


# Functions

Murat Osmanoglu

# Functions as Relations



$R(A)$  : the image of  $R$ ,  $R(A) = \{y \in B \mid (x, y) \in R, \exists x \in A\}$

Function is a relation that satisfies two conditions :


- for every element  $x$  of the domain, there is an element  $y$  in the codomain such that  $(x, y)$  is an element of the relation

Let  $R \subseteq A \times B$  be the relation,  $\forall x[(x \in A) \rightarrow (\exists y \in B \text{ s.t. } (x, y) \in R)]$


- for every element  $x$  of the domain, there is only one element  $y$  of the codomain such that  $(x, y)$  is an element of the relation

Let  $R \subseteq A \times B$  be the relation,  $\forall x[((x, y_1) \in R \wedge (x, y_2) \in R) \rightarrow (y_1 = y_2)]$


# Definition



# Definition




# Definition



- $f$  assigns every element of  $A$  to exactly one element of  $B$


# Definition



- $f$  assigns every element of  $A$  to exactly one element of  $B$

if  $(a, b) \in f$ , then  $f(a) = b$

# Definition



- $f$  assigns every element of  $A$  to exactly one element of  $B$

if  $(a, b) \in f$ , then  $f(a) = b$

preimage  
of  $b$

image  
of  $a$

# Definition

How many functions can be defined from a set  $A$  to a set  $B$  where  $|A|=n$  and  $|B|=m$  ?



# Definition

How many functions can be defined from a set  $A$  to a set  $B$  where  $|A|=n$  and  $|B|=m$  ?

- Assume  $A = \{a_1, \dots, a_n\}$  and  $B = \{b_1, \dots, b_m\}$

# Definition

How many functions can be defined from a set  $A$  to a set  $B$  where  $|A|=n$  and  $|B|=m$  ?

- Assume  $A = \{a_1, \dots, a_n\}$  and  $B = \{b_1, \dots, b_m\}$


$$f = \{(a_1, \quad), (a_2, \quad), \dots, (a_n, \quad)\}$$

# Definition

How many functions can be defined from a set  $A$  to a set  $B$  where  $|A|=n$  and  $|B|=m$  ?

- Assume  $A = \{a_1, \dots, a_n\}$  and  $B = \{b_1, \dots, b_m\}$

cannot be  $(a_1, b_1)$  and  $(a_1, b_3)$

$$f = \{(a_1, \quad), (a_2, \quad), \dots, (a_n, \quad)\}$$


# Definition

How many functions can be defined from a set  $A$  to a set  $B$  where  $|A|=n$  and  $|B|=m$  ?

- Assume  $A = \{a_1, \dots, a_n\}$  and  $B = \{b_1, \dots, b_m\}$

cannot be  $(a_1, b_1)$  and  $(a_1, b_3)$

$f = \{(a_1, \quad), (a_2, \quad), \dots, (a_n, \quad)\}$

$\uparrow \quad \uparrow \quad \uparrow$

$m \quad m \quad m$

# Definition

How many functions can be defined from a set  $A$  to a set  $B$  where  $|A|=n$  and  $|B|=m$  ?

- Assume  $A = \{a_1, \dots, a_n\}$  and  $B = \{b_1, \dots, b_m\}$

$$f = \{(a_1, \quad), (a_2, \quad), \dots, (a_n, \quad)\}$$

$\uparrow \qquad \uparrow \qquad \qquad \uparrow$   
 $m \qquad m \qquad \qquad m$

cannot be  $(a_1, b_1)$  and  $(a_1, b_3)$

$$m^n = |B|^{|A|} \text{ functions}$$

# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .

# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .

$$\forall a \forall b [f(a) = f(b) \rightarrow a = b]$$

$$\text{or } \forall a \forall b [a \neq b \rightarrow f(a) \neq f(b)]$$


# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .

$$\forall a \forall b [f(a) = f(b) \rightarrow a = b]$$

$$\text{or } \forall a \forall b [a \neq b \rightarrow f(a) \neq f(b)]$$






# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .

$$\forall a \forall b [f(a) = f(b) \rightarrow a = b]$$

$$\text{or } \forall a \forall b [a \neq b \rightarrow f(a) \neq f(b)]$$



# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .

# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

$$\forall x_1, x_2 \in \mathbb{R}, f(x_1) = f(x_2)$$

# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

$$\forall x_1, x_2 \in \mathbb{R}, f(x_1) = f(x_2) \rightarrow 3x_1 + 1 = 3x_2 + 1$$

# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{R}, \quad f(x_1) = f(x_2) &\rightarrow 3x_1 + 1 = 3x_2 + 1 \\ &\rightarrow x_1 = x_2\end{aligned}$$

# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{R}, \quad f(x_1) = f(x_2) &\rightarrow 3x_1 + 1 = 3x_2 + 1 \\ &\rightarrow x_1 = x_2\end{aligned}$$

- Determine whether the function  $f(x) = x^4 - x^2$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{R}, f(x_1) = f(x_2) &\rightarrow 3x_1 + 1 = 3x_2 + 1 \\ &\rightarrow x_1 = x_2\end{aligned}$$

- Determine whether the function  $f(x) = x^4 - x^2$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

$$\forall x_1, x_2 \in \mathbb{R}, x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)$$



# Definition

## One-to-One

- Let  $f : A \rightarrow B$ . A function is called one-to-one (or injective) if and only if  $f(a) = f(b)$  implies  $a = b$ .
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{R}, f(x_1) = f(x_2) &\rightarrow 3x_1 + 1 = 3x_2 + 1 \\ &\rightarrow x_1 = x_2\end{aligned}$$

- Determine whether the function  $f(x) = x^4 - x^2$  ( $f: \mathbb{R} \rightarrow \mathbb{R}$ ) is a one-to-one function or not.

$$\forall x_1, x_2 \in \mathbb{R}, x_1 \neq x_2 \rightarrow f(x_1) \neq f(x_2)$$

for  $x_1 = 1$  and  $x_2 = -1$ ,  $x_1 \neq x_2$  but  $f(x_1) = f(x_2)$

# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$

# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$


$$\forall b \exists a [f(a) = b]$$

# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$

$$\forall b \exists a [f(a) = b]$$




# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$

$$\forall b \exists a [f(a) = b]$$



# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$

# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{Q} \rightarrow \mathbb{Q}$ ) is a onto function or not.

# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{Q} \rightarrow \mathbb{Q}$ ) is a onto function or not.

$$\forall b \in \mathbb{Q}, f(a) = b$$



# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{Q} \rightarrow \mathbb{Q}$ ) is a onto function or not.

$$\forall b \in \mathbb{Q}, f(a) = b \leftrightarrow 3a + 1 = b$$

# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{Q} \rightarrow \mathbb{Q}$ ) is a onto function or not.

$$\begin{aligned}\forall b \in \mathbb{Q}, f(a) = b &\leftrightarrow 3a + 1 = b \\ &\leftrightarrow a = \frac{b-1}{3}\end{aligned}$$

# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{Q} \rightarrow \mathbb{Q}$ ) is a onto function or not.

$$\begin{aligned}\forall b \in \mathbb{Q}, f(a) = b &\leftrightarrow 3a + 1 = b \\ &\leftrightarrow a = \frac{b-1}{3}\end{aligned}$$

Since  $a = \frac{b-1}{3} \in \mathbb{Q}$ ,  $f$  is onto

# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{Q} \rightarrow \mathbb{Q}$ ) is a onto function or not.

$$\begin{aligned}\forall b \in \mathbb{Q}, f(a) = b &\leftrightarrow 3a + 1 = b \\ &\leftrightarrow a = \frac{b-1}{3}\end{aligned}$$

Since  $a = \frac{b-1}{3} \in \mathbb{Q}$ ,  $f$  is onto

- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{Z} \rightarrow \mathbb{Z}$ ) is a onto function or not.

# Definition

## Onto

- Let  $f : A \rightarrow B$ . A function is called onto (or surjective) if  $f(A)=B$ , i.e. for all  $b \in B$ , there is at least one  $a \in A$  such that  $f(a) = b$
- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{Q} \rightarrow \mathbb{Q}$ ) is a onto function or not.

$$\begin{aligned}\forall b \in \mathbb{Q}, f(a) = b &\leftrightarrow 3a + 1 = b \\ &\leftrightarrow a = \frac{b-1}{3}\end{aligned}$$

Since  $a = \frac{b-1}{3} \in \mathbb{Q}$ ,  $f$  is onto

- Determine whether the function  $f(x) = 3x + 1$  ( $f: \mathbb{Z} \rightarrow \mathbb{Z}$ ) is a onto function or not.

for  $5 \in \mathbb{Z}$ , there is no integer  $x \in \mathbb{Z}$  such that  $f(x) = 5$ .

# Definition

## Bijection

- If a function both one-to-one and onto, it is called bijection.

# Definition

## Bijection

- If a function both one-to-one and onto, it is called bijection.
- the identity function  $f(x) = x$  ( $f: A \rightarrow A$ ) is a bijection

# Definition

## Bijection

- If a function both one-to-one and onto, it is called bijection.
- the identity function  $f(x) = x$  ( $f: A \rightarrow A$ ) is a bijection

$$\forall x_1, x_2 \in A, f(x_1) = f(x_2) \rightarrow x_1 = x_2$$



# Definition


## Bijection

- If a function both one-to-one and onto, it is called bijection.
- the identity function  $f(x) = x$  ( $f: A \rightarrow A$ ) is a bijection

$$\forall x_1, x_2 \in A, f(x_1) = f(x_2) \rightarrow x_1 = x_2$$

$$\forall a \in A, f(a) = a, \text{ the preimage of } a \text{ is itself}$$

# Inverse




$$f: A \rightarrow B$$

$$f(a) = b$$

$$f^{-1}: B \rightarrow A$$

$$f^{-1}(b) = a$$

# Inverse




$$f: A \rightarrow B$$


$$f(a) = b$$

$$f^{-1}: B \rightarrow A$$

$$f^{-1}(b) = a$$



# Inverse




$$f: A \rightarrow B$$

$$f(a) = b$$


$$f^{-1}: B \rightarrow A$$

$$f^{-1}(b) = a$$



$$f(A) \neq B$$

# Inverse




$$f: A \rightarrow B$$


$$f(a) = b$$

$$f^{-1}: B \rightarrow A$$

$$f^{-1}(b) = a$$




$$f(A) \neq B$$



$$f(x) = f(y) = a$$

# Inverse




$$f: A \rightarrow B$$


$$f(a) = b$$

$$f^{-1}: B \rightarrow A$$

$$f^{-1}(b) = a$$




$$f(A) \neq B$$



$$f(x) = f(y) = a$$

$$f^{-1}(a) = x \text{ and } f^{-1}(a) = y$$

# Inverse





$$f: A \rightarrow B$$

$$f(a) = b$$

$$f^{-1}: B \rightarrow A$$

$$f^{-1}(b) = a$$



If  $f$  is a bijection, then  $f^{-1}$  can be defined,  
i.e.  $f$  is invertible

$= y$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible



# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = x + 1$ ,  $f$  is invertible ?

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = x + 1$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2)$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = x + 1$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow x_1 + 1 = x_2 + 1$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = x + 1$ ,  $f$  is invertible ?

$$\begin{aligned} \forall x_1, x_2 \in \mathbb{Z}, \quad f(x_1) = f(x_2) &\rightarrow x_1 + 1 = x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)} \end{aligned}$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = x + 1$ ,  $f$  is invertible ?

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) &\rightarrow x_1 + 1 = x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)}\end{aligned}$$

$$\forall y \in \mathbb{Z}, f(x) = y$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = x + 1$ ,  $f$  is invertible ?

$$\begin{aligned} \forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) &\rightarrow x_1 + 1 = x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)} \end{aligned}$$

$$\forall y \in \mathbb{Z}, f(x) = y \leftrightarrow x + 1 = y$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = x + 1$ ,  $f$  is invertible ?

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{Z}, \quad f(x_1) = f(x_2) &\rightarrow x_1 + 1 = x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)}\end{aligned}$$

$$\begin{aligned}\forall y \in \mathbb{Z}, \quad f(x) = y &\leftrightarrow x + 1 = y \\ &\leftrightarrow x = y - 1 \in \mathbb{Z} \text{ (onto)}\end{aligned}$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = x + 1$ ,  $f$  is invertible ?

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) &\rightarrow x_1 + 1 = x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)}\end{aligned}$$

$$\begin{aligned}\forall y \in \mathbb{Z}, f(x) = y &\leftrightarrow x + 1 = y \\ &\leftrightarrow x = y - 1 \in \mathbb{Z} \text{ (onto)}\end{aligned}$$

$$f^{-1}(x) = x - 1$$



# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = 2x + 1$ ,  $f$  is invertible ?

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = 2x + 1$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2)$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = 2x + 1$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 + 1 = 2x_2 + 1$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = 2x + 1$ ,  $f$  is invertible ?

$$\begin{aligned} \forall x_1, x_2 \in \mathbb{Z}, \quad f(x_1) = f(x_2) &\rightarrow 2x_1 + 1 = 2x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)} \end{aligned}$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = 2x + 1$ ,  $f$  is invertible ?

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) &\rightarrow 2x_1 + 1 = 2x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)}\end{aligned}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z} f(x) = y$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = 2x + 1$ ,  $f$  is invertible ?

$$\begin{aligned} \forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) &\rightarrow 2x_1 + 1 = 2x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)} \end{aligned}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z} f(x) = y \leftrightarrow 2x + 1 = y$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = 2x + 1$ ,  $f$  is invertible ?

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) &\rightarrow 2x_1 + 1 = 2x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)}\end{aligned}$$

$$\begin{aligned}\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z} f(x) = y &\leftrightarrow 2x + 1 = y \\ &\leftrightarrow x = \frac{y-1}{2}\end{aligned}$$



# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$ , defined as  $f(x) = 2x + 1$ ,  $f$  is invertible ?

$$\begin{aligned}\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) &\rightarrow 2x_1 + 1 = 2x_2 + 1 \\ &\rightarrow x_1 = x_2 \text{ (one-to-one)}\end{aligned}$$

$$\begin{aligned}\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z} f(x) = y &\leftrightarrow 2x + 1 = y \\ &\leftrightarrow x = \frac{y-1}{2}\end{aligned}$$

but for some  $y \in \mathbb{Z}$ ,  $x = \frac{y-1}{2} \notin \mathbb{Z}$  (not onto)

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2)$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

$$\begin{aligned} \forall x_1, x_2 \in \mathbb{Z}, \quad f(x_1) = f(x_2) &\rightarrow 2x_1 - 1 = 2x_2 - 1 \\ &\rightarrow x_1 = x_2 \end{aligned}$$

# Inverse

- If a function both one-to-one and onto, it is called bijection.  
If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible

- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2)$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2$$



# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k, \exists k \in \mathbb{Z},$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k, \exists k \in \mathbb{Z}, \text{ then } f(x) = y$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k, \exists k \in \mathbb{Z}, \text{ then } f(x) = y \leftrightarrow -2x = y$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k, \exists k \in \mathbb{Z}, \text{ then } f(x) = y \leftrightarrow -2x = y \\ \leftrightarrow x = -\frac{y}{2} = -k \in \mathbb{Z}$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k, \exists k \in \mathbb{Z}, \text{ then } f(x) = y \leftrightarrow -2x = y \\ \leftrightarrow x = -\frac{y}{2} = -k \in \mathbb{Z}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k + 1, \exists k \in \mathbb{Z},$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k, \exists k \in \mathbb{Z}, \text{ then } f(x) = y \leftrightarrow -2x = y \\ \leftrightarrow x = -\frac{y}{2} = -k \in \mathbb{Z}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k + 1, \exists k \in \mathbb{Z}, \\ \text{then } f(x) = y$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k, \exists k \in \mathbb{Z}, \text{ then } f(x) = y \leftrightarrow -2x = y \\ \leftrightarrow x = -\frac{y}{2} = -k \in \mathbb{Z}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k + 1, \exists k \in \mathbb{Z}, \\ \text{then } f(x) = y \leftrightarrow 2x - 1 = y$$



# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible ?

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k, \exists k \in \mathbb{Z}, \text{ then } f(x) = y \leftrightarrow -2x = y \\ \leftrightarrow x = -\frac{y}{2} = -k \in \mathbb{Z}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k + 1, \exists k \in \mathbb{Z}, \\ \text{then } f(x) = y \leftrightarrow 2x - 1 = y \\ \leftrightarrow x = \frac{y+1}{2} = k + 1 \in \mathbb{Z}$$

# Inverse

- If a function both one-to-one and onto, it is called bijection. If  $f$  is a bijection, then  $f^{-1}$  can be defined, i.e.  $f$  is invertible
- $f: \mathbb{Z} \rightarrow \mathbb{N}$ , defined as  $f(x) = \begin{cases} 2x - 1 & \text{if } x > 0 \\ -2x & \text{if } x \leq 0 \end{cases}$ ,  $f$  is invertible?


$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow 2x_1 - 1 = 2x_2 - 1 \\ \rightarrow x_1 = x_2$$

$$\forall x_1, x_2 \in \mathbb{Z}, f(x_1) = f(x_2) \rightarrow -2x_1 = -2x_2 \\ \rightarrow x_1 = x_2 \text{ (one-to-one)}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k, \exists k \in \mathbb{Z}, \text{ then } f(x) = y \leftrightarrow -2x = y \\ \leftrightarrow x = -\frac{y}{2} = -k \in \mathbb{Z}$$

$$\forall y \in \mathbb{Z}, \exists x \in \mathbb{Z}, \text{ if } y = 2k + 1, \exists k \in \mathbb{Z}, \\ \text{then } f(x) = y \leftrightarrow 2x - 1 = y \\ \leftrightarrow x = \frac{y+1}{2} = k + 1 \in \mathbb{Z} \\ \text{(onto)}$$


# Composition



$$f: A \rightarrow B \text{ and } g: B \rightarrow C$$

$$g \circ f: A \rightarrow C$$

# Composition




$$f: A \rightarrow B \text{ and } g: B \rightarrow C$$

$$g \circ f: A \rightarrow C$$

$$f(a) = b \text{ and } g(b) = c$$

# Composition




$$f: A \rightarrow B \text{ and } g: B \rightarrow C$$

$$g \circ f: A \rightarrow C$$

$$f(a) = b \text{ and } g(b) = c$$

$$g \circ f(a)$$

# Composition




$$f: A \rightarrow B \text{ and } g: B \rightarrow C$$

$$g \circ f: A \rightarrow C$$

$$f(a) = b \text{ and } g(b) = c$$

$$g \circ f(a) = g(f(a))$$

# Composition




$$f: A \rightarrow B \text{ and } g: B \rightarrow C$$

$$g \circ f: A \rightarrow C$$

$$f(a) = b \text{ and } g(b) = c$$

$$g \circ f(a) = g(f(a)) = g(b)$$

# Composition



$$f: A \rightarrow B \text{ and } g: B \rightarrow C$$


$$g \circ f: A \rightarrow C$$

$$f(a) = b \text{ and } g(b) = c$$

$$g \circ f(a) = g(f(a)) = g(b) = c$$



# Composition



$$f: A \rightarrow B \text{ and } g: B \rightarrow C$$

$$g \circ f: A \rightarrow C$$

$$f(a) = b \text{ and } g(b) = c$$

$$g \circ f(a) = g(f(a)) = g(b) = c$$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x)$   
 $f \circ g(x)$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x))$   
 $f \circ g(x)$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1)$   
 $f \circ g(x)$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,

$$f(x) = 3x + 1 \text{ and } g(x) = 2x - 1$$

$$g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$$

$$f \circ g(x)$$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,

$$f(x) = 3x + 1 \quad \text{and} \quad g(x) = 2x - 1$$

$$g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$$

$$f \circ g(x) = f(g(x))$$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,

$$f(x) = 3x + 1 \quad \text{and} \quad g(x) = 2x - 1$$

$$g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$$

$$f \circ g(x) = f(g(x)) = f(2x - 1)$$



# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,

$$f(x) = 3x + 1 \quad \text{and} \quad g(x) = 2x - 1$$

$$g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$$

$$f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y)$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y))$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = y$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y,$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y,$   
 $f^{-1} \circ f(x)$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y,$   
 $f^{-1} \circ f(x) = f^{-1}(f(x))$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y,$   
 $f^{-1} \circ f(x) = f^{-1}(f(x)) = f^{-1}(y)$



# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y,$   
 $f^{-1} \circ f(x) = f^{-1}(f(x)) = f^{-1}(y) = x,$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,

$$f(x) = 3x + 1 \quad \text{and} \quad g(x) = 2x - 1$$

$$g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$$

$$f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$$

- $f: A \rightarrow B$

$$f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y, \quad f \circ f^{-1} = I_B$$

$$f^{-1} \circ f(x) = f^{-1}(f(x)) = f^{-1}(y) = x, \quad f^{-1} \circ f = I_A$$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y, \quad f \circ f^{-1} = I_B$   
 $f^{-1} \circ f(x) = f^{-1}(f(x)) = f^{-1}(y) = x, \quad f^{-1} \circ f = I_A$
- If  $f$  and  $g$  are one-to-one, then  $f \circ g$  is also one-to-one.

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y, \quad f \circ f^{-1} = I_B$   
 $f^{-1} \circ f(x) = f^{-1}(f(x)) = f^{-1}(y) = x, \quad f^{-1} \circ f = I_A$
- If  $f$  and  $g$  are one-to-one, then  $f \circ g$  is also one-to-one.  
 $\forall x_1, x_2 \in A, f \circ g(x_1) = f \circ g(x_2)$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y, \quad f \circ f^{-1} = I_B$   
 $f^{-1} \circ f(x) = f^{-1}(f(x)) = f^{-1}(y) = x, \quad f^{-1} \circ f = I_A$
- If  $f$  and  $g$  are one-to-one, then  $f \circ g$  is also one-to-one.

$$\forall x_1, x_2 \in A, f \circ g(x_1) = f \circ g(x_2) \rightarrow f(g(x_1)) = f(g(x_2))$$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,

$$f(x) = 3x + 1 \text{ and } g(x) = 2x - 1$$

$$g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$$

$$f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$$

- $f: A \rightarrow B$

$$f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y, \quad f \circ f^{-1} = I_B$$

$$f^{-1} \circ f(x) = f^{-1}(f(x)) = f^{-1}(y) = x, \quad f^{-1} \circ f = I_A$$

- If  $f$  and  $g$  are one-to-one, then  $f \circ g$  is also one-to-one.

$$\begin{aligned} \forall x_1, x_2 \in A, f \circ g(x_1) = f \circ g(x_2) &\rightarrow f(g(x_1)) = f(g(x_2)) \\ &\rightarrow g(x_1) = g(x_2) \text{ (f is one-to-one)} \end{aligned}$$

# Composition

- $f, g: \mathbb{Z} \rightarrow \mathbb{Z}$ ,  
 $f(x) = 3x + 1$  and  $g(x) = 2x - 1$   
 $g \circ f(x) = g(f(x)) = g(3x + 1) = 2(3x + 1) - 1 = 6x + 1$   
 $f \circ g(x) = f(g(x)) = f(2x - 1) = 3(2x - 1) + 1 = 6x - 2$
- $f: A \rightarrow B$   
 $f \circ f^{-1}(y) = f(f^{-1}(y)) = f(x) = y, \quad f \circ f^{-1} = I_B$   
 $f^{-1} \circ f(x) = f^{-1}(f(x)) = f^{-1}(y) = x, \quad f^{-1} \circ f = I_A$
- If  $f$  and  $g$  are one-to-one, then  $f \circ g$  is also one-to-one.

$$\begin{aligned} \forall x_1, x_2 \in A, f \circ g(x_1) = f \circ g(x_2) &\rightarrow f(g(x_1)) = f(g(x_2)) \\ &\rightarrow g(x_1) = g(x_2) \text{ (f is one-to-one)} \\ &\rightarrow x_1 = x_2 \text{ (g is one-to-one)} \end{aligned}$$

# Floor and Ceiling Functions



# Floor and Ceiling Functions

- **floor function** of a real number  $x$  : is the largest integer that is less than or equal to  $x$ , denoted by  $\lfloor x \rfloor$ .

# Floor and Ceiling Functions

- **floor function** of a real number  $x$  : is the largest integer that is less than or equal to  $x$ , denoted by  $\lfloor x \rfloor$ .

$$\lfloor 1/5 \rfloor = 0, \lfloor -1/5 \rfloor = -1, \lfloor 3,56 \rfloor = 3, \lfloor -3,56 \rfloor = -4$$

# Floor and Ceiling Functions

- **floor function** of a real number  $x$  : is the largest integer that is less than or equal to  $x$ , denoted by  $\lfloor x \rfloor$ .

$$\lfloor 1/5 \rfloor = 0, \lfloor -1/5 \rfloor = -1, \lfloor 3.56 \rfloor = 3, \lfloor -3.56 \rfloor = -4$$

$$\lfloor x \rfloor = n \text{ if } n \leq x < n + 1 \quad \text{or} \quad \lfloor x \rfloor = n \text{ if } x - 1 \leq n < x$$

# Floor and Ceiling Functions

- **floor function** of a real number  $x$  : is the largest integer that is less than or equal to  $x$ , denoted by  $\lfloor x \rfloor$ .

$$\lfloor 1/5 \rfloor = 0, \lfloor -1/5 \rfloor = -1, \lfloor 3.56 \rfloor = 3, \lfloor -3.56 \rfloor = -4$$

$$\lfloor x \rfloor = n \text{ if } n \leq x < n + 1 \quad \text{or} \quad \lfloor x \rfloor = n \text{ if } x - 1 \leq n < x$$

- **ceiling function** of a real number  $x$  : is the smallest integer that is greater than or equal to  $x$ , denoted by  $\lceil x \rceil$ .

# Floor and Ceiling Functions

- **floor function** of a real number  $x$  : is the largest integer that is less than or equal to  $x$ , denoted by  $\lfloor x \rfloor$ .

$$\lfloor 1/5 \rfloor = 0, \lfloor -1/5 \rfloor = -1, \lfloor 3.56 \rfloor = 3, \lfloor -3.56 \rfloor = -4$$

$$\lfloor x \rfloor = n \text{ if } n \leq x < n + 1 \quad \text{or} \quad \lfloor x \rfloor = n \text{ if } x - 1 \leq n < x$$

- **ceiling function** of a real number  $x$  : is the smallest integer that is greater than or equal to  $x$ , denoted by  $\lceil x \rceil$ .

$$\lceil 1/5 \rceil = 1, \lceil -1/5 \rceil = 0, \lceil 3.56 \rceil = 4, \lceil -3.56 \rceil = -3$$

# Floor and Ceiling Functions

- **floor function** of a real number  $x$  : is the largest integer that is less than or equal to  $x$ , denoted by  $\lfloor x \rfloor$ .

$$\lfloor 1/5 \rfloor = 0, \lfloor -1/5 \rfloor = -1, \lfloor 3.56 \rfloor = 3, \lfloor -3.56 \rfloor = -4$$

$$\lfloor x \rfloor = n \text{ if } n \leq x < n + 1 \quad \text{or} \quad \lfloor x \rfloor = n \text{ if } x - 1 \leq n < x$$

- **ceiling function** of a real number  $x$  : is the smallest integer that is greater than or equal to  $x$ , denoted by  $\lceil x \rceil$ .

$$\lceil 1/5 \rceil = 1, \lceil -1/5 \rceil = 0, \lceil 3.56 \rceil = 4, \lceil -3.56 \rceil = -3$$

$$\lceil x \rceil = n \text{ if } n - 1 < x \leq n \quad \text{or} \quad \lceil x \rceil = n \text{ if } x \leq n < x + 1$$

# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

# Floor and Ceiling Functions


- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$   
assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$



# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $[2x] = [x] + [x + 1/2]$


assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$


# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$



$$0 \leq \varepsilon < \frac{1}{2}$$


$$\frac{1}{2} \leq \varepsilon < 1$$

# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$


$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$

$$\frac{1}{2} \leq \varepsilon < 1$$

# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$

$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$


$$2n = n + n$$

$$\frac{1}{2} \leq \varepsilon < 1$$

# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$

$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$

$$2n = n + n$$


$$\frac{1}{2} \leq \varepsilon < 1$$

$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$

# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$

$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$

$$2n = n + n$$

$$\frac{1}{2} \leq \varepsilon < 1$$


$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$

$$2n + 1 = n + n + 1$$

# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$

$$\begin{aligned}\lfloor 2n + 2\varepsilon \rfloor &= \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor \\ 2n &= n + n\end{aligned}$$

$$\frac{1}{2} \leq \varepsilon < 1$$


$$\begin{aligned}\lfloor 2n + 2\varepsilon \rfloor &= \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor \\ 2n + 1 &= n + n + 1\end{aligned}$$

- determine whether  $\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$  for all  $x, y \in \mathbb{R}$ .

# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$

$$\begin{aligned}\lfloor 2n + 2\varepsilon \rfloor &= \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor \\ 2n &= n + n\end{aligned}$$

$$\frac{1}{2} \leq \varepsilon < 1$$

$$\begin{aligned}\lfloor 2n + 2\varepsilon \rfloor &= \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor \\ 2n + 1 &= n + n + 1\end{aligned}$$

- determine whether  $\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$  for all  $x, y \in \mathbb{R}$ .


assume  $0 < x, y < \frac{1}{2}$



# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$

$$\begin{aligned}\lfloor 2n + 2\varepsilon \rfloor &= \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor \\ 2n &= n + n\end{aligned}$$

$$\frac{1}{2} \leq \varepsilon < 1$$

$$\begin{aligned}\lfloor 2n + 2\varepsilon \rfloor &= \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor \\ 2n + 1 &= n + n + 1\end{aligned}$$


- determine whether  $\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$  for all  $x, y \in \mathbb{R}$ .

assume  $0 < x, y < \frac{1}{2}$ , then  $x + y < 1$ .

# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$

$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$

$$2n = n + n$$

$$\frac{1}{2} \leq \varepsilon < 1$$

$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$

$$2n + 1 = n + n + 1$$

- determine whether  $\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$  for all  $x, y \in \mathbb{R}$ .


assume  $0 < x, y < \frac{1}{2}$ , then  $x + y < 1$ .

$$\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$$

# Floor and Ceiling Functions

- show that if  $x$  is a real number, then  $\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$

assume  $x = n + \varepsilon$  where  $n$  is integer and  $0 \leq \varepsilon < 1$


$$0 \leq \varepsilon < \frac{1}{2}$$

$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$

$$2n = n + n$$

$$\frac{1}{2} \leq \varepsilon < 1$$

$$\lfloor 2n + 2\varepsilon \rfloor = \lfloor n + \varepsilon \rfloor + \lfloor n + \varepsilon + 1/2 \rfloor$$

$$2n + 1 = n + n + 1$$

- determine whether  $\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$  for all  $x, y \in \mathbb{R}$ .

assume  $0 < x, y < \frac{1}{2}$ , then  $x + y < 1$ .

$$\lfloor x + y \rfloor = \lfloor x \rfloor + \lfloor y \rfloor$$

$$1 \neq 1 + 1$$

# Sequences

**Definition :** A sequence is a function from  $\mathbb{N}$  (or  $\mathbb{Z}^+$ ) to a set  $S$ , denoted by  $\{a_n\}$  where  $a_n$  is the general term of the sequence.

# Sequences

**Definition :** A sequence is a function from  $\mathbb{N}$  (or  $\mathbb{Z}^+$ ) to a set  $S$ , denoted by  $\{a_n\}$  where  $a_n$  is the general term of the sequence.

1, 4, 7, 10, 13, ...

# Sequences

**Definition :** A sequence is a function from  $\mathbb{N}$  (or  $\mathbb{Z}^+$ ) to a set  $S$ , denoted by  $\{a_n\}$  where  $a_n$  is the general term of the sequence.

$$1, 4, 7, 10, 13, \dots \quad \{3n + 1\}$$

# Sequences

**Definition :** A sequence is a function from  $\mathbb{N}$  (or  $\mathbb{Z}^+$ ) to a set  $S$ , denoted by  $\{a_n\}$  where  $a_n$  is the general term of the sequence.

$$1, 4, 7, 10, 13, \dots \quad \{3n + 1\}$$

$$0, 1, 3, 7, 15, \dots$$

# Sequences

**Definition :** A sequence is a function from  $\mathbb{N}$  (or  $\mathbb{Z}^+$ ) to a set  $S$ , denoted by  $\{a_n\}$  where  $a_n$  is the general term of the sequence.

$$1, 4, 7, 10, 13, \dots \quad \{3n + 1\}$$

$$0, 1, 3, 7, 15, \dots \quad \{2^n - 1\}$$



# Sequences

**Definition :** A sequence is a function from  $\mathbb{N}$  (or  $\mathbb{Z}^+$ ) to a set  $S$ , denoted by  $\{a_n\}$  where  $a_n$  is the general term of the sequence.

$$1, 4, 7, 10, 13, \dots \quad \{3n + 1\}$$

$$0, 1, 3, 7, 15, \dots \quad \{2^n - 1\}$$

- $a_n = \frac{1}{n}$

# Sequences

**Definition :** A sequence is a function from  $\mathbb{N}$  (or  $\mathbb{Z}^+$ ) to a set  $S$ , denoted by  $\{a_n\}$  where  $a_n$  is the general term of the sequence.

$$1, 4, 7, 10, 13, \dots \quad \{3n + 1\}$$

$$0, 1, 3, 7, 15, \dots \quad \{2^n - 1\}$$

- $a_n = \frac{1}{n} \quad a_1 = 1, a_2 = \frac{1}{2}, a_3 = \frac{1}{3}, \dots$

-

# Sequences

**Definition :** A sequence is a function from  $\mathbb{N}$  (or  $\mathbb{Z}^+$ ) to a set  $S$ , denoted by  $\{a_n\}$  where  $a_n$  is the general term of the sequence.

$$1, 4, 7, 10, 13, \dots \quad \{3n + 1\}$$

$$0, 1, 3, 7, 15, \dots \quad \{2^n - 1\}$$

- $a_n = \frac{1}{n} \quad a_1 = 1, a_2 = \frac{1}{2}, a_3 = \frac{1}{3}, \dots$

- $a_n = \frac{1}{3^{n+2}}$

# Sequences

**Definition :** A sequence is a function from  $\mathbb{N}$  (or  $\mathbb{Z}^+$ ) to a set  $S$ , denoted by  $\{a_n\}$  where  $a_n$  is the general term of the sequence.

$$1, 4, 7, 10, 13, \dots \quad \{3n + 1\}$$

$$0, 1, 3, 7, 15, \dots \quad \{2^n - 1\}$$

- $a_n = \frac{1}{n} \quad a_1 = 1, a_2 = \frac{1}{2}, a_3 = \frac{1}{3}, \dots$
- $a_n = \frac{1}{3^{n+2}} \quad a_0 = \frac{1}{2}, a_1 = \frac{1}{5}, a_2 = \frac{1}{11}, \dots$

# Sequences

Geometric Sequence :


# Sequences

Geometric Sequence :

$$a, ar, ar^2, \dots, ar^n, \dots$$


# Sequences

## Geometric Sequence :



# Sequences

## Geometric Sequence :




$$a_n = (-1)^n$$

$1, -1, 1, -1, \dots$



# Sequences

## Geometric Sequence :



$$a_n = (-1)^n$$


1, -1, 1, -1, ...

$$a_n = 2.3^n$$

2, 2.3, 2.9, 2.27, ...

# Sequences

## Geometric Sequence :



$$a_n = (-1)^n$$

1, -1, 1, -1, ...

$$a_n = 2.3^n$$

2, 2.3, 2.9, 2.27, ...

$$a_n = 3. (1/2)^n$$

3, 3/2, 3/4, 3/8, ...

# Sequences

Arithmetic Sequence :


# Sequences

Arithmetic Sequence :

$$a, a + d, a + 2d, \dots, a + n.d, \dots$$


# Sequences

Arithmetic Sequence :



# Sequences

## Arithmetic Sequence :




$$a_n = 1 + n$$

1, 2, 3, 4, ...

# Sequences

## Arithmetic Sequence :



$$a_n = 1 + n$$


$$1, 2, 3, 4, \dots$$

$$a_n = 2 - 4n$$

$$2, -2, -6, -10, \dots$$

# Sequences

## Arithmetic Sequence :



$$a_n = 1 + n$$

1, 2, 3, 4, ...

$$a_n = 2 - 4n$$

2, -2, -6, -10, ...

$$a_n = -1 + 8n$$

-1, 7, 15, 23, ...



# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$   
 $\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$

$$\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$$

$$\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$

$$\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$$

$$\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$$

- $S = \{2, 3, 4\}, \quad \sum_{x \in S} x^3$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$   
 $\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$   
 $\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$
- $S = \{2, 3, 4\}, \quad \sum_{x \in S} x^3 = 2^3 + 3^3 + 4^3 = 99$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$   
 $\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$   
 $\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$
- $S = \{2, 3, 4\}, \quad \sum_{x \in S} x^3 = 2^3 + 3^3 + 4^3 = 99$
- $\sum c f(x) = c \sum f(x)$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$   
 $\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$   
 $\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$
- $S = \{2, 3, 4\}, \quad \sum_{x \in S} x^3 = 2^3 + 3^3 + 4^3 = 99$
- $\sum c f(x) = c \sum f(x)$   
 $\sum (f(x) + g(x)) = \sum f(x) + \sum g(x)$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$

$$\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$$

$$\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$$

- $S = \{2, 3, 4\}, \quad \sum_{x \in S} x^3 = 2^3 + 3^3 + 4^3 = 99$

- $\sum c f(x) = c \sum f(x)$

$$\sum (f(x) + g(x)) = \sum f(x) + \sum g(x)$$

$$\sum_{i=m}^n f(i) = \sum_{i=m}^k f(i) + \sum_{i=k+1}^n f(i)$$



# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$   
 $\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$   
 $\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$
- $S = \{2, 3, 4\}, \quad \sum_{x \in S} x^3 = 2^3 + 3^3 + 4^3 = 99$
- $\sum c f(x) = c \sum f(x)$   
 $\sum (f(x) + g(x)) = \sum f(x) + \sum g(x)$   
 $\sum_{i=m}^n f(i) = \sum_{i=m}^k f(i) + \sum_{i=k+1}^n f(i)$
- $\sum_{i=1}^n i$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$   
 $\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$   
 $\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$
- $S = \{2, 3, 4\}, \quad \sum_{x \in S} x^3 = 2^3 + 3^3 + 4^3 = 99$
- $\sum c f(x) = c \sum f(x)$   
 $\sum (f(x) + g(x)) = \sum f(x) + \sum g(x)$   
 $\sum_{i=m}^n f(i) = \sum_{i=m}^k f(i) + \sum_{i=k+1}^n f(i)$
- $\sum_{i=1}^n i = 1 + 2 + \dots + \frac{n}{2} + \left(\frac{n}{2} + 1\right) + \dots + (n-1) + n$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$   
 $\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$   
 $\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$
- $S = \{2, 3, 4\}, \quad \sum_{x \in S} x^3 = 2^3 + 3^3 + 4^3 = 99$
- $\sum c f(x) = c \sum f(x)$   
 $\sum (f(x) + g(x)) = \sum f(x) + \sum g(x)$   
 $\sum_{i=m}^n f(i) = \sum_{i=m}^k f(i) + \sum_{i=k+1}^n f(i)$
- $\sum_{i=1}^n i = 1 + 2 + \dots + \frac{n}{2} + \left(\frac{n}{2} + 1\right) + \dots + (n-1) + n$   
 $= (n+1) + (n+1) + \dots + (n+1)$

# Summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \dots + a_{n-1} + a_n$   
 $\sum_{i=0}^{\infty} a_i = a_0 + a_1 + \dots + a_n + \dots$   
 $\sum_{i=2}^5 (i^2 - 1) = 4 - 1 + 9 - 1 + 16 - 1 + 25 - 1 = 50$
- $S = \{2, 3, 4\}, \quad \sum_{x \in S} x^3 = 2^3 + 3^3 + 4^3 = 99$
- $\sum c f(x) = c \sum f(x)$   
 $\sum (f(x) + g(x)) = \sum f(x) + \sum g(x)$   
 $\sum_{i=m}^n f(i) = \sum_{i=m}^k f(i) + \sum_{i=k+1}^n f(i)$
- $\sum_{i=1}^n i = 1 + 2 + \dots + \frac{n}{2} + \left(\frac{n}{2} + 1\right) + \dots + (n-1) + n$   
 $= (n+1) + (n+1) + \dots + (n+1)$   
 $= \frac{n}{2}(n+1)$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\sum_{i=0}^n (a + id)$$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\sum_{i=0}^n (a + id) = \sum_{i=0}^n a + \sum_{i=0}^n id$$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i\end{aligned}$$



# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

- $a, ar, ar^2, \dots, ar^n$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

- $a, ar, ar^2, \dots, ar^n$

$$S_n = \sum_{i=0}^n ar^i$$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

- $a, ar, ar^2, \dots, ar^n$

$$S_n = \sum_{i=0}^n ar^i \rightarrow rS_n = r \sum_{i=0}^n ar^i$$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

- $a, ar, ar^2, \dots, ar^n$

$$S_n = \sum_{i=0}^n ar^i \rightarrow rS_n = r \sum_{i=0}^n ar^i = \sum_{i=0}^n ar^{i+1}$$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

- $a, ar, ar^2, \dots, ar^n$

$$\begin{aligned}S_n = \sum_{i=0}^n ar^i &\rightarrow rS_n = r \sum_{i=0}^n ar^i = \sum_{i=0}^n ar^{i+1} \\ rS_n &= \sum_{i=1}^{n+1} ar^i\end{aligned}$$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

- $a, ar, ar^2, \dots, ar^n$

$$\begin{aligned}S_n = \sum_{i=0}^n ar^i &\rightarrow rS_n = r \sum_{i=0}^n ar^i = \sum_{i=0}^n ar^{i+1} \\ rS_n &= \sum_{i=1}^{n+1} ar^i = \sum_{i=1}^n ar^i + ar^{n+1}\end{aligned}$$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

- $a, ar, ar^2, \dots, ar^n$

$$\begin{aligned}S_n &= \sum_{i=0}^n ar^i \rightarrow rS_n = r \sum_{i=0}^n ar^i = \sum_{i=0}^n ar^{i+1} \\ rS_n &= \sum_{i=1}^{n+1} ar^i = \sum_{i=1}^n ar^i + ar^{n+1} \\ rS_n &= \sum_{i=0}^n ar^i + ar^{n+1} - a\end{aligned}$$



# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

- $a, ar, ar^2, \dots, ar^n$

$$\begin{aligned}S_n &= \sum_{i=0}^n ar^i \rightarrow rS_n = r \sum_{i=0}^n ar^i = \sum_{i=0}^n ar^{i+1} \\ rS_n &= \sum_{i=1}^{n+1} ar^i = \sum_{i=1}^n ar^i + ar^{n+1} \\ rS_n &= \sum_{i=0}^n ar^i + ar^{n+1} - a \\ rS_n &= S_n + ar^{n+1} - a\end{aligned}$$

# Summations

- $a, a + d, a + 2d, \dots, a + n.d$

$$\begin{aligned}\sum_{i=0}^n (a + id) &= \sum_{i=0}^n a + \sum_{i=0}^n id \\ &= \sum_{i=0}^n a + d \sum_{i=0}^n i \\ &= (n + 1)a + d \frac{n(n+1)}{2}\end{aligned}$$

- $a, ar, ar^2, \dots, ar^n$

$$\begin{aligned}S_n &= \sum_{i=0}^n ar^i \rightarrow rS_n = r \sum_{i=0}^n ar^i = \sum_{i=0}^n ar^{i+1} \\ rS_n &= \sum_{i=1}^{n+1} ar^i = \sum_{i=1}^n ar^i + ar^{n+1} \\ rS_n &= \sum_{i=0}^n ar^i + ar^{n+1} - a \\ rS_n &= S_n + ar^{n+1} - a \rightarrow S_n = \frac{ar^{n+1} - a}{r - 1}\end{aligned}$$

# Recurrence Relations

- sometimes the elements of the sequence are defined recursively in terms of previous and the initial elements of the sequence

# Recurrence Relations

- sometimes the elements of the sequence are defined recursively in terms of previous and the initial elements of the sequence

$$a_0 = 1, a_1 = 5, a_2 = 13, a_3 = 29, a_4 = ?$$

# Recurrence Relations

- sometimes the elements of the sequence are defined recursively in terms of previous and the initial elements of the sequence

$$a_0 = 1, a_1 = 5, a_2 = 13, a_3 = 29, a_4 = ?$$

$$a_1 = 2a_0 + 3 = 5$$

# Recurrence Relations

- sometimes the elements of the sequence are defined recursively in terms of previous and the initial elements of the sequence

$$a_0 = 1, a_1 = 5, a_2 = 13, a_3 = 29, a_4 = ?$$

$$a_1 = 2a_0 + 3 = 5$$

$$a_2 = 2a_1 + 3 = 13$$

# Recurrence Relations

- sometimes the elements of the sequence are defined recursively in terms of previous and the initial elements of the sequence

$$a_0 = 1, a_1 = 5, a_2 = 13, a_3 = 29, a_4 = ?$$

$$a_1 = 2a_0 + 3 = 5$$

$$a_2 = 2a_1 + 3 = 13$$

$$a_3 = 2a_2 + 3 = 29$$

# Recurrence Relations

- sometimes the elements of the sequence are defined recursively in terms of previous and the initial elements of the sequence

$$a_0 = 1, a_1 = 5, a_2 = 13, a_3 = 29, a_4 = ?$$

$$a_1 = 2a_0 + 3 = 5$$

$$a_2 = 2a_1 + 3 = 13$$

$$a_3 = 2a_2 + 3 = 29$$

$$a_4 = 2a_3 + 3 = 61$$



# Recurrence Relations

- sometimes the elements of the sequence are defined recursively in terms of previous and the initial elements of the sequence

$$a_0 = 1, a_1 = 5, a_2 = 13, a_3 = 29, a_4 = ?$$

$$a_1 = 2a_0 + 3 = 5$$

$$a_2 = 2a_1 + 3 = 13$$

$$a_3 = 2a_2 + 3 = 29$$

$$a_4 = 2a_3 + 3 = 61$$

**Definition :** an equation that express the general term of the sequence in terms of previous terms.

# Recurrence Relations

- sometimes the elements of the sequence are defined recursively in terms of previous and the initial elements of the sequence

$$a_0 = 1, a_1 = 5, a_2 = 13, a_3 = 29, a_4 = ?$$

$$a_1 = 2a_0 + 3 = 5$$

$$a_2 = 2a_1 + 3 = 13$$

$$a_3 = 2a_2 + 3 = 29$$

$$a_4 = 2a_3 + 3 = 61$$

**Definition :** an equation that express the general term of the sequence in terms of previous terms. A sequence is called a solution of a recurrence relation if its terms satisfy the recurrence relation.

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3 \cdot 5$$

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3 \cdot 5$$

$$a_2 = 45 = 3 \cdot (3 \cdot 5)$$

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3 \cdot 5$$

$$a_2 = 75 = 3 \cdot (3 \cdot 5)$$

$$a_3 = 225 = 3 \cdot (3 \cdot (3 \cdot 5))$$

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3 \cdot 5$$

$$a_2 = 75 = 3 \cdot (3 \cdot 5)$$

$$a_3 = 225 = 3 \cdot (3 \cdot (3 \cdot 5))$$

$$\vdots$$

$$a_n = 3^n 5$$

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3.5$$

$$a_2 = 75 = 3.(3.5)$$

$$a_3 = 225 = 3.(3.(3.5))$$

$\vdots$

$$a_n = 3^n 5 ; \text{ the unique solution of the given recurrence relation}$$



# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3.5$$

$$a_2 = 75 = 3.(3.5)$$

$$a_3 = 225 = 3.(3.(3.5))$$

$\vdots$

$$a_n = 3^n 5 ; \text{ the unique solution of the given recurrence relation}$$

- $a_{n+1} = d.a_n, a_0 = A$  where  $d$  is constant

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3.5$$

$$a_2 = 75 = 3.(3.5)$$

$$a_3 = 225 = 3.(3.(3.5))$$

$\vdots$

$$a_n = 3^n 5 ; \text{ the unique solution of the given recurrence relation}$$

- $a_{n+1} = d.a_n, a_0 = A$  where  $d$  is constant

the solution of the recurrence relation will be  $a_n = A.d^n$

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3.5$$

$$a_2 = 75 = 3.(3.5)$$

$$a_3 = 225 = 3.(3.(3.5))$$

$\vdots$

$$a_n = 3^n 5 ; \text{ the unique solution of the given recurrence relation}$$

- $a_{n+1} = d.a_n, a_0 = A$  where  $d$  is constant

the solution of the recurrence relation will be  $a_n = A.d^n$

- solve the recurrence relation  $a_{n+1} = 7.a_n$  where  $n \geq 1$  and  $a_2 = 98$

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3.5$$

$$a_2 = 75 = 3.(3.5)$$

$$a_3 = 225 = 3.(3.(3.5))$$

$\vdots$

$$a_n = 3^n 5 ; \text{ the unique solution of the given recurrence relation}$$

- $a_{n+1} = d.a_n, a_0 = A$  where  $d$  is constant

the solution of the recurrence relation will be  $a_n = A.d^n$

- solve the recurrence relation  $a_{n+1} = 7.a_n$  where  $n \geq 1$  and  $a_2 = 98$

$$a_2 = A.7^2 \rightarrow$$

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3.5$$

$$a_2 = 75 = 3.(3.5)$$

$$a_3 = 225 = 3.(3.(3.5))$$

$\vdots$

$$a_n = 3^n 5 ; \text{ the unique solution of the given recurrence relation}$$

- $a_{n+1} = d.a_n, a_0 = A$  where  $d$  is constant

the solution of the recurrence relation will be  $a_n = A.d^n$

- solve the recurrence relation  $a_{n+1} = 7.a_n$  where  $n \geq 1$  and  $a_2 = 98$

$$a_2 = A.7^2 \rightarrow 98 = A.49 \rightarrow A = 2$$

# Recurrence Relations

- $a_{n+1} = 3a_n, a_0 = 5$

$$a_1 = 15 = 3.5$$

$$a_2 = 75 = 3.(3.5)$$

$$a_3 = 225 = 3.(3.(3.5))$$

$\vdots$

$$a_n = 3^n 5 ; \text{ the unique solution of the given recurrence relation}$$

- $a_{n+1} = d.a_n, a_0 = A$  where  $d$  is constant

the solution of the recurrence relation will be  $a_n = A.d^n$

- solve the recurrence relation  $a_{n+1} = 7.a_n$  where  $n \geq 1$  and  $a_2 = 98$

$$a_2 = A.7^2 \rightarrow 98 = A.49 \rightarrow A = 2$$

the solution is  $a_n = 2.7^n$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$3$$

$$1 + 2$$

$$2 + 1$$

$$1 + 1 + 1$$



# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$3$$

$$1 + 2$$

$$2 + 1$$

$$1 + 1 + 1$$


- In how many different ways can  $n$  be written as a sum of positive integers ?

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$\begin{array}{c} 3 \\ 1 + 2 \\ 2 + 1 \\ 1 + 1 + 1 \end{array}$$

- In how many different ways can  $n$  be written as a sum of positive integers ?


$$\begin{array}{c} 4 \\ 1 + 3 \\ 2 + 2 \\ 1 + 1 + 2 \end{array}$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$\begin{array}{c} 3 \\ 1 + 2 \\ 2 + 1 \\ 1 + 1 + 1 \end{array}$$

- In how many different ways can  $n$  be written as a sum of positive integers?


$$\begin{array}{cc} \begin{array}{c} 4 \\ 1 + 3 \\ 2 + 2 \\ 1 + 1 + 2 \end{array} & \begin{array}{c} 3 + 1 \\ 1 + 2 + 1 \\ 2 + 1 + 1 \\ 1 + 1 + 1 + 1 \end{array} \end{array}$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$\begin{array}{c} 3 \\ 1 + 2 \\ 2 + 1 \\ 1 + 1 + 1 \end{array}$$


- In how many different ways can  $n$  be written as a sum of positive integers?


$$\begin{array}{cc} \begin{array}{c} 4 \\ 1 + 3 \\ 2 + 2 \\ 1 + 1 + 2 \end{array} & \begin{array}{c} 3 + 1 \\ 1 + 2 + 1 \\ 2 + 1 + 1 \\ 1 + 1 + 1 + 1 \end{array} \end{array}$$


- $a_4 = 2 \cdot a_3,$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:




- In how many different ways can  $n$  be written as a sum of positive integers?




- $a_4 = 2 \cdot a_3$ ,  $a_3 = 2 \cdot a_2$ , and  $a_2 = 2$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- In how many different ways can  $n$  be written as a sum of positive integers?



- $a_4 = 2 \cdot a_3$ ,  $a_3 = 2 \cdot a_2$ , and  $a_2 = 2$   
 $a_{n+1} = 2 \cdot a_n$ ,  $a_1 = 1$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$\begin{array}{ccc}
 & 3 & \\
 & 1 + 2 & \leftarrow 2 \\
 & 2 + 1 & \leftarrow 1 + 1 \\
 & 1 + 1 + 1 &
 \end{array}$$

- In how many different ways can  $n$  be written as a sum of positive integers?

$$\begin{array}{ccc}
 & 4 & 3 + 1 \\
 & 1 + 3 & 1 + 2 + 1 \\
 & 2 + 2 & 2 + 1 + 1 \\
 & 1 + 1 + 2 & 1 + 1 + 1 + 1
 \end{array}$$

- $a_4 = 2 \cdot a_3$ ,  $a_3 = 2 \cdot a_2$ , and  $a_2 = 2$   
 $a_{n+1} = 2 \cdot a_n$ ,  $a_1 = 1$   
 create a new sequence  $b_n = a_{n+1}$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$\begin{array}{ccc}
 & 3 & \\
 & 1 + 2 & \leftarrow 2 \\
 & 2 + 1 & \leftarrow 1 + 1 \\
 & 1 + 1 + 1 &
 \end{array}$$

- In how many different ways can  $n$  be written as a sum of positive integers?

$$\begin{array}{ccc}
 & 4 & 3 + 1 \\
 & 1 + 3 & 1 + 2 + 1 \\
 & 2 + 2 & 2 + 1 + 1 \\
 & 1 + 1 + 2 & 1 + 1 + 1 + 1
 \end{array}$$

- $a_4 = 2 \cdot a_3$ ,  $a_3 = 2 \cdot a_2$ , and  $a_2 = 2$   
 $a_{n+1} = 2 \cdot a_n$ ,  $a_1 = 1$   
 create a new sequence  $b_n = a_{n+1}$   
 $b_n = 2b_{n-1}$ ,  $b_0 = 1$ ;



# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$\begin{array}{ccc}
 & 3 & \\
 & 1 + 2 & \leftarrow 2 \\
 & 2 + 1 & \leftarrow 1 + 1 \\
 & 1 + 1 + 1 &
 \end{array}$$

- In how many different ways can  $n$  be written as a sum of positive integers?

$$\begin{array}{ccc}
 & 4 & 3 + 1 \\
 & 1 + 3 & 1 + 2 + 1 \\
 & 2 + 2 & 2 + 1 + 1 \\
 & 1 + 1 + 2 & 1 + 1 + 1 + 1
 \end{array}$$

- $a_4 = 2 \cdot a_3$ ,  $a_3 = 2 \cdot a_2$ , and  $a_2 = 2$


$$a_{n+1} = 2 \cdot a_n, a_1 = 1$$

create a new sequence  $b_n = a_{n+1}$


$$b_n = 2b_{n-1}, b_0 = 1; \text{ the solution will be } b_n = 2^n$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- In how many different ways can  $n$  be written as a sum of positive integers?



- $a_4 = 2 \cdot a_3$ ,  $a_3 = 2 \cdot a_2$ , and  $a_2 = 2$

$$a_{n+1} = 2 \cdot a_n, a_1 = 1$$

create a new sequence  $b_n = a_{n+1}$

$b_n = 2b_{n-1}$ ,  $b_0 = 1$ ; the solution will be  $b_n = 2^n$ ; thus  $a_n = 2^{n-1}$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$\begin{array}{ccc}
 & 3 & \\
 & 1 + 2 & \leftarrow 2 \\
 & 2 + 1 & \leftarrow 1 + 1 \\
 & 1 + 1 + 1 &
 \end{array}$$

- In how many different ways can  $n$  be written as a sum of positive integers?

$$\begin{array}{ccc}
 & 4 & 3 + 1 \\
 & 1 + 3 & 1 + 2 + 1 \\
 & 2 + 2 & 2 + 1 + 1 \\
 & 1 + 1 + 2 & 1 + 1 + 1 + 1
 \end{array}$$

- $a_4 = 2 \cdot a_3$ ,  $a_3 = 2 \cdot a_2$ , and  $a_2 = 2$

$$a_{n+1} = 2 \cdot a_n, a_1 = 1$$

create a new sequence  $b_n = a_{n+1}$

first order linear homogeneous  
recurrence relation

$b_n = 2b_{n-1}$ ,  $b_0 = 1$ ; the solution will be  $b_n = 2^n$ ; thus  $a_n = 2^{n-1}$

# Recurrence Relations

- $a_{n+1} - d \cdot a_n = 0, a_0 = A$  where  $d$  is constant.

# Recurrence Relations

- $a_{n+1} - d \cdot a_n = 0$ ,  $a_0 = A$  where  $d$  is constant.
  - first order since  $a_{n+1}$  only depends on  $a_n$  (the previous term)

# Recurrence Relations

- $a_{n+1} - d \cdot a_n = 0$ ,  $a_0 = A$  where  $d$  is constant.
  - first order since  $a_{n+1}$  only depends on  $a_n$  (the previous term)
  - linear since each variable appears in the first power and there is no product such as  $a_{n+1} \cdot a_n$

# Recurrence Relations

- $a_{n+1} - d \cdot a_n = 0$ ,  $a_0 = A$  where  $d$  is constant.
  - first order since  $a_{n+1}$  only depends on  $a_n$  (the previous term)
  - linear since each variable appears in the first power and there is no product such as  $a_{n+1} \cdot a_n$
  - homogeneous since the right hand side is 0

# Recurrence Relations

- $a_{n+1} - d \cdot a_n = 0, a_0 = A$  where  $d$  is constant.
  - first order since  $a_{n+1}$  only depends on  $a_n$  (the previous term)
  - linear since each variable appears in the first power and there is no product such as  $a_{n+1} \cdot a_n$
  - homogeneous since the right hand side is 0
- The second order linear homogeneous recurrence relation :

$$C_0 a_{n+1} + C_1 a_n + C_2 a_{n-1} = 0, a_0 = A, a_1 = B, n \geq 2$$



# Recurrence Relations

- $a_{n+1} - d \cdot a_n = 0, a_0 = A$  where  $d$  is constant.
  - first order since  $a_{n+1}$  only depends on  $a_n$  (the previous term)
  - linear since each variable appears in the first power and there is no product such as  $a_{n+1} \cdot a_n$
  - homogeneous since the right hand side is 0
- The second order linear homogeneous recurrence relation :

$$C_0 a_{n+1} + C_1 a_n + C_2 a_{n-1} = 0, a_0 = A, a_1 = B, n \geq 2$$

- The Fibonacci sequence:

$$F_{n+1} = F_n + F_{n-1}, F_0 = 1, F_2 = 1, n \geq 2$$

# Recurrence Relations

- The second order linear homogeneous recurrence relation :

$$C_0 a_{n+1} + C_1 a_n + C_2 a_{n-1} = 0, a_0 = A, a_1 = B, n \geq 2$$

# Recurrence Relations

- The second order linear homogeneous recurrence relation :

$$C_0 a_{n+1} + C_1 a_n + C_2 a_{n-1} = 0, a_0 = A, a_1 = B, n \geq 2$$

$a_{n+1} - d \cdot a_n = 0, a_0 = A$ . the solution was in the form of  $a_n = A \cdot d^n$

# Recurrence Relations

- The second order linear homogeneous recurrence relation :

$$C_0 a_{n+1} + C_1 a_n + C_2 a_{n-1} = 0, a_0 = A, a_1 = B, n \geq 2$$

$a_{n+1} - d \cdot a_n = 0, a_0 = A$ . the solution was in the form of  $a_n = A \cdot d^n$

- Similarly, we look for a solution in the form of  $a_n = c \cdot r^n$

# Recurrence Relations

- The second order linear homogeneous recurrence relation :

$$C_0 a_{n+1} + C_1 a_n + C_2 a_{n-1} = 0, a_0 = A, a_1 = B, n \geq 2$$

$a_{n+1} - d \cdot a_n = 0, a_0 = A$ . the solution was in the form of  $a_n = A \cdot d^n$

- Similarly, we look for a solution in the form of  $a_n = c \cdot r^n$

If we place it in the equation:

$$C_0 c \cdot r^{n+1} + C_1 c \cdot r^n + C_2 c \cdot r^{n-1} = 0$$

# Recurrence Relations

- The second order linear homogeneous recurrence relation :

$$C_0 a_{n+1} + C_1 a_n + C_2 a_{n-1} = 0, a_0 = A, a_1 = B, n \geq 2$$

$a_{n+1} - d \cdot a_n = 0, a_0 = A$ . the solution was in the form of  $a_n = A \cdot d^n$

- Similarly, we look for a solution in the form of  $a_n = c \cdot r^n$

If we place it in the equation:

$$C_0 c \cdot r^{n+1} + C_1 c \cdot r^n + C_2 c \cdot r^{n-1} = 0$$

$$C_0 r^2 + C_1 r + C_2 = 0 \quad (\text{characteristic equation})$$

# Recurrence Relations

- The second order linear homogeneous recurrence relation :

$$C_0 a_{n+1} + C_1 a_n + C_2 a_{n-1} = 0, a_0 = A, a_1 = B, n \geq 2$$

$a_{n+1} - d \cdot a_n = 0, a_0 = A$ . the solution was in the form of  $a_n = A \cdot d^n$

- Similarly, we look for a solution in the form of  $a_n = c \cdot r^n$

If we place it in the equation:

$$C_0 c \cdot r^{n+1} + C_1 c \cdot r^n + C_2 c \cdot r^{n-1} = 0$$

$$C_0 r^2 + C_1 r + C_2 = 0 \quad (\text{characteristic equation})$$

The solutions for the characteristic equation are called characteristic roots;  $r_1$  and  $r_2$

# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$



# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

$$r_1 = 2, r_2 = -3 \text{ (characteristic roots)}$$

# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

$$r_1 = 2, r_2 = -3 \text{ (characteristic roots)}$$

the solution will be in the form of  $a_n = c_1 2^n + c_2 (-3)^n$ .

# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

$$r_1 = 2, r_2 = -3 \text{ (characteristic roots)}$$

the solution will be in the form of  $a_n = c_1 2^n + c_2 (-3)^n$ .

$$a_0 = c_1 2^0 + c_2 (-3)^0$$

# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

$$r_1 = 2, r_2 = -3 \text{ (characteristic roots)}$$

the solution will be in the form of  $a_n = c_1 2^n + c_2 (-3)^n$ .

$$a_0 = c_1 2^0 + c_2 (-3)^0 \rightarrow -1 = c_1 + c_2$$

# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

$$r_1 = 2, r_2 = -3 \text{ (characteristic roots)}$$

the solution will be in the form of  $a_n = c_1 2^n + c_2 (-3)^n$ .

$$a_0 = c_1 2^0 + c_2 (-3)^0 \rightarrow -1 = c_1 + c_2$$

$$a_1 = c_1 2^1 + c_2 (-3)^1$$

# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

$$r_1 = 2, r_2 = -3 \text{ (characteristic roots)}$$

the solution will be in the form of  $a_n = c_1 2^n + c_2 (-3)^n$ .

$$a_0 = c_1 2^0 + c_2 (-3)^0 \rightarrow -1 = c_1 + c_2$$

$$a_1 = c_1 2^1 + c_2 (-3)^1 \rightarrow 8 = 2c_1 - 3c_2$$

# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

$$r_1 = 2, r_2 = -3 \text{ (characteristic roots)}$$

the solution will be in the form of  $a_n = c_1 2^n + c_2 (-3)^n$ .

$$a_0 = c_1 2^0 + c_2 (-3)^0 \rightarrow -1 = c_1 + c_2$$

$$a_1 = c_1 2^1 + c_2 (-3)^1 \rightarrow 8 = 2c_1 - 3c_2$$

$$c_1 + c_2 = -1$$

$$2c_1 - 3c_2 = 8$$



# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

$$r_1 = 2, r_2 = -3 \text{ (characteristic roots)}$$

the solution will be in the form of  $a_n = c_1 2^n + c_2 (-3)^n$ .

$$a_0 = c_1 2^0 + c_2 (-3)^0 \rightarrow -1 = c_1 + c_2$$

$$a_1 = c_1 2^1 + c_2 (-3)^1 \rightarrow 8 = 2c_1 - 3c_2$$

$$c_1 + c_2 = -1$$

$$2c_1 - 3c_2 = 8$$

---

$$c_1 = 1, c_2 = -2$$

# Recurrence Relations

- $a_{n+1} + a_n - 6a_{n-1} = 0, a_0 = -1, a_1 = 8, n \geq 2$

$$r^2 + r - 6 = 0 \text{ (characteristic equation)}$$

$$r_1 = 2, r_2 = -3 \text{ (characteristic roots)}$$

the solution will be in the form of  $a_n = c_1 2^n + c_2 (-3)^n$ .

$$a_0 = c_1 2^0 + c_2 (-3)^0 \rightarrow -1 = c_1 + c_2$$

$$a_1 = c_1 2^1 + c_2 (-3)^1 \rightarrow 8 = 2c_1 - 3c_2$$



$$\begin{array}{l} c_1 + c_2 = -1 \\ \underline{2c_1 - 3c_2 = 8} \\ c_1 = 1, c_2 = -2 \end{array} \longrightarrow a_n = 2^n - 2 \cdot (-3)^n$$

# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?


# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?





# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?




# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?




# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?



# Recurrence Relations


- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?





# Recurrence Relations


- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?



$$b_n =$$


# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?




# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?




# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?




# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?




# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?



# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it?



# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?



# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}$ ,  $n \geq 3$ ,  $b_1 = 1$  and  $b_2 = 2$

# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}, n \geq 3, b_1 = 1$  and  $b_2 = 2$   
 $r^2 - r - 1 = 0$  (characteristic equation)

# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}, n \geq 3, b_1 = 1$  and  $b_2 = 2$

$$r^2 - r - 1 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \frac{1+\sqrt{5}}{2}, r_2 = \frac{1-\sqrt{5}}{2} \quad (\text{characteristic roots})$$

# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}, n \geq 3, b_1 = 1$  and  $b_2 = 2$

$$r^2 - r - 1 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \frac{1+\sqrt{5}}{2}, r_2 = \frac{1-\sqrt{5}}{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^n + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^n$

# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}, n \geq 3, b_1 = 1$  and  $b_2 = 2$

$$r^2 - r - 1 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \frac{1+\sqrt{5}}{2}, r_2 = \frac{1-\sqrt{5}}{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^n + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^n$

$$b_0 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^0 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^0$$

# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}, n \geq 3, b_1 = 1$  and  $b_2 = 2$

$$r^2 - r - 1 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \frac{1+\sqrt{5}}{2}, r_2 = \frac{1-\sqrt{5}}{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^n + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^n$

$$b_0 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^0 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^0 \rightarrow 1 = c_1 + c_2$$

# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}, n \geq 3, b_1 = 1$  and  $b_2 = 2$

$$r^2 - r - 1 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \frac{1+\sqrt{5}}{2}, r_2 = \frac{1-\sqrt{5}}{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^n + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^n$

$$b_0 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^0 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^0 \rightarrow 1 = c_1 + c_2$$

$$b_1 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^1 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^1$$

# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}, n \geq 3, b_1 = 1$  and  $b_2 = 2$

$$r^2 - r - 1 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \frac{1+\sqrt{5}}{2}, r_2 = \frac{1-\sqrt{5}}{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^n + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^n$

$$b_0 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^0 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^0 \rightarrow 1 = c_1 + c_2$$

$$b_1 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^1 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^1 \rightarrow 2 = \left(\frac{1+\sqrt{5}}{2}\right) c_1 + \left(\frac{1-\sqrt{5}}{2}\right) c_2$$



# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}, n \geq 3, b_1 = 1$  and  $b_2 = 2$

$$r^2 - r - 1 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \frac{1+\sqrt{5}}{2}, r_2 = \frac{1-\sqrt{5}}{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^n + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^n$

$$b_0 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^0 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^0 \rightarrow 1 = c_1 + c_2$$

$$b_1 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^1 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^1 \rightarrow 2 = \left(\frac{1+\sqrt{5}}{2}\right) c_1 + \left(\frac{1-\sqrt{5}}{2}\right) c_2$$

$$c_1 = 1/\sqrt{5}, c_2 = -1/\sqrt{5}$$

# Recurrence Relations

- Suppose we have a  $2 \times n$  chessboard and we wish to cover it using  $2 \times 1$  and  $1 \times 2$  dominoes. In how many different ways can we cover it ?
- $b_n = b_{n-1} + b_{n-2}$ ,  $n \geq 3$ ,  $b_1 = 1$  and  $b_2 = 2$

$$r^2 - r - 1 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \frac{1+\sqrt{5}}{2}, r_2 = \frac{1-\sqrt{5}}{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^n + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^n$

$$b_0 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^0 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^0 \rightarrow 1 = c_1 + c_2$$

$$b_1 = c_1 \left(\frac{1+\sqrt{5}}{2}\right)^1 + c_2 \left(\frac{1-\sqrt{5}}{2}\right)^1 \rightarrow 2 = \left(\frac{1+\sqrt{5}}{2}\right) c_1 + \left(\frac{1-\sqrt{5}}{2}\right) c_2$$

$$c_1 = 1/\sqrt{5}, c_2 = -1/\sqrt{5} \quad \longrightarrow \quad b_n = \frac{1}{\sqrt{5}} \left( \left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \right)$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$3$$


$$1 + 2$$

$$2 + 1$$

$$1 + 1 + 1$$


# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:




# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



# Recurrence Relations


- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$\begin{array}{c} 3 \\ 1 + 1 + 1 \end{array}$$



# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

Diagram illustrating the partitions of 3 and their palindromic status:

- $3$  (red)
- $1 + 2$
- $2 + 1$
- $1 + 1 + 1$  (red)

palindrome

they are read the same from left to right, right to left

Blue arrows point from the word "palindrome" to the red numbers 3 and 1+1+1.

- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

Diagram illustrating the recursive generation of partitions for  $n=5$  from  $n=3$ :

$3$

$1 + 1 + 1$

Blue arrow points from  $1 + 1 + 1$  to the partitions of 5 below.

$5$

$2 + 1 + 2$

$1 + 3 + 1$

$1 + 1 + 1 + 1 + 1$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

3  
1 + 2  
2 + 1  
1 + 1 + 1

palindrome  
they are read  
the same from  
left to right,  
right to left

The diagram shows the four partitions of 3. Blue arrows point from the word 'palindrome' to the partitions '1 + 2', '2 + 1', and '1 + 1 + 1'. The number '3' at the top is also in red.

- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

3  
1 + 1 + 1

4  
1 + 2 + 1  
2 + 2  
1 + 1 + 1 + 1

5  
2 + 1 + 2  
1 + 3 + 1  
1 + 1 + 1 + 1 + 1

A blue arrow points from the partition '1 + 1 + 1' (under 3) to the partition '2 + 1 + 2' (under 5).

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:

$$\begin{array}{c}
 3 \\
 1 + 2 \\
 2 + 1 \\
 1 + 1 + 1
 \end{array}$$

palindrome  
they are read  
the same from  
left to right,  
right to left

- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$\begin{array}{c}
 3 \\
 1 + 1 + 1
 \end{array}$$


$$\begin{array}{c}
 4 \\
 1 + 2 + 1 \\
 2 + 2 \\
 1 + 1 + 1 + 1
 \end{array}$$

$$\begin{array}{c}
 6 \\
 2 + 2 + 2 \\
 3 + 3 \\
 2 + 1 + 1 + 2 \\
 1 + 4 + 1 \\
 1 + 1 + 2 + 1 + 1 \\
 1 + 2 + 2 + 1 \\
 1 + 1 + 1 + 1 + 1 + 1
 \end{array}$$

$$\begin{array}{c}
 5 \\
 2 + 1 + 2 \\
 1 + 3 + 1 \\
 1 + 1 + 1 + 1 + 1
 \end{array}$$

# Recurrence Relations


- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:




- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$b_n = 2b_{n-2}, n \geq 3, b_1 = 1 \text{ and } b_2 = 2$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:




- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$b_n = 2b_{n-2}, n \geq 3, b_1 = 1 \text{ and } b_2 = 2$$

$$r^2 - 2 = 0 \quad (\text{characteristic equation})$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?


$$b_n = 2b_{n-2}, n \geq 3, b_1 = 1 \text{ and } b_2 = 2$$

$$r^2 - 2 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \sqrt{2}, r_2 = -\sqrt{2} \quad (\text{characteristic roots})$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$b_n = 2b_{n-2}, n \geq 3, b_1 = 1 \text{ and } b_2 = 2$$

$$r^2 - 2 = 0 \quad (\text{characteristic equation})$$


$$r_1 = \sqrt{2}, r_2 = -\sqrt{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1(\sqrt{2})^n + c_2(-\sqrt{2})^n$



# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$b_n = 2b_{n-2}, n \geq 3, b_1 = 1 \text{ and } b_2 = 2$$

$$r^2 - 2 = 0 \quad (\text{characteristic equation})$$


$$r_1 = \sqrt{2}, r_2 = -\sqrt{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1(\sqrt{2})^n + c_2(-\sqrt{2})^n$

$$b_0 = c_1(\sqrt{2})^0 + c_2(-\sqrt{2})^0$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$b_n = 2b_{n-2}, n \geq 3, b_1 = 1 \text{ and } b_2 = 2$$

$$r^2 - 2 = 0 \quad (\text{characteristic equation})$$


$$r_1 = \sqrt{2}, r_2 = -\sqrt{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1(\sqrt{2})^n + c_2(-\sqrt{2})^n$

$$b_0 = c_1(\sqrt{2})^0 + c_2(-\sqrt{2})^0 \rightarrow 1 = c_1 + c_2$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$b_n = 2b_{n-2}, n \geq 3, b_1 = 1 \text{ and } b_2 = 2$$

$$r^2 - 2 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \sqrt{2}, r_2 = -\sqrt{2} \quad (\text{characteristic roots})$$


the solution will be in the form of  $b_n = c_1(\sqrt{2})^n + c_2(-\sqrt{2})^n$

$$b_0 = c_1(\sqrt{2})^0 + c_2(-\sqrt{2})^0 \rightarrow 1 = c_1 + c_2$$

$$b_1 = c_1(\sqrt{2})^1 + c_2(-\sqrt{2})^1$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$b_n = 2b_{n-2}, n \geq 3, b_1 = 1 \text{ and } b_2 = 2$$

$$r^2 - 2 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \sqrt{2}, r_2 = -\sqrt{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1(\sqrt{2})^n + c_2(-\sqrt{2})^n$

$$b_0 = c_1(\sqrt{2})^0 + c_2(-\sqrt{2})^0 \rightarrow 1 = c_1 + c_2$$

$$b_1 = c_1(\sqrt{2})^1 + c_2(-\sqrt{2})^1 \rightarrow 2 = (\sqrt{2})c_1 + (-\sqrt{2})c_2$$

# Recurrence Relations

- 3 can be written as a sum of positive integers in 4 different ways:



- How many different palindromes can be found for a given  $n \in \mathbb{Z}^+$  ?

$$b_n = 2b_{n-2}, n \geq 3, b_1 = 1 \text{ and } b_2 = 2$$

$$r^2 - 2 = 0 \quad (\text{characteristic equation})$$

$$r_1 = \sqrt{2}, r_2 = -\sqrt{2} \quad (\text{characteristic roots})$$

the solution will be in the form of  $b_n = c_1(\sqrt{2})^n + c_2(-\sqrt{2})^n$

$$b_0 = c_1(\sqrt{2})^0 + c_2(-\sqrt{2})^0 \rightarrow 1 = c_1 + c_2$$

$$b_1 = c_1(\sqrt{2})^1 + c_2(-\sqrt{2})^1 \rightarrow 2 = (\sqrt{2})c_1 + (-\sqrt{2})c_2$$

$$b_n = \left(\frac{1}{2} + \frac{1}{2\sqrt{2}}\right)(\sqrt{2})^n + \left(\frac{1}{2} - \frac{1}{2\sqrt{2}}\right)(-\sqrt{2})^n$$