


COUNTING I

Murat Osmanoglu

Counting



- Ali and Buse eat lunch together in a specific restaurant regularly.

Counting



'soup' XOR 'sandwich'



'soup' AND 'sandwich'

- Ali and Buse eat lunch together in a specific restaurant regularly.

Counting



'soup' XOR 'sandwich'




'soup' AND 'sandwich'

- Ali and Buse eat lunch together in a specific restaurant regularly.
- restaurant has 9 choices for soup
and 16 choices for sandwich

Counting



'soup' XOR 'sandwich'



'soup' AND 'sandwich'

- Ali and Buse eat lunch together in a specific restaurant regularly.
- restaurant has 9 choices for soup
and 16 choices for sandwich
- How many different meals can Ali order ?
- How many different meals can Buse order ?

Counting

- Ali can order either one soup among 9 different soups or one sandwich among 16 different sandwiches.

Counting

- Ali can order either one soup among 9 different soups or one sandwich among 16 different sandwiches.
- apply the Sum Rule here : $9 + 16 = 25$ different choices

Counting

- Ali can order either one soup among 9 different soups or one sandwich among 16 different sandwiches.
- apply the Sum Rule here : $9 + 16 = 25$ different choices

Sum Rule

- If A and B are disjoint finite sets, then the number of ways of choosing a single element from A or B is

Counting

- Ali can order either one soup among 9 different soups or one sandwich among 16 different sandwiches.
- apply the Sum Rule here : $9 + 16 = 25$ different choices

Sum Rule

- If A and B are disjoint finite sets, then the number of ways of choosing a single element from A or B is

$$|A \cup B| = |A| + |B|$$

Counting

- Ali can order either one soup among 9 different soups or one sandwich among 16 different sandwiches.
- apply the Sum Rule here : $9 + 16 = 25$ different choices

Sum Rule

- If A and B are disjoint finite sets, then the number of ways of choosing a single element from A or B is

$$|A \cup B| = |A| + |B|$$

- If a task can be done in one of n_1 ways or in one of n_2 ways such that none from n_1 ways is the same as any from n_2 ways, then there are $n_1 + n_2$ different ways to do the task

Counting

- Ali can order either one soup among 9 different soups or one sandwich among 16 different sandwiches.
- apply the Sum Rule here : $9 + 16 = 25$ different choices

Sum Rule

- If A and B are disjoint finite sets, then the number of ways of choosing a single element from A or B is

$$|A \cup B| = |A| + |B|$$

- If a task can be done in one of n_1 ways or in one of n_2 ways such that none from n_1 ways is the same as any from n_2 ways, then there are $n_1 + n_2$ different ways to do the task
- If A_1, A_2, \dots, A_n are mutually disjoint sets ($A_i \cap A_j = \emptyset$), then the number of ways of choosing a single element from A_1 or A_2 or ... A_n is

$$|A_1 \cup \dots \cup A_n| = |A_1| + \dots + |A_n|$$

Counting

Counting

- Buse can choose one soup among 9 different choices; and for each choice of soup, she can have one sandwich among 16 different sandwiches.
consider the meal as a pair (soup, sandwich)

Counting

- Buse can choose one soup among 9 different choices; and for each choice of soup, she can have one sandwich among 16 different sandwiches.
consider the meal as a pair (soup, sandwich)
- apply the Product Rule here : $9 \cdot 16 = 144$ different choices

Counting

- Buse can choose one soup among 9 different choices; and for each choice of soup, she can have one sandwich among 16 different sandwiches.
consider the meal as a pair (soup, sandwich)
- apply the Product Rule here : $9 \cdot 16 = 144$ different choices

Product Rule

- If A and B are finite sets, then the number of ways of choosing an element from A and an element from B is

Counting

- Buse can choose one soup among 9 different choices; and for each choice of soup, she can have one sandwich among 16 different sandwiches.
consider the meal as a pair (soup, sandwich)
- apply the Product Rule here : $9 \cdot 16 = 144$ different choices

Product Rule

- If A and B are finite sets, then the number of ways of choosing an element from A and an element from B is

$$|A \times B| = |A| \cdot |B|$$

Counting

- Buse can choose one one soup among 9 different choices; and for each choice of soup, she can have one sandwich among 16 different sandwiches.
consider the meal as a pair (soup, sandwich)
- apply the Product Rule here : $9 \cdot 16 = 144$ different choices

Product Rule

- If A and B are finite sets, then the number of ways of choosing an element from A and an element from B is

$$|A \times B| = |A| \cdot |B|$$

- Suppose that a task can be broken into a sequence of two small tasks. If there are n_1 ways to do the first task, and for each one there are n_2 ways to do the second task, then there are $n_1 \cdot n_2$ different ways to finish the task

Counting

- Buse can choose one soup among 9 different choices; and for each choice of soup, she can have one sandwich among 16 different sandwiches.
consider the meal as a pair (soup, sandwich)
- apply the Product Rule here : $9 \cdot 16 = 144$ different choices

Product Rule

- If A and B are finite sets, then the number of ways of choosing an element from A and an element from B is

$$|A \times B| = |A| \cdot |B|$$

- Suppose that a task can be broken into a sequence of two small tasks. If there are n_1 ways to do the first task, and for each one there are n_2 ways to do the second task, then there are $n_1 \cdot n_2$ different ways to finish the task
- If A_1, A_2, \dots, A_n are finite sets, then the number of ways of choosing an element from A_1, \dots , an element from A_n is

$$|A_1 \times \dots \times A_n| = |A_1| \cdot |A_2| \cdot \dots \cdot |A_n|$$

Counting

- How many bit-strings can you create with 3-digits ?

Counting

- How many bit-strings can you create with 3-digits ?

101, 001, 110, . . .

Counting

- How many bit-strings can you create with 3-digits ?

101, 001, 110, . . .

— — —

Counting

- How many bit-strings can you create with 3-digits ?

101, 001, 110, . . .

$$\underline{\quad} \underline{\quad} \underline{\quad} = 8$$

Counting

- How many bit-strings can you create with 3-digits ?

101, 001, 110, . . .

$$\underline{2} \times \underline{2} \times \underline{2} = 8$$

- What is the number of subsets of the set {a, b, c} ?

Counting

- How many bit-strings can you create with 3-digits ?

101, 001, 110, . . .

$$\underline{\quad} \times \underline{\quad} \times \underline{\quad} = 8$$

- What is the number of subsets of the set {a, b, c} ?

{a, b, c}

{a, b}

{c}

Counting

- How many bit-strings can you create with 3-digits ?

101, 001, 110, . . .

$$\underline{2} \times \underline{2} \times \underline{2} = 8$$


- What is the number of subsets of the set $\{a, b, c\}$?

$\{a, b, c\} \longrightarrow 111$


$\{a, b\} \longrightarrow 110$

$\{c\} \longrightarrow 001$


Counting



Counting




Counting



you pick one ball at a time


Counting



you pick one ball at a time

How many 3-digits numbers can you create with the picked numbers ?

Counting




you pick one ball at a time

How many 3-digits numbers can you create with the picked numbers ?

- If you leave them to the bag after you pick

Counting




you pick one ball at a time

How many 3-digits numbers can you create with the picked numbers ?

- If you leave them to the bag after you pick
- If you keep them after you pick

Counting



you pick one ball at a time

How many 3-digits numbers can you create with the picked numbers ?


- If you leave them to the bag after you pick

— — —

- If you keep them after you pick

— — —

Counting



you pick one ball at a time


How many 3-digits numbers can you create with the picked numbers ?

- If you leave them to the bag after you pick

$$\underline{\quad} \times \underline{\quad} \times \underline{\quad} = 125$$

- If you keep them after you pick

Counting



you pick one ball at a time

How many 3-digits numbers can you create with the picked numbers ?


- If you leave them to the bag after you pick

$$\underline{5} \times \underline{5} \times \underline{5} = 125$$


- If you keep them after you pick

$$\underline{5} \times \underline{4} \times \underline{3} = 60$$

Counting




Counting



In how many different ways can you go from the city A to the city B ?


Counting



In how many different ways can you go from the city A to the city B ?

$$9 + 11 + 10 = 30$$

Counting




In how many different ways can you go from the city A to the city B ?

$$9 + 11 + 10 = 30$$

In how many different ways can you go to B and come back to A ?

Counting



In how many different ways can you go from the city A to the city B ?

$$9 + 11 + 10 = 30$$

In how many different ways can you go to B and come back to A ?

$$30 \times 30 = 900$$

Counting

- You prepare a meal for your friends

Counting

- You prepare a meal for your friends
- There are 5 kinds of bagels, 7 kinds of sandwiches, 6 drinks (hot coffee, hot tea, iced tea, cola, orange juice, apple juice)

Counting

- You prepare a meal for your friends
- There are 5 kinds of bagels, 7 kinds of sandwiches, 6 drinks (hot coffee, hot tea, iced tea, cola, orange juice, apple juice)
- Bagels served with hot drinks and sandwiches served with cold drinks

Counting

- You prepare a meal for your friends
- There are 5 kinds of bagels, 7 kinds of sandwiches, 6 drinks (hot coffee, hot tea, iced tea, cola, orange juice, apple juice)
- Bagels served with hot drinks and sandwiches served with cold drinks

How many different meals can you prepare?

Counting

- You prepare a meal for your friends
- There are 5 kinds of bagels, 7 kinds of sandwiches, 6 drinks (hot coffee, hot tea, iced tea, cola, orange juice, apple juice)
- Bagels served with hot drinks and sandwiches served with cold drinks

How many different meals can you prepare?

(bagel, hot) or (sand, cold)

Counting

- You prepare a meal for your friends
- There are 5 kinds of bagels, 7 kinds of sandwiches, 6 drinks (hot coffee, hot tea, iced tea, cola, orange juice, apple juice)
- Bagels served with hot drinks and sandwiches served with cold drinks

How many different meals can you prepare?

(bagel, hot) or (sand, cold)

$$5 \times 2 \quad +$$

Counting

- You prepare a meal for your friends
- There are 5 kinds of bagels, 7 kinds of sandwiches, 6 drinks (hot coffee, hot tea, iced tea, cola, orange juice, apple juice)
- Bagels served with hot drinks and sandwiches served with cold drinks

How many different meals can you prepare?

(bagel, hot) or (sand, cold)

$$5 \times 2 \quad + \quad 7 \times 4$$

Counting

- You prepare a meal for your friends
- There are 5 kinds of bagels, 7 kinds of sandwiches, 6 drinks (hot coffee, hot tea, iced tea, cola, orange juice, apple juice)
- Bagels served with hot drinks and sandwiches served with cold drinks

How many different meals can you prepare?

(bagel, hot) or (sand, cold)

$$5 \times 2 + 7 \times 4 = 38$$

Counting

- How many different licence plates can be created if a plate contains of 2 digits indicating the city the car registered followed by 3 uppercase Turkish letters followed by 3 digits ?


Counting

- How many different licence plates can be created if a plate contains of 2 digits indicating the city the car registered followed by 3 uppercase Turkish letters followed by 3 digits ?

— — — — — — —


Counting

- How many different licence plates can be created if a plate contains of 2 digits indicating the city the car registered followed by 3 uppercase Turkish letters followed by 3 digits ?



Counting

- How many different licence plates can be created if a plate contains of 2 digits indicating the city the car registered followed by 3 uppercase Turkish letters followed by 3 digits ?



Counting

- How many different licence plates can be created if a plate contains of 2 digits indicating the city the car registered followed by 3 uppercase Turkish letters followed by 3 digits ?

$$\begin{array}{ccccccc} \underline{81} & \times & \underline{23} & \times & \underline{23} & \times & \underline{23} & \times & \underline{\quad} & \underline{\quad} & \underline{\quad} \\ \underbrace{\hspace{1.5cm}} & & \underbrace{\hspace{3.5cm}} & & & & \underbrace{\hspace{3.5cm}} & & & & \\ \text{indicating} & & \text{Turkish letters} & & & & \text{numbers} & & & & \\ \text{the city} & & & & & & & & & & \end{array}$$

Counting

- How many different licence plates can be created if a plate contains of 2 digits indicating the city the car registered followed by 3 uppercase Turkish letters followed by 3 digits ?


$$\underbrace{81}_{\text{indicating the city}} \times \underbrace{23 \times 23 \times 23}_{\text{Turkish letters}} \times \underbrace{10 \times 10 \times 10}_{\text{numbers}}$$

Counting


- How many different licence plates can be created if a plate contains of 2 digits indicating the city the car registered followed by 3 uppercase Turkish letters followed by 3 digits ?

$$\underbrace{81}_{\text{indicating the city}} \times \underbrace{23 \times 23 \times 23}_{\text{Turkish letters}} \times \underbrace{10 \times 10 \times 10}_{\text{numbers}} = 985527000$$

Counting




Counting



In how many different ways can you go from the city A to the city C ?


Counting



In how many different ways can you go from the city A to the city C ?

$$4 \times 3 + 2 = 14$$

Counting




In how many different ways can you go from the city A to the city C ?

$$4 \times 3 + 2 = 14$$

In how many different ways can you go to C and come back to A ?

Counting




In how many different ways can you go from the city A to the city C ?

$$4 \times 3 + 2 = 14$$

In how many different ways can you go to C and come back to A ?

$$14 \times 14 = 196$$

Counting



In how many different ways can you go from the city A to the city C ?


$$4 \times 3 + 2 = 14$$

In how many different ways can you go to C and come back to A ?

$$14 \times 14 = 196$$

In how many different ways can you go to C and come back to A so that you can use same route to come back?

Counting



In how many different ways can you go from the city A to the city C ?

$$4 \times 3 + 2 = 14$$

In how many different ways can you go to C and come back to A ?

$$14 \times 14 = 196$$

In how many different ways can you go to C and come back to A so that you can use same route to come back?

$$14 \times 13 = 182$$

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

— — — — —

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

$$\underline{10} \times \underline{9} \times \underline{8} \times \underline{7} \times \underline{6} = 30240$$

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

$$\underline{10} \times \underline{9} \times \underline{8} \times \underline{7} \times \underline{6} = 30240$$

How many different arrangements can we make for all students ?

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

$$\underline{10} \times \underline{9} \times \underline{8} \times \underline{7} \times \underline{6} = 30240$$

How many different arrangements can we make for all students ?

$$\underline{10} \times \underline{9} \times \underline{8} \times \underline{7} \times \dots \times \underline{1} = 3628800$$

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

$$10 \times 9 \times 8 \times 7 \times 6$$

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

$$10 \times 9 \times 8 \times 7 \times 6 \times \frac{5 \times 4 \times 3 \times 2 \times 1}{5 \times 4 \times 3 \times 2 \times 1}$$

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

$$10 \times 9 \times 8 \times 7 \times 6 \frac{\times 5 \times 4 \times 3 \times 2 \times 1}{5 \times 4 \times 3 \times 2 \times 1} = \frac{10!}{5!}$$

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

$$10 \times 9 \times 8 \times 7 \times 6 \frac{\times 5 \times 4 \times 3 \times 2 \times 1}{5 \times 4 \times 3 \times 2 \times 1} = \frac{10!}{5!}$$

- the number of different permutation of size 5 for 10 objects

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

$$10 \times 9 \times 8 \times 7 \times 6 \frac{\times 5 \times 4 \times 3 \times 2 \times 1}{5 \times 4 \times 3 \times 2 \times 1} = \frac{10!}{5!}$$

- the number of different permutation of size 5 for 10 objects
- the number of different permutation of size r for n objects

Permutation

Assume there are 10 students. We choose five of them, and make them sit together to get a picture

How many such arrangements can we make?

- assign them numbers 0, 1, 2, 3, 4, 5, 6, 7, 8, 9
- possible arrangements 13429, 60938, 19082, ...

$$10 \times 9 \times 8 \times 7 \times 6 \frac{\times 5 \times 4 \times 3 \times 2 \times 1}{5 \times 4 \times 3 \times 2 \times 1} = \frac{10!}{5!}$$

- the number of different permutation of size 5 for 10 objects
- the number of different permutation of size r for n objects

$$P(n,r) = \frac{n!}{(n-r)!}$$

Permutation

- Using the letters of the word 'COMPUTER' how many different words can you create ?

Permutation

- Using the letters of the word 'COMPUTER' how many different words can you create ?

8!

Permutation

- Using the letters of the word 'COMPUTER' how many different words can you create ?

8!

- Using the letters of the word 'COMPUTER', how many different words of length 5 can you create ?

Permutation

- Using the letters of the word 'COMPUTER' how many different words can you create ?

8!

- Using the letters of the word 'COMPUTER', how many different words of length 5 can you create ?

$$P(8, 5) = 8! / 5! = 336$$

Permutation

- Using the letters of the word 'COMPUTER' how many different words can you create ?

8!

- Using the letters of the word 'COMPUTER', how many different words of length 5 can you create ?

$$P(8, 5) = 8! / 5! = 336$$

- If repetitions are allowed, how many different words of length 5 can you create ?

Permutation

- Using the letters of the word 'COMPUTER' how many different words can you create ?

$$8!$$

- Using the letters of the word 'COMPUTER', how many different words of length 5 can you create ?

$$P(8, 5) = 8! / 5! = 336$$

- If repetitions are allowed, how many different words of length 5 can you create ?

$$8 \times 8 \times 8 \times 8 \times 8 = 32768$$

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

4!

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

~~4!~~

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

~~4!~~

BALL, BLAL, BLLA, ABLL, ALBL, ALLB
LBAL, LBLA, LABL, LALB, LLAB, LLBA

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

~~4!~~ 12

BALL, BLAL, BLLA, ABLL, ALBL, ALLB
LBAL, LBLA, LABL, LALB, LLAB, LLBA

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

~~4!~~ 12

BALL, BLAL, BLLA, ABLL, ALBL, ALLB
LBAL, LBLA, LABL, LALB, LLAB, LLBA

- Using the letters of the word 'ABARA', how many different words can you create?

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

~~4!~~ 12

BALL, BLAL, BLLA, ABLL, ALBL, ALLB
LBAL, LBLA, LABL, LALB, LLAB, LLBA

- Using the letters of the word 'ABARA', how many different words can you create?

$A_1BA_3RA_2$

- pretend they are different A's

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

~~4!~~ 12

BALL, BLAL, BLLA, ABLL, ALBL, ALLB
LBAL, LBLA, LABL, LALB, LLAB, LLBA

- Using the letters of the word 'ABARA', how many different words can you create?

$A_1BA_3RA_2$

- pretend they are different A's
- fix other letters and reorder A's

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

$$\cancel{4!} \quad 12$$

BALL, BLAL, BLLA, ABLL, ALBL, ALLB
LBAL, LBLA, LABL, LALB, LLAB, LLBA

- Using the letters of the word 'ABARA', how many different words can you create?

$A_1BA_3RA_2$ $A_2BA_3RA_1$ $A_3BA_2RA_1$
 $A_1BA_2RA_3$ $A_2BA_1RA_3$ $A_3BA_1RA_2$

- pretend they are different A's
- fix other letters and reorder A's

Permutation

- Using the letters of the word 'BALL', how many different words can you create?

$$\cancel{4!} \quad 12$$

BALL, BLAL, BLLA, ABLL, ALBL, ALLB
LBAL, LBLA, LABL, LALB, LLAB, LLBA

- Using the letters of the word 'ABARA', how many different words can you create?

$A_1BA_3RA_2$ $A_2BA_3RA_1$ $A_3BA_2RA_1$
 $A_1BA_2RA_3$ $A_2BA_1RA_3$ $A_3BA_1RA_2$

$$5! / 3! = 20$$


- pretend they are different A's
- fix other letters and reorder A's

Permutation

- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?

Permutation



- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?



A B C D E

Permutation



- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?



A	B	C	D	E
E	A	B	C	D

Permutation



- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?



A	B	C	D	E
E	A	B	C	D
D	E	A	B	C
C	D	E	A	B
B	C	D	E	A

Permutation

- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?




A	B	C	D	E
E	A	B	C	D
D	E	A	B	C
C	D	E	A	B
B	C	D	E	A

A blue bracket is drawn under the last four rows of the table.

For each circular arrangement, there are 5 linear arrangements


Permutation

- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?



$$5 \times (\# \text{ of circular}) = (\# \text{ of linear})$$


A B C D E
E A B C D
D E A B C
C D E A B
B C D E A



For each circular arrangement, there are 5 linear arrangements

Permutation


- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?



$$5 \times (\# \text{ of circular}) = (\# \text{ of linear})$$

$$(\# \text{ of circular}) = 5! / 5 = 24$$


A B C D E
E A B C D
D E A B C
C D E A B
B C D E A



For each circular arrangement, there are 5 linear arrangements


Permutation

- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?



Permutation


- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?



- fix one of them
- permute all others as in linear

Permutation

- There are 5 people : A, B, C, D, E
- They sit around a round table. How many different arrangements are possible ?



- fix one of them
- permute all others as in linear
- $4!$

Permutation


- You invite 2 couples for the dinner (3 couples at total)

Permutation

- You invite 2 couples for the dinner (3 couples at total)
- You have a around table. How many circular arrangements can you make such that no two women or no two men sit together ?


Permutation

- You invite 2 couples for the dinner (3 couples at total)
- You have a around table. How many circular arrangements can you make such that no two women or no two men sit together ?




Permutation

- You invite 2 couples for the dinner (3 couples at total)
- You have a around table. How many circular arrangements can you make such that no two women or no two men sit together ?




Permutation

- You invite 2 couples for the dinner (3 couples at total)
- You have a around table. How many circular arrangements can you make such that no two women or no two men sit together ?



Permutation


- You invite 2 couples for the dinner (3 couples at total)
- You have a around table. How many circular arrangements can you make such that no two women or no two men sit together ?



- $3 \times 2 \times 2 \times 1 \times 1 = 12$

Permutation

- You invite 2 couples for the dinner (3 couples at total)
- You have a around table. How many circular arrangements can you make such that no two women or no two men sit together ?




- $3 \times 2 \times 2 \times 1 \times 1 = 12$

- $3! \times 2! = 12$


Combinations




Combinations





Combinations



- Assume you play a game such that a player holds 3 cards.




Combinations




- Assume you play a game such that a player holds 3 cards.
- How many different hands can you create?

$$P(52,3) = 52! / (52-3)!$$

Combinations




- Assume you play a game such that a player holds 3 cards.
- How many different hands can you create?




$$P(52,3) = 52! / (52-3)!$$

Combinations




- Assume you play a game such that a player holds 3 cards.
- How many different hands can you create?




you count them as one

$$P(52,3) = 52! / (52-3)!$$

Combinations



- Assume you play a game such that a player holds 3 cards.
- How many different hands can you create?



you count them as one

$$52! / [(52-3)! \cdot 3!]$$

$$P(52,3) = 52! / (52-3)!$$

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student picks 4 questions from the first 5 and 3 questions from the last 5, how many different answer sheets can he prepare?

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student picks 4 questions from the first 5 and 3 questions from the last 5, how many different answer sheets can he prepare?

1 2 3 4 5

6 7 8 9 10

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student picks 4 questions from the first 5 and 3 questions from the last 5, how many different answer sheets can he prepare?

1 2 3 4 5

6 7 8 9 10

(,)

Combinations

The number of different selections of r elements out of n distinct objects :


$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student picks 4 questions from the first 5 and 3 questions from the last 5, how many different answer sheets can he prepare?



Combinations

The number of different selections of r elements out of n distinct objects :


$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student picks 4 questions from the first 5 and 3 questions from the last 5, how many different answer sheets can he prepare?



1 2 3 4 5 6 7 8 9 10

(,) $\binom{5}{4} \times \binom{5}{3} =$

Combinations

The number of different selections of r elements out of n distinct objects :


$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student picks 4 questions from the first 5 and 3 questions from the last 5, how many different answer sheets can he prepare?



$$\binom{5}{4} \times \binom{5}{3} = \frac{5!}{1! \cdot 4!} \times \frac{5!}{2! \cdot 3!} = 50$$

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student is required to pick at least 3 questions from the first 5, how many different answer sheets can he prepare?

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student is required to pick at least 3 questions from the first 5, how many different answer sheets can he prepare?

$$\binom{5}{3} \binom{5}{4} +$$

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student is required to pick at least 3 questions from the first 5, how many different answer sheets can he prepare?

$$\binom{5}{3} \binom{5}{4} + \binom{5}{4} \binom{5}{3}$$

Combinations

The number of different selections of r elements out of n distinct objects :

$$C(n,r) = n! / [(n-r)! \cdot r!]$$

There are 10 questions for an exam, and the student are required to choose 7 of them to answer.

- How many different answer sheets can a student prepare?

$$10! / (7! \cdot 3!) = (10 \cdot 9 \cdot 8) / 3! = 120$$

- If the student is required to pick at least 3 questions from the first 5, how many different answer sheets can he prepare?

$$\binom{5}{3} \binom{5}{4} + \binom{5}{4} \binom{5}{3} + \binom{5}{5} \binom{5}{2}$$

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

- Suppose you are playing a game with five cards.
How many different hands can you have ?

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

- Suppose you are playing a game with five cards.

How many different hands can you have ? $\binom{52}{5}$

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

- Suppose you are playing a game with five cards.

How many different hands can you have ? $\binom{52}{5}$

How many of them contains no club ?

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

- Suppose you are playing a game with five cards.

How many different hands can you have ? $\binom{52}{5}$

How many of them contains no club ? $\binom{39}{5}$

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

- Suppose you are playing a game with five cards.

How many different hands can you have ? $\binom{52}{5}$

How many of them contains no club ? $\binom{39}{5}$

How many of them contains at least two clubs ?

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

- Suppose you are playing a game with five cards.

How many different hands can you have ? $\binom{52}{5}$

How many of them contains no club ? $\binom{39}{5}$

How many of them contains at least two clubs ?

$$\binom{13}{2} \cdot \binom{39}{3}$$

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

- Suppose you are playing a game with five cards.

How many different hands can you have ? $\binom{52}{5}$

How many of them contains no club ? $\binom{39}{5}$

How many of them contains at least two clubs ?

$$\binom{13}{2} \cdot \binom{39}{3} + \binom{13}{3} \cdot \binom{39}{2}$$

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

- Suppose you are playing a game with five cards.

How many different hands can you have ? $\binom{52}{5}$

How many of them contains no club ? $\binom{39}{5}$

How many of them contains at least two clubs ?

$$\binom{13}{2} \cdot \binom{39}{3} + \binom{13}{3} \cdot \binom{39}{2} + \binom{13}{4} \cdot \binom{39}{1}$$

Combinations

- We are forming soccer teams with 6 players for a tournament from 30 boys and 25 girls. How many different 4 teams (2 for girls and 2 for boys) can you create ?

$$\binom{30}{6} \binom{24}{6} \binom{25}{6} \binom{19}{6}$$

- Suppose you are playing a game with five cards.

How many different hands can you have ? $\binom{52}{5}$

How many of them contains no club ? $\binom{39}{5}$

How many of them contains at least two clubs ?

$$\binom{13}{2} \cdot \binom{39}{3} + \binom{13}{3} \cdot \binom{39}{2} + \binom{13}{4} \cdot \binom{39}{1} + \binom{13}{5} \cdot \binom{39}{0}$$

Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$2 \times 2 \times \dots \times 2 = 2^8$$

Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$2 \times 2 \times \dots \times 2 = 2^8$$

- How many of them contains exactly three 1's ?

Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$2 \times 2 \times \dots \times 2 = 2^8$$

- How many of them contains exactly three 1's ?

— 1 — 1 1 — — —

Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$2 \times 2 \times \dots \times 2 = 2^8$$

- How many of them contains exactly three 1's ?

— 1 — 1 1 — — —

Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$2 \times 2 \times \dots \times 2 = 2^8$$

- How many of them contains exactly three 1's ?



Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$2 \times 2 \times \dots \times 2 = 2^8$$

- How many of them contains exactly three 1's ?

$$\binom{8}{3}$$



Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$2 \times 2 \times \dots \times 2 = 2^8$$

- How many of them contains exactly three 1's ?

$$\binom{8}{3}$$

Assume $x = x_1x_2\dots x_n$, the weight of a given encoding is

$$w(x) = x_1 + x_2 + \dots + x_n$$

Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$2 \times 2 \times \dots \times 2 = 2^8$$

- How many of them contains exactly three 1's ?

$$\binom{8}{3}$$

Assume $x = x_1x_2\dots x_n$, the weight of a given encoding is

$$w(x) = x_1 + x_2 + \dots + x_n$$

- How many encodings of length 8 have even weight ?

Combinations

$\Sigma = \{0, 1\}$.

Let's use this alphabet to create three digits encoding:

000, 010, 111, 011, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$2 \times 2 \times \dots \times 2 = 2^8$$

- How many of them contains exactly three 1's ?

$$\binom{8}{3}$$

Assume $x = x_1x_2\dots x_n$, the weight of a given encoding is

$$w(x) = x_1 + x_2 + \dots + x_n$$

- How many encodings of length 8 have even weight ?

$$\binom{8}{0} + \binom{8}{2} + \dots + \binom{8}{8}$$

Combinations

$\Sigma = \{0, 1, 2\}$.

Let's use this alphabet to create three digits encoding:

020, 010, 112, 211, . . .

- How many encodings of length 8 can you create using the alphabet Σ ?
- How many of them contains exactly three 1's ?

Assume $x = x_1x_2\dots x_n$, the weight of a given encoding is

$$w(x) = x_1 + x_2 + \dots + x_n$$

- How many encodings of length 8 have even weight ?

Combinations

$\Sigma = \{0, 1, 2\}$.

Let's use this alphabet to create three digits encoding:

020, 010, 112, 211, . . .

- How many encodings of length 8 can you create using the alphabet Σ ?

$$3 \times 3 \times \dots \times 3 = 3^8$$

- How many of them contains exactly three 1's ?

Assume $x = x_1x_2\dots x_n$, the weight of a given encoding is

$$w(x) = x_1 + x_2 + \dots + x_n$$

- How many encodings of length 8 have even weight ?

Combinations

$\Sigma = \{0, 1, 2\}$.

Let's use this alphabet to create three digits encoding:

020, 010, 112, 211, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$3 \times 3 \times \dots \times 3 = 3^8$$

- How many of them contains exactly three 1's ?

$$\binom{8}{3} \cdot 2^{8-3}$$

Assume $x = x_1x_2\dots x_n$, the weight of a given encoding is

$$w(x) = x_1 + x_2 + \dots + x_n$$

- How many encodings of length 8 have even weight ?

Combinations

$\Sigma = \{0, 1, 2\}$.

Let's use this alphabet to create three digits encoding:

020, 010, 112, 211, ...

- How many encodings of length 8 can you create using the alphabet Σ ?

$$3 \times 3 \times \dots \times 3 = 3^8$$

- How many of them contains exactly three 1's ?

$$\binom{8}{3} \cdot 2^{8-3}$$

Assume $x = x_1x_2\dots x_n$, the weight of a given encoding is

$$w(x) = x_1 + x_2 + \dots + x_n$$

- How many encodings of length 8 have even weight ?

$$\binom{8}{0} 2^8 + \dots + \binom{8}{8} 2^0$$

Binomial Theorem

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

$$\binom{n}{i} x^{n-i} y^i$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 7 \text{ and } i = 2$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 7 \text{ and } i = 2 \Rightarrow \binom{7}{2} x^5 y^2$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 7 \text{ and } i = 2 \Rightarrow \binom{7}{2} x^5 y^2 = \frac{7 \cdot 6}{2} x^5 y^2 = \mathbf{21} x^5 y^2$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 7 \text{ and } i = 2 \Rightarrow \binom{7}{2} x^5 y^2 = \frac{7 \cdot 6}{2} x^5 y^2 = \mathbf{21} x^5 y^2$$

- What is the coefficient of $x^{10}y^{15}$ in the expansion of $(2x - 3y)^{25}$?

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 7 \text{ and } i = 2 \Rightarrow \binom{7}{2} x^5 y^2 = \frac{7 \cdot 6}{2} x^5 y^2 = \mathbf{21} x^5 y^2$$

- What is the coefficient of $x^{10}y^{15}$ in the expansion of $(2x - 3y)^{25}$?

$$\binom{n}{i} x^{n-i} y^i$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 7 \text{ and } i = 2 \Rightarrow \binom{7}{2} x^5 y^2 = \frac{7 \cdot 6}{2} x^5 y^2 = \mathbf{21} x^5 y^2$$

- What is the coefficient of $x^{10}y^{15}$ in the expansion of $(2x - 3y)^{25}$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 25 \text{ and } i = 15$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 7 \text{ and } i = 2 \Rightarrow \binom{7}{2} x^5 y^2 = \frac{7 \cdot 6}{2} x^5 y^2 = \mathbf{21} x^5 y^2$$

- What is the coefficient of $x^{10}y^{15}$ in the expansion of $(2x - 3y)^{25}$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 25 \text{ and } i = 15 \Rightarrow \binom{25}{15} (2x)^{10} (-3y)^{15}$$

Binomial Theorem

- $(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$
 $(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of x^5y^2 in the expansion of $(x + y)^7$?

$$\binom{n}{i} x^{n-i} y^i \Rightarrow n = 7 \text{ and } i = 2 \Rightarrow \binom{7}{2} x^5 y^2 = \frac{7 \cdot 6}{2} x^5 y^2 = \mathbf{21} x^5 y^2$$

- What is the coefficient of $x^{10}y^{15}$ in the expansion of $(2x - 3y)^{25}$?

$$\begin{aligned} \binom{n}{i} x^{n-i} y^i &\Rightarrow n = 25 \text{ and } i = 15 \Rightarrow \binom{25}{15} (2x)^{10} (-3y)^{15} \\ &\Rightarrow \binom{25}{15} \mathbf{2^{10} (-3)^{15}} x^{10} y^{15} \end{aligned}$$

Binomial Theorem

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of $x_1^{n_1} x_2^{n_2} \dots x_k^{n_k}$ in the expansion of $(x_1 + x_2 + \dots + x_k)^n$?

Binomial Theorem

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of $x_1^{n_1} x_2^{n_2} \dots x_k^{n_k}$ in the expansion of $(x_1 + x_2 + \dots + x_k)^n$?

$$\binom{n}{n_1}$$

Binomial Theorem

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of $x_1^{n_1} x_2^{n_2} \dots x_k^{n_k}$ in the expansion of $(x_1 + x_2 + \dots + x_k)^n$?

$$\binom{n}{n_1} \cdot \binom{n - n_1}{n_2}$$

Binomial Theorem

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of $x_1^{n_1} x_2^{n_2} \dots x_k^{n_k}$ in the expansion of $(x_1 + x_2 + \dots + x_k)^n$?

$$\binom{n}{n_1} \cdot \binom{n - n_1}{n_2} \cdot \binom{n - n_1 - n_2}{n_3} \dots \binom{n - n_1 - \dots - n_{k-1}}{n_k}$$

Binomial Theorem

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of $x_1^{n_1} x_2^{n_2} \dots x_k^{n_k}$ in the expansion of $(x_1 + x_2 + \dots + x_k)^n$?

$$\binom{n}{n_1} \cdot \binom{n - n_1}{n_2} \cdot \binom{n - n_1 - n_2}{n_3} \dots \binom{n - n_1 - \dots - n_{k-1}}{n_k} = \frac{n!}{n_1! \dots n_k!}$$

Binomial Theorem

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of $x_1^{n_1} x_2^{n_2} \dots x_k^{n_k}$ in the expansion of $(x_1 + x_2 + \dots + x_k)^n$?

$$\binom{n}{n_1} \cdot \binom{n - n_1}{n_2} \cdot \binom{n - n_1 - n_2}{n_3} \dots \binom{n - n_1 - \dots - n_{k-1}}{n_k} = \frac{n!}{n_1! \dots n_k!}$$

- What is the coefficient of $x^3 y^2 z^2$ in the expansion of $(x + y + z)^7$?

Binomial Theorem

Let x and y be variables and n be non-negative integer, then

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \dots + \binom{n}{n} y^n$$

- What is the coefficient of $x_1^{n_1} x_2^{n_2} \dots x_k^{n_k}$ in the expansion of $(x_1 + x_2 + \dots + x_k)^n$?

$$\binom{n}{n_1} \cdot \binom{n - n_1}{n_2} \cdot \binom{n - n_1 - n_2}{n_3} \dots \binom{n - n_1 - \dots - n_{k-1}}{n_k} = \frac{n!}{n_1! \dots n_k!}$$

- What is the coefficient of $x^3 y^2 z^2$ in the expansion of $(x + y + z)^7$?

$$\frac{7!}{3! 2! 2!} = 210$$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$.

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

\emptyset

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

\emptyset

$\{1\}, \{2\}, \{3\}, \dots, \{n\}$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

\emptyset

$\{1\}, \{2\}, \{3\}, \dots, \{n\}$

$\{1, 2\}, \{1, 3\}, \{1, 4\}, \dots, \{n-1, n\}$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

\emptyset

$\{1\}, \{2\}, \{3\}, \dots, \{n\}$

$\{1, 2\}, \{1, 3\}, \{1, 4\}, \dots, \{n-1, n\}$

$\{1, 2, 3\}, \{1, 2, 4\}, \dots, \{n-2, n-1, n\}$

\dots

$\{1, 2, \dots, n\}$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

$$\emptyset \quad \binom{n}{0}$$

$$\{1\}, \{2\}, \{3\}, \dots, \{n\}$$

$$\{1, 2\}, \{1, 3\}, \{1, 4\}, \dots, \{n-1, n\}$$

$$\{1, 2, 3\}, \{1, 2, 4\}, \dots, \{n-2, n-1, n\}$$

...

$$\{1, 2, \dots, n\}$$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

$$\emptyset \quad \binom{n}{0}$$

$$\{1\}, \{2\}, \{3\}, \dots, \{n\} \quad \binom{n}{1}$$

$$\{1, 2\}, \{1, 3\}, \{1, 4\}, \dots, \{n-1, n\}$$

$$\{1, 2, 3\}, \{1, 2, 4\}, \dots, \{n-2, n-1, n\}$$

...

$$\{1, 2, \dots, n\}$$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

$$\emptyset \quad \binom{n}{0}$$

$$\{1\}, \{2\}, \{3\}, \dots, \{n\} \quad \binom{n}{1}$$

$$\{1, 2\}, \{1, 3\}, \{1, 4\}, \dots, \{n-1, n\} \quad \binom{n}{2}$$

$$\{1, 2, 3\}, \{1, 2, 4\}, \dots, \{n-2, n-1, n\}$$

...

$$\{1, 2, \dots, n\}$$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

$$\emptyset \quad \binom{n}{0}$$

$$\{1\}, \{2\}, \{3\}, \dots, \{n\} \quad \binom{n}{1}$$

$$\{1, 2\}, \{1, 3\}, \{1, 4\}, \dots, \{n-1, n\} \quad \binom{n}{2}$$

$$\{1, 2, 3\}, \{1, 2, 4\}, \dots, \{n-2, n-1, n\} \quad \binom{n}{3}$$

...

$$\{1, 2, \dots, n\} \quad \binom{n}{n}$$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

$$\emptyset \quad \binom{n}{0}$$

$$\{1\}, \{2\}, \{3\}, \dots, \{n\} \quad \binom{n}{1}$$

$$\{1, 2\}, \{1, 3\}, \{1, 4\}, \dots, \{n-1, n\} \quad \binom{n}{2}$$

$$\{1, 2, 3\}, \{1, 2, 4\}, \dots, \{n-2, n-1, n\} \quad \binom{n}{3}$$

...

$$\{1, 2, \dots, n\} \quad \binom{n}{n}$$

$|P(A)|$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

$$\emptyset \quad \binom{n}{0}$$

$$\{1\}, \{2\}, \{3\}, \dots, \{n\} \quad \binom{n}{1}$$

$$\{1, 2\}, \{1, 3\}, \{1, 4\}, \dots, \{n-1, n\} \quad \binom{n}{2}$$

$$\{1, 2, 3\}, \{1, 2, 4\}, \dots, \{n-2, n-1, n\} \quad \binom{n}{3}$$

...

$$\{1, 2, \dots, n\} \quad \binom{n}{n}$$

$$|P(A)| = \binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n}$$

Binomial Theorem

- Prove that $\sum_{i=0}^n \binom{n}{i} = 2^n$

Suppose there is a set $A = \{1, 2, \dots, n\}$. Let's write the elements of the power set $P(A)$:

$$\emptyset \quad \binom{n}{0}$$

$$\{1\}, \{2\}, \{3\}, \dots, \{n\} \quad \binom{n}{1}$$

$$\{1, 2\}, \{1, 3\}, \{1, 4\}, \dots, \{n-1, n\} \quad \binom{n}{2}$$

$$\{1, 2, 3\}, \{1, 2, 4\}, \dots, \{n-2, n-1, n\} \quad \binom{n}{3}$$

...

$$\{1, 2, \dots, n\} \quad \binom{n}{n}$$

$$|P(A)| = \binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n} = 2^n$$

Binomial Theorem

- Prove that $\sum_{k=0}^n (-1)^k \binom{n}{k} = 0$ where n is odd integer.

Binomial Theorem

- Prove that $\sum_{k=0}^n (-1)^k \binom{n}{k} = 0$ where n is odd integer.

$$= (-1)^0 \binom{n}{0} + (-1)^1 \binom{n}{1} + (-1)^2 \binom{n}{2} + (-1)^3 \binom{n}{3} + \dots + (-1)^n \binom{n}{n}$$

Binomial Theorem

- Prove that $\sum_{k=0}^n (-1)^k \binom{n}{k} = 0$ where n is odd integer.

$$= (-1)^0 \binom{n}{0} + (-1)^1 \binom{n}{1} + (-1)^2 \binom{n}{2} + (-1)^3 \binom{n}{3} + \dots + (-1)^n \binom{n}{n}$$

$$= \binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \binom{n}{3} + \dots + \binom{n}{n-1} - \binom{n}{n}$$

Binomial Theorem

- Prove that $\sum_{k=0}^n (-1)^k \binom{n}{k} = 0$ where n is odd integer.

$$= (-1)^0 \binom{n}{0} + (-1)^1 \binom{n}{1} + (-1)^2 \binom{n}{2} + (-1)^3 \binom{n}{3} + \dots + (-1)^n \binom{n}{n}$$

$$= \binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \binom{n}{3} + \dots + \binom{n}{n-1} - \binom{n}{n}$$

$$\bullet \quad \binom{n}{k} = \frac{n!}{(n-k)!k!} \quad \text{and} \quad \binom{n}{n-k} = \frac{n!}{k!(n-k)!}$$

Binomial Theorem

- Prove that $\sum_{k=0}^n (-1)^k \binom{n}{k} = 0$ where n is odd integer.

$$= (-1)^0 \binom{n}{0} + (-1)^1 \binom{n}{1} + (-1)^2 \binom{n}{2} + (-1)^3 \binom{n}{3} + \dots + (-1)^n \binom{n}{n}$$

$$= \binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \binom{n}{3} + \dots + \binom{n}{n-1} - \binom{n}{n}$$

- $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ and $\binom{n}{n-k} = \frac{n!}{k!(n-k)!}$

$$\text{Thus, } \binom{n}{k} = \binom{n}{n-k}$$

Binomial Theorem

- Prove that $\sum_{k=0}^n (-1)^k \binom{n}{k} = 0$ where n is odd integer.

$$= (-1)^0 \binom{n}{0} + (-1)^1 \binom{n}{1} + (-1)^2 \binom{n}{2} + (-1)^3 \binom{n}{3} + \dots + (-1)^n \binom{n}{n}$$

$$= \binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \binom{n}{3} + \dots + \binom{n}{n-1} - \binom{n}{n}$$

$$= \binom{n}{0} - \binom{n}{n} + \binom{n}{2} - \binom{n}{n-2} + \dots + \binom{n}{n-1} - \binom{n}{1}$$

- $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ and $\binom{n}{n-k} = \frac{n!}{k!(n-k)!}$

$$\text{Thus, } \binom{n}{k} = \binom{n}{n-k}$$

Binomial Theorem

- Prove that $\sum_{k=0}^n (-1)^k \binom{n}{k} = 0$ where n is odd integer.

$$= (-1)^0 \binom{n}{0} + (-1)^1 \binom{n}{1} + (-1)^2 \binom{n}{2} + (-1)^3 \binom{n}{3} + \dots + (-1)^n \binom{n}{n}$$

$$= \binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \binom{n}{3} + \dots + \binom{n}{n-1} - \binom{n}{n}$$

$$= \binom{n}{0} - \binom{n}{n} + \binom{n}{2} - \binom{n}{n-2} + \dots + \binom{n}{n-1} - \binom{n}{1} = 0$$

- $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ and $\binom{n}{n-k} = \frac{n!}{k!(n-k)!}$

$$\text{Thus, } \binom{n}{k} = \binom{n}{n-k}$$

Binomial Theorem

- $$\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

There is a set A such that $|A|=n+1$
Assume one of them is a

$$A = \{ \dots, a, \dots \}$$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$



$\{\dots\}$

create subsets of
A with k elements

There is a set A such that $|A|=n+1$
Assume one of them is a

$A = \{\dots, a, \dots\}$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

$\{\dots\dots\dots\}$

create subsets of
A with k elements

$\{a, \dots\dots\dots\}$

k-1 elements

$\{\dots\dots\dots\}$

k elements

There is a set A such that $|A|=n+1$
Assume one of them is a

$A = \{\dots, a, \dots\}$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

There is a set A such that $|A|=n+1$
Assume one of them is a

$\{\dots\dots\dots\}$

create subsets of
 A with k elements

$\{a, \dots\dots\dots\}$

$k-1$ elements

$\{\dots\dots\dots\}$

k elements

$A = \{\dots, a, \dots\}$

$$\begin{array}{c} \binom{0}{0} \\ \swarrow \quad \searrow \\ \binom{1}{0} \quad \binom{1}{1} \end{array}$$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

There is a set A such that $|A|=n+1$
Assume one of them is a

$\{\dots\dots\dots\}$

create subsets of
 A with k elements


$\{a, \dots\dots\dots\}$

$k-1$ elements

$\{\dots\dots\dots\}$

k elements

$A = \{\dots, a, \dots\}$



Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

There is a set A such that $|A|=n+1$
Assume one of them is a

$\{\dots\dots\}$

create subsets of
 A with k elements


$\{a, \dots\dots\}$

$k-1$ elements

$\{\dots\dots\}$

k elements

$A = \{\dots, a, \dots\}$



Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

There is a set A such that $|A|=n+1$
Assume one of them is a

$\{\dots\dots\dots\}$

create subsets of
 A with k elements


$\{a, \dots\dots\dots\}$

$k-1$ elements

$\{\dots\dots\dots\}$

k elements

$A = \{\dots, a, \dots\}$



Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

There is a set A such that $|A|=n+1$
Assume one of them is a

$\{\dots\dots\dots\}$

create subsets of
 A with k elements


$\{a, \dots\dots\dots\}$

$k-1$ elements

$\{\dots\dots\dots\}$

k elements

$A = \{\dots, a, \dots\}$



Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

There is a set A such that $|A|=n+1$
Assume one of them is a

$\{\dots\dots\dots\}$

create subsets of
 A with k elements



$\{a, \dots\dots\dots\}$

$k-1$ elements

$\{\dots\dots\dots\}$

k elements

$A = \{\dots, a, \dots\}$



Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

Prove that $\sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

Prove that $\sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$

$$= \binom{1}{1} + \binom{2}{1} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

Prove that $\sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$

$$= \binom{2}{2} + \binom{2}{1} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$\binom{1}{1} = \binom{2}{2}$$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

Prove that $\sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$

$$= \binom{2}{2} + \binom{2}{1} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

Prove that $\sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$

$$= \binom{2}{2} + \binom{2}{1} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$= \binom{3}{2} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

Prove that $\sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$

$$= \binom{2}{2} + \binom{2}{1} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$= \binom{3}{2} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$= \binom{4}{2} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

Prove that $\sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$

$$= \binom{2}{2} + \binom{2}{1} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$= \binom{3}{2} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$= \binom{4}{2} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$= \binom{n}{2} + \binom{n}{1}$$

Binomial Theorem

- $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

Prove that $\sum_{k=1}^n \binom{k}{1} = \binom{n+1}{2}$

$$= \binom{2}{2} + \binom{2}{1} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$= \binom{3}{2} + \binom{3}{1} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$= \binom{4}{2} + \dots + \binom{n-1}{1} + \binom{n}{1}$$

$$= \binom{n}{2} + \binom{n}{1} = \binom{n+1}{2}$$