

# Mathematical Induction

Murat Osmanoglu

# Definition

- To prove  $P(n)$  is true for all positive integers  $n$ ,

# Definition

- To prove  $P(n)$  is true for all positive integers  $n$ ,
  - verify that  $P(1)$  is true (Basic Step)

# Definition

- To prove  $P(n)$  is true for all positive integers  $n$ ,
  - verify that  $P(1)$  is true (Basic Step)
  - prove that the implication  $P(k) \rightarrow P(k + 1)$  is true for all  $k \in \mathbb{Z}^+$  (Inductive Step)

# Definition


- To prove  $P(n)$  is true for all positive integers  $n$ ,
  - verify that  $P(1)$  is true (Basic Step)
  - prove that the implication  $P(k) \rightarrow P(k + 1)$  is true for all  $k \in \mathbb{Z}^+$  (Inductive Step)

$$[P(1) \wedge \forall k P(k) \rightarrow P(k + 1)] \rightarrow \forall n P(n)$$

# Definition

- To prove  $P(n)$  is true for all positive integers  $n$ ,
  - verify that  $P(1)$  is true (Basic Step)
  - prove that the implication  $P(k) \rightarrow P(k + 1)$  is true for all  $k \in \mathbb{Z}^+$  (Inductive Step)

$$[P(1) \wedge \forall k P(k) \rightarrow P(k + 1)] \rightarrow \forall n P(n)$$



# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3



# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^3 - k$  is divisible by 3

# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^3 - k$  is divisible by 3

$$[k^3 - k = 3a, \exists a \in \mathbb{Z}]$$

# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^3 - k$  is divisible by 3

$$[k^3 - k = 3a, \exists a \in \mathbb{Z}] \rightarrow (k + 1)^3 - (k + 1)$$

# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^3 - k$  is divisible by 3

$$[k^3 - k = 3a, \exists a \in \mathbb{Z}] \rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1$$

# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^3 - k$  is divisible by 3

$$\begin{aligned} [k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \end{aligned}$$

# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^3 - k$  is divisible by 3

$$\begin{aligned}[k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3k^2 + 3k\end{aligned}$$

# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^3 - k$  is divisible by 3

$$\begin{aligned} [k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3k^2 + 3k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3(k^2 + k) \end{aligned}$$



# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^3 - k$  is divisible by 3

$$\begin{aligned} [k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3k^2 + 3k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3(k^2 + k) \\ &\rightarrow (k + 1)^3 - (k + 1) = 3a + 3b, \exists a, b \in \mathbb{Z} \end{aligned}$$

# Proofs

- Prove that  $\forall x \in \mathbb{Z}^+, x^3 - x$  is divisible by 3

Basic Step  $P(1) : 1^3 - 1 = 0$  is divisible by 3

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^3 - k$  is divisible by 3

$$\begin{aligned} [k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3k^2 + 3k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3(k^2 + k) \\ &\rightarrow (k + 1)^3 - (k + 1) = 3a + 3b, \exists a, b \in \mathbb{Z} \\ &\rightarrow (k + 1)^3 - (k + 1) \text{ is divisible by 3} \end{aligned}$$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$  is divisible by 57

# Proofs

- Prove that  $\forall n \in N, 7^{n+2} + 8^{2n+1}$  is divisible by 57

Basic Step  $P(0) : 7^2 + 8 = 57$  is divisible by 57

# Proofs

- Prove that  $\forall n \in N, 7^{n+2} + 8^{2n+1}$  is divisible by 57

Basic Step  $P(0) : 7^2 + 8 = 57$  is divisible by 57

Inductive Step  $P(k) \rightarrow P(k + 1)$

# Proofs

- Prove that  $\forall n \in N, 7^{n+2} + 8^{2n+1}$  is divisible by 57

Basic Step  $P(0) : 7^2 + 8 = 57$  is divisible by 57

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $7^{k+2} + 8^{2k+1}$  is divisible by 57

# Proofs

- Prove that  $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$  is divisible by 57

Basic Step  $P(0) : 7^2 + 8 = 57$  is divisible by 57

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $7^{k+2} + 8^{2k+1}$  is divisible by 57

$$[7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}]$$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$  is divisible by 57

Basic Step  $P(0) : 7^2 + 8 = 57$  is divisible by 57

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $7^{k+2} + 8^{2k+1}$  is divisible by 57

$$[7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] \rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1}$$



# Proofs

- Prove that  $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$  is divisible by 57

Basic Step  $P(0) : 7^2 + 8 = 57$  is divisible by 57

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $7^{k+2} + 8^{2k+1}$  is divisible by 57

$$\begin{aligned} [7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 7 \cdot 8^{2k+1} + 57 \cdot 8^{2k+1} \end{aligned}$$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$  is divisible by 57

Basic Step  $P(0) : 7^2 + 8 = 57$  is divisible by 57

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $7^{k+2} + 8^{2k+1}$  is divisible by 57

$$\begin{aligned} [7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 7 \cdot 8^{2k+1} + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7(7^{k+2} + 8^{2k+1}) + 57 \cdot 8^{2k+1} \end{aligned}$$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$  is divisible by 57

Basic Step  $P(0) : 7^2 + 8 = 57$  is divisible by 57

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $7^{k+2} + 8^{2k+1}$  is divisible by 57

$$\begin{aligned} [7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 7 \cdot 8^{2k+1} + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7(7^{k+2} + 8^{2k+1}) + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 57a + 57b, \exists a, b \in \mathbb{Z} \end{aligned}$$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$  is divisible by 57

Basic Step  $P(0) : 7^2 + 8 = 57$  is divisible by 57

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $7^{k+2} + 8^{2k+1}$  is divisible by 57

$$\begin{aligned} [7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 7 \cdot 8^{2k+1} + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7(7^{k+2} + 8^{2k+1}) + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 57a + 57b, \exists a, b \in \mathbb{Z} \\ &\rightarrow 7^{k+3} + 8^{2k+3} \text{ is divisible by } 57 \end{aligned}$$

# Proofs

- Prove that if  $\forall n \in \mathbb{Z}^+$ , then  $1 + 2 + \dots + n = n \cdot (n + 1) / 2$

# Proofs

- Prove that if  $\forall n \in \mathbb{Z}^+$ , then  $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step  $P(1) : 1 = 1 \cdot 2/2$

# Proofs

- Prove that if  $\forall n \in \mathbb{Z}^+$ , then  $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step  $P(1) : 1 = 1 \cdot 2/2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

# Proofs

- Prove that if  $\forall n \in \mathbb{Z}^+$ , then  $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step  $P(1) : 1 = 1 \cdot 2/2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + \dots + k = k \cdot (k + 1)/2$



# Proofs

- Prove that if  $\forall n \in \mathbb{Z}^+$ , then  $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step  $P(1) : 1 = 1 \cdot 2/2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + \dots + k = k \cdot (k + 1)/2$

$[1 + 2 + \dots + k = k \cdot (k + 1)/2]$

# Proofs

- Prove that if  $\forall n \in \mathbb{Z}^+$ , then  $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step  $P(1) : 1 = 1 \cdot 2/2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + \dots + k = k \cdot (k + 1)/2$

$$[1 + 2 + \dots + k = k \cdot (k + 1)/2] \rightarrow \left[ 1 + 2 + \dots + (k + 1) = k \cdot \frac{k+1}{2} + k + 1 \right]$$

# Proofs

- Prove that if  $\forall n \in \mathbb{Z}^+$ , then  $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step  $P(1) : 1 = 1 \cdot 2/2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + \dots + k = k \cdot (k + 1)/2$

$$\begin{aligned} [1 + 2 + \dots + k = k \cdot (k + 1)/2] &\rightarrow [1 + 2 + \dots + (k + 1) = k \cdot \frac{k+1}{2} + k + 1] \\ &\rightarrow [1 + 2 + \dots + (k + 1) = \frac{k(k+1) + 2(k+1)}{2}] \end{aligned}$$

# Proofs

- Prove that if  $\forall n \in \mathbb{Z}^+$ , then  $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step  $P(1) : 1 = 1 \cdot 2/2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + \dots + k = k \cdot (k + 1)/2$

$$[1 + 2 + \dots + k = k \cdot (k + 1)/2] \rightarrow [1 + 2 + \dots + (k + 1) = k \cdot \frac{k+1}{2} + k + 1]$$

$$\rightarrow [1 + 2 + \dots + (k + 1) = \frac{k(k+1) + 2(k+1)}{2}]$$

$$\rightarrow [1 + 2 + \dots + (k + 1) = \frac{(k+1)(k+2)}{2}]$$

# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

- $1 = 1$        $1 + 3 = 4$        $1 + 3 + 5 = 9$        $1 + 3 + 5 + 7 = 16$

# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

- $$\begin{array}{cccc} 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ 1^2 & 2^2 & 3^2 & 4^2 \end{array}$$

# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

- $1 = 1$                        $1 + 3 = 4$                        $1 + 3 + 5 = 9$                        $1 + 3 + 5 + 7 = 16$   
 $1^2$                                    $2^2$                                    $3^2$                                    $4^2$   
 $1 + 3 + \dots + (2n - 1) = n^2$



# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

- $1 = 1$                    $1 + 3 = 4$                    $1 + 3 + 5 = 9$                    $1 + 3 + 5 + 7 = 16$   
 $1^2$                            $2^2$                            $3^2$                            $4^2$   
 $1 + 3 + \dots + (2n - 1) = n^2$

Basic Step  $P(1) : 1 = 1^2$

# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

- $1 = 1$                    $1 + 3 = 4$                    $1 + 3 + 5 = 9$                    $1 + 3 + 5 + 7 = 16$   
 $1^2$                            $2^2$                            $3^2$                            $4^2$   
 $1 + 3 + \dots + (2n - 1) = n^2$

Basic Step  $P(1) : 1 = 1^2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

- $1 = 1$                        $1 + 3 = 4$                        $1 + 3 + 5 = 9$                        $1 + 3 + 5 + 7 = 16$   
 $1^2$                                    $2^2$                                    $3^2$                                    $4^2$   
 $1 + 3 + \dots + (2n - 1) = n^2$

Basic Step  $P(1) : 1 = 1^2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 3 + \dots + (2k - 1) = k^2$

# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

- $$\begin{array}{ccccccc} 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ 1^2 & 2^2 & 3^2 & 4^2 \\ 1 + 3 + \dots + (2n - 1) = n^2 \end{array}$$

Basic Step  $P(1) : 1 = 1^2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + \dots + (2k - 1) = k^2$

$$[1 + 2 + \dots + (2k - 1) = k^2]$$

# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

- $$\begin{array}{ccccccc} 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ 1^2 & 2^2 & 3^2 & 4^2 \\ 1 + 3 + \dots + (2n - 1) = n^2 \end{array}$$

Basic Step  $P(1) : 1 = 1^2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + \dots + (2k - 1) = k^2$

$$[1 + 2 + \dots + (2k - 1) = k^2] \rightarrow [1 + 2 + \dots + (2k - 1) + (2k + 1) = k^2 + 2k + 1]$$

# Proofs

Conjecture a formula for the sum of the first  $n$  positive odd integers, then prove your conjecture using mathematical induction

- $$\begin{array}{ccccccc} 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ 1^2 & 2^2 & 3^2 & 4^2 \\ 1 + 3 + \dots + (2n - 1) = n^2 \end{array}$$

Basic Step  $P(1) : 1 = 1^2$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + \dots + (2k - 1) = k^2$

$$[1 + 2 + \dots + (2k - 1) = k^2] \rightarrow [1 + 2 + \dots + (2k - 1) + (2k + 1) = k^2 + 2k + 1]$$

$$\rightarrow [1 + 2 + \dots + (2k - 1) + (2k + 1) = (k + 1)^2]$$

# Proofs

- Prove that if  $n \in \mathbb{N}$ , then  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

# Proofs

- Prove that if  $n \in \mathbb{N}$ , then  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step  $P(1) : 1 = 2^{0+1} - 1$



# Proofs

- Prove that if  $n \in \mathbb{N}$ , then  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step  $P(1) : 1 = 2^{0+1} - 1$

Inductive Step  $P(k) \rightarrow P(k + 1)$

# Proofs

- Prove that if  $n \in \mathbb{N}$ , then  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step  $P(1) : 1 = 2^{0+1} - 1$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

# Proofs

- Prove that if  $n \in \mathbb{N}$ , then  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step  $P(1) : 1 = 2^{0+1} - 1$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

$$[1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1]$$

# Proofs

- Prove that if  $n \in \mathbb{N}$ , then  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step  $P(1) : 1 = 2^{0+1} - 1$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

$$[1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1] \rightarrow [1 + 2 + 2^2 + \dots + 2^k + 2^{k+1} = 2^{k+1} - 1 + 2^{k+1}]$$

# Proofs

- Prove that if  $n \in \mathbb{N}$ , then  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step  $P(1) : 1 = 2^{0+1} - 1$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

$$[1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1] \rightarrow [1 + 2 + 2^2 + \dots + 2^k + 2^{k+1} = 2^{k+1} - 1 + 2^{k+1}]$$

$$\rightarrow [1 + 2 + 2^2 + \dots + 2^{k+1} = 2 \cdot 2^{k+1} - 1]$$

# Proofs

- Prove that if  $n \in \mathbb{N}$ , then  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step  $P(1) : 1 = 2^{0+1} - 1$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

$$[1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1] \rightarrow [1 + 2 + 2^2 + \dots + 2^k + 2^{k+1} = 2^{k+1} - 1 + 2^{k+1}]$$

$$\rightarrow [1 + 2 + 2^2 + \dots + 2^{k+1} = 2 \cdot 2^{k+1} - 1]$$

$$\rightarrow [1 + 2 + 2^2 + \dots + 2^{k+1} = 2^{k+1+1} - 1]$$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1} - a}{r - 1}$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1} - a}{r-1}$

Basic Step  $P(0)$  :



# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1} - a}{r-1}$

Basic Step  $P(0) : a = \frac{ar^{0+1} - a}{r-1}$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1} - a}{r-1}$

Basic Step  $P(0) : a = \frac{ar^{0+1} - a}{r-1} = \frac{a(r-1)}{r-1} = a$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1} - a}{r-1}$

Basic Step  $P(0) : a = \frac{ar^{0+1} - a}{r-1} = \frac{a(r-1)}{r-1} = a$

Inductive Step  $P(k) \rightarrow P(k+1)$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1} - a}{r-1}$

Basic Step  $P(0) : a = \frac{ar^{0+1} - a}{r-1} = \frac{a(r-1)}{r-1} = a$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $a + ar + \dots + ar^k = \frac{ar^{k+1} - a}{r-1}$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1}-a}{r-1}$

Basic Step  $P(0) : a = \frac{ar^{0+1}-a}{r-1} = \frac{a(r-1)}{r-1} = a$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1}$

$$\left[ a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1} \right]$$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1}-a}{r-1}$

Basic Step  $P(0) : a = \frac{ar^{0+1}-a}{r-1} = \frac{a(r-1)}{r-1} = a$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1}$

$$\left[ a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1} \right] \rightarrow \left[ a + ar + \dots + ar^k + ar^{k+1} = \frac{ar^{k+1}-a}{r-1} + ar^{k+1} \right]$$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1}-a}{r-1}$

Basic Step  $P(0) : a = \frac{ar^{0+1}-a}{r-1} = \frac{a(r-1)}{r-1} = a$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1}$

$$\begin{aligned} \left[ a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1} \right] &\rightarrow \left[ a + ar + \dots + ar^k + ar^{k+1} = \frac{ar^{k+1}-a}{r-1} + ar^{k+1} \right] \\ &\rightarrow \left[ a + ar + \dots + ar^{k+1} = \frac{ar^{k+1}-a+ar^{k+1}(r-1)}{r-1} \right] \end{aligned}$$

# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1}-a}{r-1}$

Basic Step  $P(0) : a = \frac{ar^{0+1}-a}{r-1} = \frac{a(r-1)}{r-1} = a$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1}$

$$\begin{aligned} \left[ a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1} \right] &\rightarrow \left[ a + ar + \dots + ar^k + ar^{k+1} = \frac{ar^{k+1}-a}{r-1} + ar^{k+1} \right] \\ &\rightarrow \left[ a + ar + \dots + ar^{k+1} = \frac{ar^{k+1}-a+ar^{k+1}(r-1)}{r-1} \right] \\ &\rightarrow \left[ a + ar + \dots + ar^{k+1} = \frac{ar^{k+1}-a+ar^{k+2}-ar^{k+1}}{r-1} \right] \end{aligned}$$



# Proofs

- Prove that  $\forall n \in \mathbb{N}, \sum_{i=0}^n ar^i = a + ar + \dots + ar^n = \frac{ar^{n+1}-a}{r-1}$

Basic Step  $P(0) : a = \frac{ar^{0+1}-a}{r-1} = \frac{a(r-1)}{r-1} = a$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e  $a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1}$

$$\left[ a + ar + \dots + ar^k = \frac{ar^{k+1}-a}{r-1} \right] \rightarrow \left[ a + ar + \dots + ar^k + ar^{k+1} = \frac{ar^{k+1}-a}{r-1} + ar^{k+1} \right]$$

$$\rightarrow \left[ a + ar + \dots + ar^{k+1} = \frac{ar^{k+1}-a+ar^{k+1}(r-1)}{r-1} \right]$$

$$\rightarrow \left[ a + ar + \dots + ar^{k+1} = \frac{ar^{k+1}-a+ar^{k+2}-ar^{k+1}}{r-1} \right]$$

$$\rightarrow \left[ a + ar + \dots + ar^{k+1} = \frac{ar^{k+2}-a}{r-1} \right]$$

# Proofs

- Prove that for every integer  $n \geq 4$ ,  $2^n < n!$

# Proofs

- Prove that for every integer  $n \geq 4$ ,  $2^n < n!$

Basic Step  $P(4)$ :  $2^4 = 16 < 4! = 24$

# Proofs

- Prove that for every integer  $n \geq 4$ ,  $2^n < n!$

Basic Step  $P(4) : 2^4 = 16 < 4! = 24$

Inductive Step  $P(k) \rightarrow P(k + 1)$

# Proofs

- Prove that for every integer  $n \geq 4$ ,  $2^n < n!$

Basic Step  $P(4) : 2^4 = 16 < 4! = 24$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $2^k < k!$

# Proofs

- Prove that for every integer  $n \geq 4$ ,  $2^n < n!$

Basic Step  $P(4) : 2^4 = 16 < 4! = 24$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $2^k < k!$

$[2^k < k!]$

# Proofs

- Prove that for every integer  $n \geq 4$ ,  $2^n < n!$

Basic Step  $P(4)$  :  $2^4 = 16 < 4! = 24$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $2^k < k!$

$$[2^k < k!] \rightarrow [2^{k+1} = 2 \cdot 2^k < 2 \cdot k!]$$

# Proofs

- Prove that for every integer  $n \geq 4$ ,  $2^n < n!$

Basic Step  $P(4)$  :  $2^4 = 16 < 4! = 24$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $2^k < k!$

$$[2^k < k!] \rightarrow [2^{k+1} = 2 \cdot 2^k < 2 \cdot k!] \rightarrow [2^{k+1} < 2 \cdot k!]$$



# Proofs

- Prove that for every integer  $n \geq 4$ ,  $2^n < n!$

Basic Step  $P(4)$  :  $2^4 = 16 < 4! = 24$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $2^k < k!$

$$[2^k < k!] \rightarrow [2^{k+1} = 2 \cdot 2^k < 2 \cdot k!] \rightarrow [2^{k+1} < 2 \cdot k! < (k+1) \cdot k!]$$

# Proofs

- Prove that for every integer  $n \geq 4$ ,  $2^n < n!$

Basic Step  $P(4)$  :  $2^4 = 16 < 4! = 24$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $2^k < k!$

$$\begin{aligned} [2^k < k!] &\rightarrow [2^{k+1} = 2 \cdot 2^k < 2 \cdot k!] \rightarrow [2^{k+1} < 2 \cdot k! < (k+1) \cdot k!] \\ &\rightarrow [2^{k+1} < (k+1)!] \end{aligned}$$

# Proofs

- Prove that for every integer  $n \geq 3$ ,  $n^2 - 7n + 12$  is nonnegative integer

# Proofs

- Prove that for every integer  $n \geq 3$ ,  $n^2 - 7n + 12$  is nonnegative integer

Basic Step  $P(3)$  :  $3^2 - 7 \cdot 3 + 12 = 0$  is a nonnegative integer

# Proofs

- Prove that for every integer  $n \geq 3$ ,  $n^2 - 7n + 12$  is nonnegative integer

Basic Step  $P(3)$  :  $3^2 - 7 \cdot 3 + 12 = 0$  is a nonnegative integer

Inductive Step  $P(k) \rightarrow P(k + 1)$

# Proofs

- Prove that for every integer  $n \geq 3$ ,  $n^2 - 7n + 12$  is nonnegative integer

Basic Step  $P(3)$  :  $3^2 - 7 \cdot 3 + 12 = 0$  is a nonnegative integer

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^2 - 7k + 12$  is a nonnegative integer

# Proofs

- Prove that for every integer  $n \geq 3$ ,  $n^2 - 7n + 12$  is nonnegative integer

Basic Step  $P(3)$  :  $3^2 - 7 \cdot 3 + 12 = 0$  is a nonnegative integer

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^2 - 7k + 12$  is a nonnegative integer

$$(k + 1)^2 - 7(k + 1) + 12$$

# Proofs

- Prove that for every integer  $n \geq 3$ ,  $n^2 - 7n + 12$  is nonnegative integer

Basic Step  $P(3)$  :  $3^2 - 7 \cdot 3 + 12 = 0$  is a nonnegative integer

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^2 - 7k + 12$  is a nonnegative integer

$$(k + 1)^2 - 7(k + 1) + 12 = k^2 + 2k + 1 - 7k - 7 + 12$$



# Proofs

- Prove that for every integer  $n \geq 3$ ,  $n^2 - 7n + 12$  is nonnegative integer

Basic Step  $P(3)$  :  $3^2 - 7 \cdot 3 + 12 = 0$  is a nonnegative integer

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^2 - 7k + 12$  is a nonnegative integer

$$\begin{aligned}(k + 1)^2 - 7(k + 1) + 12 &= k^2 + 2k + 1 - 7k - 7 + 12 \\ &= k^2 - 7k + 12 + 2k - 6\end{aligned}$$

# Proofs


- Prove that for every integer  $n \geq 3$ ,  $n^2 - 7n + 12$  is nonnegative integer

Basic Step  $P(3)$  :  $3^2 - 7 \cdot 3 + 12 = 0$  is a nonnegative integer

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^2 - 7k + 12$  is a nonnegative integer

$$\begin{aligned}(k + 1)^2 - 7(k + 1) + 12 &= k^2 + 2k + 1 - 7k - 7 + 12 \\ &= \underline{k^2 - 7k + 12} + 2k - 6\end{aligned}$$



from the assumption,  
this part is nonnegative

# Proofs

- Prove that for every integer  $n \geq 3$ ,  $n^2 - 7n + 12$  is nonnegative integer

Basic Step  $P(3)$  :  $3^2 - 7 \cdot 3 + 12 = 0$  is a nonnegative integer

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e  $k^2 - 7k + 12$  is a nonnegative integer

$$(k + 1)^2 - 7(k + 1) + 12 = k^2 + 2k + 1 - 7k - 7 + 12$$

$$= \underbrace{k^2 - 7k + 12}_{\text{from the assumption, this part is nonnegative}} + \underbrace{2k - 6}_{\text{since } n \geq 3, 2k - 6 \geq 0}$$

from the assumption,  
this part is nonnegative

since  $n \geq 3$ ,  $2k - 6 \geq 0$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1]$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step  $P(k) \rightarrow P(k + 1)$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $H_1 + \dots + H_k = (k + 1)H_k - k$



# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $H_1 + \dots + H_k = (k + 1)H_k - k$

$$[H_1 + \dots + H_k = (k + 1)H_k - k]$$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $H_1 + \dots + H_k = (k + 1)H_k - k$

$$[H_1 + \dots + H_k = (k + 1)H_k - k] \rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}]$$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $H_1 + \dots + H_k = (k + 1)H_k - k$

$$\begin{aligned} [H_1 + \dots + H_k = (k + 1)H_k - k] &\rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}] \\ &\rightarrow \left[ H_1 + \dots + H_{k+1} = (k + 1)\left(H_k - \frac{1}{k+1} + \frac{1}{k+1}\right) - k + H_{k+1} \right] \end{aligned}$$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $H_1 + \dots + H_k = (k + 1)H_k - k$

$$\begin{aligned} [H_1 + \dots + H_k = (k + 1)H_k - k] &\rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}] \\ &\rightarrow \left[ H_1 + \dots + H_{k+1} = (k + 1)\left(H_k - \frac{1}{k+1} + \frac{1}{k+1}\right) - k + H_{k+1} \right] \\ &\rightarrow \left[ H_1 + \dots + H_{k+1} = (k + 1)\left(H_{k+1} - \frac{1}{k+1}\right) - k + H_{k+1} \right] \end{aligned}$$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $H_1 + \dots + H_k = (k + 1)H_k - k$

$$[H_1 + \dots + H_k = (k + 1)H_k - k] \rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}]$$

$$\rightarrow \left[ H_1 + \dots + H_{k+1} = (k + 1)\left(H_k - \frac{1}{k+1} + \frac{1}{k+1}\right) - k + H_{k+1} \right]$$

$$\rightarrow \left[ H_1 + \dots + H_{k+1} = (k + 1)\left(H_{k+1} - \frac{1}{k+1}\right) - k + H_{k+1} \right]$$

$$\rightarrow [H_1 + \dots + H_{k+1} = (k + 1)H_{k+1} - 1 - k + H_{k+1}]$$

# Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that  $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step  $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $H_1 + \dots + H_k = (k + 1)H_k - k$

$$\begin{aligned} [H_1 + \dots + H_k = (k + 1)H_k - k] &\rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}] \\ &\rightarrow \left[ H_1 + \dots + H_{k+1} = (k + 1)\left(H_k - \frac{1}{k+1} + \frac{1}{k+1}\right) - k + H_{k+1} \right] \\ &\rightarrow \left[ H_1 + \dots + H_{k+1} = (k + 1)\left(H_{k+1} - \frac{1}{k+1}\right) - k + H_{k+1} \right] \\ &\rightarrow [H_1 + \dots + H_{k+1} = (k + 1)H_{k+1} - 1 - k + H_{k+1}] \\ &\rightarrow [H_1 + \dots + H_{k+1} = (k + 2)H_{k+1} - (k + 1)] \end{aligned}$$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$



# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in \mathbb{N}$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in \mathbb{N}$



if  $b > 0$ ,

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in N$



if  $b > 0$ ,  $k + 1 = a.3 + b.8 + 1$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in \mathbb{N}$



if  $b > 0$ ,  $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in \mathbb{N}$



if  $b > 0$ ,  $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$



# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in \mathbb{N}$



if  $b > 0$ ,  $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

if  $b = 0$ ,  $k + 1 = a.3 + 1$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in N$



if  $b > 0$ ,  $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

if  $b = 0$ ,  $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in N$



if  $b > 0$ ,  $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

if  $b = 0$ ,  $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

$$k + 1 = (a - 5).3 + 2.8$$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in \mathbb{N}$



if  $b > 0$ ,  $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

if  $b = 0$ ,  $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

$$k + 1 = (a - 5).3 + 2.8$$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in N$



if  $b > 0$ ,  $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

$P(k - 8)$

if  $b = 0$ ,  $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

$$k + 1 = (a - 5).3 + 2.8$$

$P(k - 15)$

# Proofs

- For every integer  $n \geq 14$ ,  $n$  can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step  $P(4) : 14 = 2.3 + 1.8$

Inductive Step  $P(k) \rightarrow P(k + 1)$

assume that  $P(k)$  is true, i.e.  $k = a.3 + b.8, \exists a, b \in \mathbb{N}$

if  $b > 0$ ,  $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

$P(k - 8)$

if  $b = 0$ ,  $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

$$k + 1 = (a - 5).3 + 2.8$$

$P(k - 15)$

$$[P(k - 8) \wedge P(k - 15)] \rightarrow P(k + 1)$$

# Strong Induction

- To prove  $P(n)$  is true for all positive integers  $n$ ,
  - verify that  $P(1)$  is true (Basic Step)

# Strong Induction

- To prove  $P(n)$  is true for all positive integers  $n$ ,
  - verify that  $P(1)$  is true (Basic Step)
  - prove that the implication

$$[P(1) \wedge P(2) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$$

for all  $k \in \mathbb{Z}^+$  (Inductive Step)



# Strong Induction

- Prove that for every integer  $n \geq 2$ ,  $n$  can be written as the product of primes

# Strong Induction

- Prove that for every integer  $n \geq 2$ ,  $n$  can be written as the product of primes

Basic Step  $P(2)$  is true, i.e. 2 can be written as the product of primes

# Strong Induction

- Prove that for every integer  $n \geq 2$ ,  $n$  can be written as the product of primes

Basic Step  $P(2)$  is true, i.e. 2 can be written as the product of primes

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

# Strong Induction

- Prove that for every integer  $n \geq 2$ ,  $n$  can be written as the product of primes

Basic Step  $P(2)$  is true, i.e. 2 can be written as the product of primes

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $i$  can be written as the product of primes, then

# Strong Induction

- Prove that for every integer  $n \geq 2$ ,  $n$  can be written as the product of primes

Basic Step  $P(2)$  is true, i.e. 2 can be written as the product of primes

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $i$  can be written as the product of primes, then

if  $(k + 1)$  is prime, then  $P(k + 1)$  is true

# Strong Induction

- Prove that for every integer  $n \geq 2$ ,  $n$  can be written as the product of primes

Basic Step  $P(2)$  is true, i.e. 2 can be written as the product of primes

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $i$  can be written as the product of primes, then

if  $(k + 1)$  is prime, then  $P(k + 1)$  is true

if  $(k + 1)$  is composite, then  $k + 1 = a.b$ , where  $2 \leq a \leq b < k + 1$ .

# Strong Induction

- Prove that for every integer  $n \geq 2$ ,  $n$  can be written as the product of primes

Basic Step  $P(2)$  is true, i.e. 2 can be written as the product of primes

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $i$  can be written as the product of primes, then

if  $(k + 1)$  is prime, then  $P(k + 1)$  is true

if  $(k + 1)$  is composite, then  $k + 1 = a.b$ , where  $2 \leq a \leq b < k + 1$ .  
Since  $a, b < k + 1$ ,  $P(a)$  and  $P(b)$  are true from the assumption,

# Strong Induction

- Prove that for every integer  $n \geq 2$ ,  $n$  can be written as the product of primes

Basic Step  $P(2)$  is true, i.e. 2 can be written as the product of primes

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $i$  can be written as the product of primes, then

if  $(k + 1)$  is prime, then  $P(k + 1)$  is true

if  $(k + 1)$  is composite, then  $k + 1 = a.b$ , where  $2 \leq a \leq b < k + 1$ .

Since  $a, b < k + 1$ ,  $P(a)$  and  $P(b)$  are true from the assumption, i.e.  $a$  and  $b$  can be written as the product of primes.



# Strong Induction

- Prove that for every integer  $n \geq 2$ ,  $n$  can be written as the product of primes

Basic Step  $P(2)$  is true, i.e. 2 can be written as the product of primes

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $i$  can be written as the product of primes, then

if  $(k + 1)$  is prime, then  $P(k + 1)$  is true

if  $(k + 1)$  is composite, then  $k + 1 = a.b$ , where  $2 \leq a \leq b < k + 1$ .

Since  $a, b < k + 1$ ,  $P(a)$  and  $P(b)$  are true from the assumption, i.e.  $a$  and  $b$  can be written as the product of primes.


Thus,  $k + 1 = a.b$  can also be written as the product of primes.

# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?



# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?





# Strong Induction

- Consider a puzzle. How do we assemble a puzzle?



# Strong Induction



- Consider a puzzle. How do we assemble a puzzle ?



- Show that no matter which move we make,  $n-1$  moves required to assemble a puzzle with  $n$  pieces.

# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?





- Show that no matter which move we make,  $n-1$  moves required to assemble a puzzle with  $n$  pieces.

Basic Step  $P(1)$  is true, i.e. no move required for just 1 piece

# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?





- Show that no matter which move we make,  $n-1$  moves required to assemble a puzzle with  $n$  pieces.

Basic Step  $P(1)$  is true, i.e. no move required for just 1 piece

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?



- Show that no matter which move we make,  $n-1$  moves required to assemble a puzzle with  $n$  pieces.

Basic Step  $P(1)$  is true, i.e. no move required for just 1 piece



Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e a puzzle with  $i$  pieces can be assembled with  $i-1$  moves



# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?



- Show that no matter which move we make,  $n-1$  moves required to assemble a puzzle with  $n$  pieces.

Basic Step  $P(1)$  is true, i.e. no move required for just 1 piece



Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e a puzzle with  $i$  pieces can be assembled with  $i-1$  moves

$k+1$  pieces

# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?




- Show that no matter which move we make,  $n-1$  moves required to assemble a puzzle with  $n$  pieces.

Basic Step  $P(1)$  is true, i.e. no move required for just 1 piece

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e a puzzle with  $i$  pieces can be assembled with  $i-1$  moves

$k+1$  pieces



# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?




- Show that no matter which move we make,  $n-1$  moves required to assemble a puzzle with  $n$  pieces.

Basic Step  $P(1)$  is true, i.e. no move required for just 1 piece

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e a puzzle with  $i$  pieces can be assembled with  $i-1$  moves



# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?




- Show that no matter which move we make,  $n-1$  moves required to assemble a puzzle with  $n$  pieces.

Basic Step  $P(1)$  is true, i.e. no move required for just 1 piece



Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e a puzzle with  $i$  pieces can be assembled with  $i-1$  moves



# Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?




- Show that no matter which move we make,  $n-1$  moves required to assemble a puzzle with  $n$  pieces.

Basic Step  $P(1)$  is true, i.e. no move required for just 1 piece

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e a puzzle with  $i$  pieces can be assembled with  $i-1$  moves



# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$



# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $F(i) > \alpha^{i-2}$

# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $F(i) > \alpha^{i-2}$

for  $P(k+1) : F(k+1) = F(k) + F(k-1)$

# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $F(i) > \alpha^{i-2}$

for  $P(k+1) : F(k+1) = F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3}$

# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $F(i) > \alpha^{i-2}$

for  $P(k+1) : F(k+1) = F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3}$   
 $= \alpha \cdot \alpha^{k-3} + \alpha^{k-3}$

# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $F(i) > \alpha^{i-2}$

for  $P(k+1) : F(k+1) = F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3}$   
 $= \alpha \cdot \alpha^{k-3} + \alpha^{k-3}$   
 $= (\alpha + 1) \cdot \alpha^{k-3}$

# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $F(i) > \alpha^{i-2}$

$$\begin{aligned} \text{for } P(k+1) : F(k+1) &= F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3} \\ &= \alpha \cdot \alpha^{k-3} + \alpha^{k-3} \\ &= (\alpha + 1) \cdot \alpha^{k-3} \end{aligned}$$

$\alpha = \frac{1+\sqrt{5}}{2}$  is a solution of the equation  $\alpha^2 - \alpha - 1 = 0$ . Thus,  $\alpha^2 = \alpha + 1$

# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $F(i) > \alpha^{i-2}$

$$\begin{aligned} \text{for } P(k+1) : F(k+1) &= F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3} \\ &= \alpha \cdot \alpha^{k-3} + \alpha^{k-3} \\ &= (\alpha + 1) \cdot \alpha^{k-3} = \alpha^2 \cdot \alpha^{k-3} \end{aligned}$$

$\alpha = \frac{1+\sqrt{5}}{2}$  is a solution of the equation  $\alpha^2 - \alpha - 1 = 0$ . Thus,  $\alpha^2 = \alpha + 1$



# Strong Induction

- Prove that for every integer  $n \geq 3$ ,  $F(n) > \alpha^{n-2}$  where  $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence :  $F(1) = 1, F(2) = 1$ , and  $F(n) = F(n-1) + F(n-2)$

Basic Step  $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step  $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that  $P(i)$  is true for all  $i$  such that  $2 \leq i \leq k$ , i.e  $F(i) > \alpha^{i-2}$

$$\begin{aligned} \text{for } P(k+1) : F(k+1) &= F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3} \\ &= \alpha \cdot \alpha^{k-3} + \alpha^{k-3} \\ &= (\alpha + 1) \cdot \alpha^{k-3} = \alpha^2 \cdot \alpha^{k-3} \end{aligned}$$

$$F(k+1) > \alpha^{k-1}$$

$\alpha = \frac{1+\sqrt{5}}{2}$  is a solution of the equation  $\alpha^2 - \alpha - 1 = 0$ . Thus,  $\alpha^2 = \alpha + 1$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1, F(1)^2 + F(2)^2 = 2, F(1)^2 + F(2)^2 + F(3)^2 = 6,$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1, F(1)^2 + F(2)^2 = 2, F(1)^2 + F(2)^2 + F(3)^2 = 6,$   
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15,$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1, F(1)^2 + F(2)^2 = 2, F(1)^2 + F(2)^2 + F(3)^2 = 6,$   
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15,$   
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40,$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$ ,  $F(1)^2 + F(2)^2 = 2$ ,  $F(1)^2 + F(2)^2 + F(3)^2 = 6$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$ ,  
1.1                      1.2                      2.3                      3.5                      5.8

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1, F(1)^2 + F(2)^2 = 2, F(1)^2 + F(2)^2 + F(3)^2 = 6,$   
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15,$   
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40,$   
1.1                      1.2                      2.3                      3.5                      5.8
- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1), \text{ where } n \geq 2$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$ ,  $F(1)^2 + F(2)^2 = 2$ ,  $F(1)^2 + F(2)^2 + F(3)^2 = 6$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$ ,  
1.1                      1.2                      2.3                      3.5                      5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$ , where  $n \geq 2$

Basic Step  $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$



# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$ ,  $F(1)^2 + F(2)^2 = 2$ ,  $F(1)^2 + F(2)^2 + F(3)^2 = 6$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$ ,  
1.1                      1.2                      2.3                      3.5                      5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$ , where  $n \geq 2$

Basic Step  $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step  $P(k) \rightarrow P(k+1)$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$ ,  $F(1)^2 + F(2)^2 = 2$ ,  $F(1)^2 + F(2)^2 + F(3)^2 = 6$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$ ,  
1.1                      1.2                      2.3                      3.5                      5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$ , where  $n \geq 2$

Basic Step  $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e.  $\sum_{j=1}^k F(j)^2 = F(k) \cdot F(k+1)$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$ ,  $F(1)^2 + F(2)^2 = 2$ ,  $F(1)^2 + F(2)^2 + F(3)^2 = 6$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$ ,  
1.1                      1.2                      2.3                      3.5                      5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$ , where  $n \geq 2$

Basic Step  $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e.  $\sum_{j=1}^k F(j)^2 = F(k) \cdot F(k+1)$

for  $P(k+1) : F(1)^2 + \dots + F(k)^2 + F(k+1)^2$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$ ,  $F(1)^2 + F(2)^2 = 2$ ,  $F(1)^2 + F(2)^2 + F(3)^2 = 6$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$ ,  
1.1                      1.2                      2.3                      3.5                      5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$ , where  $n \geq 2$

Basic Step  $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e.  $\sum_{j=1}^k F(j)^2 = F(k) \cdot F(k+1)$

for  $P(k+1) : F(1)^2 + \dots + F(k)^2 + F(k+1)^2 = F(k) \cdot F(k+1) + F(k+1)^2$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$ ,  $F(1)^2 + F(2)^2 = 2$ ,  $F(1)^2 + F(2)^2 + F(3)^2 = 6$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$ ,  
1.1                      1.2                      2.3                      3.5                      5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$ , where  $n \geq 2$

Basic Step  $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e.  $\sum_{j=1}^k F(j)^2 = F(k) \cdot F(k+1)$

for  $P(k+1) : F(1)^2 + \dots + F(k)^2 + F(k+1)^2 = F(k) \cdot F(k+1) + F(k+1)^2$   
 $= F(k+1)(F(k) + F(k+1))$

# Strong Induction

Conjecture a formula for the sum of the squares of the first  $n$  terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$ ,  $F(1)^2 + F(2)^2 = 2$ ,  $F(1)^2 + F(2)^2 + F(3)^2 = 6$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$ ,  
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$ ,  
1.1                      1.2                      2.3                      3.5                      5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$ , where  $n \geq 2$

Basic Step  $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step  $P(k) \rightarrow P(k+1)$

assume that  $P(k)$  is true, i.e.  $\sum_{j=1}^k F(j)^2 = F(k) \cdot F(k+1)$

$$\begin{aligned} \text{for } P(k+1) : F(1)^2 + \dots + F(k)^2 + F(k+1)^2 &= F(k) \cdot F(k+1) + F(k+1)^2 \\ &= F(k+1)(F(k) + F(k+1)) \\ &= F(k+1)F(k+2) \end{aligned}$$