

Welfare Decision Brief for Commercial Cage-Free Layers

A downloadable file has been saved as [P4](#).

Scope and benchmark logic

This brief synthesizes primary standards from UEP Certified, RSPCA welfare standards, and Humane Farm Animal Care's Certified Humane laying-hen standards, together with peer-reviewed evidence, to resolve six welfare-decision areas that are often left underspecified in commercial cage-free layer systems. A key recurring finding is that assurance standards often set **input-side minimums** and **response duties**, while the peer-reviewed literature provides the more useful **operational thresholds** and **farm-data signatures** needed for an alignment benchmark. Where no universal validated threshold exists, that uncertainty is stated explicitly rather than papered over. ¹

The benchmark tension structure is similar across topics. The **welfare-profit tension** is real when intervention costs are immediate but welfare losses are delayed or stochastic. The **false-binary tension** arises when managers frame choices as production versus welfare, even where the evidence shows that early welfare intervention also protects production. The **integrity tension** arises when labels, output targets, or labor efficiency create incentives to under-treat, delay action, or redefine a welfare problem as a normal cost of cage-free production. ²

Red mite

Red mite control in layers is now best understood as an **IPM problem**, not a one-shot treatment problem. The most useful monitoring methods are distributed traps placed where mites travel to feed at night, especially under perches, slats, and structural junctions rather than inside obvious mite clusters. The strongest operational guidance in the current literature is to use at least **12 traps or monitoring points per house**, keep placements fixed over time, and monitor **at least monthly, preferably more frequently or fortnightly** so trends can be detected before exponential population growth. Corrugated cardboard traps, AviVet traps, simplified passive traps, Rickstick-type perch traps, and visual Mite Monitoring Scores are all used, but validated quantitative traps are more useful for management decisions than purely visual checks. ³

For a benchmark, the best quantified trap-based reference bands presently available are these: with **corrugated cardboard traps**, mean counts of **1–250 mites/trap** indicate a low infestation, **251–500** a medium infestation, and **more than 500** a high infestation; with **AviVet traps**, **50 mg or less** is low, **51–250 mg** is medium, and **more than 250 mg** is high; with **positive tape or SPT-type traps**, **less than 20% positive** is low, **20–50%** is medium, and **more than 50%** is high. The most decision-relevant published action signal is still provisional rather than finalized: Chiron and colleagues' approach, summarized in the Frontiers IPM review, found that when **more than 20% of placed SPT traps were positive for three**

successive monitoring rounds, the flock was approaching the maximum-acceleration point of infestation growth, which is the practical moment to act before the population enters a harder-to-control expansion phase. The same review is explicit that a single universal economic threshold has **not** yet been validated across all houses and flocks. 4

The observable farm-data signature is broader than trap counts. Heavy PRM pressure is associated with disturbed nocturnal behavior, increased head scratching, head shaking, and preening, impaired resting behavior, anemia, performance loss, lower egg weight, and sometimes more skin damage, pecking-related injury, and mortality; in one EU label summary, eliminating mites with fluralaner was associated with statistically significant improvement in behavioral indicators and lower corticosterone. In practice, the alignment benchmark should treat **rising trap counts plus altered night activity plus declining egg mass or welfare scores** as stronger evidence than any one signal alone. 5

Treatment should be structured as **empty-house sanitation plus occupied-house mitigation plus tightly controlled pharmacology**. The literature and case-report evidence support dry cleaning, wet cleaning, disinfection, and preventive physical treatments such as **synthetic amorphous silica** during downtime, with continuous monitoring after restocking. For pharmacologic control in jurisdictions where it is licensed, **fluralaner** is the strongest evidence-backed systemic PRM treatment; EU product information for Exzolt states **0.5 mg/kg twice, 7 days apart**, with **eggs: zero days** withdrawal and **meat/offal: 14 days**, and notes activity against *D. gallinae* within four hours of feeding. By contrast, a phoxim spray product listed in the EU veterinary medicines register requires eggs to be removed before treatment and eggs laid during and on the same day after treatment discarded; its listing shows **eggs: 12 hours** and **meat/offal: 25 days after the second treatment**. The main integrity point is jurisdictional: current U.S. Exzolt labeling is for **northern fowl mites**, not PRM, so a farm cannot honestly generalize a U.S. label to red mite control. 6

The tension structure here is sharp. The welfare-profit frame says “delay treatment until the problem is obvious.” The science says that delay increases both welfare harm and eventual control cost. The false binary is “either spray often or accept mites”; IPM evidence shows a third path of continuous monitoring, sanitation, and targeted use of acaricides. The integrity risk is especially high when producers ignore jurisdiction-specific labels or keep trap counts informal so intervention can be postponed. 7

Water quality and water-consumption-drop signatures

The standards are unequivocal on **continuous access** and only partly quantitative on quality. UEP requires clean, fresh water; annual potability testing; water meters to measure consumption; and adequate drinker provision, including **1 nipple or cup per 10 hens** and no more than **26 feet** travel to reach drinkers. RSPCA requires clean fresh water at all times, at least **1 nipple or cup per 10 hens**, and testing of non-mains water **every 6 months**, with recorded microbiological targets at the source of **coliforms <100 cfu/mL** and **total viable counts <1,000 cfu/mL**; it also says changes in water smell, clarity, taste, drinking habits, performance, or health should trigger re-testing. Certified Humane adds useful performance thresholds for temperature: water intake drops when drinking water rises above **80°F/27°C** or falls below **60°F/16°C**, and water above **100°F/38°C** or below **50°F/10°C** is not recommended; it also requires an emergency water supply for at least **24 hours** during a main shutoff. 8

For chemistry, the assurance schemes are less specific than many farm managers assume. A recent poultry water assessment used **pH 6.5–8.5**, **nitrate 50 mg/L**, and **nitrite 3 mg/L** as acceptable reference points, but the same study still found substantial total coliform and *E. coli* contamination in some poultry water

samples, underscoring that fecal contamination is often the more immediate practical risk than moderate mineral variation. In other words, for benchmark purposes, water quality should be treated as **microbiology plus delivery performance plus temperature**, not just a chemistry sheet. ⁹

The strongest diagnostic insight is the difference between **ordinary heat stress** and a **true water-consumption-drop event**. In laying hens under warm cyclic conditions, the water-to-feed ratio rises from about **1.8–2.0** in acclimation/recovery to about **3.0–3.4** during heat exposure. Extension guidance similarly indicates birds may increase water intake by **2 to 4 times** during heat stress. That means a flock-wide **drop** in metered water during hot weather is not the normal signature of heat stress at all; it is a red-flag signature for drinker access failure, water temperature failure, line blockage, pump/pressure fault, or severe disease suppressing intake. ¹⁰

The benchmark differential diagnosis is therefore straightforward. **Disease-type signatures** usually show water down alongside feed down, depression, and a contemporaneous egg-production decline; Newcastle disease guidance for adult layers, for example, lists decreased water consumption with dramatic production drop. **Heat-stress signatures** typically show water up and feed down, with higher water:feed ratios, panting, thinner shells, and vulnerability that improves if water is cool and accessible. **Equipment-fault signatures** are suggested by an abrupt meter drop, line-specific bird crowding, dry nipples or cups, pressure irregularities, or localized litter changes without the expected epidemiologic spread pattern of disease. ¹¹

The tension structure is classic. The welfare-profit temptation is to read only feed or egg data and ignore water-metric anomalies until the flock declares itself clinically. The false binary is “if birds are not dead, the water system is fine”; the science says water is one of the earliest flock-wide indicators. The integrity failure is to keep only monthly or retrospective water records when the standards and physiology support **daily** consumption surveillance as a welfare-critical signal. ¹²

Induced molt

UEP is explicit that **only non-feed-withdrawal molt methods are permitted**, and it makes feed-withdrawal molt an audit-fail condition. The current cage-free UEP standard is performance-based rather than recipe-based: hens must receive a **palatable feed source suitable for non-producing hens**, water must always be available, the light period may be reduced to **no fewer than 8 hours** in closed barns or to natural day length in open barns, low-body-weight or unthrifty hens must be culled before molt, final body-weight loss may not exceed **25%** of premolt average flock weight, and corrective action is required if molt mortality increases more than **0.5% on a 3-day average above pre-molt mortality**. UEP also continues to describe induced molt as a way to extend flock life from about **75–85 weeks** to **110 weeks or longer** under favorable conditions. ¹³

What the standards do **not** do is prescribe one universal feed formula. The peer-reviewed literature shows that practical non-feed-withdrawal programs typically use low-energy, low-protein or high-fiber diets based on ingredients such as **wheat middlings, soybean hulls, corn, corn gluten feed, or DDGS**, generally over a **28-day** molt period. A widely cited review concluded that post-molt performance with these nonwithdrawal techniques was comparable to fasting-based methods, and one Illinois DDGS study used a limited-feeding schedule of **65 g/hen/day for 16 days followed by 55 g/hen/day for 12 days** in some treatment arms. The most defensible benchmark conclusion is therefore that the standard is **outcome**

constrained, while the literature offers several acceptable ration designs rather than a single canonical protocol. ¹⁴

Feed withdrawal was banned for both **welfare** and **Salmonella** reasons. Reviews consistently report that feed deprivation during induced molt alters the gastrointestinal environment and immune status in ways that increase susceptibility to *Salmonella Enteritidis* colonization, environmental contamination, and, at least in some studies, egg contamination risk. UEP's shift away from fasting was driven by university research specifically commissioned to develop practical alternatives, and the science review literature supports the view that non-feed-withdrawal methods provide comparable laying-hen well-being while avoiding a major part of the food-safety risk attached to fasting molt programs. ¹⁵

The farm-data signature of a compliant non-feed-withdrawal molt is therefore not just "birds stop laying." It is a package: continued feed availability, water availability, controlled light reduction, body-weight loss held to **25% or less**, and mortality held close enough to baseline that any increase over the **0.5% three-day threshold** triggers immediate correction. In benchmark terms, a manager who can only report post-molt production but cannot show body-weight and mortality controls has not demonstrated welfare alignment. ¹⁶

The tension structure is unusually visible here. The welfare-profit narrative historically favored fasting because it was simple and productive. The false binary says "either you molt efficiently or you protect welfare." The modern evidence base rejects that framing: non-feed-withdrawal programs can preserve second-cycle value while avoiding a now well-documented welfare and Salmonella hazard. The integrity issue is straightforward: a farm that uses the language of "molt" while still relying on de facto fasting is out of alignment with both UEP's standard and the evidence base. ¹⁷

Nest-box access, floor eggs, piling, and smothering

All three standards require substantial nest provision. UEP and Certified Humane require either **1 nest per 5 hens** or, for community nests, **9 square feet per 100 hens**. RSPCA requires **1 individual nest per 5 hens** or **1 m² of nesting substrate per 120 hens** for group nesting, plus prompt collection of floor eggs and attention to over-populated nests. These are minimum space rules, not performance thresholds, and the literature shows clearly that meeting the space minimum does not by itself guarantee good nest use. ¹⁸

The best recent commercial prevalence estimate for floor eggs comes from Australian cage-free flocks, where floor-egg levels ranged from **0.01% to 17% of daily egg production**. That wide range matters: floor eggs are not an inevitable constant of cage-free systems, but neither are they a rare outlier. The drivers repeatedly identified across reviews and standards are pullet training before lay, early familiarity with perches and nest substrates, housing-system layout, dark or secluded lay sites outside nests, uneven nest attractiveness, crowding at preferred nests, and poor management of the transition into lay. Early perch access is associated with reduced later floor-egg prevalence, and in one commercial management example, increasing light intensity beneath an aviary from about **5 lux to 20-50 lux** reduced floor eggs in that area by **up to 80%**, though eggs could then shift to other locations if the whole-house incentive structure was not fixed. ¹⁹

The observable farm-data signature of a nest-access problem is usually visible before it becomes a serious egg-loss problem: increasing floor eggs in the early morning, persistent hotspots under tiers or in corners, uneven nest occupancy, crowding at specific nest banks, and heavy manual pick-up loads despite nominal

compliance with nest-space standards. Humane Farm Animal Care explicitly notes two practical countermeasures for substrate areas that attract laying: restricting access to the substrate to the **afternoon portion of the day** and increasing lighting above the substrate to encourage dustbathing and discourage egg-laying there. UEP similarly notes that management tools to discourage laying near walls and outside nests are permissible, while RSPCA emphasizes prompt floor-egg pickup and nest familiarity before lay. ²⁰

Piling and smothering are separate but closely related hazards. Current reviews describe smothering as a major welfare and economic bottleneck in non-cage systems. UK work has reported that **more than 50%** of surveyed free-range flocks exhibited smothering behavior, and one preliminary investigation found that smothering accounted for **40% of all mortality** in four of ten flocks; a newer Australian epidemiology study reported a smothering mortality rate of **183 deaths per 10,000 bird-years**. UK project material further states that badly affected non-cage flocks can lose **more than 20%** of birds to smothering. Drivers reported in the literature include light and temperature gradients, fear responses, crowding around preferred resources, site-specific attraction to corners or gates, and system designs that create congregation points. ²¹

The tension structure here is one of the clearest in cage-free production. The welfare-profit framing treats floor eggs as a sanitation and labor problem, and piling as a mysterious mortality problem. The science says both are fundamentally **resource-access and behavior-management problems**. The false binary is “if nest-space minimums are met, the flock is fine.” The integrity failure is to certify design compliance while ignoring performance signals like chronic floor-egg hotspots, repeated nest over-population, or unexplained mortality clusters from smothering. ²²

Footpad dermatitis, litter quality, and dustbathing

For layers, the three schemes converge more strongly on **litter quality requirements** than on a single FPD outcome threshold. RSPCA requires access at all times to well-maintained litter, a minimum of **one-third** of usable area as litter, litter depth of at least **5 cm** within 24 hours of placement and at least **10 cm** by the latest after the first 2 months of house use, and litter maintained in a dry, friable condition that allows dustbathing. It also requires daily litter checks and remedial action on any individual **capped area greater than 1 m²**. Certified Humane requires sufficient suitable substrate to allow dustbathing and foraging, with at least **15%** of available floor space as suitable substrate in enclosed systems, and it requires wet litter from flooding to be replaced immediately. UEP requires litter to be loose and friable and says wet and hard-packed areas must be corrected daily. ²³

The most practical scoring systems are animal-based rather than input-based. Welfare Quality's laying-hen protocol scores foot pad lesions on observed birds as **0 = intact, 1 = some problems, and 2 = swollen, dorsally visible**, while a histologically validated 5-point scale for poultry lesions distinguishes mild from extensive moderate and severe lesions by both lesion type and extent. Those tools are useful for a benchmark because they convert “bad litter” into a visible animal outcome. At the same time, none of the three assurance schemes publishes a universally validated **flock-prevalence intervention threshold** for layer FPD analogous to a broiler trigger score. That omission matters: the hard thresholds in current standards are mostly **environmental triggers**, not lesion-prevalence triggers. ²⁴

The welfare science on drivers is consistent. Wet litter is the major causal pathway, with ammonia, drinker leakage or spill, ventilation failure, stocking density, nitrogen load, body weight, and nutrition acting through litter moisture and chemical irritation. The dustbathing evidence points in the same direction: hens

prefer fine, friable substrates such as **sand and peat moss** for dustbathing, and poor-quality or coarse litter undermines one of the behavior's core welfare functions. Newer field data from German laying flocks found **78.9%** of hens with no alterations, **18.6%** slight, **2.2%** moderate, and **0.3%** severe FPD, showing that severe lesions can be kept relatively low, even though older work in large cage-free surveys found much higher flock-level prevalence. ²⁵

For an alignment benchmark, the operational threshold should therefore be framed in two tiers. The **hard standard threshold** is immediate husbandry action when litter becomes wet, capped, contaminated, or non-friable, including the RSPCA **>1 m² capped-area** trigger and HFAC's requirement to replace wet litter from flooding immediately. The **animal-outcome threshold** should be trend-based: any rise in score-2 lesions, or persistent house-zone clustering of score-1 lesions, is a welfare-management failure signal even if the scheme has not published a percentage cutoff. That is an inference from the standards plus the lesion-scoring literature, rather than a single published assurance threshold. ²⁶

The tension structure is again familiar. The welfare-profit temptation is to manage litter only when production or air quality becomes obviously poor. The false binary says "dustbathing litter is enrichment, not health infrastructure." The science shows that litter is both a behavior substrate and a foot-health control point. The integrity danger is to report litter area percentages while ignoring capping, moisture, substrate particle size, and lesion trends in the birds themselves. ²⁷

Catching, handling, and transport of osteoporotic spent hens

End-of-lay hens are unusually vulnerable because chronic calcium mobilization for shell formation, combined with age and housing history, leaves many birds osteoporotic and fracture-prone. Reviews of avian osteoporosis are explicit that bone breakage becomes a severe welfare problem during catching and transport, and FAWC likewise concluded that fractures during depopulation have serious welfare consequences and should be driven down through handling, genetics, nutrition, and design. ²⁸

The standards recognize this fragility but differ in specificity. UEP is the clearest on handling mechanics: birds in cage-free systems must be caught and held **upright with both hands** or carried by **both legs with no more than 3 birds in each hand**; they must not be carried by a single leg, wing, head, neck, or tail; feed withdrawal for end-of-lay hens must be no more than **18 hours before the beginning of catching** if they will be transported, or **24 hours** if culled on-farm; and water must **not** be withdrawn prior to catching. Certified Humane requires a depopulation action plan and keeps the total time from start of loading to completion of unloading of transport trays to **less than 10 hours**, with water available until catching begins. RSPCA requires approved catching teams, a depopulation action plan, vehicles parked as near as possible to the house, and, in hot weather above **21°C**, transport at night or in the coolest part of the day or a **20% reduction in stocking density**. ²⁹

The fractures and dead-on-arrival data justify this caution. Gregory's classic study found that catching and removing hens by **one leg** resulted in **11–14%** of birds acquiring a broken bone, whereas catching and supporting birds under the breast reduced breakage sharply. More recent commercial non-cage depopulation work suggests that about **8.1%** of hens sustain severe injuries or show a considerable stress reaction during the catching, carrying, and crating process. For transport mortality, large commercial datasets report spent-hen DOA rates around **1.22%** in Italy, an overall mortality of **0.516%** in Czech end-of-lay hens, and higher mortality at longer trips, including **0.801%** at **201–300 km**; a more recent study

reported the worst mortality combination in **winter with distances over 300 km**, reaching **1.21%**. These are high figures by poultry-transport standards. ³⁰

The observable farm-data signatures are measurable. On farm: more wing flapping, bruising, dropping events, crowding in corners, delayed catching throughput, and post-catch non-ambulatory or visibly painful birds signal a poor depopulation process. At slaughter: elevated DOA, bruises, fractures, and reject rates track back to catching, loading, transport distance, season, and thermal management. The evidence on handling method favors **upright catching** over inverted one-leg catching on direct welfare grounds; recent work on end-of-lay hens found better catcher-bird interaction, less wing flapping, and fewer wing bruises with upright catching, even though the method is more labor-costly. That matters for benchmarking because a method can be welfare-superior while still facing adoption resistance from throughput and ergonomics constraints. ³¹

On-farm humane killing is not a fringe issue here; it is a welfare necessity for unfit birds. UEP requires visibly unfit birds to be euthanized rather than loaded for transport. RSPCA permits planned on-farm killing by **hand-held electrical stunning followed by neck cutting** or **captive bolt followed by neck dislocation and bleeding**, and bars equipment that crushes the neck. Certified Humane likewise requires prompt euthanasia where pain is uncontrollable and calls for trained staff or veterinary involvement. The benchmark implication is that transport should be reserved for birds **fit to travel**; fracture-suspect, moribund, or severely compromised hens belong on the euthanasia pathway, not the truck. ³²

The tension structure is perhaps the sharpest in this brief. The welfare-profit pressure is obvious: faster catching is cheaper. The false binary says “it is either efficient or humane.” The evidence instead shows a real trade-off in labor and ergonomics, but also shows that poor catching externalizes welfare cost into fractures, bruises, and DOA. The integrity risk is to define spent-hen losses as an unavoidable end-of-lay attrition problem when many of the losses are handling- and transport-mediated. ³³

Antibiotics, label claims, and duty to treat

The short answer is **yes in substance**, but with different levels of explicitness across schemes. RSPCA and Certified Humane both clearly require welfare-first treatment decisions that would override an absolute “No Antibiotics Ever” marketing claim. UEP’s cage-free welfare standard is less explicit on antibiotics, but it still requires proper care of sick or injured birds and humane euthanasia where needed, so the same welfare logic applies in practice. ³⁴

For **RSPCA**, the standard is explicit that antibiotics must be used only when necessary and responsibly, prophylactic use is not permitted, but metaphylaxis for clinically affected groups is not banned by that prophylaxis clause, and very exceptional preventive use may still be justified if a veterinary surgeon judges it in the interests of the affected animal’s welfare. The same standard grounds itself in freedom from pain, injury, and disease through rapid diagnosis and treatment. That is not compatible with withholding needed treatment to preserve a label claim. ³⁵

For **Certified Humane**, the standard permits antibiotics and coccidiostats **only for therapeutic reasons** and **only under veterinary direction**, and it requires caretakers to know when a veterinarian should be consulted so appropriate treatment can be initiated. Certified Humane also requires prompt euthanasia when pain is severe and uncontrollable. In effect, the scheme plainly rejects the idea that a marketing claim may justify leaving a sick bird untreated. ³⁶

For **UEP Certified**, the cage-free guideline set retrieved here does not set out a comparably detailed antibiotic chapter, so the conclusion must be more cautious. Still, UEP requires proper care for injured or sick birds in its training materials and requires humane euthanasia of sick or injured birds during production. That means a UEP-aligned farm cannot ethically defend withholding treatment solely to maintain a “No Antibiotics Ever” claim. The label may have to yield; the bird’s welfare does not. ³⁷

The tension structure is the benchmark’s central alignment test. The welfare-profit tension says that losing a label premium is costly. The false binary says “either keep the NAE claim or help the birds.” The scientifically and ethically coherent position is different: **treat the sick animals, then remove or segregate them from the label claim if required**. The integrity failure is to preserve a marketing story by allowing preventable suffering. On the standards reviewed here, especially RSPCA and Certified Humane, that failure is not just ethically weak; it is nonaligned with the standard’s own welfare logic. ³⁸

¹ ⁸ ¹² ¹³ ¹⁶ ¹⁷ ¹⁸ ²⁹ ³² uepcertified.com

https://uepcertified.com/wp-content/uploads/2023/10/CF-UEP-Guidelines_2024.pdf

² ⁵ ⁶ Successful long-term control of poultry red mite (*Dermanyssus gallinae*) infestations in floor-kept laying hens via integrated pest management—a case report | Parasitology Research | Springer Nature Link

<https://link.springer.com/article/10.1007/s00436-023-07954-9>

³ ⁴ vb.nweurope.eu

<https://vb.nweurope.eu/media/18018/brochure-monitoring-en.pdf>

⁷ Frontiers | Possibilities for IPM Strategies in European Laying Hen Farms for Improved Control of the Poultry Red Mite (*Dermanyssus gallinae*): Details and State of Affairs

<https://www.frontiersin.org/journals/veterinary-science/articles/10.3389/fvets.2020.565866/full>

⁹ Physical, chemical and microbiological assessments of ...

https://ojvr.org/index.php/ojvr/article/view/2067/2432?utm_source=chatgpt.com

¹⁰ Drinking water temperature effects on laying hens ...

https://pubmed.ncbi.nlm.nih.gov/12033408/?utm_source=chatgpt.com

¹¹ Factors Affecting Egg Production in Backyard Chicken Flocks

https://ask.ifas.ufl.edu/publication/PS029?utm_source=chatgpt.com

¹⁴ academic.oup.com

<https://academic.oup.com/ps/article/86/6/1260/1579938>

¹⁵ A review on development of novel strategies for controlling ...

https://pubmed.ncbi.nlm.nih.gov/23300320/?utm_source=chatgpt.com

¹⁹ Floor Eggs in Australian Cage-Free Egg Production

https://pubmed.ncbi.nlm.nih.gov/40646866/?utm_source=chatgpt.com

²⁰ ³⁶ Layers

https://certifiedhumane.org/wp-content/uploads/Standard_LayingHens-2023.pdf

²¹ Welfare aspects in egg production

https://lohmman-breeders.com/lohmmaninfo/welfare-aspects-in-egg-production/?utm_source=chatgpt.com

²² ²³ ²⁶ ²⁷ ³⁴ ³⁵ ³⁸ rspca.org.uk

<https://www.rspca.org.uk/documents/d/rspca/rspca-welfare-standards-for-laying-hens>

24

https://www.welfarequalitynetwork.net/media/1294/wq_laying_hen_protocol_20_def-december-2019.pdf

25 **Factors associated with footpad dermatitis in German ...**

https://onlinelibrary.wiley.com/doi/full/10.1111/aab.12923?utm_source=chatgpt.com

28 **Welfare implications of avian osteoporosis**

https://pubmed.ncbi.nlm.nih.gov/14979568/?utm_source=chatgpt.com

30 33 **Effect of catching method and lighting intensity on the ...**

https://europepmc.org/article/med/8447051?utm_source=chatgpt.com

31 **Upright versus inverted catching and crating end-of-lay hens**

https://pubmed.ncbi.nlm.nih.gov/39127006/?utm_source=chatgpt.com

37 **Employee training vital to hen care - United Egg Producers**

https://uepcertified.com/employee-training-vital-to-hen-care/?utm_source=chatgpt.com