

CHEMISTRY IN CESM2

EIRIK ROLLAND ENGER

December 7, 2022




Colorado State University

MOTIVATION


STRATOSPHERIC SUDDEN WARMINGS

Tropospheric polar vortex versus stratospheric polar vortex



climate.gov


STRATOSPHERIC SUDDEN WARMINGS



climate.gov

- Defined as a wind reversal (eastward) at 10 hPa (~ 25 km), 60° N
- Big improvement from including updated parametrizations of turbulent mountain stress (TMS), surface stress due to unresolved topography
- A lack of stratospheric internal variability without a high-top atmosphere


STRATOSPHERIC SUDDEN WARMINGS



[3] Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". 2019

EVOLUTION OF THE OZONE LAYER

- WACCM6 is able to reproduce the evolution of the ozone layer (also SH polar ozone hole)
- Ozone variability in the tropical stratosphere improves on the inclusion of an internally generated quasi-biennial oscillation (QBO)



ATMOSPHERIC BLOCKING


Frequency of the meridional gradient of 500-hPa geopotential height below a threshold of $\text{GHGS} > 0$, $\text{GHGN} < -5 \text{ m/degree}$

$$\text{GHGS} = \frac{Z(\phi_0) - Z(\phi_S)}{\phi_0 - \phi_S}$$
$$\text{GHGN} = \frac{Z(\phi_N) - Z(\phi_0)}{\phi_N - \phi_0}$$

where $\phi_N = 78.75^\circ\text{N} + \Delta$, $\phi_0 = 60^\circ\text{N} + \Delta$, $\phi_S = 41.25^\circ\text{N} + \Delta$ and $\Delta = -3.75^\circ, 0^\circ, 3.75^\circ$ [1].

[1] D'Andrea et al. "Northern Hemisphere atmospheric blocking as simulated by 15 atmospheric general circulation models in the period 1979–1988". 1998


BLOCKING FREQUENCY



SEA ICE

- The September NH sea ice extent is better in WACCM6 than in CAM6
- Less downward surface SW and LW in WACCM6 due to higher LWP¹ which in turn is due to higher aerosol number.

⇒ Tropospheric aerosol chemistry impacts Arctic sea ice.



¹ liquid water path

[3] Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". 2019


IMPLEMENTATION

COMPUTATIONAL COST

Table 1: Approximate costs of running different atmosphere models
(From lecture by Mills)

Configuration	Resolution	Chemistry	Core-hours/simulation years
CAM6	1°, 32 L	CAM	3700
WACCM6	2°, 70 L	MA	5400
WACCM6	1°, 70 L	TSMLT	22 000
WACCM6-SC	1°, 70 L	SC	6000
WACCM6-SD	1°, 88 L	TSMLT	23 000
WACCM5.4	1°, 110 L	MA	20 000
WACCM5.4-SC	1°, 110 L	SC	9000

SPATIAL



cesm.ucar.edu

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions
 - MAD (Middle atmosphere with D region ion chemistry)

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions
 - MAD (Middle atmosphere with D region ion chemistry)
 - Adds 15 positive and 21 negative ions

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions
 - MAD (Middle atmosphere with D region ion chemistry)
 - Adds 15 positive and 21 negative ions
 - Thus, below 75 km electrons are no longer the main negative charge carrier

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions
 - MAD (Middle atmosphere with D region ion chemistry)
 - Adds 15 positive and 21 negative ions
 - Thus, below 75 km electrons are no longer the main negative charge carrier
- Additional thermosphere eXtension (WACCM-X)

CHEMISTRY IN TSMLT

MAM4 (*Modal Aerosol Model*), also used in CAM6, but WACCM6 adds chemistry.

- Includes the chemical families O_x , NO_x , HO_x , ClO_x and BrO_x , as well as CH_4
- Allows growth of sulfate aerosols, so the prognostic stratospheric aerosols can increase in width
- Maximum altitude of 20 km for eruptions outputting more than 3.5 Tg SO_2


MOZART (*Model for OZone And Related chemical Tracers*)

- The chemical mechanism in CESM2, available from WACCM6, but also CAM-chem
- See table 2² for a complete list of chemical reactions included in CESM2 when run with the TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) configuration.

²https://agupubs.onlinelibrary.wiley.com/action/downloadSupplement?doi=10.1029%2F2019MS001882&file=jame21103-sup-0003-2019MS001882+Table_SI-S02.pdf


STRATOSPHERIC AEROSOLS

In CAM, stratospheric aerosols are prescribed based on output from previous WACCM simulations



Aerosol optical depth from stratospheric volcanic eruption in WACCM

STRATOSPHERIC AEROSOL OPTICAL DEPTH



Stratospheric aerosol optical depth at different locations agree well

[3] Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". 2019

LUMPING

- TSMLT has 231 solution species
- Species are lumped together to reduce the computational cost
- Example: $C_{10}H_{16}$ in MOZART-4 turned into five new lumped species, with APIN, BPIN, LIMON, MYRC and BCARY giving the primary degradation rates.

SOLAR AND GEOMAGNETICS

- Photoionization and heating rates uses parametrization of Solomon and Qian (2005), with input from the $F_{10.7}$ index
- Ion-pair production rates are prescribed
- Low energy electrons included by the parametrized auroral oval model by Roble and Ridley (1994)
- Input to the model is HP, hemispheric power, related to the K_p index:

$$HP = \begin{cases} 16.82 \exp(0.32K_p) - 4.86, & K_p \leq 7 \\ 153.13 + 73.4(K_p - 7.0), & K_p > 7 \end{cases}$$

- Since WACCM3, E region ionosphere is represented with a chemistry consisting of O^+ , O_2^+ , N^+ , N_2^+ , NO^+

REFERENCES I



F. D'Andrea et al. "Northern Hemisphere atmospheric blocking as simulated by 15 atmospheric general circulation models in the period 1979–1988". In: *Climate Dynamics* 14.6 (May 1998), pp. 385–407. ISSN: 1432-0894. DOI: 10.1007/s003820050230. URL: <https://doi.org/10.1007/s003820050230>.



R. R. Garcia et al. "Simulation of secular trends in the middle atmosphere, 1950–2003". In: *Journal of Geophysical Research: Atmospheres* 112.D9 (2007). DOI: <https://doi.org/10.1029/2006JD007485>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006JD007485>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JD007485>.



A. Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". In: *Journal of Geophysical Research: Atmospheres* 124.23 (2019), pp. 12380–12403. DOI: <https://doi.org/10.1029/2019JD030943>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019JD030943>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD030943>.

REFERENCES II



Daniel R. Marsh et al. "Climate Change from 1850 to 2005 Simulated in CESM1(WACCM)". In: *Journal of Climate* 26.19 (2013), pp. 7372–7391. DOI: 10.1175/JCLI-D-12-00558.1. URL: <https://journals.ametsoc.org/view/journals/clim/26/19/jcli-d-12-00558.1.xml>.



Daniel R. Marsh et al. "Modeling the whole atmosphere response to solar cycle changes in radiative and geomagnetic forcing". In: *Journal of Geophysical Research: Atmospheres* 112.D23 (2007). DOI: <https://doi.org/10.1029/2006JD008306>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006JD008306>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JD008306>.



Michael J. Mills et al. "Global volcanic aerosol properties derived from emissions, 1990–2014, using CESM1(WACCM)". In: *Journal of Geophysical Research: Atmospheres* 121.5 (2016), pp. 2332–2348. DOI: <https://doi.org/10.1002/2015JD024290>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2015JD024290>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JD024290>.



Michael J. Mills et al. “Radiative and Chemical Response to Interactive Stratospheric Sulfate Aerosols in Fully Coupled CESM1(WACCM)”. In: *Journal of Geophysical Research: Atmospheres* 122.23 (2017), pp. 13, 061–13, 078. DOI: <https://doi.org/10.1002/2017JD027006>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2017JD027006>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027006>.



R. G. Roble and E. C. Ridley. “A thermosphere-ionosphere-mesosphere-electrodynamics general circulation model (time-GCM): Equinox solar cycle minimum simulations (30–500 km)”. In: *Geophysical Research Letters* 21.6 (1994), pp. 417–420. DOI: <https://doi.org/10.1029/93GL03391>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/93GL03391>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/93GL03391>.



Stanley C. Solomon and Liying Qian. "Solar extreme-ultraviolet irradiance for general circulation models". In: *Journal of Geophysical Research: Space Physics* 110.A10 (2005). DOI: <https://doi.org/10.1029/2005JA011160>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2005JA011160>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005JA011160>.

