

CHEMISTRY IN CESM2

EIRIK ROLLAND ENGER

December 7, 2022




Colorado State University

MOTIVATION


STRATOSPHERIC SUDDEN WARMINGS

Tropospheric polar vortex versus stratospheric polar vortex



climate.gov


STRATOSPHERIC SUDDEN WARMINGS



climate.gov

- Defined as a wind reversal (eastward) at 10 hPa (~ 25 km), 60° N
- Big improvement from including updated parametrizations of turbulent mountain stress (TMS), surface stress due to unresolved topography
- A lack of stratospheric internal variability without a high-top atmosphere


STRATOSPHERIC SUDDEN WARMINGS



[3] Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". 2019

EVOLUTION OF THE OZONE LAYER

- WACCM6 is able to reproduce the evolution of the ozone layer (also SH polar ozone hole)
- Ozone variability in the tropical stratosphere improves on the inclusion of an internally generated quasi-biennial oscillation (QBO)



[3] Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". 2019

ATMOSPHERIC BLOCKING


Frequency of the meridional gradient of 500-hPa geopotential height below a threshold of $\text{GHGS} > 0$, $\text{GHGN} < -5 \text{ m/degree}$

$$\text{GHGS} = \frac{Z(\phi_0) - Z(\phi_S)}{\phi_0 - \phi_S}$$
$$\text{GHGN} = \frac{Z(\phi_N) - Z(\phi_0)}{\phi_N - \phi_0}$$

where $\phi_N = 78.75^\circ\text{N} + \Delta$, $\phi_0 = 60^\circ\text{N} + \Delta$, $\phi_S = 41.25^\circ\text{N} + \Delta$ and $\Delta = -3.75^\circ, 0^\circ, 3.75^\circ$ [1].

[1] D'Andrea et al. "Northern Hemisphere atmospheric blocking as simulated by 15 atmospheric general circulation models in the period 1979–1988". 1998

BLOCKING FREQUENCY




[3] Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". 2019

SEA ICE

- The September NH sea ice extent is better in WACCM6 than in CAM6
- Less downward surface SW and LW in WACCM6 due to higher LWP¹ which in turn is due to higher aerosol number.

⇒ Tropospheric aerosol chemistry impacts Arctic sea ice.



¹ liquid water path

[3] Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". 2019


IMPLEMENTATION

COMPUTATIONAL COST

Table 1: Approximate costs of running different atmosphere models
(From lecture by Mills)

Configuration	Resolution	Chemistry	Core-hours/simulation years
CAM6	1°, 32 L	CAM	3700
WACCM6	2°, 70 L	MA	5400
WACCM6	1°, 70 L	TSMLT	22 000
WACCM6-SC	1°, 70 L	SC	6000
WACCM6-SD	1°, 88 L	TSMLT	23 000

SPATIAL



cesm.ucar.edu

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions
 - MAD (Middle atmosphere with D region ion chemistry)

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions
 - MAD (Middle atmosphere with D region ion chemistry)
 - Adds 15 positive and 21 negative ions

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions
 - MAD (Middle atmosphere with D region ion chemistry)
 - Adds 15 positive and 21 negative ions
 - Thus, below 75 km electrons are no longer the main negative charge carrier

CHEMISTRY VERSIONS

- Neutral chemistry model versions of WACCM6
 - TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) *default*
 - TS (troposphere and stratosphere mechanism)
 - MA (Middle atmosphere)
 - Similar the older WACCM4, available in nominal 2° only
 - Reduced set of tropospheric reactions
 - MAD (Middle atmosphere with D region ion chemistry)
 - Adds 15 positive and 21 negative ions
 - Thus, below 75 km electrons are no longer the main negative charge carrier
- Additional thermosphere eXtension (WACCM-X)

CHEMISTRY IN TSMLT

MAM4 (*Modal Aerosol Model*), also used in CAM6, but WACCM6 adds chemistry.


- Includes the chemical families O_x , NO_x , HO_x , ClO_x and BrO_x , as well as CH_4
- Allows growth of sulfate aerosols, so the prognostic stratospheric aerosols can increase in width
- Maximum altitude of 20 km for eruptions outputting more than 3.5 Tg SO_2

MOZART (*Model for OZone And Related chemical Tracers*)

- The chemical mechanism in CESM2, available from WACCM6, but also CAM-chem
- See table 2² for a complete list of chemical reactions included in CESM2 when run with the TSMLT (troposphere, stratosphere, mesosphere, lower thermosphere) configuration.

²https://agupubs.onlinelibrary.wiley.com/action/downloadSupplement?doi=10.1029%2F2019MS001882&file=jame21103-sup-0003-2019MS001882+Table_SI-S02.pdf

STRATOSPHERIC AEROSOL OPTICAL DEPTH



Stratospheric aerosol optical depth at different locations agree well

[3] Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". 2019

LUMPING

- TSMLT has 231 solution species
- Species are lumped together to reduce the computational cost
- Example: $C_{10}H_{16}$ in MOZART-4 turned into five new lumped species, with APIN, BPIN, LIMON, MYRC and BCARY giving the primary degradation rates.

SOLAR AND GEOMAGNETICS

- Photoionization and heating rates uses parametrization of Solomon and Qian (2005), with input from the $F_{10.7}$ index
- Ion-pair production rates are prescribed
- Low energy electrons included by the parametrized auroral oval model by Roble and Ridley (1994)
- Input to the model is HP, hemispheric power, related to the K_p index:

$$HP = \begin{cases} 16.82 \exp(0.32K_p) - 4.86, & K_p \leq 7 \\ 153.13 + 73.4(K_p - 7.0), & K_p > 7 \end{cases}$$

- Since WACCM3, E region ionosphere is represented with a chemistry consisting of O^+ , O_2^+ , N^+ , N_2^+ , NO^+

REFERENCES I



F. D'Andrea et al. "Northern Hemisphere atmospheric blocking as simulated by 15 atmospheric general circulation models in the period 1979–1988". In: *Climate Dynamics* 14.6 (May 1998), pp. 385–407. ISSN: 1432-0894. DOI: 10.1007/s003820050230. URL: <https://doi.org/10.1007/s003820050230>.



R. R. Garcia et al. "Simulation of secular trends in the middle atmosphere, 1950–2003". In: *Journal of Geophysical Research: Atmospheres* 112.D9 (2007). DOI: <https://doi.org/10.1029/2006JD007485>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006JD007485>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JD007485>.



A. Gettelman et al. "The Whole Atmosphere Community Climate Model Version 6 (WACCM6)". In: *Journal of Geophysical Research: Atmospheres* 124.23 (2019), pp. 12380–12403. DOI: <https://doi.org/10.1029/2019JD030943>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019JD030943>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD030943>.

REFERENCES II



Daniel R. Marsh et al. "Climate Change from 1850 to 2005 Simulated in CESM1(WACCM)". In: *Journal of Climate* 26.19 (2013), pp. 7372–7391. DOI: 10.1175/JCLI-D-12-00558.1. URL: <https://journals.ametsoc.org/view/journals/clim/26/19/jcli-d-12-00558.1.xml>.



Daniel R. Marsh et al. "Modeling the whole atmosphere response to solar cycle changes in radiative and geomagnetic forcing". In: *Journal of Geophysical Research: Atmospheres* 112.D23 (2007). DOI: <https://doi.org/10.1029/2006JD008306>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006JD008306>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JD008306>.



Michael J. Mills et al. "Global volcanic aerosol properties derived from emissions, 1990–2014, using CESM1(WACCM)". In: *Journal of Geophysical Research: Atmospheres* 121.5 (2016), pp. 2332–2348. DOI: <https://doi.org/10.1002/2015JD024290>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2015JD024290>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JD024290>.



Michael J. Mills et al. “Radiative and Chemical Response to Interactive Stratospheric Sulfate Aerosols in Fully Coupled CESM1(WACCM)”. In: *Journal of Geophysical Research: Atmospheres* 122.23 (2017), pp. 13, 061–13, 078. DOI: <https://doi.org/10.1002/2017JD027006>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2017JD027006>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027006>.



R. G. Roble and E. C. Ridley. “A thermosphere-ionosphere-mesosphere-electrodynamics general circulation model (time-GCM): Equinox solar cycle minimum simulations (30–500 km)”. In: *Geophysical Research Letters* 21.6 (1994), pp. 417–420. DOI: <https://doi.org/10.1029/93GL03391>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/93GL03391>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/93GL03391>.



Stanley C. Solomon and Liying Qian. "Solar extreme-ultraviolet irradiance for general circulation models". In: *Journal of Geophysical Research: Space Physics* 110.A10 (2005). DOI: <https://doi.org/10.1029/2005JA011160>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2005JA011160>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005JA011160>.