esc195\_exam\_2023-c59be #313 Page 1 of 26




# ESC195 - Calculus II Final Exam April 2023

Instructor: J. W. Davis

								] 	;					
	<u> </u>	L	L	L			<u> </u>	<u> </u>	<u> </u>	 	 L. <u>.</u>	l	L	L
Last	nam∈	<u> </u>												
	T	T		<u> </u>		[	Γ	Γ	1			1		
		•	-									_		
Stud	ent r	umbei	c											
	<u> </u>	T		[		I	1		<u> </u>					
	ļ	İ			•						Ì		ļ	

Closed book, no aid sheets, no calculators There are 12 questions; each question is worth 10 marks.





#313 Page 2 of 26

#313 Page 3 of 26



1. Evaluate the integrals:

a) 
$$\int \frac{\ln x}{x^2} dx$$

$$b) \int \frac{x^3}{\sqrt{9+x^2}} \, dx$$

a) 
$$\int \frac{\ln x}{x^2} dx$$
 b)  $\int \frac{x^3}{\sqrt{9+x^2}} dx$  c)  $\int \frac{x}{(x+4)(2x-1)} dx$ 

#### 4A28C1AA-038E-454C-A7DF-3CD4A9A139B3



esc195\_exam\_2023-c59be #313 Page 4 of 26

esc195\_exam\_2023-c59be #313 Page 5 of 26



- 2. Sketch the curves and regions indicated and find an integral(s) representing the area of the region. Do not evaluate the integrals.
  - (a) Inside one loop of the curve:  $r = 4\cos 3\theta$ .
  - (b) Inside both  $r^2 = \sin 2\theta$  and  $r^2 = \cos 2\theta$ .
  - (c) Inside the cardioid  $r = 1 + \cos \theta$ , but outside the circle  $r = 3\cos \theta$

#### DBD1D13E-0E96-401F-80E7-C2BF89E501A0



esc195\_exam\_2023-c59be #313-Page 6 of 26

esc195\_exam\_2023-c59be #313 Page 7 of 26



- 3. Evaluate each limit in two ways: (i) by l'Hospital's rule, (ii) by expanding the functions as Taylor series.
- a)  $\lim_{x \to 0} \frac{e^x 1}{x}$  b)  $\lim_{x \to 0} \frac{2\cos 2x 2 + 4x^2}{2x^4}$  c)  $\lim_{x \to 0} \frac{e^{-2x} 4e^{-x/2} + 3}{2x^2}$

### E63267F8-F89A-42F5-8AA9-06913F9E75C6



esc195\_exam\_2023-c59be

#313 Page 8 of 26

#313 Page 9 of 26



4. (a) Determine whether the series is convergent or divergent:

$$i) \sum_{k=1}^{\infty} k e^{-k^2}$$

ii) 
$$\sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!}$$

(b) Suppose  $a_n > 0$  and  $\frac{a_{n+1}}{a_n} \ge \frac{n}{n+1}$  for all n. Show that  $\sum_{n=1}^{\infty} a_n$  diverges.




29281830-8B92-401D-AFE4-7A606D932B2A

esc195\_exam\_2023-c59be #313 Page 10 of 26

esc195\_exam\_2023-c59be #313 Page 11 of 26



- 5. Given the series:  $\sum_{n=1}^{\infty} \frac{(2x-1)^n}{5^n \sqrt{n}}$ 
  - (a) Find the radius and interval of convergence.
  - (b) Show that the series is not convergent (conditionally or otherwise) for x < -2.






#313 Page 12 of 26

esc195\_exam\_2023-c59be #313 Page 13 of 26



6. Determine from first principles (that is by taking derivatives) the Taylor series for the function  $f(x) = \frac{1}{x^2}$  about x = 1. Express the series in  $\sum$  notation, and find the radius and interval of convergence.






esc195\_exam\_2023-c59be #313 Page 14 of 26

# #313 Page 15 of 26



7. (a) Find an equation for the tangent plane to the surface  $xy^2 + 2z^2 = 12$  at the point (1, 2, 2).

(b) The curve  $\vec{r} = \frac{1}{2}t^2\hat{i} + \frac{4}{t}\hat{j} + (\frac{1}{2}t - t^2)\hat{k}$  intersects the hyperbolic paraboloid  $x^2 - 4y^2 - 4z = 0$  at the point (2, 2, -3). Find the angle of intersection.





#313 Page 16 of 26

esc195\_exam\_2023-c59be #313 Page 17 of 26



8. Use the formal definition for the derivative of a multivariable function (the o(h) formulation) to find the gradient of: f(x, y, z) = xy + yz. Show that all remainder terms are o(h).

### F03904D0-5A51-4E6B-99BE-C88BF6F49D9A




esc195\_exam\_2023-c59be #313 Page 18 of 26

esc195\_exam\_2023-c59be #313 Page 19 of 26



9. Find the maximum and minimum values of  $f(x,y) = \frac{x-y}{1+x^2+y^2}$  on the the upper half plane,  $y \ge 0$ . Provide a sketch of the region, and identify and show the locations of all critical points.






esc195\_exam\_2023-c59be #313 Page 20 of 26

esc195\_exam\_2023-c59be #313 Page 21 of 26



10. The plane x + y + 2z = 2 intersects the paraboloid  $z = x^2 + y^2$  in an ellipse. Use Lagrange Multipliers to find the points on this ellipse that are nearest and furthest from the origin.






#313 Page 22 of 26

esc195\_exam\_2023-c59be #313 Page 23 of 26



11. Given 
$$\int_0^{2\pi} \frac{\cos \theta}{1 - y \cos \theta} d\theta = 2\pi \frac{1 - \sqrt{1 - y^2}}{y \sqrt{1 - y^2}}$$
, where  $0 < y < 1$ , find  $\int_0^{2\pi} \ln(1 - y \cos \theta) d\theta$ 





#313 Page 24 of 26

esc195\_exam\_2023-c59be #313 Page 25 of 26



12. Show that 
$$\int_0^1 x^x dx = 1 - \frac{1}{2^2} + \frac{1}{3^3} - \frac{1}{4^4} + \frac{1}{5^5} - \frac{1}{6^6} + \cdots$$

## B58C66E0-6D92-424F-B5BA-54A899638F1F



esc195\_exam\_2023-c59be