

Last name:

First name:

ID number:

---

# ECE 286

## Midterm exam

February 9, 2022  
9:30 – 10:30 am

**Circle your lecture section:**

**LEC0101 (Monday 9-10)**

**LEC0102 (Monday 1-2)**

Guidelines:

- Write your answer in the space provided for each question. Show your work and use the back side of sheets as needed.
- The exam is Type D. You may use a non-programmable calculator and a one-sided, 8.5 by 11 handwritten note sheet (prepared by you).
- Only exams written in pen will be considered for regrades.

Problem	Score
1	/10
2	/5
3	/7
Total	/22

1. Consider a coin with  $P(H) = 0.7$  and  $P(T) = 0.3$ , where  $H$  denotes ‘Heads’ and  $T$  denotes ‘Tails’. Answer parts (a)-(d) below. Justify your answers.

- (a) You flip the coin three times. Let the random variable  $X$  be the number of Heads. Answer parts (i)-(iii) below.

- i. (2 points) Find the probability mass function of  $X$ .

*Solution:*

$$f(0) = 0.3^3 = 0.027$$

$$f(1) = 3 \times 0.7 \times 0.3^2 = 0.189$$

$$f(2) = 3 \times 0.3 \times 0.7^2 = 0.441$$

$$f(3) = 0.7^3 = 0.343.$$

- ii. (2 points) Find the cumulative distribution of  $X$ .

*Solution:*

$$F(0) = 0.027$$

$$F(1) = 0.216$$

$$F(2) = 0.657$$

$$F(3) = 1.$$

- iii. (1 point) Find the probability that  $1 \leq X \leq 2$ .

*Solution:*

$$F(2) - F(0) = 0.63.$$

- (b) (2 points) Suppose you flip the coin six times. What is the probability of the sequence  $TTTTTH$ ?

*Solution:*

$$0.3^5 \times 0.7 = 0.001701$$

- (c) (3 points) Suppose you flip the coin six times. What is the probability that the sequence has exactly three heads in a row? (For example,  $THHHTT$  has exactly three heads in a row.  $HTHHTT$  and  $HHHHHH$  do not.)

*Solution:* We must compute how many sequences have three heads in a row. Let  $*$  denote  $H$  or  $T$ . We can have:

- $HHHT**$ . There are 4 such sequences, because each  $*$  can be  $H$  or  $T$ . We have:

$$0.7^3 \times 0.3 \times (0.3^2 + 2 \times 0.3 \times 0.7 + 0.7^2) = 0.1029.$$

- $THHHT*$ . There are 2 such sequences. We have:

$$0.7^3 \times 0.3^2 \times (0.3 + 0.7) = 0.03087.$$

- $*THHHT$ . There are 2 such sequences. We have:

$$0.7^3 \times 0.3^2 \times (0.3 + 0.7) = 0.03087.$$

- $**THHH$ . There are 4 such sequences. We have:

$$0.7^3 \times 0.3 \times (0.3^2 + 2 \times 0.3 \times 0.7 + 0.7^2) = 0.1029.$$

Summing the probabilities, we have 0.26754

2. A bag contains 2 red, 3 blue, and 5 green balls. You remove three balls from the bag. Answer parts (a) and (b) below.

- (a) (2 points) What is the probability of drawing first a green ball, second a red ball, and third a green ball, in this order?

*Solution: It is*

$$P(G, R, G) = \frac{5}{10} \cdot \frac{2}{9} \cdot \frac{4}{8} = 1/18 = 0.0555\dots$$

- (b) (3 points) What is the probability of drawing one red ball and two green balls, in any order?

*Solution: There are*

$$\binom{2}{1} = \frac{2!}{1!1!} = 2$$

*ways to draw one red ball. There are*

$$\binom{5}{2} = \frac{5!}{2!3!} = 10$$

*ways to draw 2 green balls. There are thus  $2 \cdot 10 = 20$  ways to draw one red and two green balls. There are*

$$\binom{10}{3} = \frac{10!}{3!7!} = 120$$

*ways to draw 3 balls of any color. Therefore, the chance of drawing 1 red and two green balls is*

$$20/120 = 1/6.$$

3. A factory produces cans of blue and green paint. The events are:

- $B$ : a can has blue paint inside.
- $G$ : a can has green paint inside.
- $C_B$ : the can is labeled blue.
- $C_G$ : the can is labeled green.

We know the following probabilities:

- 30% of cans are labeled blue, i.e.,  $P(C_B) = 0.3$ .
- The probability a can is labeled green and has green paint inside is  $P(G \cap C_G) = 0.6$ .

Answer parts (a)-(c) below.

The probability that a can contains the correct paint color is 0.9.

- (a) (2 points) If you buy a can that is labeled green, what is the probability it has green paint inside?

*Solution:* We want to know  $P(G \mid C_G)$ . From the definition of conditional probability,

$$P(G \mid C_G) = P(G \cap C_G) / P(C_G).$$

We know  $P(C_G) = 1 - P(C_B) = 0.7$ , and therefore  $P(G \mid C_G) = 0.6 / 0.7 = 6/7$ .

- (b) (2 points) If you buy a can that is labeled green, what is the probability it has blue paint inside?

*Solution:* We want to know  $P(B \mid C_G)$ . From the definition of conditional probability,

$$P(B \mid C_G) = P(B \cap C_G) / P(C_G).$$

We know that  $P(C_G) = 1 - P(C_B) = 0.7$ . The law of total probability says

$$P(C_G) = P(B \cap C_G) + P(G \cap C_G).$$

Because  $P(G \cap C_G) = 0.6$ , we know  $P(B \cap C_G) = 0.1$ . Therefore

$$P(B \mid C_G) = 0.1 / 0.7 = 1/7.$$

Alternatively, we can see that  $P(B \mid C_G) = 1 - P(G \mid C_G) = 1/7$ .

- (c) (3 points) Suppose also that the probability a can is labeled blue and has green paint inside is  $P(G \cap C_B) = 0.1$ . If a can has blue paint inside, what is the probability that the can is labeled green?

*Solution:* We want to know  $P(C_G \mid B)$ . Bayes' Rule says

$$P(C_G \mid B) = P(B \mid C_G)P(C_G) / P(B).$$

We know that  $P(C_G) = 0.7$  and  $P(B \mid C_G) = 1/7$ . We must find  $P(B)$ . We know that

$$\begin{aligned} 1 &= P(G \cap C_G) + P(G \cap C_B) + P(B \cap C_G) + P(B \cap C_B) \\ &= 0.6 + 0.1 + 0.1 + P(B \cap C_B). \end{aligned}$$

Therefore,  $P(B \cap C_B) = 0.2$ . We know that

$$\begin{aligned} P(B) &= P(B \cap C_G) + P(B \cap C_B) \\ &= 0.1 + 0.2 \\ &= 0.3. \end{aligned}$$

Therefore,

$$P(C_G \mid B) = 0.7 \cdot (1/7)/0.3 = 1/3.$$