



**University of Toronto**  
**Faculty of Applied Science and Engineering**  
**FINAL EXAMINATION – April, 2019**

**FIRST YEAR – ENGINEERING SCIENCE**

**MAT195S CALCULUS II**

**Examiners: F. Al Faisal and J. W. Davis**

First name (please write as legibly as possible within the boxes)

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Last name

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Student number

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

- Instructions:
- (1) Closed book examination; no calculators, no aids are permitted
  - (2) Answer as many questions as you can. Parts of questions may be answered.
  - (3) Do not separate or remove any pages from this exam booklet.

FOR MARKER USE ONLY					
Question	Marks	Earned	Question	Marks	Earned
1	9		7	10	
2	9		8	10	
3	9		9	12	
4	11		10	11	
5	12		11	10	
6	10		12	8	
X		X	X		X
			Total	121	



22DF936F-B249-4B4C-99CE-B99BD2EE0036


mat195-exam-2019

#181      2 of 26



- 1) Evaluate the integrals:
- a)  $\int \frac{\sqrt{x^2 - 9}}{x^3} dx$
- b)  $\int \sqrt{1 - e^x} dx$

(9 marks)



A399AA13-6853-49C0-B8FC-6E1C842175C2

mat195-exam-2019

#181      4. of 26




2) a) Find the length of the curve:  $x = 1 + 3t^2$ ,  $y = 4 + 2t^3$ ,  $0 \leq t \leq 1$

(4 marks)

b) If two parallel planes intersect a sphere, calculate the surface area of the part of the sphere that lies between the two planes. Hence, deduce that the area depends only on the radius of the sphere and the distance between the two planes, but not on where the planes intersect the sphere.

(5 marks)



1C060ABE-2DDF-4CA4-B439-D86E849E4123

mat195-exam-2019

#181          6 of 26



3) Determine whether the following integrals converge or diverge. If they converge, find their value.

a)  $\int_1^{\infty} \frac{dx}{\sqrt{x}(x+1)}$

b)  $\int_1^e \frac{dx}{x(\ln x)^2}$

c)  $\int_2^{\infty} \frac{x}{\sqrt{x^4-1}} dx$

(9 marks)



10A8E3BB-4D05-4C03-B87F-81C4C9E33B98

mat195-exam-2019


#181      8 of 26





- 4) Sketch the region indicated, and find an integral representing the area of the region. Do not evaluate the integrals.
- a) The region enclosed by one loop of the curve  $r = \sin 3\theta$ .
  - b) The region that lies inside both  $r = \sin 2\theta$  and  $r = \cos 2\theta$ .
  - c) The region that lies inside the circle  $r = 2 \sin \theta$  but outside  $r = \frac{3}{2} - \sin \theta$ .

(11 marks)



5EEEDF4F-A9BA-449F-AACC-B7F0844415A5

mat195-exam-2019


#181      10 of 26



5) Determine whether the following series converge or diverge:

a)  $\sum_{n=1}^{\infty} \frac{n^3}{(\ln 2)^n}$    b)  $\sum_{n=1}^{\infty} \left(1 + \frac{2}{n}\right)^n$    c)  $\sum_{k=2}^{\infty} \frac{1}{k(1 + \ln k)}$    d)  $\sum_{n=1}^{\infty} \frac{n^{\left(n + \frac{1}{n}\right)}}{\left(n + \frac{1}{n}\right)^n}$

(12 marks)



A3C134A2-A6AB-4051-ACFE-AD9E339E962B

mat195-exam-2019

#181      12 of 26



- 6) a) Suppose that the power series  $\sum c_n(x-a)^n$  satisfies  $c_n \neq 0$  for all  $n$ . Show that if  $\lim_{n \rightarrow \infty} \left| \frac{c_n}{c_{n+1}} \right|$  exists, then it is equal to the radius of convergence of the power series.

(5 marks)

- b) Determine the radius and interval of convergence for the series:  $\sum_{k=2}^{\infty} \frac{(x+2)^k}{2^k \ln k}$

(5 marks)



BDE79349-F919-42D2-A30B-DB337D9FA844

mat195-exam-2019

#181      14 of 26




- 7) If the function  $f(x) = (1 - 2\mu x + x^2)^{-1/2}$  is expanded in a Maclaurin series in  $x$ :

$$\frac{1}{\sqrt{1 - 2\mu x + x^2}} = \sum_{n=0}^{\infty} P_n(\mu) x^n$$

The coefficients  $P_n(\mu)$  are called the *Legendre polynomials*.

- Find  $P_0(\mu)$ ,  $P_1(\mu)$ ,  $P_2(\mu)$  and  $P_3(\mu)$ . (The coefficients may be found in several ways, including by direct differentiation, and by expanding  $f$  as a binomial series.)
- Show that in the cases where  $\mu = \pm 1$ , the coefficients can be evaluated using the geometric series.

(10 marks)



96AC304C-74BA-412E-B55D-0D1F68ABFD71

mat195-exam-2019

#181          16 of 26





8) On our first midterm test, we defined the gamma function  $\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt$ , for  $z > 0$ .

a) Let  $I(s) = \int_0^\infty \frac{x^s}{e^x - 1} dx$ ,  $s > 0$ . Show that  $I(s) = \Gamma(s+1) \cdot \sum_{n=1}^\infty \frac{1}{n^{s+1}}$

b) Find  $\int_0^\infty \frac{x}{e^x - 1} dx$  by evaluating  $I(1)$ . Express your answer in the form  $\pi^2/k$ .

Hint 1: On our first midterm test we saw that  $\Gamma(4) = 3!$ . One can similarly show in general that  $\Gamma(n+1) = n!$  for any positive integer  $n$ . You may use this result without proof.

Hint 2: A result from the second midterm test may be helpful in part b).

(10 marks)



647B51F0-7540-44DA-B8E4-7EB1D5A7B412

mat195-exam-2019


#181          18 of 26



- 9) The motion of a particle is given by:  $\vec{r}(t) = t\hat{i} + t^2\hat{j} + \ln(1+t)\hat{k}$ , for  $t > 0$ . Determine the tangential and normal components of acceleration of this particle, and the curvature of its path at time  $t = 1$ .

Hint: There are several ways of solving this problem without finding the derivative of the unit tangent vector.

(12 marks)



938BF7C0-568B-4B2B-BC2D-B80418137067

mat195-exam-2019

#181      20 of 26



- 10) a) Find the unit direction vector(s)  $\hat{u}$  for which the directional derivative for  $f(x, y) = ye^{-xy}$  at the point  $(0, 2)$  has value 1. That is, find  $\hat{u}$  so that  $D_{\hat{u}}f(0, 2) = 1$ .

(6 marks)

- b) Are there any points on the hyperboloid  $x^2 + 4y^2 - z^2 = 4$  where the tangent plane is parallel to the plane  $2x + 2y + z = 1$ ? If so, find them.

(5 marks)



DB49B59F-D928-4A1F-B1F9-D1A05EAB648


mat195-exam-2019

#181      22 of 26



- 11) Find the maximum value of  $f(x, y) = xy - x^3y^2$  on the square  $0 \leq x \leq 1$ ,  $0 \leq y \leq 1$ . Provide a sketch of the region, and identify and show the locations of all critical points.

(10 marks)



1E9150AF-D83F-47C5-B91A-BC762F7A39E0

mat195-exam-2019


#181      24 of 26





- 12) Use Lagrange multipliers to find the extreme values of the function  $f(x, y, z) = xz + xy$  subject to the two constraints:  $x^2 + z^2 = 2$  and  $xy = 1$ .

(8 marks)



45A565C5-8F65-4C5C-88AF-AC758E7EA1B6

mat195-exam-2019

#181      26 of 26