



UNIVERSITY OF TORONTO
Faculty of Applied Science and Engineering

AER210 Midterm Test #1
Vector Calculus & Fluid Mechanics

Instructor: Alex Bercik

October 19, 2023

First name (please write as legibly as possible within the boxes)

S	O	L	U	T	I	O	N	S							
---	---	---	---	---	---	---	---	---	--	--	--	--	--	--	--

Last name

Y															
---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Student ID number

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

This exam contains 20 pages (including this cover page and a 2-page formula sheet at the back) and 8 questions. The total number of points is 65.

This exam will be scanned and graded using Crowdmark. Therefore, please write legibly, within the margins, and make a note to the grader if you need to continue your work on a separate page (otherwise, your work will not be seen by the grader). Good luck!

Distribution of Marks

Question:	1	2	3	4	5	6	7	8	Total
Points:	10	15	6	6	4	5	11	8	65
Score:									



5E9B45D7-9405-4020-90EE-A29DB6DB2E1B

midterm-1-b1d18

#235 Page 2 of 20

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.



1. (10 points) True or False. Circle the following statements that are true. You may circle only the letter before the statement, rather than the entire statement. Remember, in order for a statement to be true, it must *always* be true.

- (a) In a conservative force vector field, the work done by the force in moving an object from any point a to point b is the same regardless of the path taken.

TRUE: THM in lecture 13

- (b) Let $f(x, y) : \Omega \rightarrow \mathbb{R}$ be some function defined on a rectangular domain $\Omega = [a, b] \times [c, d]$. Then

$$\int_{\Omega} f(x, y) d\Omega = \int_c^d \left(\int_a^b f(x, y) dx \right) dy = \int_a^b \left(\int_c^d f(x, y) dy \right) dx .$$

False: No guarantee Fubini's THM holds/applies! e.g. is F integrable?

- (c) For two vector fields \mathbf{F} and \mathbf{G} , if $\nabla \times \mathbf{F} = \nabla \times \mathbf{G}$, then $\mathbf{F} = \mathbf{G}$.

False: Curl is not unique. e.g. $\vec{\nabla} \times (\vec{F} + \vec{\nabla} \phi) = \vec{\nabla} \times \vec{F}$

- (d) If f and $|f|$ are both Riemann integrable functions on Ω , then $\left| \int_{\Omega} f d\Omega \right| \leq \int_{\Omega} |f| d\Omega$.

TRUE: THM in lecture 3 (Monotonicity)

- (e) If the closed line integral $\oint_{\gamma} \mathbf{F} \cdot d\ell$ of a vector field \mathbf{F} around a specific path γ is found to be zero, then \mathbf{F} is conservative.

False: Must be zero for all paths

- (f) Given a vector field $\mathbf{F} : \Omega \rightarrow \mathbb{R}^n$ defined on a domain $\Omega \subseteq \mathbb{R}^n$, if there exists a C^1 scalar field $\phi : \Omega \rightarrow \mathbb{R}$ such that $\mathbf{F} = \nabla \phi$, then \mathbf{F} is conservative.

TRUE: Definition of conservative, lecture 13

- (g) Let $f(x, y) : \Omega \rightarrow \mathbb{R}$ be a bounded, continuous, but not differentiable function defined on the unit circle $\Omega = \{x^2 + y^2 \leq 1\}$. Any Riemann sum of f with respect to partition P must converge to the same finite value in the limit $\|P\| \rightarrow 0$. *(defn of integral)*

TRUE: THM in lecture 3: bounded & $C^0 \Rightarrow$ integrable \Rightarrow all Riemann sums converge

- (h) Let $f(x, y, z)$ be some integrable scalar field, and γ be some path in \mathbb{R}^3 with its reverse path denoted $-\gamma$. Then $\int_{\gamma} f(x, y, z) ds = - \int_{-\gamma} f(x, y, z) ds$.

False: $\int_{\gamma} f ds = \int_{-\gamma} f ds$. Standard result stated many times in lecture.

- (i) The third-order Taylor polynomial of a function $f(x, y) : \mathbb{R}^2 \rightarrow \mathbb{R}$ about a point (a, b) is unique.

TRUE: Taylor polynomials are unique! Stressed in lecture 9, on P.S., etc.

- (j) The following transformation $T : (r, \theta) \rightarrow (x, y)$ is not a diffeomorphism.

$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \end{aligned} \quad \text{where } T : [0, \infty) \times [0, 2\pi] \rightarrow \mathbb{R}^2$$

\approx bijective
differentiable
invertible

TRUE: not bijective! should be $T : (0, \infty) \times [0, 2\pi] \rightarrow \mathbb{R}^2$

importance of bounds stressed in lecture 11, on P.S.



4EB9F077-4923-4799-BBE4-69E02B3BC86F

midterm-1-b1d18

#235 Page 4 of 20

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.



2. Determine the line integrals $\int_{\gamma} \mathbf{F} \cdot d\ell$ for the given vector fields \mathbf{F} and paths γ .

- (a) (5 points) $\mathbf{F} = zy \hat{x} + x \hat{y} + z^2x \hat{z}$ along the straight line between $(0, 0, 0)$ and $(2, 3, 2)$.
- (b) (5 points) $\mathbf{F} = (2xy + \cos(x)) \hat{x} + x^2 \hat{y}$ along the line $\{(x, y) : x^2 + y^2 = 4\}$ in the counterclockwise direction, starting point $a = (2, 0)$ and finishing at point $b = (0, 2)$.
- (c) (5 points) $\mathbf{F} = (2y + \cos(y)) \hat{x} + (x + e^{\sqrt{y}}) \hat{y}$ around the region enclosed by $y = x^2$ and $x = y^2$ in the counterclockwise direction.

$\frac{\partial F_x}{\partial y} = z \neq \frac{\partial F_y}{\partial x} = 1$
 Note: \vec{F} not conservative \rightarrow No Gr THM
 & γ not closed \rightarrow No Stol THM

a) let $\vec{r}(t) = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix}t$, $t \in [0, 1]$

$$\Rightarrow \vec{r}'(t) = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix}, \quad \vec{F}(\vec{r}(t)) = \begin{pmatrix} zy \\ x \\ z^2x \end{pmatrix} = \begin{pmatrix} 6t^2 \\ 2t \\ 8t^3 \end{pmatrix}$$

$$\begin{aligned} \Rightarrow \int_{\gamma} \vec{F} \cdot d\vec{\ell} &= \int_0^1 \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt \\ &= \int_0^1 \begin{pmatrix} 6t^2 \\ 2t \\ 8t^3 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix} dt = \int_0^1 12t^2 + 6t + 16t^3 dt \\ &= 4t^3 + 3t^2 + 4t^4 \Big|_0^1 = 4+3+4 = \boxed{11} \end{aligned}$$

b) $\frac{\partial F_x}{\partial y} = 2x$, $\frac{\partial F_y}{\partial x} = 2x$, domain Ω of \vec{F} is \mathbb{R}^2
 \rightarrow simply connected

$$\frac{\partial F_x}{\partial y} = \frac{\partial F_y}{\partial x} \quad \& \quad \Omega \text{ simply connected} \quad \stackrel{\text{THM}}{\Rightarrow} \quad \vec{F} \text{ conservative}$$

Find the potential Ψ s.t. $\vec{F} = \nabla \Psi$

Continued on next page...



This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.

b) (cont'd) $\frac{\partial \varphi}{\partial x} = F_x, \quad \frac{\partial \varphi}{\partial y} = F_y$

$$\varphi = \int F_y dy = \int x^2 dy = x^2 y + g(x)$$

$$\frac{\partial \varphi}{\partial x} = 2xy + g'(x) = F_x = 2xy + \cos(x) \Rightarrow g'(x) = \cos(x)$$

$$\Rightarrow g(x) = \sin(x)$$


$$\therefore \varphi(x, y) = x^2 y + \sin(x)$$

Since \vec{F} is conservative, can use gradient THM

$$\int_{\gamma} \vec{F} \cdot d\vec{l} = \varphi(\vec{b}) - \varphi(\vec{a}) = \varphi\left(\begin{pmatrix} 0 \\ 2 \end{pmatrix}\right) - \varphi\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}\right) = (0 + \sin(0)) - (0 + \sin(2))$$

$$= -\sin(2)$$

c) $\frac{\partial F_x}{\partial y} = 2 \quad \frac{\partial F_y}{\partial x} = 1 \quad \Rightarrow$ Not conservative, but closed!
i.e. Can use Green's THM



- ✓ regular region (closed, bounded, no isolated pts)
- ✓ piecewise-smooth boundary
- ✓ $\vec{F}: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is C^1 (at least on $y \geq 0$)
- ✓ positive orientation (CCW)

$$\oint \vec{F} \cdot d\vec{l}$$

$$\oint_L \left(\frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right) dx = \int_0^1 \int_{x^2}^{\sqrt{x}} -1 dy dx = + \int_0^1 x^2 - \sqrt{x} dx = \left[\frac{x^3}{3} - \frac{2}{3} x^{3/2} \right]_0^1 = +\frac{1}{3} - \frac{2}{3} = \boxed{-\frac{1}{3}}$$




3. (6 points) Consider the triple integral $\int_0^4 \int_0^{4-z} \int_0^{\sqrt{y}} f(x, y, z) dx dy dz$.

(a) Rewrite the integral in the form $\int_?^? \int_?^? \int_?^? f(x, y, z) dz dy dx$ with the proper bounds.

(b) Rewrite the integral in the form $\int_?^? \int_?^? \int_?^? f(x, y, z) dy dx dz$ with the proper bounds.

$$\int_0^4 \int_0^{4-z} \int_0^{\sqrt{y}} \Rightarrow \begin{aligned} 0 &\leq z \leq 4 \\ 0 &\leq y \leq 4-z \\ 0 &\leq x \leq \sqrt{y} \\ \Rightarrow 0 &\leq x^2 \leq y \end{aligned}$$




a) $0 \leq z \leq 4-y$

$x^2 \leq y \leq 4$

$0 \leq x \leq 2$

$$\int_0^2 \int_{x^2}^4 \int_0^{4-y} f(x, y, z) dz dy dx$$




b) $x^2 \leq y \leq 4-z$

$0 \leq x \leq \sqrt{4-z}$

$0 \leq z \leq 4$

$$\int_0^4 \int_0^{\sqrt{4-z}} \int_{x^2}^{4-z} f(x, y, z) dy dx dz$$





BOF4BD60-B273-4737-B0B1-DFE12B4ACCB9

midterm-1-b1d18

#235 Page 8 of 20

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.



4. (a) (3 points) Consider a vector field $\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with magnitude $f(\mathbf{x}) = |\mathbf{F}(\mathbf{x})|$, and a path γ with parametrization $\mathbf{r}(t) : I \subset \mathbb{R} \rightarrow \mathbb{R}^n$. Show that if \mathbf{F} is everywhere aligned with γ , i.e. it points in the same direction as tangent vectors of $\mathbf{r}(t)$ everywhere along the path, then the line integral of \mathbf{F} reduces to the line integral of $f(\mathbf{x})$. That is,

$$\int_{\gamma} \mathbf{F} \cdot d\ell = \int_{\gamma} f(\mathbf{x}) ds$$

- (b) (3 points) Consider \mathbb{R}^2 . Show that if $\mathbf{F} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ points purely in the azimuthal direction, i.e. $\mathbf{F} = f(x, y)\hat{\theta}$, and we take the line integral along a circle of radius R in the counterclockwise direction from $\theta = \theta_i$ to $\theta = \theta_f$ then the line integral of \mathbf{F} reduces to

$$\int_{\gamma} \mathbf{F} \cdot d\ell = R \int_{\theta_i}^{\theta_f} f(x(r, \theta), y(r, \theta)) d\theta$$

a) (done in class, lecture 12, and Review Demo session)

$$\begin{aligned}
 \vec{F} \text{ aligned with } \gamma &\Rightarrow \vec{F}(\vec{r}(t)) = F(\vec{r}(t)) \hat{T}(\vec{r}(t)) \\
 &= F(\vec{r}(t)) \frac{\vec{r}'(t)}{\|\vec{r}'(t)\|} \quad \text{tangent vector to } \gamma \\
 \Rightarrow \int_{\gamma} \vec{F} \cdot d\vec{\ell} &= \int_{t_i}^{t_f} \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt \\
 &= \int_{t_i}^{t_f} F(\vec{r}(t)) \frac{\vec{r}'(t) \cdot \vec{r}'(t)}{\|\vec{r}'(t)\|} dt \\
 &= \int_{t_i}^{t_f} F(\vec{r}(t)) \frac{\|\vec{r}'(t)\|^2}{\|\vec{r}'(t)\|} dt \\
 &= \int_{t_i}^{t_f} F(\vec{r}(t)) \underbrace{\|\vec{r}'(t)\|}_{ds} dt = \int_{\gamma} F(\vec{x}) ds \quad \square
 \end{aligned}$$

b) on next page



This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.

b) Posted as note on Piazza (@26)

take polar coordinates $\Rightarrow \vec{F} = F\hat{\theta}$, $d\vec{r} = dr\hat{r} + r d\theta \hat{\theta}$
 $\Rightarrow \vec{F} \cdot d\vec{r} = Fr d\theta$ since $\hat{\theta} \cdot \hat{r} = 0$

$$\begin{aligned}\therefore \int_{\gamma} \vec{F} \cdot d\vec{r} &= \int_{\gamma} F(r) r(\theta) d\theta = \int_{\theta_i}^{\theta_f} F(R) R d\theta \quad \text{since } r=R \text{ along } \gamma \\ &= R \int_{\theta_i}^{\theta_f} F(x(R, \theta), y(R, \theta)) d\theta \quad \square\end{aligned}$$

OR

Compute directly using $\hat{\theta} = -\sin\theta \hat{x} + \cos\theta \hat{y}$

$$\Rightarrow \vec{F} = \begin{bmatrix} -F\sin\theta \\ F\cos\theta \end{bmatrix}, \quad \vec{r}(\theta) = \begin{bmatrix} R\cos\theta \\ R\sin\theta \end{bmatrix}, \quad \vec{r}'(\theta) = \begin{bmatrix} -R\sin\theta \\ R\cos\theta \end{bmatrix}$$
$$\begin{aligned}\therefore \int_{\gamma} \vec{F} \cdot d\vec{r} &= \int_{\theta_i}^{\theta_f} \vec{F}(\vec{r}(\theta)) \cdot \vec{r}'(\theta) d\theta = \int_{\theta_i}^{\theta_f} \begin{bmatrix} -F\sin\theta \\ F\cos\theta \end{bmatrix} \cdot \begin{bmatrix} -R\sin\theta \\ R\cos\theta \end{bmatrix} d\theta \\ &= \int_{\theta_i}^{\theta_f} RF(\sin^2\theta + \cos^2\theta) d\theta, \quad \sin^2\theta + \cos^2\theta = 1 \\ &= R \int_{\theta_i}^{\theta_f} F(x(R, \theta), y(R, \theta)) d\theta \quad \square\end{aligned}$$



5. (4 points) Given the vector field $\mathbf{F} = y \hat{x} - x \hat{y}$, show that the line integral around a piecewise smooth, simple, closed, and bounded curve γ in the x - y plane in the clockwise direction satisfies

$$\oint_{\gamma} \mathbf{F} \cdot d\ell = 2A$$

where A is the area enclosed by the curve γ .

- $\vec{\mathbf{F}}$ is C^1 on \mathbb{R}^2
 - Simple, closed, bounded $\gamma \Rightarrow$ regular region Ω
 - piecewise-smooth boundary $\gamma = \partial\Omega$
- \Rightarrow Green's THM applies

$$\frac{\partial F_x}{\partial y} = 1, \quad \frac{\partial F_y}{\partial x} = -1, \quad \text{Note: CW direction is negative orientation!}$$

$$\begin{aligned} \therefore \oint_{\gamma} \vec{\mathbf{F}} \cdot d\ell &= - \int_{\Omega} \left(\frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right) d\Omega \\ &= - \int_{\Omega} -2 \, d\Omega \\ &= 2 \int_{\Omega} d\Omega \\ &= 2A \quad \text{since } A = \int_{\Omega} d\Omega \quad \square \end{aligned}$$



ADA756AB-E228-4722-84E9-1EA80C4D9F28

midterm-1-b1d18

#235 Page 12 of 20

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.



6. (5 points) Let $\mathbf{F} : \Omega \rightarrow \mathbb{R}^2$ and $\mathbf{G} : \Omega \rightarrow \mathbb{R}^2$ be vector fields over a domain $\Omega \subset \mathbb{R}^2$. Prove or disprove (e.g. find a counterexample to) the following statement.

If $\mathbf{F} + \mathbf{G}$ is conservative, then \mathbf{F} and \mathbf{G} are also conservative.

(similar to HW problems Lec 13)

The statement is False.

Proof by counter-example:

$$\text{let } \vec{F} = y\hat{x} - x\hat{y}$$

$$\vec{G} = -y\hat{x} + x\hat{y} + \nabla \psi \text{ for some } C^1 \psi \text{ on } \Omega$$

by previous Q5, \vec{F} and \vec{G} are not conservative

$$\left[\frac{\partial F_x}{\partial y} = 1 \neq -1 = \frac{\partial F_y}{\partial x}, \quad \frac{\partial G_{xc}}{\partial y} = -1 + \frac{\partial^2 \psi}{\partial x \partial y} \neq 1 + \frac{\partial^2 \psi}{\partial y \partial x} = \frac{\partial G_y}{\partial x} \right]$$

$$\begin{aligned} \text{but } \vec{F} + \vec{G} &= (y-y)\hat{x} + (-x+x)\hat{y} + \nabla \psi \\ &= \nabla \psi \end{aligned}$$

i.e. $\vec{F} + \vec{G}$ is conservative by definition \square

i.e. we showed,

$(\vec{F} + \vec{G} \text{ is conservative, but neither } \vec{F} \text{ nor } \vec{G} \text{ are conservative})$



FD77125A-1B50-4601-AE3E-63F01B35E5F4

midterm-1-b1d18

#235 Page 14 of 20

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.




7. Consider the volume Ω in \mathbb{R}^3 bounded by $y = x^2 + z^2$ and the plane $y = 4$.

(a) (5 points) Evaluate the integral $\int_{\Omega} \sqrt{x^2 + z^2} d\Omega$.

(b) (6 points) Compute the total surface area $\Gamma = \partial\Omega$ of the volume Ω

a) Without loss of generality, swap y & z

$$\text{i.e. } z = x^2 + y^2, \quad z = 4, \quad \underbrace{\int_{\Omega} \sqrt{x^2 + y^2} d\Omega}$$



$= \int_{\Omega} r d\Omega$ cylindrical coordinates!

$$\therefore \int_{\Omega} \sqrt{x^2 + y^2} d\Omega = \int_0^4 \int_0^{2\pi} \int_0^{\sqrt{z}} r \cdot r dr d\theta dz$$

$$= 2\pi \int_0^4 \frac{1}{3} r^3 \Big|_0^{\sqrt{z}} dz$$

$$= \frac{2\pi}{3} \int_0^4 z^{3/2} dz = \frac{2\pi}{3} \cdot \frac{2}{5} z^{5/2} \Big|_0^4$$


$$2, 4, 8, 16, 32, 64, 128$$

$$= \frac{4\pi}{15} 2^5 = \boxed{\frac{128\pi}{15}}$$

b) on next page

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.

b) Surface Area = $\int_{\text{paraboloid}} + \int_{\text{top cap}}$



$$\begin{aligned} \int_{\text{paraboloid}} &= \int_{\Omega} \sqrt{\left(\frac{\partial F}{\partial x}\right)^2 + \left(\frac{\partial F}{\partial y}\right)^2 + 1} d\Omega, \quad F(x, y) = z = x^2 + y^2 \\ &= \int_{\Omega} \sqrt{(2x)^2 + (2y)^2 + 1} d\Omega = \int_{\Omega} \sqrt{4(x^2 + y^2) + 1} d\Omega \\ &= \int_{\Omega} \sqrt{4r^2 + 1} d\Omega \quad \text{use polar coordinates! } r^2 = x^2 + y^2 \\ &= \int_0^{2\pi} \int_0^2 \sqrt{4r^2 + 1} r dr d\theta \quad (x, y) \rightarrow (r, \theta) \\ &= 2\pi \int_1^{17} \sqrt{u} \cdot \frac{1}{8} du \quad \text{let } u = 4r^2 + 1 \quad r=0 \Rightarrow u=1 \\ &\qquad \qquad \qquad du = 8r dr \quad r=2 \Rightarrow u=17 \\ &= \frac{\pi}{4} \cdot \frac{2}{3} u^{3/2} \Big|_1^{17} \\ &= \boxed{\frac{\pi}{6} (17^{3/2} - 1)} \approx 5.76 \end{aligned}$$

$$\int_{\text{top cap}} = \pi r^2 = \pi (2)^2 = \boxed{4\pi}$$

$$\therefore \text{Surface Area} = \boxed{4\pi + \frac{\pi}{6} (17^{3/2} - 1)}$$




8. (8 points) Consider a 2D plate defined in the x - y plane with mass density

$$\rho(x, y) = (x^2 + y^2) e^{|xy|} \frac{\text{kg}}{\text{m}^2},$$

where x and y have units of m. Suppose we are only interested in the region of this plate Ω found in the 1st quadrant, bounded between the curves $x^2 - y^2 = 1$, $y^2 = x^2 - 4$, $y = 0$, and


$y = \frac{1}{2x}$. Compute the total mass of the plate contained in the region Ω .

$$\underbrace{y > 0}_{y > 0}$$



$$\begin{aligned} \text{let } u &= x^2 - y^2 \\ v &= y/x \end{aligned} \rightarrow$$

$$\begin{aligned} \Rightarrow \rho(x, y) &= (x^2 + y^2) e^{|v|} \\ &= (x^2 + y^2) e^v \text{ since } v \geq 0 \end{aligned}$$



$$\vec{T}: (u, v) \rightarrow (x, y) \quad \text{and} \quad \vec{T}^{-1}: (x, y) \rightarrow (u, v)$$

$$D_{uv} \vec{T}^{-1} = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} = \begin{pmatrix} 2x & -2y \\ y & x \end{pmatrix} \rightarrow \vec{T}^{-1} \text{ is differentiable (C')} \& \text{continuous}$$

$$\det(D_u \vec{T}) = \frac{1}{\det(D_x \vec{T}^{-1})} = \frac{1}{\begin{vmatrix} 2x & -2y \\ y & x \end{vmatrix}} = \frac{1}{2x^2 + 2y^2} = \frac{1}{2(x^2 + y^2)} > 0 \text{ since } x > 0, y \geq 0$$

$\therefore \vec{T}^{-1}$ is C' , determinants > 0 , \vec{T}^{-1} bijective (^{no proof required}) \Rightarrow invertible

$\Rightarrow \vec{T}$ & \vec{T}^{-1} are diffeomorphisms \Rightarrow use change of variables

$$\begin{aligned} \int_{\Omega} \rho(x, y) d\Omega &= \int_0^4 \int_1^4 (x^2 + y^2) e^v \cdot \frac{1}{2(x^2 + y^2)} du dv \\ &= \int_1^4 du \cdot \frac{1}{2} \int_0^{1/2} e^v dv = \frac{3}{2} e^v \Big|_0^{1/2} = \boxed{\frac{3}{2} (\sqrt{e} - 1)} \end{aligned}$$



2959EC20-5F81-40BF-841E-92A278036507

midterm-1-b1d18

#235 Page 18 of 20

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.



Useful Formulas

Vector Identities

$$\begin{aligned}\nabla(fg) &= f(\nabla g) + g(\nabla f) \\ \nabla(f \cdot g) &= f \times (\nabla \times g) + g \times (\nabla \times f) + (f \cdot \nabla)g + (g \cdot \nabla)f \\ \nabla(fg) &= f(\nabla \cdot g) + g \cdot (\nabla f) \\ \nabla \times (fg) &= f(\nabla \times g) - g \times (\nabla f) \\ \nabla \times (f \times g) &= (g \cdot \nabla)f - (f \cdot \nabla)g + f(\nabla \cdot g) - g(\nabla \cdot f) \\ \nabla \cdot (\nabla \times f) &= 0 \\ \nabla \times (\nabla f) &= 0, \quad \text{where } f \text{ is the scalar potential of the conservative vector field } \nabla f\end{aligned}$$

(Note: the following are *not* rigorous mathematical statements!)

Change of Variables

Let $\mathbf{T} : \tilde{\Omega} \rightarrow \Omega$ be a diffeomorphism (invertible & differentiable bijection). Then

$$\int_{\Omega} f(x) d\Omega = \int_{\tilde{\Omega} = \mathbf{T}^{-1}(\Omega)} f(\mathbf{T}(u)) |\det(D_u \mathbf{T})| d\tilde{\Omega}$$

where $D_u \mathbf{T}$ is the Jacobian matrix of \mathbf{T} with respect to variables $u = \mathbf{T}^{-1}(x)$.

Gradient Theorem (Fundamental Theorem of Calculus for line integrals)

Let γ be a continuous curve which starts at a and ends at point b . Then

$$\int_{\gamma} (\nabla f) \cdot d\ell = f(b) - f(a)$$

Green's Theorem

Let $\Omega \subset \mathbb{R}^2$ be a regular region with boundary $\partial\Omega = \gamma$. Then

$$\oint_{\gamma} \mathbf{F} \cdot d\ell = \int_{\Omega} \left(\frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right) d\Omega$$

Cartesian Coordinates (x, y, z)

Line element: $d\ell = dx \hat{x} + dy \hat{y} + dz \hat{z}$

Volume element: $d\Omega = dx dy dz$

$$\text{Gradient: } \nabla f = \frac{\partial f}{\partial x} \hat{x} + \frac{\partial f}{\partial y} \hat{y} + \frac{\partial f}{\partial z} \hat{z}$$

$$\text{Divergence: } \nabla \cdot \mathbf{f} = \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} + \frac{\partial f_z}{\partial z}$$

$$\text{Curl: } \nabla \times \mathbf{f} = \left(\frac{\partial f_z}{\partial y} - \frac{\partial f_y}{\partial z} \right) \hat{x} + \left(\frac{\partial f_x}{\partial z} - \frac{\partial f_z}{\partial x} \right) \hat{y} + \left(\frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) \hat{z}$$

**Spherical Coordinates (r, θ, ϕ)**

$$\begin{aligned}x &= r \cos \theta \sin \phi & \hat{x} &= \cos \theta \sin \phi \hat{r} - \sin \theta \hat{\theta} + \cos \theta \cos \phi \hat{\phi} \\y &= r \sin \theta \sin \phi & \hat{y} &= \sin \theta \sin \phi \hat{r} + \cos \theta \hat{\theta} + \sin \theta \cos \phi \hat{\phi} \\z &= r \cos \phi & \hat{z} &= \cos \phi \hat{r} - \sin \phi \hat{\phi}\end{aligned}$$

$$\begin{aligned}r &= \sqrt{x^2 + y^2 + z^2} & \hat{r} &= \cos \theta \sin \phi \hat{x} + \sin \theta \sin \phi \hat{y} + \cos \phi \hat{z} \\\theta &= \tan^{-1} \left(\frac{y}{x} \right) & \hat{\theta} &= -\sin \theta \hat{x} + \cos \theta \hat{y} \\\phi &= \cos^{-1} \left(\frac{z}{r} \right) & \hat{\phi} &= \cos \theta \cos \phi \hat{x} + \sin \theta \cos \phi \hat{y} - \sin \phi \hat{z}\end{aligned}$$

Line element: $d\ell = dr \hat{r} + r \sin \phi d\theta \hat{\theta} + r d\phi \hat{\phi}$ Volume element: $d\Omega = r^2 \sin \phi dr d\theta d\phi$

Gradient: $\nabla f = \frac{\partial f}{\partial r} \hat{r} + \frac{1}{r \sin \phi} \frac{\partial f}{\partial \theta} \hat{\theta} + \frac{1}{r} \frac{\partial f}{\partial \phi} \hat{\phi}$

Divergence: $\nabla \cdot \mathbf{f} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 f_r) + \frac{1}{r \sin \phi} \frac{\partial f_\theta}{\partial \theta} + \frac{1}{r \sin \phi} \frac{\partial}{\partial \phi} (\sin \phi f_\phi)$

Curl:
$$\begin{aligned}\nabla \times \mathbf{f} = & \frac{1}{r \sin \phi} \left(\frac{\partial}{\partial \phi} (\sin \phi f_\theta) - \frac{\partial f_\phi}{\partial \theta} \right) \hat{r} \\& + \frac{1}{r} \left(\frac{\partial}{\partial r} (rf_\phi) - \frac{\partial f_r}{\partial \phi} \right) \hat{\theta} + \frac{1}{r} \left(\frac{1}{\sin \phi} \frac{\partial f_r}{\partial \theta} - \frac{\partial}{\partial r} (rf_\theta) \right) \hat{\phi}\end{aligned}$$

Cylindrical Coordinates (r, θ, z)

$$\begin{aligned}x &= r \cos \theta & \hat{x} &= \cos \theta \hat{r} - \sin \theta \hat{\theta} \\y &= r \sin \theta & \hat{y} &= \sin \theta \hat{r} + \cos \theta \hat{\theta} \\z &= z & \hat{z} &= \hat{z}\end{aligned}$$

$$\begin{aligned}r &= \sqrt{x^2 + y^2} & \hat{r} &= \cos \theta \hat{x} + \sin \theta \hat{y} \\\theta &= \tan^{-1} \left(\frac{y}{x} \right) & \hat{\theta} &= -\sin \theta \hat{x} + \cos \theta \hat{y}\end{aligned}$$

Line element: $d\ell = dr \hat{r} + r d\theta \hat{\theta} + dz \hat{z}$ Volume element: $d\Omega = r dr d\theta dz$

Gradient: $\nabla f = \frac{\partial f}{\partial r} \hat{r} + \frac{1}{r} \frac{\partial f}{\partial \theta} \hat{\theta} + \frac{\partial f}{\partial z} \hat{z}$

Divergence: $\nabla \cdot \mathbf{f} = \frac{1}{r} \frac{\partial}{\partial r} (rf_r) + \frac{1}{r} \frac{\partial f_\theta}{\partial \theta} + \frac{\partial f_z}{\partial z}$

Curl:
$$\nabla \times \mathbf{f} = \left(\frac{1}{r} \frac{\partial f_z}{\partial \theta} - \frac{\partial f_\theta}{\partial z} \right) \hat{r} + \left(\frac{\partial f_r}{\partial z} - \frac{\partial f_z}{\partial r} \right) \hat{\theta} + \frac{1}{r} \left(\frac{\partial}{\partial r} (rf_\theta) - \frac{\partial f_r}{\partial \theta} \right) \hat{z}$$