



**UNIVERSITY OF TORONTO**  
**Faculty of Applied Science and Engineering**

**AER210 Midterm Test #2**  
**Vector Calculus & Fluid Mechanics**

Instructor: Alex Bercik

November 30, 2023

First name (please write as legibly as possible within the boxes)

S	O	L	U	T	I	O	N	S									
---	---	---	---	---	---	---	---	---	--	--	--	--	--	--	--	--	--

Last name

B	E	R	C	I	K											
---	---	---	---	---	---	--	--	--	--	--	--	--	--	--	--	--

Student ID number

:	)															
---	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

This exam contains 22 pages (including this cover page and a 3-page formula sheet at the back) and 8 questions. You may remove the formula sheet. The total number of points is 50.

This exam will be scanned and graded using Crowdmark. Therefore, please write legibly, within the margins, and **make a note to the grader if you need to continue your work on a separate page** (otherwise, your work will not be seen by the grader). Good luck!

**Distribution of Marks**

Question:	1	2	3	4	5	6	7	8	Total
Points:	6	6	3	4	6	8	12	5	50
Score:									



C5F3F330-FA93-4A69-BABE-D1103F918ACD

midterm-2-42ddd

#47 Page 2 of 22

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.



1. Provide short answers to the following questions.

- (a) (1 point) What is a Newtonian fluid?
- (b) (1 point) What determines whether the continuum approach is valid or not?
- (c) (1 point) Find the dimensions of dynamic viscosity in an MLT system.
- (d) (1 point) Two spheres of equal diameter are submerged in water. One is made of steel, while the other is made of wood. Assuming the ratio of densities is  $\rho_{\text{steel}} = 12\rho_{\text{wood}}$ , what is the ratio of the buoyant forces  $F_{\text{steel}}$  and  $F_{\text{wood}}$ ?
- (e) (2 points) How does the dynamic viscosity of air change with temperature, and why?

a) A fluid where shear stress is proportional to deformation rate OR normal velocity gradient  
i.e.  $\tau = \mu \frac{d\gamma}{dt}$  or  $\tau_{yx} = \mu \frac{\partial u}{\partial y}$

b) Continuum Approach is valid when ratio of microscopic length scale (mean free path) to macroscopic length scale is  $\ll 1$ .  
i.e. Knudsen number  $K_n = \frac{\lambda}{L} \ll 1$

$$c) [\mu] = [\tau] / \left[ \frac{\partial u}{\partial y} \right] = \left[ \frac{\text{Force}}{\text{Area}} \right] / \left[ \frac{\text{velocity}}{\text{length}} \right] = \frac{M}{TL}$$

d)  $\vec{F}_B = W_{\text{fluid displaced}} = \rho_{\text{fluid}} V_{\text{object}}$ . Same radius  $\Rightarrow$  Same volume  
Same  $V_{\text{object}} \Rightarrow$  Same  $F_B \Rightarrow F_{\text{steel}} = F_{\text{wood}}$

e)  $\mu$  increases with temperature due to increased particle collisions (1 pt for relationship, 1 pt for reason)



2F8A5209-C823-411A-87BD-0535465E3264

midterm-2-42ddd

#47 Page 4 of 22

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.



- Page 5



77B14AA8-0ACF-4B1B-A26D-401CE50779FA

midterm-2-42ddd

#47 Page 6 of 22

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.




3. (3 points) A manometer containing water with density  $\rho_w$  and mercury with density  $\rho_m$  is connected to a tank containing air at an internal pressure  $p$ , the other end being open to atmospheric pressure  $p_{\text{atm}}$ . Derive an expression relating  $p$  to  $p_{\text{atm}}$ .

$$P_{\text{Air}} = P_A \quad \text{since } \gamma_{\text{air}} \ll \gamma_{\text{water}}$$

$$P_A = P_B - \rho_w g h_1$$

$$P_B = P_C - \rho_w g (h_3 - h_2)$$

$$P_C = P_{\text{atm}} + \rho_m g h_4$$



$$\Rightarrow P = P_{\text{Air}} = P_{\text{atm}} + \rho_m g h_4 - \rho_w g (h_3 - h_2) - \rho_w g h_1$$

$$\Rightarrow \boxed{P = P_{\text{atm}} + \rho_m g h_4 - \rho_w g (h_3 - h_2 + h_1)}$$



2E13661D-6995-4FD0-A534-DF59CDE56845

midterm-2-42ddd

#47 Page 8 of 22

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.





Fluid

4. (4 points) Show that for an incompressible ~~flow~~ ( $\nabla \cdot \mathbf{v} = 0$ ), assuming both the momentum field  $\rho \mathbf{v}$  and pressure field  $p$  are  $C^1$ , then

$$\int_{\Omega} \nabla p \cdot \rho \mathbf{v} \, d\Omega = \oint_{\partial\Omega} p \rho \mathbf{v} \cdot \hat{\mathbf{n}} \, d\Gamma$$

whenever  $\Omega$  is a regular region with piecewise-smooth boundary  $\partial\Omega$ .

incompressible Fluid  $\Rightarrow \vec{\nabla} \cdot \vec{v} = 0$  and  $\rho$  constant

Not a Fluids question, just a calculus problem!

product rule  
(or vector identity)

$$\nabla p \cdot \rho \vec{v} = \nabla \cdot (p \rho \vec{v}) - p \rho (\underbrace{\nabla \cdot \vec{v}}_{=0})$$

possible because all  $C^1$

$$\begin{aligned} \Rightarrow \int_{\Omega} (\nabla p \cdot \rho \vec{v}) \, d\Omega &= \int_{\Omega} \nabla \cdot (p \rho \vec{v}) \, d\Omega \\ &= \oint_{\partial\Omega} p \rho \vec{v} \cdot \hat{\mathbf{n}} \, d\Gamma \end{aligned}$$

by Divergence THM.

We can apply it since  $p \rho \vec{v}$  is  $C^1$  and  $\Omega$  is a regular region w/ piecewise-smooth bdy.



07504852-5076-4787-B4F5-205F125265A0


midterm-2-42ddd

#47 Page 10 of 22

This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.



5. (6 points) The concrete seawall in the figure below has a parabolic surface given by  $z = \frac{x^2}{2}$ , a width (into the page) of  $w = 2\text{m}$ , and restrains seawater ( $\rho = 1000 \frac{\text{kg}}{\text{m}^3}$ ) at a depth of  $8\text{m}$ . Determine the moment with respect to the axis through the point A acting on the seawall due to the seawater. Use  $g = -9.8 \frac{\text{m}}{\text{s}^2} \hat{z}$ .



let O be the origin @ base of seawall

parametrize surface  $\vec{r}_0 = \begin{pmatrix} x \\ y \\ \frac{x^2}{2} \end{pmatrix}$   
 ↑  
 from origin O to surface


Surface vectors  $\frac{\partial \vec{r}_0}{\partial x} = \begin{pmatrix} 1 \\ 0 \\ x \end{pmatrix}$  &  $\frac{\partial \vec{r}_0}{\partial y} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$

⇒ scaled normal  $\vec{n} = \frac{\partial \vec{r}_0}{\partial x} \times \frac{\partial \vec{r}_0}{\partial y} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ 1 & 0 & x \\ 0 & 1 & 0 \end{vmatrix} = \begin{pmatrix} -x \\ 0 \\ 1 \end{pmatrix}$

Check direction: @  $x=0$   $\vec{n} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \hat{z}$  ✓ points outward to surface

distance from axis to surface  $\vec{r}_A = \begin{pmatrix} x-6 \\ 0 \\ \frac{x^2}{2} \end{pmatrix}$

∴  $M_y = \left[ \int_{\Gamma} \vec{r}_A \times d\vec{F} \right]_y = \left[ \int_{\Gamma} (\vec{r}_A \times -p\hat{n}) d\Gamma \right]_y$  ← can easily verify only y-comp. non-zero



$$z = \frac{x^2}{2} = 8$$

when  $x^2 = 4$

⇒  $x = 4$  →

$$= \int_{\Gamma} \left[ \begin{pmatrix} x-6 \\ 0 \\ \frac{x^2}{2} \end{pmatrix} \times \begin{pmatrix} x \\ 0 \\ -1 \end{pmatrix} \right]_y (p g (8-z)) dx dy$$

$$= \int_0^w dy \int_0^4 \left[ \left( \frac{x^2}{2} \right)(x) - (x-6)(-1) \right] p g \left( 8 - \frac{x^2}{2} \right) dx$$

$$= w p g \int_0^4 \left[ 8 \frac{x^3}{2} + 8x - 48 - \frac{x^5}{4} - \frac{x^3}{2} + 3x^2 \right] dx$$

$$= w p g \left[ 8 \frac{x^4}{8} + 8 \frac{x^2}{2} - 48x - \frac{x^6}{24} - \frac{x^4}{8} + 3 \frac{x^3}{3} \right]_0^4$$

$$= w p g (-32/3)$$




This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.

$$\Rightarrow \vec{M} = M_y \hat{y} = \rho g (-32/3) = (2)(1000)(9.8) (-32/3)$$

$$= -209066.6 \hat{y} \text{ N}\cdot\text{m} \quad (\text{or } \text{kg} \cdot \frac{\text{m}^2}{\text{s}^2})$$

Alternatively, use graphical approach. BUT Not very easy!



$$F^L = (\frac{1}{2} \rho g 8)(8)(w) = 32 \rho g w \quad \text{by pressure prism}$$

$$W = -\rho g w A = -\rho g w \int_0^4 (8 - \frac{x^2}{2}) dx = -\rho g w (32 - \frac{4^3}{3})$$

$$= -\rho g w \frac{64}{3}$$


$$\text{then } F_x^R = -F^L, \quad F_z^R = -W$$

to balance moments,  $F_x^R$  must act at same height as  $F^L$

$F_z^R$  must act at same  $x$  as  $W$ ,  $x_{CG}$

i.e. moments acting on wall will not be same as applying  $F^R$  at cp

$F^R$  acts at point  $x$  with normal  $\vec{n} = \begin{pmatrix} -x \\ 0 \\ 1 \end{pmatrix}$  (scaled) ← from integration approach



by similar triangles,  $\tan(\theta) = \frac{1}{x} = \frac{W}{F^L}$

$$\Rightarrow x = \frac{F^L}{W} = \frac{32 \rho g w}{64/3 \rho g w} = \frac{3}{2}$$

$$\therefore \vec{M} = \vec{r} \times F^R = (z F_x^R - x F_z^R) \hat{y} \quad \text{with } F_z^R = W$$

$$= \rho g w \left( \left( \frac{8}{3} \right) (32) - \left( \frac{3}{2} - 6 \right) \left( -\frac{64}{3} \right) \right) \hat{y} = -209066.6 \hat{y}$$

Same  $z$  as  $F^L$   
 $\Rightarrow$  centroid of pressure prism  
 $\Rightarrow z = h/3 = 8/3$





6. An open container in the form of a hemisphere of radius  $R$ , as depicted in the diagram below, is fully filled with a liquid of density  $\rho$  in hydrostatic equilibrium. Assume  $\mathbf{g} = -g\hat{z}$ .


- (a) (5 points) By direct integration of the pressure forces acting on the surface of the container, find the net resultant pressure force  $\mathbf{F}_p$  acting on the container.

Note: the use of any method other than the direct integration of the pressure forces will not receive full credit.

- ~~(3 points)~~ Prove that regardless of the container's shape,  $\mathbf{F}_p$  depends only on the container's volume and always points in the  $-\hat{z}$  direction.

$$\vec{F}_p = - \int_{\Gamma} p \hat{n} d\Gamma \quad \text{surface integral of pressure}$$

parametrize surface with  $\vec{r} = \begin{pmatrix} R \cos \theta \sin \varphi \\ R \sin \theta \sin \varphi \\ -R \cos \varphi \end{pmatrix}$   
 $0 \leq \theta < 2\pi, \quad 0 \leq \varphi \leq \pi/2$



note: could alternatively use  $z = R \cos \varphi$  if  $\pi/2 \leq \varphi \leq \pi$

$$\frac{\partial \vec{r}}{\partial \theta} = \begin{pmatrix} -R \sin \theta \sin \varphi \\ R \cos \theta \sin \varphi \\ 0 \end{pmatrix} \quad \frac{\partial \vec{r}}{\partial \varphi} = \begin{pmatrix} R \cos \theta \cos \varphi \\ R \sin \theta \cos \varphi \\ R \sin \varphi \end{pmatrix} \quad \frac{\partial \vec{r}}{\partial \theta} \times \frac{\partial \vec{r}}{\partial \varphi} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ -R \sin \theta \sin \varphi & R \cos \theta \sin \varphi & 0 \\ R \cos \theta \cos \varphi & R \sin \theta \cos \varphi & R \sin \varphi \end{vmatrix}$$

and  $p(z) = \rho g(z) = \rho g(R \cos \varphi)$  \* ignore  $P_{atm}$  or use  $P_{atm} = 0$

$$\therefore \vec{F}_p = - \int \rho g R \cos \varphi \hat{n} d\Gamma$$

$$= - \int_0^{\pi/2} \int_0^{2\pi} \rho g R \cos \varphi \begin{pmatrix} \cos \theta \sin \varphi \\ \sin \theta \sin \varphi \\ + \cos \varphi \end{pmatrix} R^2 \sin \varphi d\theta d\varphi$$

do each component:

$$F_p^x = + \rho g R^3 \underbrace{\int_0^{\pi/2} \cos \varphi \sin^2 \varphi d\varphi}_{=1/3} \underbrace{\int_0^{2\pi} \cos \theta d\theta}_{=0} = 0$$

$$F_p^y = + \rho g R^3 \underbrace{\int_0^{\pi/2} \cos \varphi \sin^2 \varphi d\varphi}_{=1/3} \underbrace{\int_0^{2\pi} \sin \theta d\theta}_{=0} = 0$$

$$= R^2 \sin \varphi \begin{pmatrix} \cos \theta \sin \varphi \\ \sin \theta \sin \varphi \\ -\cos \varphi \end{pmatrix}$$

consistent with what we know about spherical coordinates!

check sign:  $\hat{n} = \begin{pmatrix} \cos \theta \sin \varphi \\ \sin \theta \sin \varphi \\ -\cos \varphi \end{pmatrix}$

@  $\varphi = \pi \quad \hat{n} = \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}$  -ve of what we want!

take  $-\hat{n}$

conf'd...



This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.

and

$$F_p^z = -\rho g R^3 \underbrace{\int_0^{\pi/2} \cos^2 \varphi \sin \varphi d\varphi}_{=1/3} \underbrace{\int_0^{2\pi} d\theta}_{=2\pi} = -\frac{2}{3}\pi \rho g R^3$$

$$\therefore \vec{F}_p = -\frac{2}{3}\pi \rho g R^3 \hat{z} \quad \left( -\pi R^2 P_{\text{atm}} \hat{z} \right) \quad \begin{matrix} \swarrow \text{if used } P_{\text{atm}} \neq 0 \\ \frac{1}{2} \cdot \frac{4}{3}\pi R^3 \end{matrix}$$

Note: consistent with idea that Force = Weight =  $\frac{1}{2} \text{Vol(sphere)} \cdot \rho g$

↑ Only  $\frac{1}{5}$  marks to be awarded if answered with this method.

If Divergence theorem is used, i.e.  $\int_{\partial V} p \hat{n} d\Gamma = \int_V \nabla p dV$   
(and then hydrostatics)

then only award  $\frac{3}{5}$  marks.

(this was the originally intended purpose of part b, but was removed for time's sake)

$$\begin{aligned} &= \int_V \rho \vec{g} dV \\ &= \rho \vec{g} \int_V dV \\ &= \rho \vec{g} \frac{2}{3}\pi R^3 \end{aligned}$$






7. A U-tube of bottom width  $\ell = 0.5\text{m}$  is filled with water such that the liquid reaches an initial height  $H = 0.25\text{m}$  in both tubes. The tube is then rotated with a constant angular velocity  $\omega$  about the offset vertical axis shown in the figure,  $0.1\text{m}$  away from the inner tube. The water in the tubes eventually reach different heights  $h_1$  and  $h_2$  in the inner and outer tubes, respectively.

(a) (7 points) Assuming that no spillage occurs, find the angular velocity  $\omega$  required to make the fluid in the inner tube drop to a height of  $h_1 = 0\text{m}$ .

Hint: The magnitude of centripetal acceleration is  $r\omega^2$ . Find an expression for the free surface, then use a conservation of mass argument relating  $h_1$  and  $h_2$  to  $H = 0.25$  to eliminate any unknown constants. Use  $g = -9.81\hat{z}$ .

(b) (5 points) Suppose the relevant variables for this problem are  $g, \omega, H, \ell$ , and the distance from the inner tube to the axis,  $a$  (in this case  $a = 0.1\text{m}$ ). If our dependent variable is  $\omega$ , find appropriate Pi terms required to non-dimensionalize this problem (Note: radians are treated as dimensionless).

a) Very similar to Newton's bucket!

$$\rho \vec{a} = -\nabla p - \rho g \hat{z} \xrightarrow{\text{cylindrical coords}} \rho \begin{bmatrix} -r\omega^2 \\ 0 \\ 0 \end{bmatrix} = - \begin{bmatrix} \frac{\partial p}{\partial r} \\ \frac{1}{r} \frac{\partial p}{\partial \theta} \\ \frac{\partial p}{\partial z} \end{bmatrix} - \rho g \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$


$$\Rightarrow \frac{\partial p}{\partial r} = \rho r \omega^2, \quad \frac{\partial p}{\partial \theta} = 0, \quad \frac{\partial p}{\partial z} = -\rho g$$

integrate in  $z$ :  $p(r, z) = \int -\rho g dz = -\rho g z + f(r)$

differentiate in  $r$ :  $\frac{\partial p}{\partial r} = f'(r) = \rho r \omega^2 \Rightarrow f(r) = \frac{\rho}{2} r^2 \omega^2 + C$

$$\therefore p(r, z) = \frac{1}{2} \rho r^2 \omega^2 - \rho g z + C$$

remove one point if total differential  $dp = \frac{\partial p}{\partial r} dr + \frac{\partial p}{\partial z} dz$  is used. NOT correct, as explained on PS2

To find  $C$ , argue that  $H + H = h_1 + h_2$  (conservation of mass!)

$$\Rightarrow \text{if } z_s \text{ is s.t. } p(r, z_s) = p_{\text{atm}} = \frac{1}{2} \rho r^2 \omega^2 - \rho g z_s + C$$

$$\Rightarrow z_s = \underbrace{\frac{C - p_{\text{atm}}}{\rho g}}_{= \tilde{C} \leftarrow \text{constant!}} + \frac{1}{2} \frac{1}{g} r^2 \omega^2$$

$$h_1 = z_s(r=0.1) = \tilde{C} + \frac{1}{2} \frac{1}{g} (0.1)^2 \omega^2$$

$$h_2 = z_s(r=0.4) = \tilde{C} + \frac{1}{2} \frac{1}{g} (0.4)^2 \omega^2$$

$$\Rightarrow 2H = h_1 + h_2 = 2\tilde{C} + \frac{1}{2} \frac{1}{g} \omega^2 (0.1^2 + 0.4^2)$$

$$\Rightarrow \tilde{C} = \underbrace{H}_{= 0.25} - \frac{1}{4} \frac{1}{g} \omega^2 \underbrace{(0.1^2 + 0.4^2)}_{= 0.17}$$

Cont'd...



This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.

$$\therefore h_1 = 0 = z_s(r=0.1) = H - \frac{1}{4g}\omega^2(0.1^2 + 0.4^2) + \frac{1}{2}\frac{1}{g}\omega^2(0.1)^2$$

$$= H + \frac{1}{4}\frac{1}{g}\omega^2(0.1)^2 - \frac{1}{4g}\omega^2(0.4^2)$$

$$\Rightarrow \underbrace{H}_{0.25} = \frac{1}{4}\frac{1}{g}\omega^2(0.4^2 - 0.1^2) \Rightarrow \omega = \sqrt{\frac{4g(0.25)}{(0.4)^2 - (0.1)^2}} = 8.087 \text{ s}^{-1}$$

$$\text{general form } \omega^2 = \frac{4gH}{(l-a)^2 - a^2} = \frac{4gH}{l^2 - 2la}$$

b) Variables are  $g, \omega, H, l, a$

use  $M, L, T$  system:  $[g] = \frac{L}{T^2}$ ,  $[\omega] = \frac{1}{T}$ ,  $[H] = L$   
 $[l] = L$ ,  $[a] = L$

3 primary dimensions, but only 2 reference dimensions,  $L$  &  $T$

$\Rightarrow$  by Buckingham Pi Theorem,

$$\begin{aligned} \# \text{ of Pi terms} &= \# \text{ of variables} - \# \text{ of reference dimensions} \\ &= 5 - 2 = 3 \end{aligned}$$

Choose 2 repeating variables:  $g, l$

$\omega$  is dependent variable, thus we need  $g$  to get  $T$

(could choose  $H$  or  $a$  instead but can not choose  $\omega$  & must choose  $g$ )

Can form  $\pi$  terms with  $[\pi_1] = \cancel{M^0} L^0 T^0 = [\omega][g]^a[l]^b \rightarrow$

$$[\pi_2] = \cancel{M^0} L^0 T^0 = [H][g]^a[l]^b \rightarrow \text{solve for } a, b$$

$$[\pi_3] = \cancel{M^0} L^0 T^0 = [a][g]^a[l]^b \rightarrow$$

OR... cont'd...





This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.

Q7 cont'd:

OR From visual inspection can easily see

$$\pi_1 = \omega \sqrt{\frac{l}{g}}, \quad \pi_2 = \frac{H}{l}, \quad \pi_3 = \frac{a}{l}$$

(swap  $l$  with  $H$  &  $a$  to get all other possible combinations)

(not part of question)

Note: we can use this to write a non-dimensional form of our answer


$$\omega^2 = \frac{4gH}{l^2 - 2la} \rightarrow \omega^2 \frac{l}{g} = \frac{4 \left( \frac{H}{l} \right)}{1 - 2 \left( \frac{a}{l} \right)}$$



8. (5 points) Consider the vector field  $\mathbf{F} = x\hat{x} - y\hat{y} + 2x\hat{z}$  and the two surfaces  $\Gamma_1$  and  $\Gamma_2$  formed by the intersection of the sphere  $x^2 + y^2 + z^2 = R^2$  with the horizontal plane  $z = k$  at some height  $-R < k < R$ . The curve at which they intersect is  $\gamma$  and goes in the counterclockwise direction as looking from above. Show that the surface integral of  $\mathbf{F}$  through  $\Gamma_1$  is the opposite of that through  $\Gamma_2$ , i.e.

$$\int_{\Gamma_1} \mathbf{F} \cdot \hat{n} d\Gamma = - \int_{\Gamma_2} \mathbf{F} \cdot \hat{n} d\Gamma$$

Can solve either with Stokes' THM  
or Divergence THM



Stokes THM:  $\int_{\Gamma} (\nabla \times \vec{A}) \cdot \hat{n} d\Gamma = \oint_{\gamma=\partial\Gamma} \vec{A} \cdot d\vec{\ell}$

idea: if  $\nabla \times \vec{A} = \vec{F}$  we can apply Stokes' THM.

Does such an  $\vec{A}$  exist? Yes! Infinitely many.

e.g. let  $\vec{A} = \begin{bmatrix} 0 \\ x^2 \\ xy \end{bmatrix}$ , then  $\nabla \times \vec{A} = \begin{bmatrix} x \\ -y \\ 2x \end{bmatrix} = \vec{F} \checkmark$

$\vec{A}$  is  $C^1$ ,  $\Gamma_1$  &  $\Gamma_2$  are bounded smooth surfaces w/ boundaries  $\gamma$  &  $-\gamma$

$$\Rightarrow \int_{\Gamma_1} \vec{F} \cdot \hat{n} d\Gamma = \int_{\Gamma_1} (\nabla \times \vec{A}) \cdot \hat{n} d\Gamma = \oint_{\gamma} \vec{A} \cdot d\vec{\ell} \quad \leftarrow \text{vectors!}$$

$$\int_{\Gamma_2} \vec{F} \cdot \hat{n} d\Gamma = \int_{\Gamma_2} (\nabla \times \vec{A}) \cdot \hat{n} d\Gamma = \oint_{-\gamma} \vec{A} \cdot d\vec{\ell} = - \oint_{\gamma} \vec{A} \cdot d\vec{\ell}$$

$$\therefore \int_{\Gamma_1} \vec{F} \cdot \hat{n} d\Gamma = - \int_{\Gamma_2} \vec{F} \cdot \hat{n} d\Gamma \text{ as desired.}$$

Divergence THM:  $\int_{\Omega} (\nabla \cdot \vec{F}) d\Omega = \oint_{\partial\Omega} \vec{F} \cdot \hat{n} d\Gamma$

idea: the divergence inside the sphere is equal to the flux out of  $\Gamma_1 + \Gamma_2$   
if  $\nabla \cdot \vec{F} = 0$  then, those fluxes must be opposite

cont'd...



This page is intentionally left blank for continued work. Please make a note to the grader on the main question page if you wish them to grade any work on this page.

$$\text{check: } \nabla \cdot \vec{F} = \nabla \cdot \begin{bmatrix} x \\ -y \\ 2z \end{bmatrix} = 1 - 1 + 0 = 0 \quad \checkmark$$

$\vec{F}$  is  $C^1$ , the sphere  $\Omega$  is a regular region with smooth boundary  $\partial\Omega = \Gamma_1 + \Gamma_2$

$$\Rightarrow 0 = \int_{\Omega} \nabla \cdot \vec{F} \, d\Omega = \oint_{\partial\Omega} \vec{F} \cdot \hat{n} \, d\Gamma = \int_{\Gamma_1} \vec{F} \cdot \hat{n} \, d\Gamma + \int_{\Gamma_2} \vec{F} \cdot \hat{n} \, d\Gamma$$

$$\Rightarrow \int_{\Gamma_1} \vec{F} \cdot \hat{n} \, d\Gamma = - \int_{\Gamma_2} \vec{F} \cdot \hat{n} \, d\Gamma \quad \text{as desired.}$$

Alternative proof w/ Divergence THM:

instead pick regions  $\Omega_{\text{top}}$  = top of sphere

$\Omega_{\text{bot}}$  = bottom of sphere

$\Gamma_{\text{mid}}$  = intersection surface w/ plane

$$\begin{aligned} \Rightarrow \int_{\Omega_{\text{top}}} \nabla \cdot \vec{F} \, d\Omega &= \int_{\Gamma_1} \vec{F} \cdot \hat{n} \, d\Gamma + \int_{\Gamma_{\text{mid}}} \vec{F} \cdot \hat{n} \, d\Gamma \\ \int_{\Omega_{\text{bot}}} \nabla \cdot \vec{F} \, d\Omega &= \int_{\Gamma_2} \vec{F} \cdot \hat{n} \, d\Gamma + \int_{\Gamma_{\text{mid}}} \vec{F} \cdot (-\hat{n}) \, d\Gamma \end{aligned}$$

points down  
same  $\hat{n}$   
(different "out" direction)

$$\text{since } \nabla \cdot \vec{F} = 0 \text{ as before, } - \int_{\Gamma_{\text{mid}}} \vec{F} \cdot \hat{n} \, d\Gamma = \int_{\Gamma_1} \vec{F} \cdot \hat{n} \, d\Gamma$$

$$\int_{\Gamma_{\text{mid}}} \vec{F} \cdot \hat{n} \, d\Gamma = \int_{\Gamma_2} \vec{F} \cdot \hat{n} \, d\Gamma$$

$$\Rightarrow \int_{\Gamma_1} \vec{F} \cdot \hat{n} \, d\Gamma = - \int_{\Gamma_2} \vec{F} \cdot \hat{n} \, d\Gamma \quad \text{as desired.}$$



## Useful Formulas

### Vector Identities

$$\begin{aligned}
 \mathbf{f} \cdot (\mathbf{g} \times \mathbf{h}) &= \mathbf{g} \cdot (\mathbf{h} \times \mathbf{f}) = \mathbf{h} \cdot (\mathbf{f} \times \mathbf{g}) \\
 \mathbf{f} \times (\mathbf{g} \times \mathbf{h}) &= \mathbf{g}(\mathbf{f} \cdot \mathbf{h}) - \mathbf{h}(\mathbf{f} \cdot \mathbf{g}) \\
 \nabla(fg) &= f(\nabla g) + g(\nabla f) \\
 \nabla(\mathbf{f} \cdot \mathbf{g}) &= \mathbf{f} \times (\nabla \times \mathbf{g}) + \mathbf{g} \times (\nabla \times \mathbf{f}) + (\mathbf{f} \cdot \nabla)\mathbf{g} + (\mathbf{g} \cdot \nabla)\mathbf{f} \\
 \nabla \cdot (\mathbf{f}\mathbf{g}) &= f(\nabla \cdot \mathbf{g}) + \mathbf{g} \cdot (\nabla f) \\
 \nabla \cdot (\mathbf{f} \times \mathbf{g}) &= \mathbf{g} \cdot (\nabla \times \mathbf{f}) - \mathbf{f} \cdot (\nabla \times \mathbf{g}) \\
 \nabla \times (\mathbf{f}\mathbf{g}) &= f(\nabla \times \mathbf{g}) - \mathbf{g} \times (\nabla f) \\
 \nabla \times (\mathbf{f} \times \mathbf{g}) &= (\mathbf{g} \cdot \nabla)\mathbf{f} - (\mathbf{f} \cdot \nabla)\mathbf{g} + \mathbf{f}(\nabla \cdot \mathbf{g}) - \mathbf{g}(\nabla \cdot \mathbf{f}) \\
 \nabla \cdot (\nabla \times \mathbf{f}) &= 0 \\
 \nabla \times (\nabla f) &= 0, \quad \text{where } f \text{ is the scalar potential of the conservative vector field } \nabla f \\
 \nabla \times (\nabla \times \mathbf{f}) &= \nabla(\nabla \cdot \mathbf{f}) - \nabla^2 \mathbf{f}, \quad \text{where } \nabla^2 \mathbf{f} \text{ is the vector Laplacian}
 \end{aligned}$$

(Note: the following are *not* rigorous mathematical statements!)

### Change of Variables

Let  $\mathbf{T} : \tilde{\Omega} \rightarrow \Omega$  be a diffeomorphism (invertible & differentiable bijection). Then

$$\int_{\Omega} f(\mathbf{x}) d\Omega = \int_{\tilde{\Omega}=\mathbf{T}^{-1}(\Omega)} f(\mathbf{T}(\mathbf{u})) |\det(D_{\mathbf{u}}\mathbf{T})| d\tilde{\Omega}$$

where  $D_{\mathbf{u}}\mathbf{T}$  is the Jacobian matrix of  $\mathbf{T}$  with respect to variables  $\mathbf{u} = \mathbf{T}^{-1}(\mathbf{x})$ .

### Gradient Theorem (Fundamental Theorem of Calculus for line integrals)

Let  $\gamma$  be a continuous curve which starts at  $\mathbf{a}$  and ends at point  $\mathbf{b}$ . Then

$$\int_{\gamma} (\nabla f) \cdot d\ell = f(\mathbf{b}) - f(\mathbf{a})$$

### Divergence Theorem (Gauss' Theorem)

Let  $\Omega$  be a regular region with piecewise-smooth boundary  $\Gamma = \partial\Omega$  and outward normal  $\hat{\mathbf{n}}$ . Then

$$\int_{\Omega} \nabla \cdot \mathbf{f} d\Omega = \oint_{\partial\Omega} (\mathbf{f} \cdot \hat{\mathbf{n}}) d\Gamma$$

### Stokes' Theorem (Curl Theorem, Baby Stokes' Theorem)

Let  $\Gamma$  be a bounded piecewise-smooth surface (2D area) with closed boundary  $\partial\Gamma = \gamma$ . Then

$$\int_{\Gamma} \underbrace{(\nabla \times \mathbf{f}) \cdot \hat{\mathbf{n}}}_{(\nabla \times \vec{f}) \cdot \hat{n}} d\Gamma = \oint_{\partial\Gamma} \mathbf{f} \cdot d\ell$$



## Buckingham Pi Theorem

(# of Pi terms) = (# of variables) - (min. # of reference dimensions)

## Fluids Equations

$$\rho \mathbf{a} = -\nabla p + \rho \mathbf{g} \quad , \quad \tau_{yx} = \mu \frac{\partial v_x}{\partial y} \quad , \quad \Delta p = \rho g \Delta z$$

$$Re = \frac{\rho |v| L}{\mu} \quad , \quad Ma = \frac{|v|}{c} \quad , \quad c = \sqrt{\gamma RT} \quad , \quad p = \rho RT$$

## Useful Integrals

$$\int_0^{2\pi} \cos \theta \sin \theta \, d\theta = 0 \quad , \quad \int_0^{\pi} \cos \theta \sin \theta \, d\theta = 0 \quad , \quad \int_0^{\pi/2} \cos \theta \sin \theta \, d\theta = \frac{1}{2}$$

$$\int_0^{2\pi} \cos \theta \sin^2 \theta \, d\theta = 0 \quad , \quad \int_0^{\pi} \cos \theta \sin^2 \theta \, d\theta = 0 \quad , \quad \int_0^{\pi/2} \cos \theta \sin^2 \theta \, d\theta = \frac{1}{3}$$

$$\int_0^{2\pi} \cos^2 \theta \sin \theta \, d\theta = 0 \quad , \quad \int_0^{\pi} \cos^2 \theta \sin \theta \, d\theta = 0 \quad , \quad \int_0^{\pi/2} \cos^2 \theta \sin \theta \, d\theta = \frac{1}{3}$$

$$\int \cos^2 \theta \, d\theta = \frac{1}{2}\theta + \frac{1}{4}\sin 2\theta + C \quad , \quad \int \sin^2 \theta \, d\theta = \frac{1}{2}\theta - \frac{1}{4}\sin 2\theta + C$$

## Trigonometric Identities

$$\sin(\theta \pm \frac{\pi}{2}) = \pm \cos \theta \quad , \quad \sin(\theta + \pi) = -\sin \theta \quad , \quad \cos(\theta + \pi) = -\cos \theta$$

$$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta \quad , \quad \cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta \quad , \quad \cos(2\theta) = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta$$

## Cartesian Coordinates $(x, y, z)$

Line element:  $d\ell = dx \hat{x} + dy \hat{y} + dz \hat{z}$

Volume element:  $d\Omega = dx \, dy \, dz$

Gradient:  $\nabla f = \frac{\partial f}{\partial x} \hat{x} + \frac{\partial f}{\partial y} \hat{y} + \frac{\partial f}{\partial z} \hat{z}$

Divergence:  $\nabla \cdot \mathbf{f} = \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} + \frac{\partial f_z}{\partial z}$

Curl:  $\nabla \times \mathbf{f} = \left( \frac{\partial f_z}{\partial y} - \frac{\partial f_y}{\partial z} \right) \hat{x} + \left( \frac{\partial f_x}{\partial z} - \frac{\partial f_z}{\partial x} \right) \hat{y} + \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) \hat{z}$

Cross Product:  $\mathbf{a} \times \mathbf{b} = (a_y b_z - a_z b_y) \hat{x} + (a_z b_x - a_x b_z) \hat{y} + (a_x b_y - a_y b_x) \hat{z}$

**Spherical Coordinates**  $(r, \theta, \phi)$ 

$$x = r \cos \theta \sin \phi \quad \hat{x} = \cos \theta \sin \phi \hat{r} - \sin \theta \hat{\theta} + \cos \theta \cos \phi \hat{\phi}$$

$$y = r \sin \theta \sin \phi \quad \hat{y} = \sin \theta \sin \phi \hat{r} + \cos \theta \hat{\theta} + \sin \theta \cos \phi \hat{\phi}$$

$$z = r \cos \phi \quad \hat{z} = \cos \phi \hat{r} - \sin \phi \hat{\phi}$$

$$r = \sqrt{x^2 + y^2 + z^2} \quad \hat{r} = \cos \theta \sin \phi \hat{x} + \sin \theta \sin \phi \hat{y} + \cos \phi \hat{z}$$

$$\theta = \tan^{-1} \left( \frac{y}{x} \right) \quad \hat{\theta} = -\sin \theta \hat{x} + \cos \theta \hat{y}$$

$$\phi = \cos^{-1} \left( \frac{z}{r} \right) \quad \hat{\phi} = \cos \theta \cos \phi \hat{x} + \sin \theta \cos \phi \hat{y} - \sin \phi \hat{z}$$

$$\text{Line element: } d\ell = dr \hat{r} + r \sin \phi d\theta \hat{\theta} + r d\phi \hat{\phi}$$

$$\text{Volume element: } d\Omega = r^2 \sin \phi dr d\theta d\phi$$

$$\text{Gradient: } \nabla f = \frac{\partial f}{\partial r} \hat{r} + \frac{1}{r \sin \phi} \frac{\partial f}{\partial \theta} \hat{\theta} + \frac{1}{r} \frac{\partial f}{\partial \phi} \hat{\phi}$$

$$\text{Divergence: } \nabla \cdot \mathbf{f} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 f_r) + \frac{1}{r \sin \phi} \frac{\partial f_\theta}{\partial \theta} + \frac{1}{r \sin \phi} \frac{\partial}{\partial \phi} (\sin \phi f_\phi)$$

$$\begin{aligned} \text{Curl: } \nabla \times \mathbf{f} = & \frac{1}{r \sin \phi} \left( \frac{\partial}{\partial \phi} (\sin \phi f_\theta) - \frac{\partial f_\phi}{\partial \theta} \right) \hat{r} \\ & + \frac{1}{r} \left( \frac{\partial}{\partial r} (r f_\phi) - \frac{\partial f_r}{\partial \phi} \right) \hat{\theta} + \frac{1}{r} \left( \frac{1}{\sin \phi} \frac{\partial f_r}{\partial \theta} - \frac{\partial}{\partial r} (r f_\theta) \right) \hat{\phi} \end{aligned}$$

**Cylindrical Coordinates**  $(r, \theta, z)$ 

$$x = r \cos \theta \quad \hat{x} = \cos \theta \hat{r} - \sin \theta \hat{\theta}$$

$$y = r \sin \theta \quad \hat{y} = \sin \theta \hat{r} + \cos \theta \hat{\theta}$$

$$z = z \quad \hat{z} = \hat{z}$$

$$r = \sqrt{x^2 + y^2} \quad \hat{r} = \cos \theta \hat{x} + \sin \theta \hat{y}$$

$$\theta = \tan^{-1} \left( \frac{y}{x} \right) \quad \hat{\theta} = -\sin \theta \hat{x} + \cos \theta \hat{y}$$

$$\text{Line element: } d\ell = dr \hat{r} + r d\theta \hat{\theta} + dz \hat{z}$$

$$\text{Volume element: } d\Omega = r dr d\theta dz$$

$$\text{Gradient: } \nabla f = \frac{\partial f}{\partial r} \hat{r} + \frac{1}{r} \frac{\partial f}{\partial \theta} \hat{\theta} + \frac{\partial f}{\partial z} \hat{z}$$

$$\text{Divergence: } \nabla \cdot \mathbf{f} = \frac{1}{r} \frac{\partial}{\partial r} (r f_r) + \frac{1}{r} \frac{\partial f_\theta}{\partial \theta} + \frac{\partial f_z}{\partial z}$$

$$\text{Curl: } \nabla \times \mathbf{f} = \left( \frac{1}{r} \frac{\partial f_z}{\partial \theta} - \frac{\partial f_\theta}{\partial z} \right) \hat{r} + \left( \frac{\partial f_r}{\partial z} - \frac{\partial f_z}{\partial r} \right) \hat{\theta} + \frac{1}{r} \left( \frac{\partial}{\partial r} (r f_\theta) - \frac{\partial f_r}{\partial \theta} \right) \hat{z}$$