



DO NOT WRITE ABOVE THIS LINE

First name (please write as legibly as possible within the boxes)

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Last name

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Student ID number

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

UNIVERSITY OF TORONTO

FACULTY OF APPLIED SCIENCE AND ENGINEERING

ESC103F – Engineering Mathematics and Computation

Final Exam

December 17, 2022

Instructor – Professor W.R. Cluett



This is a closed book test. No calculators are permitted.

All six questions are of equal value.

This test contains 30 pages including the cover page 1, this information page 2, and page 30 that is for rough work only. The test is printed two-sided.

Do not tear any pages from this test.

Present complete solutions in the space provided.

Given information:

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta$$

$$\begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} \times \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} y_1 z_2 - z_1 y_2 \\ -(x_1 z_2 - z_1 x_2) \\ x_1 y_2 - y_1 x_2 \end{bmatrix}$$

$$\text{proj}_{\vec{v}} \vec{u} = \frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|^2} \vec{v}$$

The formula for the inverse of a 2x2 matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is given by:

$$\frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

The normal system of equations corresponding to $A\vec{x} = \vec{b}$ is given by:

$$(A^T A)\vec{x}_{LS} = A^T \vec{b}$$

Euler's method for solving a first order differential equation $y'(t) = f(t, y(t))$ is given by:

$$y_{n+1} = y_n + \Delta t f(t_n, y_n)$$

For higher order systems that can be expressed in the vector-matrix form $Z' = AZ$, Euler's method is given by:

$$Z_{n+1} = Z_n + \Delta t AZ_n$$



DO NOT WRITE ABOVE THIS LINE

Q1: Consider matrix A given below,

$$A = \begin{bmatrix} 2 & 4 & 6 & 4 \\ 2 & 5 & 7 & 6 \\ 2 & 3 & 5 & 2 \end{bmatrix}$$

a) Express $\begin{bmatrix} 6 \\ 7 \\ 5 \end{bmatrix}$ as a linear combination of $\begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 4 \\ 5 \\ 3 \end{bmatrix}$.

b) Express $\begin{bmatrix} 4 \\ 6 \\ 2 \end{bmatrix}$ as a linear combination of $\begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 4 \\ 5 \\ 3 \end{bmatrix}$.



52987A67-116A-41C8-89A0-81C585890B0B

final-exam-fa79c

#58 Page 4 of 30

DO NOT WRITE ABOVE THIS LINE

- c) Using Gaussian elimination, find the reduced row echelon form (R_0) of matrix A .



DO NOT WRITE ABOVE THIS LINE

- d) For the homogeneous system $A\vec{x} = \vec{0}$, find the special solution(s). (As a check on your solution, remember that $n - r$ corresponds to the number of special solutions where n is the number of unknowns and r is the rank of matrix A .)



2192B18F-2921-4065-8E06-E8FD4DB75F54

final-exam-fa79c

#58 Page 6 of 30

DO NOT WRITE ABOVE THIS LINE

- e) Factor matrix $A = CR$ by determining matrices C and R .



DO NOT WRITE ABOVE THIS LINE

Q2: Consider matrix A given below,

$$A = \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}$$

- a) By inspection, can you determine a nonzero vector \vec{x} such that $A\vec{x} = \vec{0}$?
- b) Using Gaussian elimination, determine if the inverse of matrix A (A^{-1}) exists.



224B1DB6-3B62-4582-A42B-D855B9174AF6

final-exam-fa79c

#58 Page 8 of 30

DO NOT WRITE ABOVE THIS LINE

- c) Using Gaussian elimination, find the vector solution to $A\vec{x} = \vec{0}$ and give a geometric interpretation of the solution.



DO NOT WRITE ABOVE THIS LINE

d) Note: this part is not related to parts a-c.

Suppose matrix B is square ($n \times n$) and invertible. Let us say that you replace row 2 of matrix B by row 2 plus row 1 and this produces matrix C . Is matrix C invertible? Explain your answer. If matrix C is invertible, how could you find C^{-1} from B^{-1} ? Explain your answer.



963165ED-B73C-45F6-A5B5-BB761DE60F17

final-exam-fa79c

#58 Page 10 of 30

DO NOT WRITE ABOVE THIS LINE

Q3: Consider the following three (x, y) data points that have been collected, $(-1, 7), (1, 7), (2, 21)$. We wish to fit a straight line to this data of the form $y = cx + d$.

- a) Formulate this problem in the form $A\vec{x} = \vec{b}$, where $\vec{x} = \begin{bmatrix} c \\ d \end{bmatrix}$.



DO NOT WRITE ABOVE THIS LINE

- b) Determine a scalar equation that describes the column space of matrix A .



6CB003C4-1B64-409C-9AA8-16184983668E

final-exam-fa79c

#58 Page 12 of 30

DO NOT WRITE ABOVE THIS LINE

- c) Determine if the vector $\begin{bmatrix} 7 \\ 7 \\ 21 \end{bmatrix}$ lies in the column space of matrix A .



DO NOT WRITE ABOVE THIS LINE

- d) Find the projection of the vector $\begin{bmatrix} 7 \\ 7 \\ 21 \end{bmatrix}$ in the column space of matrix A , i.e. what we will

refer to as $A\vec{x}_{LS}$, by making use of the scalar equation that describes the column space of matrix A as determined in part b.



0670957D-D57C-42EA-9EC8-097549E987CC

final-exam-fa79c

#58 Page 14 of 30

DO NOT WRITE ABOVE THIS LINE

- e) From the vector $A\vec{x}_{LS}$ found in part d, find the values for $\vec{x}_{LS} = \begin{bmatrix} c \\ d \end{bmatrix}$.



DO NOT WRITE ABOVE THIS LINE

Q4: Consider the system of linear equations,

$$x + 4y - 2z = 1$$

$$x + 7y - 6z = 6$$

$$0x + 3y + qz = t$$

The goal is to find solutions to this system.

- a) Begin by formulating and defining the problem in the form $A\vec{x} = \vec{b}$, where $\vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$.



364E1F2A-E0E6-41F9-8808-E116A45D56CF

final-exam-fa79c

#58 Page 16 of 30

DO NOT WRITE ABOVE THIS LINE

- b) Determine values for q and t that result in matrix A not being invertible.



DO NOT WRITE ABOVE THIS LINE

- c) Determine values for q and t that result in infinite solutions.



2AD07AD9-1F0A-4915-ADE7-B6665129CADD

final-exam-fa79c

#58 Page 18 of 30

DO NOT WRITE ABOVE THIS LINE

- d) Give the vector equation representing the solution for the case with infinite solutions as determined in part c.



DO NOT WRITE ABOVE THIS LINE

e) Give the row picture of the infinite solutions to the system of linear equations from part c.


f) Find the solution when $z = 1$.




DO NOT WRITE ABOVE THIS LINE

Q5: One of the problems EngSci students get asked in another first year course, CIV102F, is to solve for the forces in the members of a Warren truss, like the one shown below:

2. Calculate the joint loads and solve for the forces in the members of the Warren truss shown below. All members have the same length. Select the lightest square HSS which can be used for all of the truss members. Use $E = 200,000$ MPa and $\sigma_y = 350$ MPa.



After converting the 70 kN/m load to discrete "joint loads", we get:



Let \overrightarrow{AB} denote the force in kN carried in the member which connects joints A and B , with tension denoted as positive and compression denoted as negative. Treat the leftmost joint as letter A and the rightmost joint as letter M . There are 13 joints that produce two equations each from the x and y static force balances at each joint for a total of 26 equations.

For the purpose of this question on the ESC103F final exam, we want to express the system of equations that need to be solved in the form $A\vec{x} = \vec{b}$. In the vector of unknowns \vec{x} , if we start with \overrightarrow{AB} and end with \overrightarrow{LM} , you end up with a vector of 23 unknowns:

$$\vec{x} = \begin{bmatrix} \overrightarrow{AB} \\ \overrightarrow{AC} \\ \overrightarrow{BC} \\ \overrightarrow{BD} \\ \overrightarrow{CD} \\ \vdots \\ \overrightarrow{LM} \end{bmatrix}$$



DO NOT WRITE ABOVE THIS LINE

- a) What is the size of matrix A ? What is the shape of matrix A (square, tall and thin, short and wide)? What is the size of vector \vec{b} ?

- b) After applying Gaussian elimination to the augmented matrix $[A|\vec{b}]$ in order to obtain $[R_0|\vec{d}]$, the reduced row echelon form of matrix A is of this form,

$$R_0 = \begin{bmatrix} I_{23 \times 23} \\ \text{row of zeros} \\ \text{row of zeros} \\ \text{row of zeros} \end{bmatrix}$$

In other words, a 23×23 identity matrix appears on the top of R_0 , followed by three rows of zeros. What is the rank of matrix A ? Explain your answer.



F1F7D452-27A7-4BE0-A02F-349FBFFCAA60

final-exam-fa79c

#58 Page 22 of 30

DO NOT WRITE ABOVE THIS LINE

- c) Based on the specific form of R_0 given in part b, what are the possible outcomes for solutions to $A\vec{x} = \vec{b}$, i.e., no solution, one solution, infinite solutions? Explain your answer.



DO NOT WRITE ABOVE THIS LINE

- d) After applying Gaussian elimination to the augmented matrix $[A|\vec{b}]$ to obtain $[R_0|\vec{d}]$, vector \vec{d} is given by,

$$\vec{d} = \begin{bmatrix} -1010 \\ 505 \\ 1010 \\ -1010 \\ -606 \\ 1313 \\ 606 \\ -1617 \\ -202 \\ 1718 \\ 202 \\ -1819 \\ 202 \\ 1718 \\ -202 \\ -1617 \\ 606 \\ 1313 \\ -606 \\ -1010 \\ 1010 \\ 505 \\ -1010 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Given the vector \vec{d} shown above, how many solutions are there to $A\vec{x} = \vec{b}$? Explain your answer.



CD38D258-674D-49F7-A5B4-1BD67DF5B4CE

final-exam-fa79c

#58 Page 24 of 30

DO NOT WRITE ABOVE THIS LINE

- e) What is the solution for \overrightarrow{LM} , the force carried in the member which connects joints L and M ?



DO NOT WRITE ABOVE THIS LINE

Q6: From physics, we know that a spring-mass system can be described by the following second order differential equation,

$$mx'' + kx = 0$$

with mass m , spring constant k , and $x(t)$ representing the displacement of the mass from its equilibrium position.

For a particular spring-mass system, $m = \frac{1}{16}$ and $k = 4$. Therefore,

$$\frac{1}{16}x'' + 4x = 0$$

or,

$$x'' + 64x = 0$$

The initial conditions for this particular spring-mass system are given as follows,

$$x(0) = 0 \text{ and } x'(0) = -16$$

The exact analytical solution for this initial value problem (IVP) is given by,

$$x(t) = -2\sin(8t)$$

- a) Show that this analytical solution satisfies both the second order differential equation and the initial conditions.



60CCE806-3979-4A73-9604-6834DC06B5D5

final-exam-fa79c

#58 Page 26 of 30

DO NOT WRITE ABOVE THIS LINE

- b) Formulate this problem by defining $Z(t) = \begin{bmatrix} x(t) \\ x'(t) \end{bmatrix}$ and $Z' = AZ$. Determine matrix A and $Z(0)$.



DO NOT WRITE ABOVE THIS LINE

- c) If you were to solve this IVP numerically using Euler's Method (EM), determine two separate difference equations, one for x_{n+1} and one for x'_{n+1} , assuming a step size of Δt .



9CD762B7-39A0-426F-8B25-96DE68B649A4

final-exam-fa79c

#58 Page 28 of 30

DO NOT WRITE ABOVE THIS LINE

- d) Consider solving this problem using EM with two different step sizes, $\Delta t = 0.005$ and $\Delta t = 0.01$. Calculate two numerical estimates of the displacement $x(t)$ at $t = 0.01$, one using $\Delta t = 0.005$ and one using $\Delta t = 0.01$.



DO NOT WRITE ABOVE THIS LINE

- e) Consider a different version of the Improved Euler's method studied in ESC103F. Write the difference equation(s) for updating vector Z_{n+1} for a method that uses the slope estimate at the midpoint of the interval instead of the average of the left and right slope estimates.



7D6ECA80-033D-4974-9E7D-FDCAC7D9E8CA

final-exam-fa79c

#58 Page 30 of 30

DO NOT WRITE ABOVE THIS LINE

Rough work only on this page.