

Applied Epidemiology I: Tables and interpreting results

Enoch Yi-Tung Chen

Department of Medical Epidemiology and Biostatistics, Karolinska Institutet

November 26, 2020

Acknowledgements

This course material is based on my learning from Anastasia Lam's teachings in last year's Applied Epidemiology I lab sessions, and readings from *Epidemiology* by Gordis [1], *A First Course in Probability and Statistics* by Goldsman and Goldsman [2], *Principles of Biostatistics* by Pagano and Gauvreau [3], and *Biostatistics I* by Gabriel and Frumento [4]. I especially want to thank Marlene Stratmann for reviewing the slides and Prof. Paul Dickman for providing me with suggestions to improving the teaching.

① Tables

- One-way tables
- Two by two tables
- Stata tool for Epidemiology

② Basic Epidemiology terms

- Rate vs. proportion
- Risk, risk difference, risk ratio
- Odds, odds ratio

③ Interpreting results

- Principles
- Ratio $>$ or < 1
- More examples

④ Calculate ratios using Stata

- Risk ratio
- Odds ratio
- Incidence rate ratio

⑤ References

Tables: One-way tables

- We use cancer data still.
sysuse cancer, clear
keep if drug ==1 | drug == 2

Tables: One-way tables

- We use cancer data still.
sysuse cancer, clear
keep if drug ==1 | drug == 2
- One-way table of frequencies with mean and sd of age
. table died, contents(freq mean age sd age)

1 if				
patient				
died		Freq.	mean(age)	sd(age)
-----+-----				
0		9	55.1111	5.487359
1		25	56.88	6.227091

Tables: One-way tables

- One-way table of frequencies

```
. tabulate died
```

1 if			
patient			
died	Freq.	Percent	Cum.
-----+-----			
0	9	26.47	26.47
1	25	73.53	100.00
-----+-----			
Total	34	100.00	

Tables: One-way tables

- One-way tables of frequencies each variable specified

. tab1 died drug

-> tabulation of died

1 if patient died	Freq.	Percent	Cum.
0	9	26.47	26.47
1	25	73.53	100.00
Total	34	100.00	

-> tabulation of drug

Drug type (1=placebo)	Freq.	Percent	Cum.
0	14	41.18	41.18
1	20	58.82	100.00
Total	34	100.00	

Tables: Two by two tables

- 2 by 2 table for drug and died with relative frequency by column or row

```
. tabulate died drug, col row
```

1 if patient Drug type (1=placebo) died	0		1	Total
0	8	1		9
	88.89	11.11		100.00
	57.14	5.00		26.47
1	6	19		25
	24.00	76.00		100.00
	42.86	95.00		73.53
Total	14	20		34
	41.18	58.82		100.00
	100.00	100.00		100.00

Tables: Two by two tables

- 2 by 2 table with chi-square test and fisher's exact test

```
. tabulate died drug, col row chi2 exact
```

1 if			
patient Drug type (1=placebo)			
died	0	1	Total
-----+-----			
0	8	1	9
	88.89	11.11	100.00
	57.14	5.00	26.47
-----+-----			
1	6	19	25
	24.00	76.00	100.00
	42.86	95.00	73.53
-----+-----			
Total	14	20	34
	41.18	58.82	100.00
	100.00	100.00	100.00

```
Pearson chi2(1) = 11.5039 Pr = 0.001
```

```
Fisher's exact = 0.001
```

Tables: Two by two tables

- 2 by 2 table with chi-square test and fisher's exact test

- How to interpret the results?

```
. tabulate died drug, col row chi2 exact
```

1 if			
patient Drug type (1=placebo)			
died	0	1	Total
-----+-----			
0	8	1	9
	88.89	11.11	100.00
	57.14	5.00	26.47
-----+-----			
1	6	19	25
	24.00	76.00	100.00
	42.86	95.00	73.53
-----+-----			
Total	14	20	34
	41.18	58.82	100.00
	100.00	100.00	100.00

```
Pearson chi2(1) = 11.5039 Pr = 0.001
```

```
Fisher's exact = 0.001
```

Tables: Two by two tables

- 2 by 2 table with chi-square test and fisher's exact test

```
. tabulate died drug, col row chi2 exact
1 if |
patient | Drug type (1=placebo)
died |      0      1 |      Total
-----+-----+-----
0 |      8      1 |      9
  |  88.89  11.11 |  100.00
  |  57.14   5.00 |  26.47
-----+-----+-----
1 |      6     19 |     25
  |  24.00  76.00 |  100.00
  |  42.86  95.00 |  73.53
-----+-----+-----
Total |     14     20 |     34
  |  41.18  58.82 |  100.00
  |  100.00 100.00 |  100.00
```

```
Pearson chi2(1) = 11.5039   Pr = 0.001
Fisher's exact =           0.001
```

- How to interpret the results?
- Chi-square test: testing the association between two binary variables.

Tables: Two by two tables

- 2 by 2 table with chi-square test and fisher's exact test

```
. tabulate died drug, col row chi2 exact
1 if |
patient | Drug type (1=placebo)
died |      0      1 |      Total
-----+-----+-----
0 |      8      1 |      9
  | 88.89  11.11 | 100.00
  | 57.14   5.00 | 26.47
-----+-----+-----
1 |      6     19 |     25
  | 24.00  76.00 | 100.00
  | 42.86  95.00 | 73.53
-----+-----+-----
Total |     14     20 |     34
  | 41.18  58.82 | 100.00
  | 100.00 100.00 | 100.00
```

```
Pearson chi2(1) = 11.5039   Pr = 0.001
Fisher's exact =           0.001
```

- How to interpret the results?
- Chi-square test: testing the association between two binary variables.
- Using placebo has association with that the patients died or not.

Tables: Two by two tables

- 2 by 2 tables stratified by sex

```
. bysort sex: tab died drug, col row chi2
```

```
-> sex = 0
```

1 if patient	Drug type (1=placebo)		
died	0	1	Total

0	6	1	7
	85.71	14.29	100.00
	75.00	9.09	36.84

1	2	10	12
	16.67	83.33	100.00
	25.00	90.91	63.16

Total	8	11	19
	42.11	57.89	100.00
	100.00	100.00	100.00

Pearson chi2(1) = 8.6466 Pr = 0.003

```
-> sex = 1
```

1 if patient	Drug type (1=placebo)		
died	0	1	Total

0	2	0	2
	100.00	0.00	100.00
	33.33	0.00	13.33

1	4	9	13
	30.77	69.23	100.00
	66.67	100.00	86.67

Total	6	9	15
	40.00	60.00	100.00
	100.00	100.00	100.00

Pearson chi2(1) = 3.4615 Pr = 0.063

Tables: Stata tool for Epidemiology

- How to use Stata to generate risk ratios (relative risk) and odds ratios?

Tables: Stata tool for Epidemiology

- How to use Stata to generate risk ratios (relative risk) and odds ratios?
- A useful tool in Stata's default function can be found at
- Statistics - Epidemiology and related - Tables for epidemiologists

Stata/IC 16.1 File Edit View Data Graphics **Statistics** User Window Help

Tables: Stata tool for Epidemiology

- How to use Stata to generate risk ratios (relative risk) and odds ratios?
- A useful tool in Stata's default function can be found at
- Statistics - Epidemiology and related - Tables for epidemiologists

Stata/IC 16.1 File Edit View Data Graphics **Statistics** User Window Help

- But before demonstrating how this works, a recapture on basic epi terms!

Basic Epidemiology terms: Rate vs. proportion

Rate

Proportion

Risk

Basic Epidemiology terms: Rate vs. proportion

Rate

- Incidence (rate): $\frac{\text{no. of diseased}}{\text{total person-time}}$

Proportion

Risk

Basic Epidemiology terms: Rate vs. proportion

Rate

- Incidence (rate): $\frac{\text{no. of diseased}}{\text{total person-time}}$
- Mortality (rate): $\frac{\text{no. of deaths}}{\text{total person-time}}$

Proportion

Risk

Basic Epidemiology terms: Rate vs. proportion

Rate

- Incidence (rate): $\frac{\text{no. of diseased}}{\text{total person-time}}$
- Mortality (rate): $\frac{\text{no. of deaths}}{\text{total person-time}}$
- Hazard (rate): in survival analysis, usually hazard is defined as mortality rate.

Proportion

Risk

Basic Epidemiology terms: Rate vs. proportion

Rate

- Incidence (rate): $\frac{\text{no. of diseased}}{\text{total person-time}}$
- Mortality (rate): $\frac{\text{no. of deaths}}{\text{total person-time}}$
- Hazard (rate): in survival analysis, usually hazard is defined as mortality rate.

Proportion

- Cumulative incidence: **over a time period**, $\frac{\text{no. of new cases of the disease}}{\text{no. of initially disease-free persons}}$

Risk

Basic Epidemiology terms: Rate vs. proportion

Rate

- Incidence (rate): $\frac{\text{no. of diseased}}{\text{total person-time}}$
- Mortality (rate): $\frac{\text{no. of deaths}}{\text{total person-time}}$
- Hazard (rate): in survival analysis, usually hazard is defined as mortality rate.

Proportion

- Cumulative incidence: **over a time period**, $\frac{\text{no. of new cases of the disease}}{\text{no. of initially disease-free persons}}$
- Point prevalence: **at a specified time**, $\frac{\text{no. of diseased}}{\text{no. of persons}}$

Risk

Basic Epidemiology terms: Rate vs. proportion

Rate

- Incidence (rate): $\frac{\text{no. of diseased}}{\text{total person-time}}$
- Mortality (rate): $\frac{\text{no. of deaths}}{\text{total person-time}}$
- Hazard (rate): in survival analysis, usually hazard is defined as mortality rate.

Proportion

- Cumulative incidence: **over a time period**, $\frac{\text{no. of new cases of the disease}}{\text{no. of initially disease-free persons}}$
- Point prevalence: **at a specified time**, $\frac{\text{no. of diseased}}{\text{no. of persons}}$
- Period prevalence: **over a time period**, $\frac{\text{no. of diseased}}{\text{no. of persons}}$

Risk

Basic Epidemiology terms: Rate vs. proportion

Rate

- Incidence (rate): $\frac{\text{no. of diseased}}{\text{total person-time}}$
- Mortality (rate): $\frac{\text{no. of deaths}}{\text{total person-time}}$
- Hazard (rate): in survival analysis, usually hazard is defined as mortality rate.

Proportion

- Cumulative incidence: **over a time period**, $\frac{\text{no. of new cases of the disease}}{\text{no. of initially disease-free persons}}$
- Point prevalence: **at a specified time**, $\frac{\text{no. of diseased}}{\text{no. of persons}}$
- Period prevalence: **over a time period**, $\frac{\text{no. of diseased}}{\text{no. of persons}}$
- Survival (proportion/probability) ~~rate~~:
$$\frac{\text{no. of alive persons (since diagnosis)}}{\text{no. of initially disease-free persons (since diagnosis)}}$$

Risk

Basic Epidemiology terms: Rate vs. proportion

Rate

- Incidence (rate): $\frac{\text{no. of diseased}}{\text{total person-time}}$
- Mortality (rate): $\frac{\text{no. of deaths}}{\text{total person-time}}$
- Hazard (rate): in survival analysis, usually hazard is defined as mortality rate.

Proportion

- Cumulative incidence: **over a time period**, $\frac{\text{no. of new cases of the disease}}{\text{no. of initially disease-free persons}}$
- Point prevalence: **at a specified time**, $\frac{\text{no. of diseased}}{\text{no. of persons}}$
- Period prevalence: **over a time period**, $\frac{\text{no. of diseased}}{\text{no. of persons}}$
- Survival (proportion/probability) ~~rate~~:
$$\frac{\text{no. of alive persons (since diagnosis)}}{\text{no. of initially disease-free persons (since diagnosis)}}$$

Risk

- Is risk a rate or a proportion?

Basic Epidemiology terms: Risk, risk difference, risk ratio

Risk: the proportion (probability) of an event, e.g., death.

Basic Epidemiology terms: Risk, risk difference, risk ratio

Risk: the proportion (probability) of an event, e.g., death.

- Estimates of risk: cumulative incidence, cumulative hazard

Basic Epidemiology terms: Risk, risk difference, risk ratio

Risk: the proportion (probability) of an event, e.g., death.

- Estimates of risk: cumulative incidence, cumulative hazard
- E.g., in survival analysis,

Cumulative hazard = 1 – Survival proportion = Cumulative probability of death

$$F(t) = 1 - S(t) = P(T \leq t)$$

Basic Epidemiology terms: Risk, risk difference, risk ratio

Risk: the proportion (probability) of an event, e.g., death.

- Estimates of risk: cumulative incidence, cumulative hazard
- E.g., in survival analysis,

Cumulative hazard = 1 – Survival proportion = Cumulative probability of death

$$F(t) = 1 - S(t) = P(T \leq t)$$

- Risk difference: the difference of the probabilities of an event between the exposed group and non-exposed group

Basic Epidemiology terms: Risk, risk difference, risk ratio

Risk: the proportion (probability) of an event, e.g., death.

- Estimates of risk: cumulative incidence, cumulative hazard
- E.g., in survival analysis,

Cumulative hazard = 1 – Survival proportion = Cumulative probability of death

$$F(t) = 1 - S(t) = P(T \leq t)$$

- Risk difference: the difference of the probabilities of an event between the exposed group and non-exposed group
- Risk ratio (relative risk): the ratio of the probabilities of an event between the exposed group and non-exposed group

Basic Epidemiology terms: Risk, risk difference, risk ratio

	Female (Exposed)	Male (Unexposed)	Total
shiba (Case)	2	2	4
guinea pig (Noncase)	2	1	3
Total	4	3	7

Epidemiologists love two by two tables!

Basic Epidemiology terms: Risk, risk difference, risk ratio

	Female (Exposed)	Male (Unexposed)	Total
shiba (Case)	2	2	4
guinea pig (Noncase)	2	1	3
Total	4	3	7

Epidemiologists love two by two tables!

- Risk difference between females having shiba and males having shiba = $\widehat{p}_F - \widehat{p}_M = 2/4 - 2/3 = -0.16667$
- Interpretation: Females have 16.67 % lower risk of having shiba than males.

Basic Epidemiology terms: Risk, risk difference, risk ratio

	Female (Exposed)	Male (Unexposed)	Total
shiba (Case)	2	2	4
guinea pig (Noncase)	2	1	3
Total	4	3	7

Epidemiologists love two by two tables!

- Risk difference between females having shiba and males having shiba
$$= \widehat{p}_F - \widehat{p}_M = 2/4 - 2/3 = -0.16667$$
- Interpretation: Females have 16.67 % lower risk of having shiba than males.
- Risk ratio between females having shiba and males having shiba
$$= \widehat{p}_F \div \widehat{p}_M = 2/4 \div 2/3 = 0.75.$$
- Interpretation: The RR of females having shiba is 0.75 times than males having shiba.

Basic Epidemiology terms: Odds, odds ratio

Odds: the ratio between those having and not having an outcome.

$$Odds = \frac{p}{1 - p}$$

Basic Epidemiology terms: Odds, odds ratio

Odds: the ratio between those having and not having an outcome.

$$Odds = \frac{p}{1 - p}$$

- Odds ratio measures the association between an exposure and an outcome.

$$\text{Odds ratio (OR)} = \frac{Odds_{exposed}}{Odds_{unexposed}}$$

Basic Epidemiology terms: Odds, odds ratio

Odds: the ratio between those having and not having an outcome.

$$Odds = \frac{p}{1 - p}$$

- Odds ratio measures the association between an exposure and an outcome.

$$\text{Odds ratio (OR)} = \frac{Odds_{exposed}}{Odds_{unexposed}}$$

- E.g., the OR is 0.5, which indicates that there is a 50% decrease in the odds of having an outcome among the exposed compared to the unexposed.

Basic Epidemiology terms: Odds, odds ratio

Odds: the ratio between those having and not having an outcome.

$$Odds = \frac{p}{1 - p}$$

- Odds ratio measures the association between an exposure and an outcome.

$$\text{Odds ratio (OR)} = \frac{Odds_{\text{exposed}}}{Odds_{\text{unexposed}}}$$

- E.g., the OR is 0.5, which indicates that there is a 50% decrease in the odds of having an outcome among the exposed compared to the unexposed.
- Why there is no odds difference?

Basic Epidemiology terms: Odds, odds ratio

	Female (Exposed)	Male (Unexposed)	Total
shiba (Case)	2	2	4
guinea pig (Noncase)	2	1	3
Total	4	3	7

Basic Epidemiology terms: Odds, odds ratio

	Female (Exposed)	Male (Unexposed)	Total
shiba (Case)	2	2	4
guinea pig (Noncase)	2	1	3
Total	4	3	7

- The odds of having shiba among females is

$$\begin{aligned}\widehat{odds}_F &= \frac{p(\text{having shiba}|\text{female})}{p(\text{having guinea pig}|\text{female})} \\ &= \frac{(2/4)}{(2/4)} = 1\end{aligned}$$

- The odds of having shiba among males is 2 (calculation ignored).

Basic Epidemiology terms: Odds, odds ratio

	Female (Exposed)	Male (Unexposed)	Total
shiba (Case)	2	2	4
guinea pig (Noncase)	2	1	3
Total	4	3	7

- OR of having shiba (females to males)
- $OR = \frac{Odds_f}{Odds_m} = \frac{1}{2}$

- The odds of having shiba among females is

$$\begin{aligned}\widehat{odds}_f &= \frac{p(\text{having shiba}|\text{female})}{p(\text{having guinea pig}|\text{female})} \\ &= \frac{(2/4)}{(2/4)} = 1\end{aligned}$$

- The odds of having shiba among males is 2 (calculation ignored).

Basic Epidemiology terms: Odds, odds ratio

	Female (Exposed)	Male (Unexposed)	Total
shiba (Case)	2	2	4
guinea pig (Noncase)	2	1	3
Total	4	3	7

- The odds of having shiba among females is

$$\begin{aligned}\widehat{odds}_f &= \frac{p(\text{having shiba}|\text{female})}{p(\text{having guinea pig}|\text{female})} \\ &= \frac{(2/4)}{(2/4)} = 1\end{aligned}$$

- The odds of having shiba among males is 2 (calculation ignored).

- OR of having shiba (females to males)
- $OR = \frac{Odds_f}{Odds_m} = \frac{1}{2}$
- Interpretation: there is a 50% decrease in the odds of having shiba among females compared to males. Higher odds of shiba ownership among males than females!
- It seems that females instead love guinea pigs more.

Interpreting results: Principles

- When describing a ratio, it can ideally be illustrated by
 1. Exposed group
 2. Ratio (exact value, higher or lower percentage)
 3. Outcome
 4. Unexposed

Interpreting results: Principles

- When describing a ratio, it can ideally be illustrated by
 1. Exposed group
 2. Ratio (exact value, higher or lower percentage)
 3. Outcome
 4. Unexposed
- Example:
 1. Females have a RR of 0.75 having shiba compared to males.

Interpreting results: Principles

- When describing a ratio, it can ideally be illustrated by
 1. Exposed group
 2. Ratio (exact value, higher or lower percentage)
 3. Outcome
 4. Unexposed
- Example:
 1. Females have a RR of 0.75 having shiba compared to males.
 2. Females have a 50% decrease in the odds of having shiba compared to males.

Interpreting results: Ratio $>$ or $<$ 1

Ratio

Interpreting results: Ratio $>$ or $<$ 1

Ratio

- As ratio $<$ 1,
 - $(1 - \text{RR/OR}) \times 100\%$
 - E.g., $\text{RR} = 0.75$, $(1 - 0.75) \times 100\% = 25\%$
 - 25% lower risk

Interpreting results: Ratio $>$ or $<$ 1

Ratio

- As ratio $<$ 1,
 - $(1 - \text{RR/OR}) \times 100\%$
 - E.g., $\text{RR} = 0.75$, $(1 - 0.75) \times 100\% = 25\%$
 - 25% lower risk
- As ratio $>$ 1,
 - $(\text{RR/OR} - 1) \times 100\%$
 - E.g., $\text{OR} = 2.05$, $(2.05 - 1) \times 100\% = 105\%$
 - The odds is 2 times higher.
 - Twice the odds

Interpreting results: More examples

Diabetes Is a Risk Factor for Pulmonary Tuberculosis: A Case-Control Study from Mwanza, Tanzania (Faurholt-Jepsen, 2011)

	OR (95% C.I.)	OR (95% C.I.)	OR (95% C.I.)
		Model 1	Model 2
	Unadjusted	Adjusted for age, sex, socio-demography ²	Model 1 + AGP ³
HIV negative (n = 770)			
Glucose intolerance status ¹			
normal glucose tolerance	ref.	ref.	ref.
IFG/IGT	2.26 (1.50;3.41)	2.34 (1.52;3.61)	2.65 (1.00;7.06)
diabetes	2.15 (1.35;3.42)	2.14 (1.32;3.46)	4.23 (1.54;11.57)

Interpreting results: More examples

Diabetes Is a Risk Factor for Pulmonary Tuberculosis: A Case-Control Study from Mwanza, Tanzania (Faurholt-Jepsen, 2011)

	OR (95% C.I.)	OR (95% C.I.)	OR (95% C.I.)
		Model 1	Model 2
	Unadjusted	Adjusted for age, sex, socio-demography ²	Model 1 + AGP ³
HIV negative (n = 770)			
Glucose intolerance status ¹			
normal glucose tolerance	ref.	ref.	ref.
IFG/IGT	2.26 (1.50;3.41)	2.34 (1.52;3.61)	2.65 (1.00;7.06)
diabetes	2.15 (1.35;3.42)	2.14 (1.32;3.46)	4.23 (1.54;11.57)

1. People with diabetes had a higher odds of TB (OR 2.15, 95% CI: 1.35-3.42) relative to people without diabetes.

Interpreting results: More examples

Diabetes Is a Risk Factor for Pulmonary Tuberculosis: A Case-Control Study from Mwanza, Tanzania (Faurholt-Jepsen, 2011)

	OR (95% C.I.)	OR (95% C.I.)	OR (95% C.I.)
		Model 1	Model 2
	Unadjusted	Adjusted for age, sex, socio-demography ²	Model 1 + AGP ³
HIV negative (n = 770)			
Glucose intolerance status ¹			
normal glucose tolerance	ref.	ref.	ref.
IFG/IGT	2.26 (1.50;3.41)	2.34 (1.52;3.61)	2.65 (1.00;7.06)
diabetes	2.15 (1.35;3.42)	2.14 (1.32;3.46)	4.23 (1.54;11.57)

1. People with diabetes had a higher odds of TB (OR 2.15, 95% CI: 1.35-3.42) relative to people without diabetes.
2. Having diabetes was associated with more than a 2-fold increase (OR: 2.15, 95% CI: 1.35; 3.42) in the odds of TB compared to not having diabetes.

Interpreting results: More examples

Bidirectional association between physical activity and symptoms of anxiety and depression: the Whitehall II study (Azevedo Da Silva, 2012)

Table 3 Cross-sectional associations between physical activity at recommended levels and anxiety and/or depression symptoms at phase 1 (1985–1988) (N = 9,309)

	OR (CI 95 %)	<i>P</i> value
<i>Anxiety symptoms</i>		
Model 1		
Physical activity		
Yes	0.71 (0.54, 0.91)	0.01
No	1 (reference)	
Model 2		
Physical activity		
Yes	0.71 (0.55, 0.93)	0.01
No	1 (reference)	
<i>Depression symptoms</i>		
Model 1		
Physical activity		
Yes	0.63 (0.48, 0.81)	<0.001
No	1 (reference)	
Model 2		
Physical activity		
Yes	0.63 (0.49, 0.82)	0.001
No	1 (reference)	

Interpreting results: More examples

Bidirectional association between physical activity and symptoms of anxiety and depression: the Whitehall II study (Azevedo Da Silva, 2012)

Table 3 Cross-sectional associations between physical activity at recommended levels and anxiety and/or depression symptoms at phase 1 (1985–1988) (N = 9,309)

	OR (CI 95 %)	<i>P</i> value
<i>Anxiety symptoms</i>		
Model 1		
Physical activity		
Yes	0.71 (0.54, 0.91)	0.01
No	1 (reference)	
Model 2		
Physical activity		
Yes	0.71 (0.55, 0.93)	0.01
No	1 (reference)	
<i>Depression symptoms</i>		
Model 1		
Physical activity		
Yes	0.63 (0.48, 0.81)	<0.001
No	1 (reference)	
Model 2		
Physical activity		
Yes	0.63 (0.49, 0.82)	0.001
No	1 (reference)	

1. Patients who conducted recommended levels of physical activity had a 29% lower odds of anxiety (OR: 0.71, 95% CI: 0.54-0.91) and a 37% lower odds of depression (OR: 0.63, 95% CI: 0.48-0.81) relative to those who did not.

Interpreting results: More examples

Bidirectional association between physical activity and symptoms of anxiety and depression: the Whitehall II study (Azevedo Da Silva, 2012)

Table 3 Cross-sectional associations between physical activity at recommended levels and anxiety and/or depression symptoms at phase 1 (1985–1988) (N = 9,309)

	OR (CI 95 %)	<i>P</i> value
<i>Anxiety symptoms</i>		
Model 1		
Physical activity		
Yes	0.71 (0.54, 0.91)	0.01
No	1 (reference)	
Model 2		
Physical activity		
Yes	0.71 (0.55, 0.93)	0.01
No	1 (reference)	
<i>Depression symptoms</i>		
Model 1		
Physical activity		
Yes	0.63 (0.48, 0.81)	<0.001
No	1 (reference)	
Model 2		
Physical activity		
Yes	0.63 (0.49, 0.82)	0.001
No	1 (reference)	

1. Patients who conducted recommended levels of physical activity had a 29% lower odds of anxiety (OR: 0.71, 95% CI: 0.54-0.91) and a 37% lower odds of depression (OR: 0.63, 95% CI: 0.48-0.81) relative to those who did not.
2. Our results showed that individuals who practiced recommended levels of physical activity were less likely to have anxiety (OR: 0.71, 95% CI: 0.54-0.91) and depression (OR: 0.63, 95% CI: 0.48-0.81) in comparison with those who did not.

Calculate ratios using Stata: Risk ratio

- Finally we come back to Stata again!

Calculate ratios using Stata: Risk ratio

- Finally we come back to Stata again!
- cs case exposed

```
. cs died drug
```

	Drug type [1=placebo]		
	Exposed	Unexposed	Total
Cases	19	6	25
Noncases	1	8	9
Total	20	14	34
Risk	.95	.4285714	.7352941
	Point estimate		[95% Conf. Interval]
Risk difference	.5214286		.245166 .7976911
Risk ratio	2.216667		1.200631 4.092525
Attr. frac. ex.	.5488722		.1671043 .7556521
Attr. frac. pop	.4171429		
+-----			
chi2(1) = 11.50 Pr>chi2 = 0.0007			

Calculate ratios using Stata: Odds ratio

- cs case exposed, or
. cs died drug, or

	Drug type [1=placebo]			
	Exposed	Unexposed	Total	
Cases	19	6	25	
Noncases	1	8	9	
Total	20	14	34	
Risk	.95	.4285714	.7352941	
	Point estimate		[95% Conf. Interval]	
Risk difference	.5214286		.245166	.7976911
Risk ratio	2.216667		1.200631	4.092525
Attr. frac. ex.	.5488722		.1671043	.7556521
Attr. frac. pop	.4171429			
Odds ratio	25.33333		3.189793	. (Cornfield
+-----+				

Calculate ratios using Stata: Incidence rate ratio

- ir case exposed studytime

```
. ir died drug studytime
```

```
Incidence-rate comparison
```

	Drug type [1=placebo]			
	Exposed	Unexposed	Total	
1 if patient die	19	6	25	
Months to death	180	209	389	
Incidence rate	.1055556	.0287081	.0642674	
	Point estimate		[95% Conf. Interval]	
Inc. rate diff.	.0768474		.0241182	.1295766
Inc. rate ratio	3.676852		1.411772	11.24864 (exact)
Attr. frac. ex.	.7280282		.2916701	.9111003 (exact)
Attr. frac. pop	.5533014			

```
Mid p-values for tests of incidence-rate difference:
```

```
Adj Pr(Exposed 1 if patient die <= 19) = 0.9985 (lower one-sided)
```

```
Adj Pr(Exposed 1 if patient die >= 19) = 0.0015 (upper one-sided)
```

```
Two-sided p-value = 0.0031
```

References

1. Gordis L. *Epidemiology*. Philadelphia, PA: Elsevier/Saunders, 2014. ISBN 9781455737338.
2. David Goldsman PG. *A First Course in Probability and Statistics*. Georgia Institute of Technology, 2020.
3. Marcello Pagano KG. *Principles of Biostatistics*. Cengage Learning, Inc, 2000. ISBN 0534229026.
4. Erin Gabriel PF. Epidemiology PhD program, Karolinska Institutet, 2020.