
MATH 2023 Fall 2021

Multivariable Calculus

Written By: Ljm

Chapter 15 Vector Field


1 Intro. to Vector Field

So far, we have learned two kinds of functions involving vector:

- $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}$: for each t , provides a *position* vector $\langle x(t), y(t), z(t) \rangle$, so this is a (parametric) curve.
- $z = f(\mathbf{r}) = f(x_1, x_2, \dots, x_n)$: for a given vector \mathbf{r} , this gives a real number, so this is a function of *several variables*. This is also a **scalar field** since for any point \mathbf{r} in **field**, it gives a scalar value.

Now we are looking at **vector-valued** function \mathbf{F} of a vector \mathbf{r} , i.e., $\mathbf{F}(\mathbf{r})$. This is a **vector field**, which means for any point \mathbf{r} in **field**, it gives a vector $\mathbf{F}(\mathbf{r})$.

You can consider a world map showing the *speed* and *direction* of wind.



You can see that in a 2D map(like above), if we put a vector on each point, the vector must have same dimension as the map, i.e., all vectors must also be 2D vectors.


$$\mathbf{F}(\mathbf{r}) = \begin{cases} (F_1(\mathbf{r}), F_2(\mathbf{r})) & \mathbf{r} = (x, y) & 2D \\ (F_1(\mathbf{r}), F_2(\mathbf{r}), F_3(\mathbf{r})) & \mathbf{r} = (x, y, z) & 3D \\ (F_1(\mathbf{r}), F_2(\mathbf{r}), \dots, F_n(\mathbf{r})) & \mathbf{r} = (x_1, x_2, \dots, x_n) & nD \end{cases}$$

Summary: *dimension of \mathbf{F} must be the same as \mathbf{r} .*

[Example.] Assume a vector field: $\mathbf{F}(x, y) = \frac{y\mathbf{i} - x\mathbf{j}}{\sqrt{x^2 + y^2}}$.

[Solution.] Notice that $\|\mathbf{F}\| = \frac{y^2}{x^2 + y^2} + \frac{x^2}{x^2 + y^2} = 1$, all vectors $\mathbf{F}(x, y)$ are unit vectors. Moreover, let $\mathbf{r} = (x, y)$, then $\mathbf{r} \cdot \mathbf{F} = 0$, so $\mathbf{r} \perp \mathbf{F}$.

So all vectors are unit vectors tangent to circles centered at the origin with radius $\sqrt{x^2 + y^2}$.



2 Divergence and Curl

Recall that the **gradient operator** is a *vector operator*:

$$\nabla = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right) \quad (\text{a vector})$$

If $\mathbf{F}(\mathbf{r}) = F_1(\mathbf{r})\mathbf{i} + F_2(\mathbf{r})\mathbf{j} + F_3(\mathbf{r})\mathbf{k}$, then we define:

- **divergence** of \mathbf{F} , written $\text{div } \mathbf{F}$:

$$\text{div } \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$$

- **curl** of \mathbf{F} , written $\text{curl } \mathbf{F}$:

$$\text{curl } \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix} = \left(\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z} \right) \mathbf{i} - \left(\frac{\partial F_3}{\partial x} - \frac{\partial F_1}{\partial z} \right) \mathbf{j} + \left(\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) \mathbf{k}$$

This example shows basic computation of **divergence** and **curl**.

[**Example.**] Let $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ and $\mathbf{u} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$, where a, b and c are constants, show that

- (a) $\nabla \cdot \mathbf{r} = 3$
- (b) $\nabla \times \mathbf{r} = \mathbf{0}$
- (c) $\nabla \cdot (\mathbf{u} \times \mathbf{r}) = 0$
- (d) $\nabla \times (\mathbf{u} \times \mathbf{r}) = 2\mathbf{u}$.

[**Solution.**] (a) $\nabla \cdot \mathbf{r} = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (x\mathbf{i} + y\mathbf{j} + z\mathbf{k}) = \frac{\partial x}{\partial x} + \frac{\partial y}{\partial y} + \frac{\partial z}{\partial z} = 3$

$$(b) \nabla \times \mathbf{r} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & y & z \end{vmatrix} = \mathbf{0}$$

$$(c) \mathbf{u} \times \mathbf{r} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a & b & c \\ x & y & z \end{vmatrix} = (bz - cy)\mathbf{i} - (az - cx)\mathbf{j} + (ay - bx)\mathbf{k}$$

$$\therefore \nabla \cdot (\mathbf{u} \times \mathbf{r}) = \frac{\partial}{\partial x}(bz - cy) - \frac{\partial}{\partial y}(az - cx) + \frac{\partial}{\partial z}(ay - bx) = 0$$


$$(d) \nabla \times (\mathbf{u} \times \mathbf{r}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ bz - cy & -az + cx & ay - bx \end{vmatrix} = 2(a\mathbf{i} + b\mathbf{j} + c\mathbf{k}) = 2\mathbf{u}$$

2.1 Interpretation of Divergence

Imagine water in a bath tank, if the **velocity** of water at any point of the tank is given by

$$\mathbf{u}(\mathbf{r}) = u_1(\mathbf{r})\mathbf{i} + u_2(\mathbf{r})\mathbf{j} + u_3(\mathbf{r})\mathbf{k}$$


then **net outward flux per unit volume** is $\text{div } \mathbf{u} = \nabla \cdot \mathbf{u}$.



Moreover,

- If more water comes inside, then $\text{div } \mathbf{u} < 0$
- If more water comes outside, then $\text{div } \mathbf{u} > 0$
- If the amount of water comes inside equals to comes outside, then $\text{div } \mathbf{u} = 0$

This page proves the interpretation of divergence.



Imagine the box with volume $\Delta V = \Delta x \Delta y \Delta z$, firstly consider faces **I** and **II**, the total flux *out of* faces **I** and **II**, as shown above, is:

$$\begin{aligned}
 & [u_1(x + \Delta x, y, z) - u_1(x, y, z)] \Delta y \Delta z \\
 &= \frac{[u_1(x + \Delta x, y, z) - u_1(x, y, z)]}{\Delta x} \Delta x \Delta y \Delta z \\
 &= \frac{\partial u_1}{\partial x} \Delta x \Delta y \Delta z, \quad (\text{in the limit of } \Delta x \rightarrow 0)
 \end{aligned}$$

Similarly, the two faces in the y - and z - direction contribute

$$\frac{\partial u_2}{\partial y} \Delta x \Delta y \Delta z, \quad \frac{\partial u_3}{\partial z} \Delta x \Delta y \Delta z$$

Hence net outward flux is:


$$\left(\frac{\partial u_1}{\partial x} + \frac{\partial u_2}{\partial y} + \frac{\partial u_3}{\partial z} \right) \cdot \Delta V$$

Therefore outward flux *per unit volume* is $\nabla \cdot \mathbf{u}$.

2.2 Interpretation of Curl

Curl is something related to rotation. Consider a small object flying in strong wind, where the speed and direction of wind can be treated as a vector field \mathbf{F} . If the object locates at position \mathbf{r} , then its rotation has some relation with curl \mathbf{F} .


Actually, the object will rotate about the direction $\nabla \times \mathbf{F}(\mathbf{r})$ (direction is determined by right-hand rule), and with speed $\omega = \frac{1}{2} \|\nabla \times \mathbf{F}(\mathbf{r})\|$.



The rest of this page prove the relation above.

Consider a disk in xy -plane, in y direction, the differential velocity *normal to* Δx is:

$$u_2(x + \Delta x) - u_2(x) = \frac{\partial u_2}{\partial x} \Delta x$$



Recall that $v = \omega r$, so the angular velocity is $\omega_1 = \frac{\partial u_2}{\partial x}$

Similarly, in the y -direction, (notice the negative sign)

$$-u_1(x, y + \Delta y) + u_1(x, y) = -\frac{\partial u_1}{\partial y} \Delta y, \quad \omega_2 = -\frac{\partial u_1}{\partial y}$$

Thus the *averaged angular velocity* is: $\omega = \frac{1}{2} \left(\frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} \right)$

The curl of this vector field is:

$$\text{curl } \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u_1 & u_2 & 0 \end{vmatrix} = \left(\frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} \right) \mathbf{k} = 2\omega \mathbf{k}$$

Thus prove the result.


Below example is used to explain the meaning of curl, it's the same example in intro.

[Example.] Assume a vector field: $\mathbf{F}(x, y) = \frac{y\mathbf{i} - x\mathbf{j}}{\sqrt{x^2 + y^2}}$.

[Solution.]

$$\begin{aligned} \vec{\omega} = \nabla \times \mathbf{F} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{y}{\sqrt{x^2 + y^2}} & \frac{-x}{\sqrt{x^2 + y^2}} & 0 \end{vmatrix} \\ &= \left[-\frac{\partial}{\partial x} \left(\frac{x}{\sqrt{x^2 + y^2}} \right) - \frac{\partial}{\partial y} \left(\frac{y}{\sqrt{x^2 + y^2}} \right) \right] \mathbf{k} \\ &= \mathbf{0} \end{aligned}$$

Consider a small object in the vector field, it doesn't rotate (since $\text{curl } \mathbf{F} = \mathbf{0}$ everywhere), it just move in circular, along the vector field.



This definition is optional.

Laplacian Operator

$$\begin{aligned}\nabla^2 &= \nabla \cdot \nabla = \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot \left(\mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \\ &= \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}\end{aligned}$$

∇^2 is a **scalar** differential operator. Note that

$$\begin{aligned}\nabla^2 f &= \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \\ \nabla^2 \mathbf{F} &= \nabla^2 F_1 \mathbf{i} + \nabla^2 F_2 \mathbf{j} + \nabla^2 F_3 \mathbf{k}\end{aligned}$$

2.3 Vector differential identities

Let ϕ, ψ are scalar fields and \mathbf{F} and \mathbf{G} are vector fields, then

$$(a) \nabla(\phi\psi) = \phi\nabla\psi + \psi\nabla\phi$$

$$(b) \nabla \cdot (\phi\mathbf{F}) = \nabla\phi \cdot \mathbf{F} + \phi(\nabla \cdot \mathbf{F})$$

$$(c) \nabla \times (\phi\mathbf{F}) = \nabla\phi \times \mathbf{F} + \phi(\nabla \times \mathbf{F})$$

$$(d) \nabla \cdot (\mathbf{F} \times \mathbf{G}) = (\nabla \times \mathbf{F}) \cdot \mathbf{G} - \mathbf{F} \cdot (\nabla \times \mathbf{G})$$

$$(e) \nabla \cdot (\nabla \times \mathbf{F}) = 0$$

$$(f) \nabla \times (\nabla\phi) = 0$$

$$(g) \nabla \times (\nabla \times \mathbf{F}) = \nabla(\nabla \cdot \mathbf{F}) - \nabla^2\mathbf{F}$$

[Proof.]