

Section 5.2 Confidence Intervals Using Normal Distributions

E. Nordmoe

Math 261

Outline

- Confidence interval using a normal distribution

Example: Hearing Loss in Teenagers

New York Times [report](#) on a study of hearing loss in teenagers:

- What proportion of Americans aged 12 to 19 have some hearing loss?
 - Give a 95% confidence interval.

Confidence Intervals Using Normal Distributions

If a bootstrap distribution is bell-shaped, a P% confidence interval can be found as the interval containing the middle P% of the normal distribution with mean equal to the observed sample statistic and standard deviation equal to the standard error of the statistic:

$$N(\text{sample statistic}, SE)$$

Confidence Intervals Using the Standard Normal


- If a statistic is normally distributed, we find a P% confidence interval for the parameter using

$$\text{statistic} \pm z^* \text{SE}$$

where z^* is the *critical value* with area P% between $-z^*$ and z^* in the standard normal distribution.

- Extends SE method to confidence levels beyond just 95%.

Use StatKey to Find the Critical Value z^*



Chap 5 Summary: General Formulas

Confidence interval

$$\text{Sample Statistic} \pm z^* \text{SE}$$

Hypothesis test statistic

$$\frac{\text{Sample statistic} - \text{Null parameter}}{\text{SE}}$$

Looking ahead

- For now, the SE comes from **resampling** methods (randomization or bootstrap).
- Beginning next class, we use classical model-based **formulas** to compute SE.