Egg Layer Production Guide

Comprehensive Rearing & Laying Cycle Schedule for 100 Hens (Day 1 to 72 Weeks)

FLOCK SIZE: 100 HENS

REARING PHASE FEED	LAY PHASE DAILY FEED	EXPECTED EGG YIELD	PEAK LAYING RATE
490 kg	**11.0–11.5 kg**	**30,000+ Eggs**	**92% – 96%**
Weeks 1–17 (20 Bags)	110g–115g per hen/day	Per 100 hens (72-wk cycle)	Reached at Wk 26–30

1. TOP COMMERCIAL LAYER BREEDS & PERFORMANCE

BREED NAME	TYPE	EGG COLOR	ANNUAL YIELD	PEAK LAY	KEY TRAITS & SUITABILITY
Lohmann Brown	Hybrid	Dark Brown	320 – 350 eggs	95%	Industry standard. Superior feed efficiency, calm nature, consistent shell strength.
Hy-Line Brown	Hybrid	Rich Brown	320 – 340 eggs	95%	Excellent feather retention, disease resistance, robust in varying climates.
Lohmann LSL White	Hybrid	White	330 – 360 eggs	96%	Lowest feed consumption (~105g/day). Highest egg count for commercial white egg sales.
Amberlink	Hybrid	Medium Brown	300 – 320 eggs	92%	High heat tolerance, excellent foraging traits. Ideal for free-range / pasture systems.
Black Australorp	Dual Breed	Light Brown	250 – 280 eggs	85%	Hardy heritage breed. Provides excellent meat value at end of laying cycle.

2. FEED SCHEDULE & CONSUMPTION (100 HENS)

PHASE	AGE PERIOD	FEED TYPE	CRUDE PROTEIN	DAILY INTAKE / BIRD	PHASE TOTAL (100 BIRDS)
Chick Starter	Week 1 – 6 (Days 1–42)	Chick Starter Mash	20.0% CP	15g – 30g	63.0 kg (3 Bags)
Pullet Grower	Week 7 – 15 (Days 43–105)	Pullet Grower Mash	15.0% CP	45g – 60g	315.0 kg (13 Bags)
Pre-Lay Transition	Week 16 – 17 (Days 106–119)	Pre-Lay Mash (High Calcium)	17.5% CP	75g – 85g	112.0 kg (5 Bags)
Layer Phase 1	Week 18 – 45 (Peak Lay)	Layer Mash 1	17.0% CP	110g	2,156.0 kg (87 Bags)
Layer Phase 2	Week 46 – 72+ (Late Lay)	Layer Mash 2	15.5% CP	115g	2,173.5 kg (87 Bags)

3. HEALTH, VACCINATION & MANAGEMENT PROGRAM

AGE / STAGE	VACCINATION / MEDICATION SCHEDULE	LIGHTING & NUTRITION FOCUS	KEY MANAGEMENT ACTION
Days 1 – 7	**Day 1:** Glucose + Electrolytes **Day 2–5:** Chick Starter Multivitamins **Day 7:** Newcastle (ND) B1	Natural daylight only. Fresh water twice daily.	Brooder temp at 33–35°C. Paper feeding for easy feed pickup.
Days 8 – 28	**Day 14:** Gumboro (IBD) Vaccine **Day 21:** ND LaSota Booster **Day 28:** Anti-coccidial treatment	Maintain 20% CP Starter. Clean, fresh water continuously.	Gradually decrease temperature 2°C weekly down to 22°C. Expand pen.
Weeks 7 – 15	**Week 9:** ND / IB Booster **Week 10:** Fowl Pox (Wing Web) **Week 12:** Deworming (Piperazine/Levamisole)	Switch to Grower Mash. Do not overfeed to prevent early fat deposition.	Monitor body weight uniformity. Aim for 1.2–1.4 kg target weight at Week 16.
Weeks 16 – 17	**Week 16:** Final ND + IB Oil Vaccine **Week 17:** Pre-lay multivitamins + Calcium	Switch to Pre-Lay Mash. Increase day length by 1hr/week.	Install nest boxes (1 box per 4 hens). Transfer hens to laying house.
Weeks 18 – 72+	**Every 8–10 Wks:** Layer-safe Deworming **Monthly:** Vitamin C + Electrolytes (Stress)	**16 Hours Light/Day**. Provide free-choice Oyster Shell grit.	Collect eggs 3x daily. Maintain clean nest box bedding to avoid dirty eggs.

CRITICAL SUCCESS FACTORS FOR COMMERCIAL LAYER OPERATIONS

- **Lighting Rule:** Never decrease light duration during the laying phase. 16 hours of light triggers hormone production needed for continuous egg laying.