

Lines on cubic surfaces, elliptic surfaces and the E_6 lattice

T. Shioda, *Weierstrass Transformations and Cubic Surfaces*

Enrique Acosta

Department of Mathematics
University of Arizona

May 2009

Cubic Surfaces

Theorem


Any smooth cubic surface in \mathbb{P}^3 contains exactly 27 lines.

Cubic Surfaces


Theorem

Any smooth cubic surface in \mathbb{P}^3 contains exactly 27 lines.

However...the lines are usually not defined over the reals.




5 real lines



7 real lines

Cubic Surfaces




The Clebsch Diagonal Cubic

$$x^3 + y^3 + z^3 + w^3 - (x + y + z + w)^3 = 0$$

Blowing up

- ▶ Given any smooth algebraic surface, replace a point by a line (the **Exceptional Line**): a copy of \mathbb{P}^1 .
- ▶ Each point on this line corresponds a tangent direction at the point.




Cubic Surfaces and blow up's

- ▶ Every smooth cubic in \mathbb{P}^3 is isomorphic to \mathbb{P}^2 blown up at six points (not all on a conic and no three on a line).
- ▶ The 27 lines are given by:

Cubic Surfaces and blow up's

- ▶ Every smooth cubic in \mathbb{P}^3 is isomorphic to \mathbb{P}^2 blown up at six points (not all on a conic and no three on a line).
- ▶ The 27 lines are given by:

6: The six exceptional lines.




Cubic Surfaces and blow up's

- ▶ Every smooth cubic in \mathbb{P}^3 is isomorphic to \mathbb{P}^2 blown up at six points (not all on a conic and no three on a line).
- ▶ The 27 lines are given by:

6: The six exceptional lines.


15: The strict transform of the lines connecting any two of the six points.



Cubic Surfaces and blow up's

- ▶ Every smooth cubic in \mathbb{P}^3 is isomorphic to \mathbb{P}^2 blown up at six points (not all on a conic and no three on a line).
- ▶ The 27 lines are given by:

- 6: The six exceptional lines.
- 15: The strict transform of the lines connecting any two of the six points.
- 6: The strict transform of the 6 (unique) conics through five of the six points.



Two natural questions

Question 1

Given a cubic smooth surface, ¿Can you find the equations of the 27 lines?

Two natural questions

Question 1

Given a cubic smooth surface, ¿Can you find the equations of the 27 lines?

Question 2

Given six points in \mathbb{P}^2 , ¿Can find the equation of the corresponding smooth cubic surface and the 27 lines in it?

Question 1:

¿Can you find the equations of the 27 lines?

Very difficult!...the lines are usually not defined over the field of definition of the surface.

Question 1:

Can you find the equations of the 27 lines?

Very difficult!...the lines are usually not defined over the field of definition of the surface.

Example (Shioda)

The minimum field extension of \mathbb{Q} where all the 27 lines of the cubic surface

$$y^2 + 2yz = x^3 + x + xz^2 + z + z^2 + 1$$

are defined is the splitting field of a polynomial of degree 27. The degree of this extension is 51,840.

Question 1:

¿Can you find the equations of the 27 lines?

Very difficult!...the lines are usually not defined over the field of definition of the surface.

Example (Shioda)

The minimum field extension of \mathbb{Q} where all the 27 lines of the cubic surface

$$y^2 + 2yz = x^3 + x + xz^2 + z + z^2 + 1$$

are defined is the splitting field of a polynomial of degree 27. The degree of this extension is 51,840.

More Generally (Harris)

There are no explicit equations for the 27 lines of a general cubic surface.

Question 2:

Given the six points you blow up, ¿ Can you find the equation of the resulting smooth cubic surface and the 27 lines in it?

Question 2: The Cubic resulting from the Blow-up

Question 2: The Cubic resulting from the Blow-up

- ▶ The space of plane cubics is 9-dimensional (10 projective coefficients).

Question 2: The Cubic resulting from the Blow-up

- ▶ The space of plane cubics is 9-dimensional (10 projective coefficients).
- ▶ Assuming the six points are not on a conic and no three are on a line, the space of cubics through the six points is 3-dimensional ($9 - 6 = 3$), i.e., the conditions are independent.

Question 2: The Cubic resulting from the Blow-up

- ▶ The space of plane cubics is 9-dimensional (10 projective coefficients).
- ▶ Assuming the six points are not on a conic and no three are on a line, the space of cubics through the six points is 3-dimensional ($9 - 6 = 3$), i.e., the conditions are independent.
- ▶ Let f_0, f_1, f_2, f_3 be a basis.

Question 2: The Cubic resulting from the Blow-up

- ▶ The space of plane cubics is 9-dimensional (10 projective coefficients).
- ▶ Assuming the six points are not on a conic and no three are on a line, the space of cubics through the six points is 3-dimensional ($9 - 6 = 3$), i.e., the conditions are independent.
- ▶ Let f_0, f_1, f_2, f_3 be a basis. The equation of any plane cubic through the six points is a linear combination of the f_i .

Question 2: The Cubic resulting from the Blow-up

- ▶ The space of plane cubics is 9-dimensional (10 projective coefficients).
- ▶ Assuming the six points are not on a conic and no three are on a line, the space of cubics through the six points is 3-dimensional ($9 - 6 = 3$), i.e., the conditions are independent.
- ▶ Let f_0, f_1, f_2, f_3 be a basis. The equation of any plane cubic through the six points is a linear combination of the f_i .
- ▶ Define the rational map $\mathbb{P}^2 \rightarrow \mathbb{P}^3$

$$[x : y : z] \mapsto [f_0(x, y, z) : \dots : f_3(x, y, z)]$$

Question 2: The Cubic resulting from the Blow-up

- ▶ The space of plane cubics is 9-dimensional (10 projective coefficients).
- ▶ Assuming the six points are not on a conic and no three are on a line, the space of cubics through the six points is 3-dimensional ($9 - 6 = 3$), i.e., the conditions are independent.
- ▶ Let f_0, f_1, f_2, f_3 be a basis. The equation of any plane cubic through the six points is a linear combination of the f_i .
- ▶ Define the rational map $\mathbb{P}^2 \rightarrow \mathbb{P}^3$

$$[x : y : z] \mapsto [f_0(x, y, z) : \dots : f_3(x, y, z)]$$

- ▶ The map is not defined at the six points since all the f_i give zero, but it extends to a map from the blowup of \mathbb{P}^2 at those points to \mathbb{P}^3 which is in fact an ISOMORPHISM.

Question 2: The Cubic resulting from the Blow-up

- ▶ The image is a cubic surface by arguments involving how many times a general line in \mathbb{P}^3 intersects the surface.

Question 2: The Cubic resulting from the Blow-up

- ▶ The image is a cubic surface by arguments involving how many times a general line in \mathbb{P}^3 intersects the surface.
- ▶ The equation for the surface is never given, and there is no apparent way to find it in general.

Question 2: The Cubic resulting from the Blow-up

- ▶ The image is a cubic surface by arguments involving how many times a general line in \mathbb{P}^3 intersects the surface.
- ▶ The equation for the surface is never given, and there is no apparent way to find it in general.

The Article

T. Shioda, *Weierstrass Transformations and Cubic Surfaces*, 1994.

- ▶ Explicit equation for the cubic in terms of the equations of the blown-up points.
- ▶ The construction involves the E_6 lattice and its dual E_6^*

The answer

- Let P_1, \dots, P_6 be six points in \mathbb{P}^2 not on a conic and no three on a line.

The answer

- ▶ Let P_1, \dots, P_6 be six points in \mathbb{P}^2 not on a conic and no three on a line.
- ▶ By a linear change of coordinates we can assume that they all lie on the cuspidal cubic $y^2 = x^3$.

The answer

- ▶ Let P_1, \dots, P_6 be six points in \mathbb{P}^2 not on a conic and no three on a line.
- ▶ By a linear change of coordinates we can assume that they all lie on the cuspidal cubic $y^2 = x^3$.
- ▶ The cuspidal cubic has a parametrization $(1/u^2, 1/u^3)$, so each P_i corresponds to a u_i .

The answer

- ▶ Let P_1, \dots, P_6 be six points in \mathbb{P}^2 not on a conic and no three on a line.
- ▶ By a linear change of coordinates we can assume that they all lie on the cuspidal cubic $y^2 = x^3$.
- ▶ The cuspidal cubic has a parametrization $(1/u^2, 1/u^3)$, so each P_i corresponds to a u_i .
- ▶ The condition on their configuration is equivalent to:
 - ▶ $u_i \neq u_j$ if $i \neq j$.
 - ▶ $\sum u_i \neq 0$
 - ▶ $u_i + u_j + u_k \neq 0$ for i, j, k distinct.

The answer

- ▶ Let P_1, \dots, P_6 be six points in \mathbb{P}^2 not on a conic and no three on a line.
- ▶ By a linear change of coordinates we can assume that they all lie on the cuspidal cubic $y^2 = x^3$.
- ▶ The cuspidal cubic has a parametrization $(1/u^2, 1/u^3)$, so each P_i corresponds to a u_i .
- ▶ The condition on their configuration is equivalent to:
 - ▶ $u_i \neq u_j$ if $i \neq j$.
 - ▶ $\sum u_i \neq 0$
 - ▶ $u_i + u_j + u_k \neq 0$ for i, j, k distinct.
- ▶ Let c_n be the n -th symmetric function in the u_i :
$$\prod (x - u_i) = x^6 - c_1 x^5 + c_2 x^4 + \dots + c_6$$

The answer

Let ε_n is the n -th symmetric function in the 27 forms:

$$a_i = \frac{c_1}{3} - u_i \quad a'_i = \frac{-2c_1}{3} - u_i \quad a''_{ij} = \frac{c_1}{3} - u_i - u_j$$

The answer

Let ε_n is the n -th symmetric function in the 27 forms:

$$a_i = \frac{c_1}{3} - u_i \quad a'_i = \frac{-2c_1}{3} - u_i \quad a''_{ij} = \frac{c_1}{3} - u_i - u_j$$

The equation of the cubic surface obtained by blowing up P_1, \dots, P_6 is

$$y^2 + 2y = x^3 + x(p_0 + p_1z + p_2z^2) + (q_0 + q_1z + q_2z^2)$$

where

The answer

Let ε_n is the n -th symmetric function in the 27 forms:

$$a_i = \frac{c_1}{3} - u_i \quad a'_i = \frac{-2c_1}{3} - u_i \quad a''_{ij} = \frac{c_1}{3} - u_i - u_j$$

The equation of the cubic surface obtained by blowing up P_1, \dots, P_6 is

$$y^2 + 2y = x^3 + x(p_0 + p_1z + p_2z^2) + (q_0 + q_1z + q_2z^2)$$

where $p_2 = \varepsilon_2/12$

The answer

Let ε_n is the n -th symmetric function in the 27 forms:

$$a_i = \frac{c_1}{3} - u_i \quad a'_i = \frac{-2c_1}{3} - u_i \quad a''_{ij} = \frac{c_1}{3} - u_i - u_j$$

The equation of the cubic surface obtained by blowing up P_1, \dots, P_6 is

$$y^2 + 2y = x^3 + x(p_0 + p_1z + p_2z^2) + (q_0 + q_1z + q_2z^2)$$

where

$$p_2 = \varepsilon_2/12$$

$$p_1 = \varepsilon_5/48$$

The answer

Let ε_n is the n -th symmetric function in the 27 forms:

$$a_i = \frac{c_1}{3} - u_i \quad a'_i = \frac{-2c_1}{3} - u_i \quad a''_{ij} = \frac{c_1}{3} - u_i - u_j$$

The equation of the cubic surface obtained by blowing up P_1, \dots, P_6 is

$$y^2 + 2y = x^3 + x(p_0 + p_1z + p_2z^2) + (q_0 + q_1z + q_2z^2)$$

where

$$p_2 = \varepsilon_2/12$$

$$p_1 = \varepsilon_5/48$$

$$q_2 = (\varepsilon_6 - 168p_2^3)/96$$

The answer

Let ε_n is the n -th symmetric function in the 27 forms:

$$a_i = \frac{c_1}{3} - u_i \quad a'_i = \frac{-2c_1}{3} - u_i \quad a''_{ij} = \frac{c_1}{3} - u_i - u_j$$

The equation of the cubic surface obtained by blowing up P_1, \dots, P_6 is

$$y^2 + 2y = x^3 + x(p_0 + p_1z + p_2z^2) + (q_0 + q_1z + q_2z^2)$$

where

$$p_2 = \varepsilon_2/12$$

$$p_1 = \varepsilon_5/48$$

$$q_2 = (\varepsilon_6 - 168p_2^3)/96$$

$$p_0 = (\varepsilon_8 - 294p_1p_2^2)/1344$$

The answer

Let ε_n is the n -th symmetric function in the 27 forms:

$$a_i = \frac{c_1}{3} - u_i \quad a'_i = \frac{-2c_1}{3} - u_i \quad a''_{ij} = \frac{c_1}{3} - u_i - u_j$$

The equation of the cubic surface obtained by blowing up P_1, \dots, P_6 is

$$y^2 + 2y = x^3 + x(p_0 + p_1z + p_2z^2) + (q_0 + q_1z + q_2z^2)$$

where

$$p_2 = \varepsilon_2/12$$

$$p_1 = \varepsilon_5/48$$

$$q_2 = (\varepsilon_6 - 168p_2^3)/96$$

$$p_0 = (\varepsilon_8 - 294p_1p_2^2)/1344$$

$$q_0 = (\varepsilon_{12} - 608p_1^2p_2 - \dots + 1248q_2^2)/17280$$

The answer

The equations for the 27 lines are also explicit:

$$x = az + b \quad \bigcap \quad y = dz + e$$

$$L_i: a = a_i.$$

$$L'_i: a = a'_i.$$

$$L'_{ij}: a = a'_{ij}.$$

$$b = \text{complicated expression in } c_1, c_2, c_3, c_4, u_i$$

In all cases:

$$d = (a^3 + ap_2)/2$$

$$e = (3a^2b - d^2 + ap_1 + bp_2 + q_2)/2$$

¿How?

How?

Theorem

The Mordell-Weil lattice of the elliptic curve

$$E : y^2 = x^3 + x(p_0 + p_1t + p_2t^2) + (q_0 + q_1t + q_2t^2 + t^4)$$

defined over $\mathbb{Q}(t)$ is isomorphic to E_6^* .

How?

Theorem

The Mordell-Weil lattice of the elliptic curve

$$E : y^2 = x^3 + x(p_0 + p_1t + p_2t^2) + (q_0 + q_1t + q_2t^2 + t^4)$$

defined over $\mathbb{Q}(t)$ is isomorphic to E_6^* . Moreover,

- The 54 minimal roots are given by 27 points $P = (x, y)$ of the form

$$x = at + b \quad y = t^2 + dt + e$$

and the 27 negatives of the points above $-P = (x, -y)$.

How?

Theorem

The Mordell-Weil lattice of the elliptic curve

$$E : y^2 = x^3 + x(p_0 + p_1t + p_2t^2) + (q_0 + q_1t + q_2t^2 + t^4)$$

defined over $\mathbb{Q}(t)$ is isomorphic to E_6^* . Moreover,

- ▶ The 54 minimal roots are given by 27 points $P = (x, y)$ of the form

$$x = at + b \quad y = t^2 + dt + e$$

and the 27 negatives of the points above $-P = (x, -y)$.

- ▶ The coefficients a, b, c, d can be given explicitly in terms of the p_i, q_j .

How?

Theorem

The Mordell-Weil lattice of the elliptic curve

$$E : y^2 = x^3 + x(p_0 + p_1t + p_2t^2) + (q_0 + q_1t + q_2t^2 + t^4)$$

defined over $\mathbb{Q}(t)$ is isomorphic to E_6^* . Moreover,

- ▶ The 54 minimal roots are given by 27 points $P = (x, y)$ of the form

$$x = at + b \quad y = t^2 + dt + e$$

and the 27 negatives of the points above $-P = (x, -y)$.

- ▶ The coefficients a, b, c, d can be given explicitly in terms of the p_i, q_j .
- ▶ These roots generate the Mordell-Weil group, and one can give six explicit points which generate the Mordell-Weil group.

How?

$$E : y^2 = x^3 + x(p_0 + p_1t + p_2t^2) + (q_0 + q_1t + q_2t^2 + t^4)$$

- ▶ The parameter t in the equation for E can be viewed as a coordinate and E itself as an affine surface.

How?

$$E : y^2 = x^3 + x(p_0 + p_1t + p_2t^2) + (q_0 + q_1t + q_2t^2 + t^4)$$

- ▶ The parameter t in the equation for E can be viewed as a coordinate and E itself as an affine surface.
- ▶ The map $y \mapsto y - t^2$ maps this surface to a cubic surface and the root vectors of the form $P = (at + b, t^2 + dt + e)$ to lines $(at + b, dt + e)$.

How?

$$E : y^2 = x^3 + x(p_0 + p_1t + p_2t^2) + (q_0 + q_1t + q_2t^2 + t^4)$$

- ▶ The parameter t in the equation for E can be viewed as a coordinate and E itself as an affine surface.
- ▶ The map $y \mapsto y - t^2$ maps this surface to a cubic surface and the root vectors of the form $P = (at + b, t^2 + dt + e)$ to lines $(at + b, dt + e)$.
- ▶ So we have a cubic surface and the 27 lines on it.

How?

$$E : y^2 = x^3 + x(p_0 + p_1t + p_2t^2) + (q_0 + q_1t + q_2t^2 + t^4)$$

- ▶ The parameter t in the equation for E can be viewed as a coordinate and E itself as an affine surface.
- ▶ The map $y \mapsto y - t^2$ maps this surface to a cubic surface and the root vectors of the form $P = (at + b, t^2 + dt + e)$ to lines $(at + b, dt + e)$.
- ▶ So we have a cubic surface and the 27 lines on it.
- ▶ Now relate this somehow to the blowup of six points in the plane....

The 28 bitangents

