




Prevalence and Characteristics of Asymmetric Hearing Loss in Korean Adults: A KNHANES 2022–2023 Analysis

PD 125

Jaeman PARK¹, Dongjun WOO², Jiyeon PARK², Gi Jung IM¹, Jae-Jun SONG¹, and Sung-Won CHAE¹


¹ Department of Otorhinolaryngology–Head and Neck Surgery, College of Medicine, Korea University, Seoul, South Korea

² College of Medicine, Korea University, Seoul, South Korea



INTRODUCTION

- Asymmetric Hearing Loss (AHL) is defined by interaural threshold difference.
- National epidemiologic data on AHL in Korean adults remain limited.
- This study evaluated prevalence and audiometric characteristics of AHL using **KNHANES 2022–2023**.
- This study aimed to estimate the **prevalence of AHL** in Korean adults aged ≥ 40 years and to **characterize** its age, frequency, laterality, and interaural difference patterns.



MATERIALS & METHODS

Definition of AHL

Absolute interaural difference ≥ 15 dB HL in 4-frequency pure-tone average (0.5, 1, 2, 4 kHz), based on the AAO–HNS criterion

Adults aged ≥ 40 years

- Exclusion: Negative 0.5, 1, 2, and 4 kHz hearing thresholds, type B or flat tympanogram

Final sample: N = 6574 [Weighted overall population: ~ 22.8 million]

AHL case subset: n = 447



RESULTS ► KEY FINDINGS



6.5%

Weighted prevalence of AHL



4 kHz

Most common asymmetry frequency



L > R

Left-sided AHL more common than right



15–20 dB HL

Largest category of interaural difference

Weighted Prevalence of AHL by Age Group (N = 6574)




Figure 1. Weighted prevalence of AHL increased with age overall. Men showed a higher prevalence than women. Dashed line indicates the total weighted prevalence of 6.5%

Weighted Mean Hearing Thresholds by Frequency (N = 6574)




Figure 2. Left-ear hearing thresholds were higher than right-ear hearing thresholds across all tested frequencies, with larger interaural differences at 2 and 4 kHz.

Frequency-Specific Asymmetry (n = 447)




Figure 3. Asymmetry was more common at higher frequencies. Proportions are not mutually exclusive and do not sum to 100%, because participants could have asymmetry at more than one frequency.

Laterality of AHL (n = 447)




Figure 4. Left-sided AHL was more common than right-sided AHL overall and in both sexes, indicating a consistent left-sided predominance across sex subgroups.

Interaural Differences by Age (n = 447)





Figure 5. Interaural differences did not differ significantly across age groups ($p = .530$), with most cases clustering within the 15–20 dB HL range. Horizontal bars and error bars indicate mean $\pm \frac{1}{2}$ SD.




CONCLUSION

In Korean adults aged ≥ 40 years, the weighted prevalence of AHL was **6.5%**. AHL prevalence increased with age overall and was higher in men than in women. AHL was characterized by **higher-frequency asymmetry**, **left-sided predominance**, and clustering near the lower end of the asymmetry range, with **most cases falling within the 15–20 dB HL range**. These findings indicate that AHL is most often expressed as relatively mild interaural asymmetry rather than marked asymmetry.



RESOURCES



SCAN FOR POSTER & REFERENCES

Access the full poster, and reference links via QR code

