

4D to 3D reduction of Seiberg duality for $SU(N)$ susy
gauge theories with adjoint matter: a partition
function approach

Carlo Sana

1 | Introduction

2 | Physics

2.1 Introduction

Supersymmetric quantum field theories enjoy an enlarged group of symmetries compared to other field theories. Since the symmetry group is a non trivial combination of internal and spacetime symmetries, they have many unexpected feature and new techniques were found to study them.

Many of the new tools found are available only for supersymmetric theories, making them the theater for exciting discoveries in physics.

2.2 Four dimensional dualities

BUUUUU

Seiberg Duality

This is the duality originally found by Seiberg [1].

POSSO DIRE ALTRO!!

The electric theory is a $SU(N_c)$ supersymmetric non-chiral gauge theory with N_f flavours global symmetry group $SU(N_f) \times SU(N_f) \times U(1)_B \times U(1)_R$. The charges of the matter content of the theory are summarized in the table below.

	$SU(N_c)$	$SU(N_f)_L$	$SU(N_f)_R$	$U(1)_B$	$U(1)_R$
Q	N_c	N_F	1	1	$\frac{N_f - N_c}{N_f}$
\tilde{Q}	$\overline{N_c}$	1	$\overline{N_F}$	-1	$\frac{N_f - N_c}{N_f}$

Table 2.1: Charge of matter content of the electric theory

The *R-Charge* is fixed by requiring that the *R-Symmetry* is non anomalous.

Da spiegare meglio.

What really happens is that $U(1)_A$ symmetry (which is anomalous) mixes with the classical (anomalous) $U(1)_{R'}$ R-symmetry. Their mixing result in a non anomalous $U(1)_R$ R-symmetry and the disappearance of $U(1)_A$.

The triangular graph corresponding to this anomaly constrains the R-charge of the quarks

imposing

$$R_{gaugino}T(\text{Ad}) + \sum_{\text{fermions } f} (R_f - 1)T(r) = 0$$

$$N_c + \frac{1}{2} 2N_f(R_Q - 1) = 0 \quad \rightarrow \quad R_Q = \frac{N_f - N_c}{N_f}$$

The magnetic theory is a theory with the same global symmetries as the electric theory, but the gauge group is now $SU(N_f - N_c)$ and in addition there are N_f^2 fields, that we will call mesons. Dual quarks will be represented as q, \tilde{q} and mesons as M_j^i . The charges for the magnetic theory are given by

	$SU(N_c)$	$SU(N_f)_L$	$SU(N_f)_R$	$U(1)_B$	$U(1)_R$
q	N_c	N_f	1	$\frac{N_c}{N-f-N_c}$	$\frac{N_c}{N_f}$
\tilde{q}	$\overline{N_c}$	1	$\overline{N_f}$	$-\frac{N_c}{N-f-N_c}$	$\frac{N_c}{N_f}$
M_j^i	1	N_f	$\overline{N_f}$	0	$2\frac{N_f-N_c}{N_f}$

Table 2.2: Charge of matter content of the magnetic theory

Kutasov-Schwimmer duality

2.3 3D dualities

Aharony duality

Kutasov-Schwimmer duality

3 | Math

4 | My work

5 | conclusions

6 | Appendix

Bibliography

- [1] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” *Nucl.Phys.*, vol. B435, pp. 129–146, 1995.
- [2] N. Seiberg, “The Power of holomorphy: Exact results in 4-D SUSY field theories,” 1994.
- [3] A. C. Davis, M. Dine, and N. Seiberg, “The Massless Limit of Supersymmetric QCD,” *Phys.Lett.*, vol. B125, p. 487, 1983.
- [4] K. A. Intriligator and N. Seiberg, “Lectures on supersymmetric gauge theories and electric - magnetic duality,” *Nucl.Phys.Proc.Suppl.*, vol. 45BC, pp. 1–28, 1996.
- [5] D. Kutasov, A. Schwimmer, and N. Seiberg, “Chiral rings, singularity theory and electric - magnetic duality,” *Nucl.Phys.*, vol. B459, pp. 455–496, 1996.
- [6] O. Aharony, S. S. Razamat, N. Seiberg, and B. Willett, “3d dualities from 4d dualities,” *JHEP*, vol. 1307, p. 149, 2013.
- [7] A. Amariti and C. Klare, “A journey to 3d: exact relations for adjoint SQCD from dimensional reduction,” 2014.
- [8] F. van de Bult, “Hyperbolic hypergeometric functions,” *Master thesis*, 2007.
- [9] F. Dolan and H. Osborn, “Applications of the Superconformal Index for Protected Operators and q-Hypergeometric Identities to N=1 Dual Theories,” *Nucl.Phys.*, vol. B818, pp. 137–178, 2009.
- [10] V. Spiridonov and G. Vartanov, “Elliptic Hypergeometry of Supersymmetric Dualities,” *Commun.Math.Phys.*, vol. 304, pp. 797–874, 2011.
- [11] K. Nii, “3d duality with adjoint matter from 4d duality,” *JHEP*, vol. 1502, p. 024, 2015.
- [12] H. Kim and J. Park, “Aharony Dualities for 3d Theories with Adjoint Matter,” *JHEP*, vol. 1306, p. 106, 2013.
- [13] F. Benini, C. Closset, and S. Cremonesi, “Comments on 3d Seiberg-like dualities,” *JHEP*, vol. 1110, p. 075, 2011.
- [14] C. Closset, T. T. Dumitrescu, G. Festuccia, Z. Komargodski, and N. Seiberg, “Comments on Chern-Simons Contact Terms in Three Dimensions,” *JHEP*, vol. 1209, p. 091, 2012.

- [15] A. Amariti and C. Klare, “Chern-Simons and RG Flows: Contact with Dualities,” *JHEP*, vol. 1408, p. 144, 2014.
- [16] O. Aharony, A. Hanany, K. A. Intriligator, N. Seiberg, and M. Strassler, “Aspects of N=2 supersymmetric gauge theories in three-dimensions,” *Nucl.Phys.*, vol. B499, pp. 67–99, 1997.