

4D to 3D reduction of Seiberg duality for $SU(N)$ susy
gauge theories with adjoint matter: a partition
function approach

Carlo Sana

1 | Physics

—— INTRODUCTION OUTLINE ——

- ~ More symmetry = more tools for studying theories
- ~ State structure: multiplet & superspace
- ~ Milder divergences
- ~ Renormalization constraints
- ~ Non renormalization theorems (perturbative)
- ~ Holomorphicity, couplings as background fields
(important smoothness of weak coupling limits, e.g. classic limit g in well defined.)
- ~ Exact results (superpotential, witten index, exact beta function)
- ~ Use of wilsonian action: no IR divergences)
- ~ Superconformal index
- ~ Localization
- ~ Moduli space

1.1 Introduction

Supersymmetric quantum field theories enjoy an enlarged group of symmetries compared to other field theories. Since the symmetry group is a non trivial combination of internal and spacetime symmetries, they have many unexpected features and new techniques were found to study them. Almost all of the new tools found are available only for supersymmetric field theories, making them the theater for many advances in physics.

In this section we will analyse the features of supersymmetric field theories that are crucial to the discovery of electric magnetic duality and its generalisations.

States and their representation

Being a symmetry between bosons and fermions, supersymmetry imposes that states are organized in multiplets containing different representations of the *Lorentz group* i.e different type of particles. Various multiplets exist and their properties depend on the number of supercharges of the theory and on what they represent e.g matter, glue or gravity.

We will introduce the multiplets that can be defined for $4d \mathcal{N} = 1$ theories and only later we will explain the differences with $3d \mathcal{N} = 2$ theories. For four dimensional theories, we can define two different multiplet that are invariant under supersymmetry transformation.

The matter or chiral multiplet contains a complex scalar (*squark*) and a Weyl fermion. It identifies the matter content of the theory.

The vector or gauge multiplet contains a Weyl fermion (*gaugino*) and a vector.

Particles in the same multiplet transform in the same representation of global or gauge symmetries. For this reason the gaugino cannot represent matter.

A representation of these multiplets on fields can be easily found using the *superspace* formalism. In this formalism it is possible to represent fields that are *off-shell*, in contrast with multiplets that we introduced previously that are *on-shell* since they represent states in Hilbert space.

Matter and gauge multiplets are represented by (anti)chiral and real superfields respectively.

General renormalization properties

A remarkable feature of supersymmetry is the constraint that the additional symmetry imposes on the renormalization properties of the theories.

One of the first aspects that brought attention to supersymmetry was that divergences of loop diagrams were milder because of the cancellation between diagrams with bosons and fermions running in the loops.

Nowadays we know powerful theorems that restrict the behaviour of supersymmetric field theories under the process of renormalization. In order to preserve supersymmetry, the renormalization process has to preserve the Hilbert space structure. For example the wave function renormalization of different *particles* inside a multiplet must be the same, otherwise the renormalized lagrangian is not supersymmetric invariant anymore.

Moreover, in the supersymmetry algebra P^2 is still a Casimir operator i.e. it commutes with every operator in the algebra: particles in the same multiplet must have the same mass. Renormalization cannot break this condition, otherwise it would break supersymmetry.

For a *Super Yang Mills* theory with $\mathcal{N} = 1$ we have the additional requirement that gV , where g is the coupling and V is the vector superfield, cannot be renormalized by symmetry considerations.

Adding more supersymmetry the wave function renormalization of the various field are even more constrained by symmetry. For example, for $\mathcal{N} = 4$ *SYM* the fields and the coupling are not renormalized at all.

Superpotential: holomorphy and non-renormalization

Other than renormalization constraints, supersymmetry provides non-renormalization theorems for certain objects, such as the superpotential. In [1] was demonstrated that the superpotential is tree-level exact, i.e. it does not receive correction in perturbation theory. However it usually receive contributions from non perturbative dynamics. It was demonstrated using *supergraph* calculations that allow to work directly with superfields using Feynman graph, keeping supersymmetry explicit. Cancellations between loops with fermions and bosons are trivial and many simplifications occur during calculations.

The demonstration is based on the fact that for general supersymmetric field theories, supergraph loops diagrams yield a term that can be written in the form

$$\int d^4x_1 \dots d^4x_n d^2\theta d^2\bar{\theta} G(x_1, \dots, x_n) F_1(x_1, \theta, \bar{\theta}) \dots F_n(x_n, \theta, \bar{\theta}) \quad (1.1)$$

where $G(x_1, \dots, x_n)$ is translationally invariant function.

The importance of this result is that is given by a single integral over full superspace ($d^2\theta d^2\bar{\theta}$) whereas the superpotential must be written as an integral in half-superspace ($d^2\theta$ only) of chiral fields. Exploiting the fact that a product of chiral fields is a chiral field, the most general form of a superpotential is the following

$$W(\lambda, \Phi) = \sum_{n=1}^{\infty} \left(\int d^2\theta \lambda_n \Phi^n + \int d^2\bar{\theta} \lambda_n^\dagger \bar{\Phi}^n \right) \quad (1.2)$$

The second term of the superpotential is added in order to give a real lagrangian after the integration in superspace. From the definition, we can see that the superpotential is holomorphic: it does not depend on conjugate fields or coupling constants.

Fifteen years later, Seiberg [2] provided a proof of this theorem using a different approach. He noted that the coupling constants λ_n can be treated as background fields, i.e. chiral superfields with no dynamics.

Using this "trick" we can assign transformation laws to the fields and the coupling constants: in this way the superpotential acquires a bigger symmetry under which the effective lagrangian has to be invariant. In the limit of weak coupling, the effective superpotential must flow to the tree-level superpotential. The holomorphicity condition of the superpotential in addition to these requirements, force the effective superpotential to be equal to the tree-level superpotential.

Exact results

Bibliography

- [1] M. T. Grisaru, W. Siegel, and M. Rocek, “Improved Methods for Supergraphs,” *Nucl.Phys.*, vol. B159, p. 429, 1979.
- [2] N. Seiberg, “Naturalness versus supersymmetric nonrenormalization theorems,” *Phys.Lett.*, vol. B318, pp. 469–475, 1993.
- [3] N. Seiberg, “The Power of holomorphy: Exact results in 4-D SUSY field theories,” 1994.
- [4] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” *Nucl.Phys.*, vol. B435, pp. 129–146, 1995.
- [5] A. C. Davis, M. Dine, and N. Seiberg, “The Massless Limit of Supersymmetric QCD,” *Phys.Lett.*, vol. B125, p. 487, 1983.
- [6] K. A. Intriligator and N. Seiberg, “Lectures on supersymmetric gauge theories and electric - magnetic duality,” *Nucl.Phys.Proc.Suppl.*, vol. 45BC, pp. 1–28, 1996.
- [7] D. Kutasov, A. Schwimmer, and N. Seiberg, “Chiral rings, singularity theory and electric - magnetic duality,” *Nucl.Phys.*, vol. B459, pp. 455–496, 1996.
- [8] O. Aharony, S. S. Razamat, N. Seiberg, and B. Willett, “3d dualities from 4d dualities,” *JHEP*, vol. 1307, p. 149, 2013.
- [9] A. Amariti and C. Klare, “A journey to 3d: exact relations for adjoint SQCD from dimensional reduction,” 2014.
- [10] F. van de Bult, “Hyperbolic hypergeometric functions,” *Master thesis*, 2007.
- [11] F. Dolan and H. Osborn, “Applications of the Superconformal Index for Protected Operators and q-Hypergeometric Identities to N=1 Dual Theories,” *Nucl.Phys.*, vol. B818, pp. 137–178, 2009.
- [12] V. Spiridonov and G. Vartanov, “Elliptic Hypergeometry of Supersymmetric Dualities,” *Commun.Math.Phys.*, vol. 304, pp. 797–874, 2011.
- [13] K. Nii, “3d duality with adjoint matter from 4d duality,” *JHEP*, vol. 1502, p. 024, 2015.
- [14] H. Kim and J. Park, “Aharony Dualities for 3d Theories with Adjoint Matter,” *JHEP*, vol. 1306, p. 106, 2013.

- [15] F. Benini, C. Closset, and S. Cremonesi, “Comments on 3d Seiberg-like dualities,” *JHEP*, vol. 1110, p. 075, 2011.
- [16] C. Closset, T. T. Dumitrescu, G. Festuccia, Z. Komargodski, and N. Seiberg, “Comments on Chern-Simons Contact Terms in Three Dimensions,” *JHEP*, vol. 1209, p. 091, 2012.
- [17] A. Amariti and C. Klare, “Chern-Simons and RG Flows: Contact with Dualities,” *JHEP*, vol. 1408, p. 144, 2014.
- [18] O. Aharony, A. Hanany, K. A. Intriligator, N. Seiberg, and M. Strassler, “Aspects of N=2 supersymmetric gauge theories in three-dimensions,” *Nucl.Phys.*, vol. B499, pp. 67–99, 1997.