

4D to 3D reduction of Seiberg duality for $SU(N)$ susy
gauge theories with adjoint matter: a partition
function approach

Carlo Sana

1 | Introduction

2 | Physics

2.1 General features of Seiberg-like dualities

2.2 Four dimensional dualities

BUUUUU

Seiberg Duality

This is the duality originally found by Seiberg [1].

POSSO DIRE ALTRO!!

The electric theory is a $SU(N_c)$ supersymmetric non-chiral gauge theory with N_f flavours global symmetry group $SU(N_f) \times SU(N_f) \times U(1)_B \times U(1)_R$. The charges of the matter content of the theory are summarized in the table below.

	$SU(N_c)$	$SU(N_f)_L$	$SU(N_f)_R$	$U(1)_B$	$U(1)_R$
Q	N_c	N_F	1	1	$\frac{N_f - N_c}{N_f}$
\tilde{Q}	$\overline{N_c}$	1	$\overline{N_F}$	-1	$\frac{N_f - N_c}{N_f}$

Table 2.1: Charge of matter content of the electric theory

The *R-Charge* is fixed by requiring that the *R-Symmetry* is non anomalous.

Da spiegare meglio.

What really happens is that $U(1)_A$ symmetry (which is anomalous) mixes with the classical (anomalous) $U(1)_{R'}$ R-symmetry. Their mixing result in a non anomalous $U(1)_R$ R-symmetry and the disappearance of $U(1)_A$.

The triangular graph corresponding to this anomaly constrains the R-charge of the quarks imposing

$$R_{gaugino}T(\text{Ad}) + \sum_{\text{fermions } f} (R_f - 1)T(r) = 0$$

$$N_c + \frac{1}{2} 2N_f(R_Q - 1) = 0 \quad \rightarrow \quad R_Q = \frac{N_f - N_c}{N_f}$$

The magnetic theory is a theory with the same global symmetries as the electric theory, but the gauge group is now $SU(N_f - N_c)$ and in addition there are N_f^2 fields, that we will call mesons. Dual quarks will be represented as q, \tilde{q} and mesons as M_j^i . The charges for the magnetic theory are given by

	$SU(N_c)$	$SU(N_f)_L$	$SU(N_f)_R$	$U(1)_B$	$U(1)_R$
q	N_c	N_F	1	$\frac{N_c}{N-f-N_c}$	$\frac{N_c}{N_f}$
\tilde{q}	$\overline{N_c}$	1	$\overline{N_F}$	$-\frac{N_c}{N-f-N_c}$	$\frac{N_c}{N_f}$
M_j^i	1	N_f	$\overline{N_f}$	0	$2\frac{N_f-N_c}{N_f}$

Table 2.2: Charge of matter content of the magnetic theory

Kutasov-Schwimmer duality

3 | Physics

3.1 General features of Seiberg-like dualities

3.2 Four dimensional dualities

BUUUUU

Seiberg Duality

This is the duality originally found by Seiberg [1].

POSSO DIRE ALTRO!!

The electric theory is a $SU(N_c)$ supersymmetric non-chiral gauge theory with N_f flavours global symmetry group $SU(N_f) \times SU(N_f) \times U(1)_B \times U(1)_R$. The charges of the matter content of the theory are summarized in the table below.

	$SU(N_c)$	$SU(N_f)_L$	$SU(N_f)_R$	$U(1)_B$	$U(1)_R$
Q	N_c	N_F	1	1	$\frac{N_f - N_c}{N_f}$
\tilde{Q}	$\overline{N_c}$	1	$\overline{N_F}$	-1	$\frac{N_f - N_c}{N_f}$

Table 3.1: Charge of matter content of the electric theory

The *R-Charge* is fixed by requiring that the *R-Symmetry* is non anomalous.

Da spiegare meglio.

What really happens is that $U(1)_A$ symmetry (which is anomalous) mixes with the classical (anomalous) $U(1)_{R'}$ R-symmetry. Their mixing result in a non anomalous $U(1)_R$ R-symmetry and the disappearance of $U(1)_A$.

The triangular graph corresponding to this anomaly constrains the R-charge of the quarks imposing

$$R_{gaugino}T(\text{Ad}) + \sum_{\text{fermions } f} (R_f - 1)T(r) = 0$$

$$N_c + \frac{1}{2} 2N_f(R_Q - 1) = 0 \quad \rightarrow \quad R_Q = \frac{N_f - N_c}{N_f}$$

The magnetic theory is a theory with the same global symmetries as the electric theory, but the gauge group is now $SU(N_f - N_c)$ and in addition there are N_f^2 fields, that we will call mesons. Dual quarks will be represented as q, \tilde{q} and mesons as M_j^i . The charges for the magnetic theory are given by

	$SU(N_c)$	$SU(N_f)_L$	$SU(N_f)_R$	$U(1)_B$	$U(1)_R$
q	N_c	N_F	1	$\frac{N_c}{N-f-N_c}$	$\frac{N_c}{N_f}$
\tilde{q}	$\overline{N_c}$	1	$\overline{N_F}$	$-\frac{N_c}{N-f-N_c}$	$\frac{N_c}{N_f}$
M_j^i	1	N_f	$\overline{N_f}$	0	$2\frac{N_f-N_c}{N_f}$

Table 3.2: Charge of matter content of the magnetic theory

Kutasov-Schwimmer duality

3.3 3D dualities

Aharony duality

Kutasov-Schwimmer duality

4 | Math

5 | My work

6 | conclusions

7 | Appendix

Bibliography

- [1] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” *Nucl.Phys.*, vol. B435, pp. 129–146, 1995.
- [2] N. Seiberg, “The Power of holomorphy: Exact results in 4-D SUSY field theories,” 1994.
- [3] A. C. Davis, M. Dine, and N. Seiberg, “The Massless Limit of Supersymmetric QCD,” *Phys.Lett.*, vol. B125, p. 487, 1983.
- [4] K. A. Intriligator and N. Seiberg, “Lectures on supersymmetric gauge theories and electric - magnetic duality,” *Nucl.Phys.Proc.Suppl.*, vol. 45BC, pp. 1–28, 1996.
- [5] D. Kutasov, A. Schwimmer, and N. Seiberg, “Chiral rings, singularity theory and electric - magnetic duality,” *Nucl.Phys.*, vol. B459, pp. 455–496, 1996.
- [6] O. Aharony, S. S. Razamat, N. Seiberg, and B. Willett, “3d dualities from 4d dualities,” *JHEP*, vol. 1307, p. 149, 2013.
- [7] A. Amariti and C. Klare, “A journey to 3d: exact relations for adjoint SQCD from dimensional reduction,” 2014.
- [8] F. van de Bult, “Hyperbolic hypergeometric functions,” *Master thesis*, 2007.
- [9] F. Dolan and H. Osborn, “Applications of the Superconformal Index for Protected Operators and q-Hypergeometric Identities to N=1 Dual Theories,” *Nucl.Phys.*, vol. B818, pp. 137–178, 2009.
- [10] V. Spiridonov and G. Vartanov, “Elliptic Hypergeometry of Supersymmetric Dualities,” *Commun.Math.Phys.*, vol. 304, pp. 797–874, 2011.
- [11] K. Nii, “3d duality with adjoint matter from 4d duality,” *JHEP*, vol. 1502, p. 024, 2015.
- [12] H. Kim and J. Park, “Aharony Dualities for 3d Theories with Adjoint Matter,” *JHEP*, vol. 1306, p. 106, 2013.
- [13] F. Benini, C. Closset, and S. Cremonesi, “Comments on 3d Seiberg-like dualities,” *JHEP*, vol. 1110, p. 075, 2011.
- [14] C. Closset, T. T. Dumitrescu, G. Festuccia, Z. Komargodski, and N. Seiberg, “Comments on Chern-Simons Contact Terms in Three Dimensions,” *JHEP*, vol. 1209, p. 091, 2012.

- [15] A. Amariti and C. Klare, “Chern-Simons and RG Flows: Contact with Dualities,” *JHEP*, vol. 1408, p. 144, 2014.
- [16] O. Aharony, A. Hanany, K. A. Intriligator, N. Seiberg, and M. Strassler, “Aspects of N=2 supersymmetric gauge theories in three-dimensions,” *Nucl.Phys.*, vol. B499, pp. 67–99, 1997.