

Armstrong Topology Solutions

George Blikas
Gregory Grant
David Hong
Grant Baker
Kristy Ong

April 2016

Contents

0	Preliminaries	7
0.1	Definitions	7
0.1.1	Functions	7
0.2	Extra Proofs and Lemmas	7
0.2.1	Group Theory & Abstract Algebra	7
0.2.2	Limit Points, Closure & Density	8
0.2.3	Separation	9
0.2.4	Compactness	10
0.2.5	Connectedness	10
0.2.6	Quotient Maps & Quotient Topology	11
0.2.7	Topological Groups	12
0.2.8	Homotopy Type	13
1	Introduction	17
1.1	Euler's Theorem	17
1.2	Topological Equivalence	17
1.3	Surfaces	17
1.4	Abstract Spaces	17
1.5	A Classification Theorem	17
1.6	Topological Invariants	17
2	Continuity	27
2.1	Open and Closed Sets	27
2.2	Continuous Functions	32
2.3	Space-Filling Curves	36
2.4	The Tietze Extension Theorem	37
3	Compactness and Connectedness	41
3.1	Closed and Bounded Subsets of \mathbb{E}^n	41
3.2	The Heine-Borel Theorem	41
3.3	Properties of Compact Spaces	43
3.4	Product Spaces	50
3.5	Connectedness	54
3.6	Joining Points by Paths	57

4	Identification Spaces	59
4.1	Constructing the Möbius Strip	59
4.2	The Identification Topology	59
4.3	Topological Groups	66
4.4	Orbit Spaces	71
5	The Fundamental Group	77
5.1	Homotopic Maps	77
5.2	Construction of the Fundamental Group	80
5.3	Calculations	83
5.4	Homotopy Type	86
5.5	Brouwer Fixed-Point Theorem	91
5.6	Separation of the Plane	95
5.7	The Boundary of a Surface	97
6	Triangulations	99
6.1	Triangulating Spaces	99
6.2	Barycentric Subdivision	106
6.3	Simplicial Approximation	106
6.4	The Edge Group of a Complex	107
6.5	Triangulating Orbit Spaces	108
6.6	Infinite Complexes	109
7	Surfaces	111
7.1	Classification	111
7.2	Triangulation and Orientation	112
7.3	Euler Characteristics	113
7.4	Surgery	114
7.5	Surface Symbols	114
8	Simplicial Homology	117
8.1	Cycles and Boundaries	117
8.2	Homology Groups	117
8.3	Examples	117
8.4	Simplicial Maps	117
8.5	Stellar Subdivision	117
8.6	Invariance	117

Acknowledgements

George B: Much thanks to all the contributors of this book. Special thanks goes to Gregory Grant, and particularly in helping establish chapter 1, section 3.4, and problems 10, 11, and 12 of section 4.2, all of which greatly sped-up the processes of this project.

Chapter 0

Preliminaries

0.1 Definitions

0.1.1 Functions

(1) Let $f : A \rightarrow B$ be a function. Then,

- f is onto if, and only if, there exists a function $g : B \rightarrow A$ such that $fg = 1_B$.
- f is one-to-one if, and only if, there exists a function $g : B \rightarrow A$ such that $gf = 1_A$ (provided A is non-empty).
- f is in 1-1 correspondence if there exists a function $g : B \rightarrow A$ such that $fg = 1_B$ and $gf = 1_A$. In this case, g is unique and is called the inverse function of f , typically denoted f^{-1} .

0.2 Extra Proofs and Lemmas

0.2.1 Group Theory & Abstract Algebra

(1) Split Exact Sequence: Given an exact sequence of abelian groups and homomorphisms,

$$0 \rightarrow G \xrightarrow{\theta} H \xrightarrow{\phi} K \rightarrow 0,$$

and a homomorphism $\psi : K \rightarrow H$, such that $\phi\psi \equiv 1_K$, we have $H \cong G \oplus K$.

Proof. Define $\alpha : G \oplus K \rightarrow H$ by $\alpha(g \oplus k) = \theta(g) + \psi(k)$: it is easy to see that α is a homomorphism. Also, α is (1-1), for if $\alpha(g \oplus k) = 0$, we have

$$0 = \phi(\theta(g) + \psi(k)) = \phi\psi(k) = k;$$

but, then $\theta(g) = 0$, so that $g = 0$ since θ is one-to-one.

Moreover, α is onto, since given $h \in H$, we have

$$\phi(h - \psi\phi(h)) = \phi(h) - \phi\psi\phi(h) = 0$$

Thus, there exists $g \in G$ such that $h - \psi\phi(h) = \theta(g)$, that is,

$$h = \theta(g) + \psi\phi(h) = \alpha(g \oplus \psi(h))$$

□

0.2.2 Limit Points, Closure & Density

- (1) Let (X, τ) be a topological space, and $A \subset X$. We show that $\overline{A} = A \cup L_A$, where L_A is the set of all accumulation points of A :

Proof. If $\overline{A} = A$, then by Theorem 2.2, pg. 29, $L_A \subset A$ and we are done. So, suppose that A is not closed. Then, A does not contain all of its limit points; consequently, $L_A \neq \emptyset$. Further, the set $B = A \cup L_A$ is closed by Theorem 2.2, pg. 29.

To conclude the proof, we show that any closed set C containing A , contains B . Indeed; Let $C \subset X$ be closed, such that $A \subset C$. Now, for $a \in L_A$ and open set O_a containing a , we have that $A \cap (O_a - \{a\}) \neq \emptyset$. But, as $A \subset C$, we have $C \cap (O_a - \{a\}) \neq \emptyset$. Therefore, $a \in C$. So, $L_A \subset C$, implying $A \cup L_A \subset C$. By Theorem 2.3, pg. 30, $\overline{A} = B = A \cup L_A$. □

- (2) Let (X, τ) be a topological space, and $A \subset X$. We prove that L_A , defined above, contains all limit points of sequences contained in A :¹

Proof. Let $\{a_n\}_{n=1}^\infty \subset A$, such that $a_n \rightarrow a$. Then, $a \in L_{\{a_n : n \in \mathbb{N}\}}$ and every neighbourhood of a contains a point of $\{a_n : n \in \mathbb{N}\} - \{a\} = B$. As $\{a_n : n \in \mathbb{N}\} \subset A$, every neighbourhood of a has a point in $A - \{a\}$; $a \in L_A$. □

- (3) Let (X, τ) be a topological space, and $A \subset X$ such that $\overline{A} = X$. We prove that $A \cap O \neq \emptyset$, for all $O \neq \emptyset$, $O \in \tau$:

Proof. Suppose that for some $O \in \tau$, $O \neq \emptyset$, $A \cap O = \emptyset$. By a previous lemma, we have $\overline{A} = (L_A - A) \cup O^c$, implying $O^c = A$. But then we have $X = O^c = \overline{O^c} = \overline{A}$, as O is open. But, contrarily, this implies that $(O^c)^c = O = X^c = \emptyset$. □

- (4) We prove that the intersection of a closed set and a compact set is always compact:

¹This assumes the general definition of limit points of a set

Proof. Let (X, τ) be a topological space. Let $H, K \subset X$, such that H is closed and K is compact. Consider $H \cap K$. Now, if $\{O_\alpha\}_\alpha$ is an open cover of $H \cap K$, then $K \subset \bigcup_{\alpha \in N} O_\alpha \cup (X - H)$. But, since H is closed, $X - H$ is open. In conclusion, as

$$H \cap K \subset \bigcup_{\alpha \in N} O_\alpha \cup (X - H)$$

such a finite subcover of $H \cap K$ exists. \square

- (5) We prove that if (X, d) is a metric space with the induced topology, then $C \subset X$ is closed if, and only if, whenever $\{a_n\}_{n=1}^\infty$ is a sequence in C , with $\{a_n\} \rightarrow L$, we have $L \in C$:

Proof.

(\implies): Suppose, to the contrary, that $\overline{C} = C$, but $L \notin C$. Thus, there exists some $\epsilon > 0$, such that $B_\epsilon(L) \cap C = \emptyset$, as L is not a limit point of C . But then, $a_n \notin B_\epsilon(L)$, for all $n \geq N$, $N \in \mathbb{N}$ is sufficiently large; A contradiction.

(\impliedby): Suppose, to the contrary, that $\overline{C} \neq C$. Then, by extra lemma², there is some $l \in L_C$, such that $L \notin C$. Thus, for each $n \in \mathbb{N}$, we pick $a_n \in B_{1/n}(L) \cap C \neq \emptyset$. Consequently, $\{a_n\}_{n=1}^\infty$ is a sequence in C such that $a_n \rightarrow L \in C$, by hypothesis; A contradiction. \square

0.2.3 Separation

- (1) We show that a compact T_2 space T_3 . Consequently, we show that it is T_4 :

Proof. Let X be the compact T_2 space. We first show that X is T_3 :

Let $A \subset X$, such that A is closed. Then A is compact. Further, let $b \in X$ such that $b \notin A$. Now, for each $a \in A$, there exists an open set O_a , and some open set O_b^a , such that $O_a \cap O_b^a = \emptyset$. It follows that $A \subset \bigcup_{a \in N} O_a \subset \bigcup_a O_a$ and that $\bigcup_{a \in N} O_a = O_A$ is open. Further, $b \in \bigcap_{a \in N} O_b^a = O_b$, is open. And by construction, we have $O_A \cap O_b = \emptyset$. So, X is T_3 .

To show that X is T_4 , let $A, B \subset X$, be disjoint and closed. Then, arguing as above, we have two disjoint open subsets $O_B \in \tau$, $O_A \in \tau$ with $O_A \cap O_B = \emptyset$. \square

²reference this

0.2.4 Compactness

- (1) We show that a homeomorphism between locally compact T_2 spaces, X, Y , extends to a homeomorphism between the Alexandroff compactifications; in other-words, locally compact homeomorphic T_2 spaces have homeomorphic one point compactifications.

Proof. Suppose that $f : X \rightarrow Y$ is the homeomorphism. Define $g : X \cup \{\infty_1\} \rightarrow Y \cup \{\infty_2\}$ as follows:

$$g(x) = \begin{cases} f(x) & x \neq \infty_1 \\ \infty_2 & x = \infty_1 \end{cases}$$

It is clear that as f is a bijection, so is g . We show that g is a homeomorphism:

Suppose that U is open in Y . Then, $g^{-1}[U] = f^{-1}[U]$, which is open in $X \cup \{\infty_1\}$. Now, if \mathcal{O} is open in $Y \cup \{\infty_2\}$, we have $\mathcal{O} = K \cup \{\infty_2\}$, where $K \subset Y$ is compact. Then,

$$g^{-1}[\mathcal{O}] = g^{-1}[K \cup \{\infty_2\}] = g^{-1}[K] \cup g^{-1}[\{\infty_2\}] = g^{-1}[K] \cup \{\infty_1\}$$

which is open in $X \cup \{\infty_1\}$.

Consequently, as g is one-to-one and onto, since f is, g is a homeomorphism by Theorem 3.7, pg. 48. \square

- (2) We show that if (X, τ) is a topological space, and $S \subset X$ is compact, then if W is a neighbourhood of S , there exists another neighbourhood G , such that

$$S \subset G \subset W$$

Proof. By exercise 21, section 3 pg. 55, and the fact that $X \cup \{y\} \cong X$, the result follows. \square

0.2.5 Connectedness

- (1) We show that if X is locally connected, then every connected component of X is open in X ; hence X is the disjoint union of its connected components:

Proof. Let $x \in Y$, where Y is a connected component of X . By definition, x is contained in some open connected subset U of X . Since Y is a maximal connected set containing x , we have $x \in U \subset Y$. This shows that Y is open in X . \square

0.2.6 Quotient Maps & Quotient Topology

- (1) We show that if $q : X \rightarrow Y$ is a quotient map³, then the topology of Y is the largest which makes q continuous:

Proof. Suppose that τ was some other topology on Y , such that q was continuous. We conclude by showing that $\tau \subset \tau_Y$:

Indeed; if $\mathcal{O} \in \tau$, then $q^{-1}[\mathcal{O}]$ is open in X . But then, $q[q^{-1}[\mathcal{O}]] = \mathcal{O} \in \tau_Y$. Thus, $\tau \subset \tau_Y$. \square

- (2) Suppose that (X, τ) is a topological space, and A a non-empty set. We show that if $f : X \rightarrow A$, then the quotient topology on A , τ_A is indeed a topology:

Proof. By definition, as noted above, \mathcal{O} is open in A , if $f^{-1}[\mathcal{O}]$ is open in X :

(a) As $f^{-1}[\emptyset] = \emptyset$, and $f^{-1}[A] = X$, $\emptyset, A \in \tau_A$.

(b) As

$$f^{-1}\left[\bigcup_{j \in J} \mathcal{O}_j\right] = \bigcup_{j \in J} f^{-1}[\mathcal{O}_j],$$

we see that $\bigcup_{j \in J} \mathcal{O}_j \in \tau_A$.

(c) Lastly,

$$f^{-1}\left[\bigcap_{j=1}^{\infty} \mathcal{O}_j\right] = \bigcap_{j=1}^{\infty} f^{-1}[\mathcal{O}_j]$$

So, $\bigcap_{j=1}^{\infty} \mathcal{O}_j \in \tau_A$.

\square

- (3) We show that projective real n -space, \mathbb{P}^n , is Hausdorff⁴:

Proof. Let $q : S^n \rightarrow \mathbb{P}^n$ be the quotient map, and suppose that $u, v \in \mathbb{P}^n$, $u \neq v$; there are such $x, y \in S^n$, such that $q^{-1}[\{u\}] = \{x, -x\}$, and $q^{-1}[\{v\}] = \{y, -y\}$. We exhibit an $\epsilon \in \mathbb{R}$ such that the open neighbourhoods of radius ϵ , centered at u, v are disjoint:

Let

$$\epsilon = \frac{1}{2} \min\{\|x - y\|, \|x + y\|\},$$

³Armstrong does not do a good job describing what the topology on Y is. A simple exercise shows that by letting \mathcal{O} be open in Y whenever $q^{-1}[\mathcal{O}]$ is open in X , we have a topology on Y ; call this τ_Y . Further, Armstrong does not do an adequate job describing what a quotient map is: $q : X \rightarrow Y$ is a quotient map if it is onto, continuous with respect to τ_Y , and such that $g^{-1}[\mathcal{O}]$ is open in X implies \mathcal{O} open in Y . We can summarize by saying that q is onto, and such that \mathcal{O} is open in Y iff $g^{-1}[\mathcal{O}]$ is open in X .

⁴Credit for the initial idea of the proof goes to Brain M. Scott. In addition, we use the notation $B(z, z')$ for the open ball of radius z' , centered at z .

and set $U = B(x, \epsilon)$, and $V = B(y, \epsilon)$. It follows from the construction of U, V , that $U, V, -U, -V$ are pairwise disjoint, and open neighbourhoods of $x, y, -x, -y \in S^n$. Moreover, as noted above, $q^{-1}(q[U]) = U \cup -U$, and $q^{-1}(q[V]) = V \cup -V$. To conclude, we show that $q[U] \cap q[V]$ are disjoint neighbourhoods for $u, v \in \mathbb{P}^n$.

Indeed;

$$\begin{aligned} q[U] \cap q[V] &= q[B(x, \epsilon) \cap S^n] \cap q[B(y, \epsilon) \cap S^n] \\ &= q[B(x, \epsilon)] \cap q[B(y, \epsilon)] \cap \mathbb{P}^n \\ &= B(u, q(\epsilon)) \cap B(v, q(\epsilon)) \cap \mathbb{P}^n \\ &= B_1 \cap B_2 \cap \mathbb{P}^n \end{aligned}$$

Now, if $B_1 \cap B_2 \neq \emptyset$, then we would have $q^{-1}[B_1 \cap B_2] = U \cap V \neq \emptyset$, contrary to assumption. Thus,

$$q[U] \cap q[V] = \emptyset \cap \mathbb{P}^n = \emptyset$$

□

0.2.7 Topological Groups

- (1) Suppose that $(G, \tau, *)$ is a topological group. Fixing $x \in G$, we show that $x\mathcal{O} \in \tau$ if, and only if, $\mathcal{O} \in \tau$:

Proof.

(\implies): Suppose that $x\mathcal{O} \in \tau$. It follows from pg. 75, that $L_{x^{-1}} : G \rightarrow G$ is a homeomorphism and an open map: thus, $L_{x^{-1}}[\mathcal{O}] \in \tau$.

(\impliedby): Suppose that $\mathcal{O} \in \tau$. Then, as noted on pg. 75, $L_x : G \rightarrow G$ is a homeomorphism and an open map: thus, $L_x[\mathcal{O}] = x\mathcal{O} \in \tau$.

□

- (2) Suppose that $(G, \tau, *)$ is a topological group. We show that \mathcal{O} is a neighbourhood of $x \in G$, if, and only if, $x^{-1}\mathcal{O}$ is a neighbourhood of e :

Proof. Suppose that $x \in G$, and without loss of generality, \mathcal{O} is an open set containing x . Then, by a previous lemma⁵, $L_{x^{-1}}[\mathcal{O}] = x^{-1}\mathcal{O} \in \tau$, and contains $x^{-1}x = e$.

By considering L_x , similar logic shows the reverse implication. □

- (3) We show that \mathbb{R} , with the Euclidean topology and addition is a topological group:

⁵reference this

Proof. The fact that $(\mathbb{R}, +)$ is a group is clear. To conclude, we show that $+ \equiv m : \mathbb{R} \rightarrow \mathbb{R}$, and $-1 \equiv i : \mathbb{R} \rightarrow \mathbb{R}$ are continuous:

Now, $i(x) = -x$, which is polynomial, and so continuous. And, as $m(x, y) = x + y = P_1(x, y) + P_2(x, y)$, where P_i is the projection mapping from \mathbb{R}^2 , m is continuous; it is the sum of two continuous functions.

A similar argument show that $(\mathbb{R}^n, \tau, +)$ is a topological group. \square

- (4) Let (G, τ, m) be a topological group. We show that the topological automorphisms of (G, τ, m) , for a subgroup of $(\text{Aut})(G)$. We denote the set of all topological automorphism of G by $\text{Aut}_\tau(G)$.

Proof. It is well known that $(\text{Aut}(G), \circ)$ is a group. We show that $(\text{Aut}_\tau(G), \circ)$ is a group:

As the composition of automorphism/homeomorphisms is another automorphism/homeomorphism, the fact that $(\text{Aut}_\tau(G), \circ)$ is closed is clear. To conclude, as $f \in \text{Aut}_\tau(G)$ is a homeomorphism, $f^{-1} \in \text{Aut}_\tau(G)$. Furthermore, $e : (G, \tau) \rightarrow (G, \tau)$ given by $e(x) = x$ is a homeomorphism. So, $e \in \text{Aut}_\tau(G)$.

By the subgroup test,

$$\text{Aut}_\tau(G) \leq \text{Aut}(G)$$

\square

- (5) Suppose that (G, τ, m) is a T_2 , finite, topological group. We claim $\tau = \mathcal{P}(G)$:

Proof. As G is T_2 , $\{x\}$ is open for each $x \in G$. Thus, as the union of open sets is open, it follows that $\tau = \mathcal{P}(G)$. \square

- (6) We claim that all topological groups of order 2 are topologically isomorphic.

Proof. Let (G, τ_G, m) be the topological group of order 2. Consider the topological group $(Z_2, \tau_{Z_2}, +)$ and the map $\phi : G \rightarrow Z_2$, given by $\phi(e_G) = 0$, and $\phi(g) = 1$. The fact that ϕ is a group isomorphism is well-known. Further, ϕ is 1-1 and onto, and ϕ^{-1} is continuous, as $\tau_G = \mathcal{P}(G)$ and $\tau_{Z_2} = \mathcal{P}(Z_2)$. This concludes the proof. \square

0.2.8 Homotopy Type

- (1) We prove that if A is a subspace of a topological space X and $G : X \times I \rightarrow X$ is deformation retract relative to A , then X and A have the same homotopy type⁶:

⁶Armstrong does not do a good job of defining a deformation retract. We use the following definition. "Strong" deformation retract: A is a 'strong' deformation retract of X iff there exists a map $D : X \times I \rightarrow X$ such that $D(a, t) = a$ for every $a \in A$, $D(x, 0) = x$ and $D(x, 1) \in A$ for all $x \in X$. See this post on SE.

Proof. Indeed: Let $f : A \rightarrow X$ be the inclusion map and $g : X \rightarrow A$ be defined by $g(x) = G(x, 1)$. We have previously shown that f and g are continuous. It is only left to show that $f \circ g \simeq 1_X$ and $g \circ f \simeq 1_A$:

Direct computation shows that

$$(g \circ f)(a) = g(f(a)) = g(a) = G(a, 1) = a,$$

while $f \circ g \simeq_G 1_X$. This completes the proof. \square

- (2) We show that any non-empty convex subset, X , of a euclidean space is homotopy equivalent to a point:

Proof. Without loss of generality, we assume that $X \subset \mathbb{E}^n$. We first show the existence of a deformation retract, $G : X \times I \hookrightarrow \{x\}$, where $x \in X$ is fixed:

Let G be defined as $G(y, t) = (1 - t)y + tx$, for all $y \in X$. Then, clearly $G(y, 0) = y$, and $G(y, 1) = x \in \{x\}$ for all $y \in X$. So, G is deformation retract. By the above, X and A are of the same homotopy type: null-homotopic. \square

- (3) We show that (X, τ) is contractible⁷ if, and only if, every map $f : X \rightarrow Y$, (Y, τ_Y) a topological space, is null-homotopic:

Proof.

(\implies): Suppose that (X, τ) is contractible to $x_0 \in X$. Let F be the contraction, and $f : X \rightarrow Y$ any map. The claim is that $f \circ F : X \times I \rightarrow Y$ is a homotopy between f , and $f(x_0)$.

Indeed; We note that the composition of two continuous functions is continuous and

$$(f \circ F)(x, 0) = f(F(x, 0)) = f(x)$$

As well as,

$$(f \circ F)(x, 1) = f(F(x, 1)) = f(x_0)$$

(\impliedby): Suppose that every map $f : X \rightarrow Y$ is null-homotopic. Then, in particular, $i : X \rightarrow X$ is null-homotopic. Thus, there exists a map $F : X \times I \rightarrow X$, such that $F(x, 0) = x = i(x)$, and $F(x, 1) = x_0 = i(x_0)$.

\square

⁷A space (X, τ) is contractible (to a point $x_0 \in X$) provided that there exists a map $F : X \times I \rightarrow X$ such that $F(x, 0) = x$ and $F(x, 1) = x_0$.

- (4) Let (X, τ) be a topological space. We show that the cone on X , CX is contractible. We consider the "cone tip" as $X \times \{0\}$:

Proof. Consider the function $\overline{H} : (X \times I) \times I \rightarrow X \times I$, given by $\overline{H}((x, v), s) = (x, v(1 - s))$. By the product topology, it follows that \overline{H} is continuous.⁸ Let $\pi : X \times I \rightarrow CX = X \times I / X \times \{0\}$ be the canonical map. The claim is that $\pi \circ \overline{H} : (X \times I) \times I \rightarrow CX$ is the desired homotopy.

Direct computation shows that, for all $x \in X$, $v \in V$,

$$\pi(\overline{H}((x, v), 0)) = \pi((x, v)) = (x, v),$$

and

$$\pi(\overline{H}((x, v), 1)) = \pi((x, 0)) = (x, 0)$$

Thus, $H \equiv \pi \circ \overline{H}$ is the desired homotopy. □

⁸For every $t \in I$, we can associate $\pi(\overline{H}((x, v), t))$ with $(c, v(1 - t), t)$, showing that the product topology of $CX \times I$ agrees with that of $(X \times I) \times I$.

Chapter 1

Introduction

1.1 Euler's Theorem

There are no exercises listed for this section.

1.2 Topological Equivalence

There are no exercises listed for this section.

1.3 Surfaces

There are no exercises listed for this section.

1.4 Abstract Spaces

There are no exercises listed for this section.

1.5 A Classification Theorem

There are no exercises listed for this section.

1.6 Topological Invariants

1. We prove that $v(T) - e(T) = 1$ for any tree T :

Proof. We proceed by induction on the number of edges in T :

- Basis: If $e(T) = 1$, then $v(T) = 2$, and so, $v(T) - e(T) = 1$.

- Hypothesis: Suppose that $v(T) - e(T) = 1$ for any tree T , such that $e(T) = n$, for some $n \in \mathbb{N}$.
- Step: Suppose that T is a tree, such that $e(T) = n + 1$. Now, removing any edge will disconnect the tree, since by definition, T contains no loops. Say that we remove an edge $e_0 \in T$, breaking up T into T_1, T_2 . Now, clearly, $v(T) = v(T_1) + v(T_2)$, and $e(T) = n + 1 = e(T_1) + e(T_2) + 1$. Thus, by the inductive hypothesis,

$$\begin{aligned} v(T) - e(T) &= v(T_1) + v(T_2) - e(T_1) - e(T_2) - 1 \\ &= 1 + 1 - 1 \\ &= 1 \end{aligned}$$

□

2. We show that inside any graph we can always find a tree which contains all the vertexes:

Proof. Let G be a finite graph. If G is already a tree, we are done. So, suppose that G is not a tree. Then, by the comments on pg. 3, G contains a (without loss of generality, minimal) loop, L . Now, we remove some edge $e_L \in L$. Now, as L is a loop, e_L does not disconnect L , and so $L - e_L$ is a tree. We continue in this way, as long as the new graph formed by removing an edge is not a tree.

To conclude, we note that this process must stop at some loop, because $e(G) < \infty$; further, we have not removed any vertexes. □

3. We prove that $v(\Gamma) - e(\Gamma) \leq 1$ for any graph Γ , with equality, precisely when Γ is a tree:

Proof. We have previously shown the equality condition. Using the above proof, we select a subtree, with the same vertexes as Γ . Call this tree Γ' . Then, $v(\Gamma') - e(\Gamma') = 1$. As Γ' was created by removing edges, and not vertexes, we have $v(\Gamma) = v(\Gamma')$, and $e(\Gamma) \leq e(\Gamma')$. So,

$$e(\Gamma) - e(\Gamma) \leq v(\Gamma') - e(\Gamma') = 1$$

□

4. We find a tree in the polyhedron of Fig. 1.3 which contains all the vertexes and construct the dual graph Γ and show that Γ contains loops:

Proof. Please refer to figures 1.1a and 1.1b □

5. Having done Problem 4, thicken both T and Γ in the polyhedron. T is a tree, so thickening it gives a disc. We investigate what happens when you thicken Γ ?



(a) The tree that contains every vertex.



(b) Blue: Dual graph. Green: loops.

Proof. Γ is basically two loops connected at a point, with some other edges connected that do not make any more loops. So thickening should produce something homeomorphic to what is shown in Problem 11 (b). \square

6. Let P be a regular polyhedron in which each face has p edges and for which q faces meet at each vertex. We use Euler's formula to prove that,

$$\frac{1}{p} + \frac{1}{q} = \frac{1}{2} + \frac{1}{e}$$

Proof. We count the number of faces of P as follows: Since each face of P has p edges, $pf = 2e$. We count the number of vertexes as follows: Since each vertex has q faces which meet on it, $qv = 2e$. In all, $f = 2e/p$, and $v = 2e/q$. Assuming that Euler's formula holds for P , we have

$$v - e + f = \frac{2e}{q} - e + \frac{2e}{p} = 2$$

And the result follows. \square


7. We deduce that there are only 5 regular (convex) polyhedra.

Proof. If P is a polyhedra, it must satisfy the above criterion. Further, we assume that $p \geq 3$, since if not, we cannot construct a polygon. We use Shlüt notation. By checking routinely, for values up to $p, q \leq 5$ we have the following set of polyhedra:


$$\{\{3, 3\}, \{3, 5\}, \{5, 3\}, \{3, 4\}, \{4, 3\}\}$$

Now, if $p \geq 6$, then we must have


$$\frac{1}{6} + \frac{1}{q} = \frac{1}{2} + \frac{1}{e}$$




(a) Tetrahedron




(b) Cube



(c) Octahedron



(d) Dodecahedron



(e) Icosahedron

Figure 1.2: All five platonic solids.

Further, as $0 \leq 1/q \leq 1/5$, we have

$$\begin{aligned} \frac{1}{6} + \frac{1}{q} &= \frac{q+6}{6q} \\ &< \frac{12}{6q} = \frac{2}{q} \\ &< \frac{1}{2} + \frac{1}{e} \end{aligned}$$

A contradiction.

Please see figures 1.2a, 1.2b, 1.2c, 1.2d, and 1.2e. □

8. We check that $v - e + f = 0$ for the polyhedron shown in Fig. 1.3 and find a polyhedron which can be deformed into a pretzel and calculate its Euler number:




Figure 1.3: "pretzel"

Proof. For the polyhedron in Fig. 1.3, $v = 20$, $e = 40$, $f = 20$. Therefore $v - e + f = 0$. Figure 1.3 is basically a donut with two holes, i.e. a "pretzel".

Figure 1.3 has 38 faces (10 on the top, 10 on the bottom, 10 inside the hole, and 8 around the outside sides). It has 76 edges (29 on the top, 29 on the bottom, 10 vertical ones inside the holes and 8 vertical ones around the outside sides), and it has 36 vertexes (18 on the top, 18 on the bottom); Therefore $v - e + f = -2$. \square

9. This is left to the reader as an exercise.
10. We find a homeomorphism from the real line to the open interval $(0, 1)$, and show that any two open intervals are homeomorphic:

Proof. We show that the real line, \mathbb{R} is homeomorphic to $(-\pi/2, \pi/2)$. This amounts to showing that $\arctan : \mathbb{R} \rightarrow (-\pi/2, \pi/2)$ is continuous, 1-1, onto and that $\tan \equiv \arctan^{-1}$ is continuous. Thus, $\mathbb{R} \simeq (-\pi/2, \pi/2)$.

To conclude the proof, we show that any open subset, $(a, b) \subset \mathbb{R}$ is homeomorphic to $(-\pi/2, \pi/2)$, $a \neq b$:

Indeed, under the mapping $f(x) = (b - a)(\frac{x}{\pi} + \frac{1}{2}) + a$, we have

$$(-\pi/2, \pi/2) \rightarrow (a, b)$$

As f is a degree 1 polynomial in \mathbb{R} , it follows that it is 1-1, onto and continuous with a continuous inverse. \square

11. This is left to the reader as an exercise.
12. We find a homeomorphism from $S^2 - N = S^2 - \{(0, 0, 1)\}$ to \mathbb{E}^2 :

Proof. As pointed out in Armstrong, we find a formula for stereographic projection. From the fact that $S^2 = \{x \in \mathbb{R}^3 : \|x\|_2 = 1\}$, and rules of calculus, any line passing through $v = (x, y, z) \in S^2 - N$ and N is of the following form:

$$p(t) = (0, 0, 1) + t(x, y, z) \equiv \langle tx, ty, 1 - t(1 - x) \rangle \quad (1.1)$$

For $t \in [0, \infty)$. As the z component of p is zero exactly when $t_0 = 1/(1-z)$, we substitute t_0 back in (1) to obtain our formula for stereographic projection:

$$\Pi(x, y, z) = \left(\frac{x}{1-z}, \frac{y}{1-z} \right)$$

It is left to show that Π is 1-1, onto, continuous, with a continuous inverse.

1-1: By component wise comparison, the result follows.

Onto: Given $(x_0, y_0) \in \mathbb{R}^2$, we have, after some simple algebra,

$$\Pi\left(\frac{x_0}{r}, \frac{y_0}{r}, \frac{x_0^2 + y_0^2}{r}\right) = (x_0, y_0),$$

where $r = 1 + x_0^2 + y_0^2$.

Continuity: As Π is a linear transformation in \mathbb{R}^2 , we know that from rules of calculus/real analysis that it is continuous.

Cont. Inverse: Π^{-1} is as outline above in the derivation of onto. For the same reasons the function is continuous.

□

13. Let $x, y \in S^2$. We find a homeomorphism from S^2 to S^2 which takes x to y . We do the same for the plane, and torus:

Proof. We assume that results about matrix groups are applicable.

- S^2 : Consider S^2 and $SO(3)$. Let $x \in S^2$. Then, by the Gram-Schmidt Orthogonalization process, there exists $X \in SO(3)$ for which x is the first column of X . Likewise for y , its corresponding matrix Y . Then, $YX^{-1} \in SO(3)$ and maps x to y . As $SO(3)$ is a group, $(YX^{-1})^{-1} \in SO(3)$. Thus, YX^{-1} is a homeomorphism which takes x to y .

- \mathbb{E}^2 : Fix $(x, y), (z, w) \in \mathbb{R}^2$. Then, the function, f that sends $(a, b) \in \mathbb{R}^2$ to

$$((x+z) - a, (y+w) - b)$$

sends (x, y) to (z, w) . As f is linear, it is continuous. Likewise, its inverse is continuous.

- Torus: Consider the torus as $S^1 \times S^1$. Then component wise examination shows that the example above proves the result.

□

14. This is left to the reader as an exercise.

15. This is left to the reader as an exercise.

16. This is left to the reader as an exercise.
17. Define $f : [0, 1) \rightarrow C$, by $f(x) = e^{2\pi ix}$. We prove that f is a continuous bijection. In addition, we find a point $x \in [0, 1)$ and a neighbourhood N , of x in $[0, 1)$, such that $f[N]$ is not a neighbourhood of $f(x)$ in C ; consequently, this means f is not a homeomorphism.

Proof. We first show that f is a continuous bijection:

Via Euler's formula,

$$e^{2\pi ix} = \cos(2\pi x) + i \sin(2\pi x), \quad \forall x \in \mathbb{R}$$

Using this fact, we show that f is 1-1 and onto.

- 1-1: Suppose that $\cos(2\pi x) + i \sin(2\pi x) = \cos(2\pi y) + i \sin(2\pi y)$ for some $x, y \in [0, 1)$. Then, by component-wise comparison of complex numbers we have $\cos(2\pi y) = \cos(2\pi x)$, iff, $2\pi y = 2\pi x$, iff $x = y$, as $x, y \in [0, 1)$; likewise for \sin . Thus, f is 1-1.
- Onto: Let $y \in C$. Then, $y = \cos(2\pi y_0) + i \sin(2\pi y_1)$ for some $(y_0, y_1) \in \mathbb{R}^2$. But as $f : [0, 1) \rightarrow C$, we have $0 \leq 2\pi y_0, 2\pi y_1 \leq 2\pi$. Consequently, f is onto.
- Continuity: By the definition of continuity in \mathbb{C} , and the fact that \cos, \sin are continuous, we have that f is continuous.

Next, consider $[0, 1/2) \subset [0, 1)$. This is an open neighbourhood in $[0, 1)$, by definition. But, $f[[0, 1/2)]$ is C intersected with the upper half of the plane minus $z = -1$. Which is not a neighbourhood of $z = 1 \in C$, because any open ball centered around $z = 1$ must contain the lower plane. Therefore, f is not a homeomorphism. \square

18. This is left to the reader as an exercise.
19. This is left to the reader as an exercise.
20. We prove that the radial projection of the tetrahedron to the sphere is a homeomorphism:

Proof. A solid proof of this relies on the fact that a circumscribed polygon with center of mass 0 is homeomorphic to S^n via radial projection. The result of which are shown in chapter 5. \square

21. Let C denote the unit circle in the complex plane and D the disc which it bounds. Given two points $x, y \in D - C$, we find a homeomorphism from D to D which interchanges x and y and leaves all points of C fixed:

Proof. This is more or less intuitively obvious. But writing down an explicit function is not so easy. First note that for any $a \in \mathbb{C}$, the function $f(z) = \frac{z-a}{1-\bar{a}z}$ takes S^1 to itself. To see this suppose $|z| = 1$. then

$$\begin{aligned} \left| \frac{z-a}{1-\bar{a}z} \right| &= \left| \frac{z-a}{\bar{z}z-\bar{a}z} \right| \\ &= \left| \frac{z-a}{(\bar{z}-\bar{a})z} \right| \\ &= \left| \frac{z-a}{\overline{z-a}} \right| \frac{1}{|z|} \\ &= 1 \end{aligned}$$

If $|a| < 1$ then the denominator never vanishes so this is a continuous function on D . Also, $f(0) = -a$, so if $|a| < 1$ then since f takes C to itself and 0 maps to $-a \in (D-C)$, f must take all of D to itself. The inverse f^{-1} is therefore also a continuous function from D to D that takes C to C . Now suppose $x, y \in \mathbb{C}$ with $|x| < 1$ and $|y| < 1$, let

$$f_1(z) = \frac{z-x}{1-\bar{x}z}$$

and

$$f_2(z) = \frac{z-ty_1}{1-\bar{t}y_1z}$$

where $y_1 = f_1(y)$ and $t = \frac{1-\sqrt{1-|y_1|^2}}{|y_1|^2}$. As shown above, both f_1 and f_2 take C to itself. Finally let

$$g(z) = ze^{i(1-|z|)\pi/(1-|x_2|)}$$


where $x_2 = f_2(0)$. Then $g(x_2) = -x_2$ and $g(-x_2) = x_2$ and $g(z) = z \forall z \in C$. Since $z \mapsto |z|$ is continuous, g is built up from sums, products and compositions of continuous functions and therefore g is continuous.

The function $f_1^{-1}f_2^{-1}gf_2f_1$ is therefore a continuous function from D to D that switches x and y and fixes C . \square

22. With C, D as above (in Problem 21), define $h : D-C \rightarrow D-C$ by

$$\begin{aligned} h(0) &= 0 \\ h(re^{i\theta}) &= r \exp \left[i \left(\theta + \frac{2\pi r}{1-r} \right) \right] \end{aligned}$$

We show that h is a homeomorphism, but that h cannot be extended to a homeomorphism from D to D and draw a picture which shows the effect of h on a diameter of D :



Proof. The function h restricted to a circle of radius r acts by rotation of $2\pi r/(1-r)$ radians. Now $2\pi r/(1-r) \rightarrow \infty$ as $r \rightarrow 1$, so as the circle radius grows towards one, it gets rotated to greater and greater angles approaching infinity. Thus intuitively it's pretty obvious this could not be extended to the boundary. See figure ??.

Now, we can think of (r, θ) as polar coordinates in \mathbb{R}^2 . And the topology on \mathbb{C} is the same as that on \mathbb{R}^2 . Thus as a function of two variables r and θ this is just a combination of continuous functions by sums, products, quotients and composition. Since the only denominator involved does not vanish for $|r| < 1$, this is a continuous function of r and θ on D which is clearly onto. Since it is a simple rotation on each circle of radius r , it is also clearly one-to-one. The inverse is evidently

$$h^{-1}(0) = 0$$

$$h^{-1}(re^{i\theta}) = r \exp \left[i \left(\theta - \frac{2\pi r}{1-r} \right) \right]$$

Now, let $r_n = \frac{n}{n+2}$ for n odd and $r_n = \frac{n-1}{n}$ for n even. Then for n odd, $\frac{r}{1-r} = \frac{n}{2}$, and for n even $\frac{r}{1-r} = n-1$. Therefore $\exp \left[i \left(\frac{2\pi r_n}{1-r_n} \right) \right]$ equals 1 if r is even and -1 if r is odd. Now, $r_n \rightarrow 1$. So if h could be extended to all of D we must have $h(1) = \lim h(r_n)$. But $h(r_n)$ does not converge, it alternates between 1 and -1 . Therefore, there is no way to extend h to C to be continuous. \square

23. Using the intuitive notion of connectedness, we argue that a circle and a circle with a spike attached cannot be homeomorphic (Fig. 1.26):

Proof. In the circle, if we remove any one point what remains is still connected. However in the circle with a spike attached there is one point we can remove that renders the space not-connected. Since this property of being able to remove a point and retain connectedness must be a topological property preserved by homeomorphism, the two spaces cannot be homeomorphic. \square

24. Let X, Y be the subspace of the plane shown in Fig. 1.27. Under the assumption that any homeomorphism from the annulus to itself must send

the points of the two boundary circles among themselves, we argue that X and Y cannot be homeomorphic:

Proof. The two points that connect the two spikes to the two boundary circles in X must go to the two points that connect the two spikes to the boundary circles in Y , because those are the only two points on the boundary circles that can be removed to result in a disconnected space, and because by assumption the circles go to the circles. Since the two points lie on the same circle in Y but on different circles in X , some part of the outer circle in Y must go to the outer circle in X and the rest must go to the inner circle in X . But then some part of the outer circle in Y must go to the interior of X . I'm not sure exactly how Armstrong expects us to prove this but it basically follows from the intermediate value theorem, applied to the two coordinates thinking of these shapes as embedded in \mathbb{R}^2 . \square

25. With X and Y as above, consider the following two subspaces of \mathbb{E}^3 :

$$X \times [0, 1] = \{(x, y, z) \mid (x, y) \in X, 0 \leq z \leq 1\},$$

$$Y \times [0, 1] = \{(x, y, z) \mid (x, y) \in Y, 0 \leq z \leq 1\}.$$

Convince yourself that if these spaces are made of rubber then they can be deformed into one another, and hence that they are homeomorphic:

Proof. With the extra dimension, the squareness can be continuously deformed so that it is a solid torus, with two flat rectangular shapes sticking off. One has both rectangles pointing out and one has one pointing out and the other pointing in. Since the torus is round, the first space made from X can be rotated at the location where the inner rectangle is a full half turn to point the rectangle out, and as parallel slices (discs) of the torus move away from where the rectangle is attached, the rotation gradually gets less and less until it becomes zero before reaching the other rectangle. In this way the inner rectangle can be rotated to point out without affecting the other rectangle and with a gradual change in rotation angle between them guaranteeing the operation is continuous. \square

26. This is left to the reader as an exercise.

27. This is left to the reader as an exercise.

Chapter 2

Continuity

2.1 Open and Closed Sets

(1) We verify each of the following for arbitrary subsets A, B of a space X :

- $\overline{A \cup B} = \overline{A} \cup \overline{B}$:

Proof. \overline{A} and \overline{B} are closed by theorem (2.3). Thus $\overline{A} \cup \overline{B}$ is closed. Now $A \subseteq \overline{A}$ and $B \subseteq \overline{B}$. Therefore $\overline{A} \cup \overline{B}$ is a closed set containing $A \cup B$. By Theorem 2.3 $\overline{A \cup B}$ is the smallest closed set containing $A \cup B$, thus it must be that $\overline{A \cup B} \subseteq \overline{A} \cup \overline{B}$. Conversely, $\overline{A \cup B}$ is a closed set that contains A , so $\overline{A \cup B} \supseteq \overline{A}$. Similarly $\overline{A \cup B} \supseteq \overline{B}$. Thus $\overline{A \cup B} \supseteq \overline{A} \cup \overline{B}$. Thus $\overline{A \cup B} = \overline{A} \cup \overline{B}$. \square

- $\overline{A \cap B} \subseteq \overline{A} \cap \overline{B}$:

Proof. \overline{A} is a closed set that contains $A \cap B$, so $\overline{A \cap B} \subseteq \overline{A}$. Likewise $\overline{A \cap B} \subseteq \overline{B}$. Thus $\overline{A \cap B} \subseteq \overline{A} \cap \overline{B}$.

To see that equality does not hold, let $A = \mathbb{Q}$ and let $B = \mathbb{R} - \mathbb{Q}$. Then $A \cap B = \emptyset$, so $\overline{A \cap B} = \emptyset$. But $\overline{A} = \mathbb{R}$ and $\overline{B} = \mathbb{R}$, so $\overline{A} \cap \overline{B} = \mathbb{R}$. \square

- $\overline{\overline{A}} = \overline{A}$:

Proof. $\overline{\overline{A}}$ is the smallest closed set containing \overline{A} by corollary (2.4), and \overline{A} is closed by theorem (2.3), that contains \overline{A} . Thus $\overline{\overline{A}} = \overline{A}$. \square

- $(A \cup B)^\circ \supseteq \overset{\circ}{A} \cup \overset{\circ}{B}$:

Proof. Let $x \in \overset{\circ}{A} \cup \overset{\circ}{B}$. Assume, without loss of generality, that $x \in \overset{\circ}{A}$. Then there is an open set $U \subseteq A$ such that $x \in A$. But then $x \in U \subseteq A \cup B$. So $x \in \overset{\circ}{A \cup B}$.

Thus

$$\overset{\circ}{A} \cup \overset{\circ}{B} \subseteq (A \cup B)^\circ$$

To see that equality does not hold, let $A = \mathbb{Q}$ and let $B = \mathbb{R} - \mathbb{Q}$. Then $\overset{\circ}{A} = \emptyset$ and $\overset{\circ}{B} = \emptyset$. And $A \cup B = \mathbb{R}$, so $(A \cup B)^\circ = \mathbb{R}$. Therefore $(A \cup B)^\circ = \mathbb{R}$ but $\overset{\circ}{A} \cup \overset{\circ}{B} = \emptyset$. \square

- $(A \cap B)^\circ = \overset{\circ}{A} \cap \overset{\circ}{B}$:

Proof. Let U be an open set in $A \cap B$. Then $U \subseteq A$ and $U \subseteq B$. Thus $(A \cap B)^\circ \subseteq \overset{\circ}{A} \cap \overset{\circ}{B}$. Conversely suppose $x \in \overset{\circ}{A} \cap \overset{\circ}{B}$. Then \exists open sets U and V s.t. $x \in U \subseteq A$ and $x \in V \subseteq B$. Then $U \cap V$ is open and $x \in U \cap V \subseteq A \cap B$. Thus $(A \cap B)^\circ \supseteq \overset{\circ}{A} \cap \overset{\circ}{B}$.

Thus

$$(A \cap B)^\circ = \overset{\circ}{A} \cap \overset{\circ}{B}$$

\square

- $(\overset{\circ}{A})^\circ = \overset{\circ}{A}$:

Proof. Clearly

$$(\overset{\circ}{A})^\circ \subseteq \overset{\circ}{A}$$

Let $x \in \overset{\circ}{A}$. Then there exists an open set U , such that $x \in U \subseteq A$. Now $\overset{\circ}{A}$ is a union of open sets so is open. Let $V = \overset{\circ}{A} \cap U$. Then $x \in V \subseteq \overset{\circ}{A}$. Therefore $x \in (\overset{\circ}{A})^\circ$ and so,

$$\overset{\circ}{A} \subseteq (\overset{\circ}{A})^\circ$$

Thus,

$$\overset{\circ}{A} = (\overset{\circ}{A})^\circ$$

\square

(2) This is left to the reader as an exercise.

(3) We specify the interior, closure and frontier of the following subsets.

$$\{(x, y) : 1 < x^2 + y^2 \leq 2\} \tag{2.1}$$

$$\mathbb{E}^2 - \{(0, t), (t, 0) : t \in \mathbb{R}\} \tag{2.2}$$

$$\mathbb{E}^2 - \{(x, \sin(1/x)) : x > 0\} \tag{2.3}$$

- $\{(x, y) : 1 < x^2 + y^2 \leq 2\} = A$:

$$\begin{aligned}\text{Fr } A &= \overline{A} \cap \overline{(\mathbb{E} - A)} \\ &= \overline{A} \cap \{ \{(x, y) : 0 \leq x^2 + y^2 \leq 1\} \cup \{(x, y) : 2 \leq x^2 + y^2\} \} \\ &= \{(x, y) : x^2 + y^2 = 1 \vee x^2 + y^2 = 2\}\end{aligned}$$

The rest are left to the reader.

- $\mathbb{E}^2 - \{(0, t), (t, 0) : t \in \mathbb{R}\} = B$:

From the fact that $\overline{\mathbb{E}^2 - B} = \mathbb{E}^2$, we have

$$\text{Fr } B = B$$

The rest are left to the reader.

- $\mathbb{E}^2 - \{(x, \sin(1/x)) : x > 0\} = C$:

From the fact that $\overline{\mathbb{E}^2 - C} = \mathbb{E}^2$, we have

$$\text{Fr } C = C$$

The rest are left to the reader as an exercise.

(4) This is left to the reader as an exercise.

(5) We show that if A is a dense subset of a space (X, τ) , and if $\mathcal{O} \in \tau$, that $\mathcal{O} \subset \overline{A \cap \mathcal{O}}$:

Proof. Suppose, to the contrary, that $\mathcal{O} \not\subset \overline{A \cap \mathcal{O}}$. Then, there exist some $x \in \mathcal{O}$, such that $x \notin \overline{A \cap \mathcal{O}}$.

As $\overline{A \cap \mathcal{O}}$ is closed, $x \in (\overline{A \cap \mathcal{O}})^c$ and so, there exists some $\mathcal{O}_x \in \tau$ such that $x \in \mathcal{O}_x$, and

$$\overline{A \cap \mathcal{O}} \cap (\mathcal{O}_x - \{x\}) = \emptyset$$

But, as $x \notin \overline{A \cap \mathcal{O}}$, we have

$$\overline{A \cap \mathcal{O}} \cap \mathcal{O}_x = \emptyset$$

and consequently, $A \cap \mathcal{O} \cap \mathcal{O}_x = \emptyset$. But then, setting $B = \mathcal{O} \cap \mathcal{O}_x$, we have $x \in B$, $B \in \tau$, but $A \cap B = \emptyset$, contrary to extra lemma; $\overline{A} \neq X$, a contradiction. \square

(6) We prove that if Y is subspace of X , and Z is a subspace of Y , that Z is a subspace of X :

Proof. The open sets in Y are exactly the sets $\mathcal{O} \cap Y$ where \mathcal{O} is open in X . The open sets in A as a subspace of Y are therefore sets of the form $A \cap (Y \cap \mathcal{O})$ where \mathcal{O} is open in X . But $A \subseteq Y$, so $A \cap (Y \cap \mathcal{O}) = A \cap \mathcal{O}$. Therefore the open sets in A as a subspace of X are exactly the same as the open sets of A as a subspace of Y . \square

- (7) Suppose that Y is a subspace of (X, τ) . We show that a subset A of Y is closed in Y if it is the intersection of Y with a closed set in X . Further, we show that we get the same result if we take the closer in Y or X :

Proof. If $A \subset Y$ is closed in Y , then $Y - A$ is open in Y . But then, by the definition of subspace topology, $Y - A = Y \cap \mathcal{O}_y$, for some open $\mathcal{O}_y \in X$. Consequently, we have

$$\begin{aligned} A &= Y - (Y \cap \mathcal{O}_y) \\ &= Y \cap (Y \cap \mathcal{O}_y)^c \\ &= Y \cap (X - \mathcal{O}_y) \end{aligned}$$

And, as $X - \mathcal{O}_y$ is closed in X , this proves the result. We note that a similar case holds in the case where A is open.

Letting \overline{A}_y and \overline{A}_x denote the closure of $A \subset Y$ in Y and X respectively, we show that $\overline{A}_y = \overline{A}_x$:

From the previous part of this proof, we have that $\overline{A}_y = Y \cap C$, where C is closed in X . Now, since C is a closed set in X containing A , we have $\overline{A}_x \subset C$, and so $Y \cap \overline{A}_x \subset \overline{A}_y = Y \cap C$.

Again by the first problem, we have $Y \cap \overline{A}_x$ is closed in Y and contains A . So,

$$\overline{A}_y \subset Y \cap \overline{A}_x \subset \overline{A}_x$$

Thus, $\overline{A}_y = \overline{A}_x$. □

- (8) Let Y be a subspace of (X, τ) . Given $A \subset Y$, we show that $A_X^\circ \subset A_Y^\circ$, and give an example when the two may not be equal:

Proof. Let $x \in A_X^\circ$. Then, there exists an open $\mathcal{O}_x \subset X$, with $x \in \mathcal{O}_x$ and $\mathcal{O}_x \subset A$. Now, since $\mathcal{O}_x \subset A \subset Y$, $x \in \mathcal{O}_x \cap Y \subset A$ and, $\mathcal{O}_x \cap Y$ is open in Y , by definition. Thus, $x \in A_Y^\circ$.

An example when they might not be equal is in the following case: Let $X = \mathbb{R}$, $Y = \mathbb{Z}$, and $A = \{0\}$. Then, $A_X^\circ = \emptyset$. But, every point of \mathbb{Z} is open in the subspace topology, and so $A_Y^\circ = \{0\}$. □

- (9) Let Y be a subspace of X . We show that if A is open (closed) in Y , and if Y is open (closed) in X , that A is open (closed) in X :

Proof. Suppose $A \subset Y \subset X$ and A open in Y . Then $A = Y \cap U$ where U is open in X . The intersection of two open sets is open, so if Y is open in X then A is open in X . Similarly, if $A \subset Y \subset X$ and A closed in Y , then $A = Y \cap C$ where C is closed in X (by exercise 7). The intersection of two closed sets is closed, so if Y is closed in X then A is closed in X . □

- (10) We show that the frontier of a set always contains the frontier of its interior, and describe the relationship between $\text{Fr}(A \cup B)$ and $\text{Fr } A, \text{Fr } B$:

Proof. Let (X, τ) be a topological space, and let $A \subset X$. We want to show that $\text{Fr } A^\circ \subset \text{Fr } A$.

Let $x \in \text{Fr } A^\circ$. Then,

$$x \in \overline{A^\circ} \cap \overline{(X - A^\circ)} = \overline{A^\circ} \cup \overline{(X - A) \cup (A - A^\circ)}$$

Now, if $x \in \overline{A^\circ}$ and $x \in \overline{X - A}$, we are done. So suppose that $x \in \overline{A^\circ}$ and $x \in \overline{(A - A^\circ)}$. But then, $x \in \overline{A^\circ} \cup \overline{(A - A^\circ)} = \overline{A}$. Thus, the result follows.

The inclusion $\text{Fr}(A \cup B) \subset \text{Fr } A \cup \text{Fr } B$ always holds, and is left to the reader. To show that the reverse inclusion does not always hold, consider $X = \mathbb{R}$, and $A = \mathbb{Q}$. Then,

$$\begin{aligned} \text{Fr}(A \cup A^c) &= \text{Fr } \mathbb{R} = \emptyset \\ &\neq \text{Fr } A \cup \text{Fr } A^c = \mathbb{R} \end{aligned}$$

□

- (11) This main part of this exercise is left to the reader. However, we do show that this topology does not have a countable base:

Proof. Let $B = \tau$ be the topology specified. Suppose, to the contrary, that $\{B_n\}_{n=1}^\infty$ is a countable base for the topology τ . Define the function $f : \mathbb{R} \rightarrow \mathbb{N}$ as follows: for each $x \in \mathbb{R}$, let $f(x) = n$, such that $B_n \subset [x, 1+x)$. Now, we show that f is 1-1 to arrive at a contradiction:

Indeed; Suppose to the contrary, without loss of generality, that $x < y$. Then, if $f(x) = f(y)$, $f(x) = [x, x+1) \subset B_{f(y)} = [y, y+1)$, which is impossible. Thus, $x = y$. □

- (12) We show that if a topological space (X, τ) has a countable base for its topology, then X contains a countable dense subset. I.e. A second countable space is separable:

Proof. Let $\{B_n\}_n$ be a countable base for τ . By the Axiom of Choice, let A be the collection of elements $\{a_i\}_i$ such that $a_i \in B_i$. The claim is that $\overline{A} = X$.

Indeed; let $\mathcal{O} \in \tau$. Then, $\mathcal{O} = \bigcup_j B_j$, where $B_j \in B$. Now, as $A = \bigcup_i x_i$, where $x_i \in B_i$, we have $A \cap \mathcal{O} \neq \emptyset$. Thus, by extra lemma, $\overline{A} = X$. □

2.2 Continuous Functions

- (1) We show that if $f : \mathbb{R} \rightarrow \mathbb{R}$ is a map, then the set of points left fixed by f is closed. Further, the kernel of f is closed.

Proof. Define $f_0(x) = f(x) - x$. The rest is left to the reader. \square

- (2) We show that the function $h(x) = \frac{e^x}{1+e^x}$ is a homeomorphism from the real line to the open interval $(0, 1)$:

Proof. Let

$$f(x) = \frac{e^x}{1+e^x}, \quad y \in (0, 1), \quad x = \ln \left(\frac{y}{1-y} \right)$$

We show that f is onto and 1-1:

- As $f(x) = y$, f is onto.
- Suppose $f(x) = f(y)$. Then,

$$\frac{e^x}{1+e^x} = \frac{e^y}{1+e^y} \text{ implies } e^x + e^{x+y} = e^y + e^{x+y}$$

Therefore $e^x = e^y$ and f is 1-1.

For $a, b \in \mathbb{R}$,

$$f^{-1}((a, b)) = \left(\ln \frac{a}{1-a}, \ln \frac{b}{1-b} \right),$$

which is open. Since the intervals (a, b) are a basis, it follows from Theorem (2.9(b)) that f is continuous. And $f((a, b)) = (f(a), f(b))$, so f takes open sets in the base to open sets. Therefore f^{-1} is continuous, and so, f is a homeomorphism. \square

- (3) Let $f : \mathbb{E} \rightarrow \mathbb{R}$ be a map and define $\Gamma_f : \mathbb{E} \rightarrow \mathbb{E}^2$ by $\Gamma_f(x) = (x, f(x))$. We show that Γ_f is continuous and that its image, with the induced topology, is homeomorphic to \mathbb{E} :

Proof. We show that Γ_f is continuous. To do this, we use the sequential criterion for continuity in \mathbb{E}^n . Let $\{x_n\}_n$ be a sequence in \mathbb{E} , such that $x_n \rightarrow x \in \mathbb{E}$. Then Γ_f is clearly continuous by the fact that f is, and component wise comparison. That is,

$$\Gamma_f(x_n) = (x_n, f(x_n)) \rightarrow (x, f(x)) = \Gamma_f(x)$$

Next, we show that $\text{Im } \Gamma_f$ is homeomorphic to \mathbb{E} :

We claim that the function $p_1 : \text{Im } \Gamma_f \rightarrow \mathbb{E}$, defined by $p_1((x, f(x))) = x$ is the desired homeomorphism.

Continuity: We have previously shown that the projection map is continuous.

1-1: This is true, by construction.

Onto: This is clear, as f a map from \mathbb{E} to \mathbb{E} .

Cont. Inverse: This is clear, as $p_1^{-1} \equiv \Gamma_f$, and we have shown Γ_f is continuous.

This proves the result.¹

□

- (4) We determine what topology on X implies that every real-valued function defined on X is continuous:

Proof. X must have the discrete topology, where every subset is open. To show this, it suffices to show points in X are open.

Indeed; Fix $x \in X$ and define $f : X \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} f(x) = 0 \\ f(y) = 1 & y \neq x \end{cases}$$

Then,

$$f^{-1}((-1/2, 1/2)) = \{x\}$$

and thus f is continuous, since $(-1/2, 1/2)$ is open and $\{x\}$ is open. □

- (5) Consider $X = \mathbb{R}$ with the co-finite topology, (\mathbb{R}, CO) . We show that $f : \mathbb{E} \rightarrow X$ defined by $f(x) = x$ is a map, but not a homeomorphism:

Proof. To see that f is continuous, let $\mathcal{O} \in CO$. Then, $X - \mathcal{O} = \mathbb{R} - \mathcal{O}$ is finite. But then, $f^{-1}[\mathcal{O}] = \mathcal{O}$, and \mathcal{O}^c is finite in \mathbb{E} . This implies that $\mathcal{O} = \overline{\mathcal{O}}$. Thus, $\mathcal{O}^c = \mathcal{O}$ is open in \mathbb{E} . The fact that f is 1-1 is clear.

Consider the inverse function of f . Note that (a, b) , $a < b$ is open in \mathbb{E} . But, $f^{-1}[(a, b)] = (a, b) = f[(a, b)]$, which is not open in X . Thus, f^{-1} is not continuous. □

- (6) Suppose $X = A_1 \cup A_2 \cup \dots$, where $A_n \subseteq \overset{\circ}{A}_{n+1}$ for each n . We show that if $f : X \rightarrow Y$ is a function such that, for each n , $f|_{A_n} : A_n \rightarrow Y$ is continuous with respect to the induced topology on A_n , then f is itself continuous:

Proof. Let $x \in X$. Then $x \in A_n$ for some n and $A_n \subseteq \overset{\circ}{A}_{n+1}$. Thus $x \in \overset{\circ}{A}_{n+1}$, and $X = \cup \overset{\circ}{A}_n$.

¹A similar argument applies to \mathbb{E}^n .

Now, let U be open in Y . Since f is continuous on A_n , by theorem (2.8), f is continuous on $\overset{\circ}{A}_n$. Consequently,

$$f^{-1}(U) \cap \overset{\circ}{A}_n = f|_{\overset{\circ}{A}_n}^{-1}(U)$$

is open in $\overset{\circ}{A}_n$. So, there exists open sets V_n such that

$$V_n \cap \overset{\circ}{A}_n = f^{-1}(U) \cap \overset{\circ}{A}_n$$

And so,

$$f^{-1}(U) = f^{-1}(U) \cap \bigcup_n \overset{\circ}{A}_n = \bigcup_n (f^{-1}(U) \cap \overset{\circ}{A}_n) = \bigcup_n (V_n \cap \overset{\circ}{A}_n),$$

which is a union of open sets and so is open. \square

- (7) Let (X, τ) be a topological space, $A \subset X$, and χ_A its characteristic function. We describe the frontier of A in terms of χ :

Proof. We note that if $\overline{(X - A)} \cap \overline{A} \neq \emptyset$, then there exists $a \in \overline{(X - A)}$, $a \in \overline{A}$. With this in mind, we claim that χ_A is continuous at $a \in X$ if, and only if, $a \notin \text{Fr } A$.

Indeed;

(\implies): Suppose, to the contrary, that χ_A is continuous at $a \in X$, but that $a \in \text{Fr } A$. Then, $a \in \overline{X - A}$, and $a \in \overline{A}$. Now, let $\mathcal{O} = (.99, 1.99) \subset \mathbb{R}$, without loss of generality. Then, $\chi_A(a) = 1 \in \mathcal{O}$. Now, since χ_A is continuous, $\chi_A^{-1}[\mathcal{O}]$ is open, and such that $a \in \chi_A^{-1}[\mathcal{O}]$. But then, $\overline{X - A} \cap \chi_A^{-1}[\mathcal{O}] \neq \emptyset$. Further, this implies that $X - A \cap \chi_A^{-1}[\mathcal{O}] \neq \emptyset$. But then, $\chi_A(a) = 0 \in \mathcal{O}$, a contradiction.

(\impliedby):

\square

- (8) We determine which of the following maps are open or closed:

- $x \mapsto e^x$, on \mathbb{R} :

Proof. Open: Let $C = f(\mathbb{R})$. Since $f(A \cup B) = f(A) \cup f(B)$, it suffices to check that f is open on a base of open sets. Let B be an open interval in \mathbb{R} . If B has length greater than 2π then $f(B)$ is all of C . So $f(B)$ is open in C . Otherwise $f(B)$ is an open arc of the circle. Also open. Thus in all cases f maps B to an open set. Since open balls are a base f must be an open map.

Not Closed: To show f is not closed, for each $n \in \mathbb{N}$ let $E_n = [2n\pi + 1/n, (2n+1)\pi - 1/n]$. Let $E = \cup_n E_n$. Suppose x is a limit point of E .

Then $(x - 1/2, x + 1/2)$ intersects at most one E_n . Thus x is a limit point of E_n . Thus $x \in E_n$ since E_n is closed. Thus E is closed. But $f(E)$ is equal to C intersected with the upper half plane $\text{im}(z) > 0$. This is an open set in C , and not closed since $z = 1$ is a limit point of $f(E)$ not in $f(E)$. Thus $f(E)$ is not closed. \square

- $f : \mathbb{E}^2 \rightarrow \mathbb{E}^2$, $f(x, y) = (x, |y|)$:

Proof. Not Open: To see f is not open, let D be the open unit disc. Let $H^+ = \{(x, y) \mid y \geq 0\}$ and $H^- = \{(x, y) \mid y \leq 0\}$. Then $f(D) = D \cap H^+$. Therefore $z = 0 \in f(D)$ and every open ball containing z intersects $H^{+c} \subset f(D)^c$. Thus $f(D)$ is not open.

Closed: Now, suppose E is closed in \mathbb{E}^2 . Let $E' = E \cap H^-$ and let E'' be E' reflected about the x -axis. Then $f^{-1}(E) = (E \cap H^+) \cup (E'')$. Now E and H^+ are closed so $E \cap H^+$ is closed. And E'' is closed because H^- is closed, so E' is closed, and reflection is a homeomorphism. Thus $f^{-1}(E)$ is closed. \square

- $z \mapsto z^3$ on \mathbb{C} :

Proof. Open: We first show it is open. Let $A_{\theta_1, \theta_2, r_1, r_2} = \{re^{i\theta} \in \mathbb{C} \mid \theta_1, \theta_2, r_1, r_2 \in [0, \infty) \text{ and } \theta_1 < \theta < \theta_2, r_1 < r < r_2\}$. Then the set $\beta = \{A_{\theta_1, \theta_2, r_1, r_2} \mid \theta_2 - \theta_1 < 2\pi \text{ and } 0 \leq r_1 < r_2\}$ form a base for the usual topology on \mathbb{C} . The sets are clearly open and the intersection of any two of them is another one. Also for any $z \in A_{\theta_1, \theta_2, r_1, r_2}$ there is an open disc D s.t. $z \in D \subset A_{\theta_1, \theta_2, r_1, r_2}$ and for any open disc D with $z \in D$ we can find $\theta_1, \theta_2, r_1, r_2$ s.t. $z \in A_{\theta_1, \theta_2, r_1, r_2} \subset D$. Now consider what happens to $A_{\theta_1, \theta_2, r_1, r_2}$ under the function f . If $\theta_2 - \theta_1 > 2\pi/3$ then $f(A_{\theta_1, \theta_2, r_1, r_2})$ is a full open annulus. Otherwise $f(A_{\theta_1, \theta_2, r_1, r_2}) = A_{3\theta_1, 3\theta_2, r_1, r_2}$ which is open. Thus f takes basic open sets in β to open sets. Since $f(A \cup B) = f(A) \cup f(B)$ for any sets A and B , it suffices to check open-ness on the base β . Thus f is an open map.

Closed: We now show f is a closed map. Let E be any closed set in \mathbb{C} . Let Q_1 be the first quadrant $\{x + iy \mid x, y \geq 0\}$. Let Q_2, Q_3, Q_4 be the other (closed) quadrants. Then Q_1 is closed, so $E \cap Q_1$ is closed. The map f restricted to Q_1 is a homeomorphism from Q_1 to its image $f|_{Q_1} : Q_1 \rightarrow f(Q_1) = Q_1 \cup Q_2 \cup Q_3$ because its inverse $re^{i\theta} \mapsto re^{i\theta/3}$ is continuous. Thus $f(E \cap Q_1)$ is closed in Q_1 . Likewise $f(E \cap Q_i)$ is closed for all $i = 1, 2, 3, 4$. Functions respect unions, thus since $E = \cup_i (E \cap Q_i)$ it follows that $f(E) = \cup_i f(E \cap Q_i)$. But each $f(E \cap Q_i)$ is closed. Thus $f(E)$ is closed. \square

- (9) We show that the *unit ball* in \mathbb{E}^n (the set of points whose coordinates satisfy $x_1^2 + \cdots + x_n^2 \leq 1$) and the *unit cube* (points whose coordinates satisfy $|x_i| \leq 1$, $1 \leq i \leq n$) are homeomorphic if they are both given the subspace topology from \mathbb{E}^n :

Proof. Note that nowhere in the proof of lemma (2.10) is it used that we are in two dimensions. The proof goes through basically without change to any finite dimension where we replace "disc" with "ball". Now, let

$$f : \mathbb{E}^n - 0 \rightarrow \mathbb{E}^n - 0$$

be given by

$$f(\mathbf{v}) = \frac{1}{\|\mathbf{v}\|} \mathbf{v}$$

Then f is continuous and the image of f is the unit sphere. Let g be f restricted to *the surface* of the unit cube. Then g is one-to-one continuous. The intersection of an open ball with the surface of the cube maps to the intersection of an open ball with the sphere. Thus g is an open map. Therefore the inverse of g must be continuous. Therefore g is a homeomorphism. By the generalization of lemma (2.10), g may be extended from the boundaries to a homeomorphism from the whole cube to the whole ball. \square

2.3 Space-Filling Curves

- (1) We find a Peano curve which fills out the unit square in \mathbb{E}^2 :

Proof. We apply lemma 2.10. By a previous exercise, we have that $\partial[0, 1]^2$ is homeomorphic to ∂S^1 . Further, the boundary of the unit triangle is homeomorphic to ∂S^1 . Therefore, by lemma 2.10, there is a homeomorphism, f , from the unit triangle to the unit disc. Let $h : [0, 1] \rightarrow T$ be the space filling curve of the triangle as mentioned in Armstrong. Then, $f \circ h$ is a continuous mapping from $[0, 1] \rightarrow [0, 1]^2$. Further, it is space filling. \square

- (2) We find an onto, continuous function from $[0, 1]$ to S^2 :

Proof. From a previous exercise, we have that $\mathbb{E}^2 \cong_f (0, 1) \times (0, 1)$. Further, we have shown that $\mathbb{E}^2 \cong_\pi S^2 - (0, 0, 1)$. Thus, we extend f to $g : [0, 1]^2 \rightarrow S^2$ by

$$g(x) = \begin{cases} f(x) & x \notin \partial[0, 1]^2 \\ (0, 0, 1) & x \in \partial[0, 1]^2 \end{cases}$$

Then, if $\mathcal{O} \subset S^2$ is open, and contains $(0, 0, 1)$, then $g^{-1}[\mathcal{O}]$ is the entire square via homeomorphism. Now, if $\mathcal{O} \subset S^2$ does not contain $(0, 0, 1)$ and is open, then $g^{-1}[\mathcal{O}] = f^{-1}[\mathcal{O}]$ which is clearly open.

Thus, $g \circ h$, where h is the triangle space-filling curve, is the desired function. \square

- (3) We determine whether or no a space-filling curve can fill out the plane:

Proof. Note, $[0, 1]$ is compact, while \mathbb{E}^2 is not. \square

- (4) We determine whether or not a space filling curve can fill out all of the unit cube in \mathbb{E}^3 :

Proof. Let $I = [0, 1]$. Let $f : I \rightarrow I \times I$ be a space filling curve. Let $g : I \times I \rightarrow I \times I \times I \times I$ be the map $(x, y) \mapsto (f(x), f(y))$ with the natural identifications $(I \times I) \times (I \times I)$ with $I \times I \times I \times I$. Let p be projection onto the first three coordinates of $I \times I \times I \times I$. Then $p \circ g \circ f$ is a continuous function from I onto I^3 . \square

- (5) To prove a rigorous proof of this, at this point, is out of the question. However, via theorems (3.3), and (3.7), it is not true.

2.4 The Tietze Extension Theorem

Throughout these exercises, we assume that (X, d) is a metric space, and that if $A \subset X$, it has the subspace metric. Further, the topology on X is the induced topology, if not stated otherwise.

- (1) We show that $d(x, A) = 0$ if, and only if, $x \in \overline{A}$:

Proof.

Suppose that $d(x, A) = 0 = \inf_{a \in A} d(x, a)$. Then, for every $\epsilon > 0$, there exists $a \in A$, such that

$$|d(x, a) - 0| = d(x, a) < \epsilon$$

Now, let \mathcal{O} be an open subset of X , $x \in \mathcal{O}$. Choose $a \in A$ such that $d(x, a) < \epsilon$. Then, $a \in \mathcal{O}$, and $\mathcal{O} \cap A \neq \emptyset$. Thus, $x \in \overline{A}$.

Suppose that $x \in \overline{A}$. Then, we do the canonical ball construction and pick a sequence $\{a_n\}_n$ of elements of A , such that $d(x, a_n) \rightarrow 0, n \rightarrow \infty$.

\square

- (2) We show that if $A, B \subset X$ are disjoint and closed, there exists disjoint open sets U, V such that $A \subset U$ and $B \subset V$:

Proof. By lemma 2.13, there exists a continuous function $f : X \rightarrow [-1, 1]$, such that $f[A] \equiv 1$, and $f[B] \equiv -1$. Let $O_1 = [-1, 0]$, and $O_2 = [0, 1]$. Then, O_1, O_2 are open in $[-1, 1]$. Thus, $f^{-1}[O_i]$ is open, as f is continuous. Further, $f^{-1}[O_1] \cap f^{-1}[O_2] = \emptyset$, and $A \subset f^{-1}[O_1], B \subset f^{-1}[O_2]$. \square

- (3) We consider what topology the discrete metric gives a space:

Proof. We first show d is a metric. It is real-valued. And clearly

$$d(x, y) \geq 0 \text{ iff } x = y$$

Also clearly $d(x, y) = d(y, x)$. Finally, the only way we could have

$$d(x, y) + d(y, z) < d(x, z)$$

is if the left-hand-side is 0. But then $x = y = z$ and so the right-hand-side is also zero. Thus d is a metric.

Since

$$\{x\} = \{y \mid d(x, y) < 1/2\},$$

the sets $\{x\}$ are open. Since every set is a union of its points, every set is open. Thus this metric gives the discrete topology. \square

- (4) We show that every closed subset of a metric space is the intersection of a countable number of open sets:

Proof. Let A be a closed subset of X . Define

$$A_n = \{x \in X : d(x, A) < \frac{1}{n}\}$$

Then, A_n is open, for each $n \in \mathbb{N}$. The claim is that $\bigcap_n A_n = A$:

Clearly, $A \subset A_n$, for each n so, $A \subset \bigcap_n A_n$. Next, suppose that $x \in \bigcap_n A_n$, but $x \notin A$. Then,

$$1 > \inf_{a \in \bigcap_n A_n} d(x, a) = \epsilon > 0$$

By the Archimedian Principal, there exists n_0 large enough so that $1/n_0 < \epsilon$. But then,

$$x \notin \bigcap_n A_n$$

Thus, $\bigcap_n A_n \subset A$.

This concludes the proof. \square

- (5) If A, B are subsets of a metric space, their *distance apart* $d(A, B)$ is the infimum of the numbers $d(x, y)$ where $x \in A$ and $y \in B$. We find two disjoint closed subsets of the plane which are zero distance apart and check that both of the closed sets which you have just found have infinite diameter:

Proof. Let A be the x -axis and B be the set

$$\{(x, 1/x) \mid x > 0\}$$

The functions $x \mapsto 0$ and $x \mapsto \frac{1}{x}$ are continuous on $(0, \infty)$ so A and B are closed by chapter 2, problem 15.

Now, let

$$\{a_n\}_{n=1}^{\infty} := (n, 0), \quad \{b_n\}_{n=1}^{\infty} := (n, 1/n)$$

Then $a_n \in A$ and $b_n \in B$ and $d(a_n, b_n) = \frac{1}{n}$ and thusly,

$$d(A, B) < \frac{1}{n}, \quad \forall n$$

Consequently, $d(A, B) = 0$. Both sets clearly have infinite diameter. \square

- (6) We show that if A is a closed subset of X , then any map $f : A \rightarrow \mathbb{E}^n$ can be extended over X :

Proof. This is the Tietze extension theorem applied component-wise. \square

- (7) We find a map from $\mathbb{E}^1 - \{0\}$ to \mathbb{E}^1 which cannot be extended over \mathbb{E}^1 :

Proof. Let $f(x) = 1/x$. Then f is continuous on $\mathbb{E}^1 - \{0\}$ by Theorem 2.9 (b) because f^{-1} of an open interval is an open interval, or the union of two open intervals. Now suppose g extends f to all of \mathbb{E} . Let $a_n = \frac{1}{n}$. Then $a_n \rightarrow 0$. Thus $g(a_n) \rightarrow g(0)$. But $g(a_n) = f(a_n) = n \rightarrow \infty$. Thus no such g can exist. \square

- (8) This is left to the reader as an exercise.
- (9) Given a map $f : X \rightarrow \mathbb{E}^{n+1} - \{0\}$ we find a map $g : X \rightarrow S^n$ which agrees with f on the set $f^{-1}(S^n)$:

Proof. Let $h : \mathbb{E}^{n+1} - \{0\} \rightarrow S^n$ be given by

$$\mathbf{v} \mapsto \frac{1}{\|\mathbf{v}\|} \mathbf{v}$$

Then h is continuous and h is the identity on S^n . Let $g = h \circ f$. Then g is continuous and agrees with f on $f^{-1}(S^n)$. \square

- (10) If X is a metric space and A closed in X , we show that a map $f : A \rightarrow S^n$ can always be extended over a *neighborhood of A* , in other words over a subset of X which is a neighborhood of each point of A . (Think of S^n as a subspace of \mathbb{E}^{n+1} and extend f to a map of X into \mathbb{E}^{n+1} . now use Problem 35.):

Proof. Following the hint we think of S^n as a subspace of \mathbb{E}^{n+1} . Then $f = (f_1, \dots, f_n)$ and

$$f_i \equiv p_i \circ f,$$

where p_i is the i -th projection. The solution to problem 32 shows p_i is continuous, so that each f_i is continuous.

By theorem (2.15), each component f_i can be extended to a function g_i on all of X such that g_i agrees with f_i on A . Then $g = (g_1, \dots, g_n)$ extends f on A to a map from X to \mathbb{E}^{n+1} .

The same argument as in problem 32 shows g is continuous. Note that

$$g^{-1}(0) \cap A = \emptyset$$

because f maps A into S^n . And $g^{-1}(0)$ is a closed set in X (theorem (2.9(e))).

Thus by problem 28 we can find *disjoint* open sets U and V in X such that

$$A \subseteq U, \quad g^{-1}(0) \subseteq V$$

Let h be the map from problem 35. Then $h \circ g$ is well-defined as long as $g(x) \neq 0$. Thus $h \circ g$ is well-defined on U . And

$$h \circ g|_U$$

agrees with f on A , since h is the identity on S^n . □

Chapter 3

Compactness and Connectedness

3.1 Closed and Bounded Subsets of \mathbb{E}^n

There are not exercises for this section.

3.2 The Heine-Borel Theorem

- (1) This is left to the reader as an exercise.
- (2) Let $S \supseteq S_1 \supseteq S_2 \supseteq \cdots$ be a nested sequence of squares in the plane whose diameters tend to zero as we proceed along the sequence. We prove that the intersection of all these squares consists of exactly one point:

Proof. Each S_n is a square so is of the form $I_n \times J_n$ for closed one-dimensional intervals. And $S_n \supset S_{n+1}$ means $I_n \supset I_{n+1}$ and $J_n \supset J_{n+1}$. Thus we can apply the one-dimensional argument given above to the I_n 's and J_n 's. We get a sequence of points x_n converging to p and y_n converging to q . So $(x_n, y_n) \in S_n$ therefore converges to (p, q) which must be in S_n for all n since S_n is closed, so $(p, q) \in \bigcap_{n=1}^{\infty} S_n$. Similarly the one-dimensional argument shows there can be only a unique x and y coordinate of anything in $\bigcap_{n=1}^{\infty} S_n$. Thus,

$$\bigcap_{n=1}^{\infty} S_n = \{(p, q)\}$$

□

- (3) We use the Heine-Borel theorem to show that an infinite subset of a closed interval must have a limit point:

Proof. By 'infinite', we assume that the author means that the cardinality of the set is non-finite and countable. We proceed by contradiction:

Let C be the closed interval. Suppose that such an infinite subset, A , of C does not have any limit points. Then, for each $x \in C$, there is an open set \mathcal{O}_x in C , such that $x \in \mathcal{O}_x$ and $\mathcal{O}_x \cap (A - \{x\}) = \emptyset$. In addition,

$$C \subset \bigcup_{x \in C} \mathcal{O}_x$$

So, $\bigcup_{x \in C} \mathcal{O}_x$ is an open cover of C , and consequently, has a finite subcover:

$$\mathcal{F} = \bigcup_{\substack{x_i \\ i \in \{1, 2, \dots, N\}}} \mathcal{O}_{x_i}$$

Then, $A \subset \mathcal{F}$ and $\mathcal{O}_{x_i} \cap A = x_i$ for each i . Thus,

$$A = \mathcal{F} \cap A = \{x_1, x_2, \dots, x_N\}$$

A contradiction. □

- (4) We rephrase the definition of compactness in terms of closed sets:¹

Proof. We claim that X is compact if, and only if, for every collection $\{C_i\}_{i \in I}$ of closed sets in X with the FIP, $\bigcap_i C_i \neq \emptyset$. Indeed:

(\implies) We proceed by contradiction: Let X be compact, and let $\{C_i\}_{i \in I}$ be a collection of closed sets with the FIP, such that, $\bigcap_i C_i = \emptyset$.

Then, by De-Morgan's Laws, as C_i^c is open,

$$\bigcup_i C_i^c = \left(\bigcap_i C_i \right)^c = \emptyset^c = X$$

Thus, there exists a finite subcover,

$$\{C_{i_1}^c, C_{i_2}^c, \dots, C_{i_n}^c\},$$

with $X = \bigcup_k C_{i_k}^c$. And, so,

$$X^c = \emptyset = \left(\bigcup_k C_{i_k}^c \right)^c = \bigcap_k C_{i_k} \neq \emptyset$$

However, this is clearly a contradiction. Thus, $\bigcap_i C_i \neq \emptyset$.

¹Definition; The finite intersection property (FIP): Let \mathcal{F} be a collection of sets. Then, \mathcal{F} has the finite intersection property if whenever $F_1, F_2, \dots, F_n \in \mathcal{F}$, $F_1 \cap F_2 \cap \dots \cap F_n \neq \emptyset$.

(\Leftarrow) Let $\mathcal{F} = \{\mathcal{O}_i\}_{i \in I}$ be an open cover of X . Then, $\{\mathcal{O}_i^c\}_{i \in I}$ is a collection of closed sets, such that

$$\bigcap_i \mathcal{O}_i^c = \left(\bigcup_i \mathcal{O}_i \right)^c = X^c = \emptyset$$

Thus, there exists a finite set $\{\mathcal{O}_1^c, \mathcal{O}_2^c, \dots, \mathcal{O}_n^c\}$, such that

$$\bigcap_{k=1,2,\dots,n} \mathcal{O}_k^c = \emptyset$$

But then,

$$\left(\bigcap_{k=1,2,\dots,n} \mathcal{O}_k^c \right)^c = \bigcup_{k=1,2,\dots,n} \mathcal{O}_k = (X^c)^c = X$$

So, \mathcal{F} contains a finite subcover.

□

3.3 Properties of Compact Spaces

(1) We determine which of the following are compact:

- $\mathbb{Q} \subset \mathbb{E}$:

Proof. By theorem (3.9), a compact subset of \mathbb{E} is closed and bounded, \mathbb{Q} is neither. □

- S^n with a finite number of points removed:

Proof. $S^n - \{p_1, \dots, p_n\}$ is not compact; by theorem (3.9), a compact subset of \mathbb{R}^{n+1} is closed and bounded. Since S^n lives in \mathbb{R}^{n+1} , the theorem applies. S^n is bounded, but $S^n - \{p_1, \dots, p_n\}$ is not closed, because one can find a sequence in S^n that converges to any of the removed points. □

- the torus with an open disc removed:

Proof. Yes, the torus with an open disc removed is compact. The torus can be embedded in \mathbb{R}^3 as a bounded subset. And since we are removing an open disc, what remains is a closed subset of the torus and therefore (by chapter 2, problem 7, page 31) is a closed subset of \mathbb{R}^3 . Therefore, by theorem (3.9), it is compact. □

- the Klein Bottle:

Proof. The Klein bottle is compact. It is the continuous image of a closed finite rectangle. By theorem (3.9) a closed finite rectangle is compact. So by theorem (3.4) the Klein bottle is compact. Alternatively, the Klein bottle can be embedded into \mathbb{R}^4 as a closed and bounded set. Therefore, it is compact. \square

- the Möbius strip with its boundary circles removed:

Proof. The Möbius strip with its boundary circle removed is not compact. Think of the strip as a subset of \mathbb{R}^3 . Then one can find a sequence of points in the strip that converge to a point on the boundary, which has been removed. Since compact sets must be closed in \mathbb{R}^3 , the Möbius with boundary removed cannot be compact. \square

- (2) We show that the Hausdorff condition cannot be relaxed in theorem (3.7):

Proof. Let $X = \{a, b\}$ a set with two points. Let X_1 be X with the discrete topology (so every subset is open). Let X_2 be X with the indiscrete topology (in other words the only open sets are X and \emptyset). Then X_2 is not Hausdorff. The function $f : X_1 \rightarrow X_2$ given by $f(a) = a$, $f(b) = b$ is one-to-one, onto and continuous. But f^{-1} is not continuous, so f is not a homeomorphism. \square

- (3) We show that Lebesgue's Lemma fails for \mathbb{E}^2 :

Proof. Let $B(0, 1)$ be the open ball of radius one, centered around the origin. And, for each $p \neq 0$ in \mathbb{E}^2 , let $B(p, 1/\|p\|_2)$ be the open ball centered at p and radius $1/\|p\|_2$. Now, let $\delta > 0$, and let $n \in \mathbb{N}$ be such that

$$\frac{1}{n} < \frac{\delta}{3}$$

Then, the open ball of radius $2\delta/3$ around the point $(n, 0)$ is not contained in any $B(p, 1/\|p\|_2)$. \square

- (4) *Lindelöf's Theorem:* We show that if X has a countable base for its topology, τ , then any open cover of X contains a countable subcover:

Proof. Suppose that \mathcal{F} is a countable base for the topology on (X, τ) . Then, $\mathcal{F} = \{F_1, F_2, \dots, F_n, \dots\}$. Let $\{\mathcal{O}_\alpha\}_\alpha$ be an open cover of X . Then, as \mathcal{F} is a base for τ ,

$$\bigcup_{\alpha} \mathcal{O}_\alpha = \bigcup_{i \in I} F_i$$

In addition, for each $x \in X$ there exists $n \in I$, and α_0 , such that

$$x \in F_n \subset \mathcal{O}_{\alpha_0}$$

Thus, let

$$\{F_{n_k}\}_k$$

be the collection of all such F_{n_k} , as described above. Then, clearly

$$\bigcup_k F_{n_k} = X$$

and, for each $F_{n_k} \in \{F_{n_k}\}_k$, there is an α_{n_k} , such that

$$F_{n_k} \subset \mathcal{O}_{\alpha_{n_k}}$$

Consequently,

$$\{\mathcal{O}_{\alpha_{n_k}}\}_{n_k}$$

is a subcover of $\{\mathcal{O}_\alpha\}_\alpha$. Finally, as I is countable, so is $\{F_{n_k}\}$ and $\{\mathcal{O}_{\alpha_{n_k}}\}$. \square

- (5) This is shown by an extra lemma, and the fact that compact subsets of T_2 spaces are closed.
- (6) Let A be a compact subset of a metric space (X, d) . We show,

- that the diameter of A , $\text{Diam } A$, is equal to $d(x, y)$ for some $x, y \in A$:

Proof. By lemma (2.13), for a fixed $y \in A$, the function defined by $f(x) = d(x, \{y\})$ is continuous. By theorem 3.10, f is bounded, and obtains its bounds on A ; for some $x_0 \in A$, $d(x, \{y\}) \leq d(x_0, \{y\})$ for all $x \in A$. Further, ranging over $y \in A$, we see that

$$\text{Diam } A = d(x_0, y_0), \quad x_0, y_0 \in A.$$

\square

- that given $x \in X$, $d(x, A) = d(x, y)$, for some $y \in A$:

Proof. This is lemma (2.13), and the fact that A is compact. \square

- that given a closed subset B , disjoint from A , that $d(A, B) > 0$:

Proof. Suppose, to the contrary, that $d(A, B) = 0$. Then, by definition, we have

$$\inf_{\substack{x \in A \\ y \in B}} d(x, y) = d(A, B) = 0$$

An application of the above shows that, actually, $d(A, B) = 0 = d(a, b)$ for some $a \in A$, and $b \in B$. But then, by properties of metrics spaces, we have $a = b$; a contraction, as $A \cap B = \emptyset$. \square

- (7) We find an example of a topological space (X, τ) and a compact subset whose closure is not compact:

Proof. Consider $\mathbb{R} = X$, and $\tau = \{\mathbb{R}, \emptyset, (-a, a) : a > 0\}$. Then, $\{0\}$ is clearly compact in X , and it is closed. But, $\{0\}^c = \mathbb{R}$ as each $(-a, a)^c = (-\infty, -a] \cup [a, \infty)$ and $0 \notin (-\infty, -a] \cup [a, \infty)$. \square

- (8) We show that the real numbers with the finite complement topology are compact, and that the reals with the half-open topology is not compact:

Proof. FC: The real numbers with finite-complement topology is compact. Let $\beta = \{U_\alpha\}$ be an open cover. Then let U_{α_0} be an element of β . Then there are only a finite number n of points that are not in U_{α_0} . We only need n more elements of β to cover everything.

HO: The real numbers with the half-open interval topology is not compact. Let β be the open cover $\{[n, n+1) \mid n \in \mathbb{Z}\}$. Then β clearly does not have a finite subcover. \square

- (9) Let $f : X \rightarrow Y$ be a closed map such that $f^{-1}(y)$, $y \in Y$ is a compact subset of X . We show that $f^{-1}[K]$ is compact whenever K is compact in Y :

Proof. This follows from the fact that

$$f^{-1}[K] = f^{-1}\left[\bigcup_{k \in K} \{k\}\right] = \bigcup_{k \in K} f^{-1}[\{k\}]$$

and the fact that the union of finite sets is finite. \square

- (10) This is left to the reader as an exercise. Hint: Use the fact that a subset of a T_2 space is T_2 , and theorem 3.7. Show that $f : X \rightarrow f(X)$ is onto.
- (11) The proof of the first part is left to the reader. However, we show

- that any closed subset of a locally compact space is locally compact:

Proof. Let (X, τ) be the locally compact space, and $A \subset X$, with $\overline{A} = A$. Let $p \in A$. Then, as X is locally compact, there exists a compact neighbourhood

$$K_X \subset X, \quad p \in K_X$$

Then, via the subspace topology, $K_X \cap A$ is a neighbourhood of p in A .

Next, let \mathcal{F} be an open cover of $K_X \cap A$. Then, we see that $\mathcal{F} \cup \{A^c\}$ is an open cover of K_X . Further, there exists a finite subcover $\mathcal{F}_1 \subset \mathcal{F}$, such that $\mathcal{F}_1 - \{A^c\}$ is a finite subcover of

$$K_X \cap A \subset \bigcup_1 \mathcal{F}_1 - \{A^c\}$$

and so $K_X \cap A$ is a compact neighbourhood of p in A . \square

- that \mathbb{Q} is not locally compact (as a subset of \mathbb{R}):

Proof. Suppose, to the contrary, that \mathbb{Q} is locally compact. Then, let $A \subset \mathbb{Q}$ be a compact neighbourhood. Then, there exists an open interval $I \subset \mathbb{R}$, with

$$I \cap \mathbb{Q} \subset A$$

Now, let $x \in I$, where x is irrational. As \mathbb{Q} is dense in \mathbb{R} , there exists a sequence of rationals in $\mathbb{Q} \cap A$ with,

$$\{q_n\} \rightarrow x \notin A, \quad n \rightarrow \infty$$

But then, A is not closed. Thus, by theorem (3.5), it cannot be compact. \square

- that local compactness is preserved under homeomorphism:

Proof. Suppose that $f : X \rightarrow Y$ is a homeomorphism, and that X is locally compact. We want to show that Y is locally compact.

Let $p \in Y$. Then, there is a locally compact neighbourhood K_p of $f^{-1}(p)$ in X . Now, let \mathcal{F} be an open cover of $f(\{K_p\})$. Then,

$$\mathcal{F}_1 = \{f^{-1}(\{F\}) : F \in \mathcal{F}\}$$

is an open cover of K_p in X . It follows that \mathcal{F}_1 has a finite subcover, \mathcal{F}_2 , and that

$$\{f(\{F\}) : F \in \mathcal{F}_2\}$$

is a finite subcover for $f(\{K_p\}) \subset Y$. \square

- (12) Suppose that X is locally compact and Hausdorff. Given $x \in X$, and a neighbourhood \mathcal{O} of x , we find a compact neighbourhood K of x which is contained in \mathcal{O} :

Proof. Let \mathcal{O} be as given. Then, since X is locally compact, there exists a compact neighbourhood V , such that $x \in V$. By the subspace topology, it follows that $x \in V \cap \mathcal{O}$ and $\mathcal{O} \cap V$ is open in V . Further, as V is compact, it is closed, by theorem 3.6. \square

- (13) Let X be a locally compact Hausdorff space which is not compact. Form a new space by adding one extra point, usually denoted by ∞ , to X and taking the open sets of $X \cup \{\infty\}$ to be those of X together with sets of the form $\{X - K\} \cup \{\infty\}$, where K is a compact subset of x . We check the axioms for a topology, and show that $X \cup \{\infty\}$ is a compact Hausdorff space which contains X as a dense subset:

Proof. Let β be the topology on X . Define

$$\beta' = \beta \cup \{U \cup \{\infty\} \mid U \subseteq X \text{ and } U^c \text{ is compact} \}$$

Clearly, $\emptyset \in \beta'$ and since \emptyset is compact,

$$X \cup \{\infty\} \in \beta'$$

It remains to show β' is closed with respect to arbitrary unions and finite intersection:

- Arbitrary Unions: Let

$$\{U_\alpha\}_{\alpha \in A} \subseteq \beta' \subset \beta'$$

We must show $\cup_{\alpha \in A} U_\alpha \in \beta'$; Write

$$\begin{aligned} A &= A_1 \cup A_2 \\ \alpha \in A_1 &\Rightarrow \infty \notin U_\alpha \\ \alpha \in A_2 &\Rightarrow \infty \in U_\alpha \end{aligned}$$

Then,

$$\{U_\alpha\}_{\alpha \in A} = \{U_\alpha\}_{\alpha \in A_1} \cup \{U_\alpha\}_{\alpha \in A_2}$$

If $A_2 = \emptyset$, then

$$\bigcup_{\alpha \in A} U_\alpha = \bigcup_{\alpha \in A_1} U_\alpha \in \beta \subseteq \beta'$$

If $A_2 \neq \emptyset$, then we must show that

$$\left(\bigcup_{\alpha \in A} U_\alpha \right)^c$$

is compact.

For $\alpha \in A_2$, U_α^c is compact: Since X is Hausdorff U_α^c is closed (theorem (3.6)). Thus

$$\bigcap_{\alpha \in A_2} U_\alpha^c$$

is closed. Now, fix any $\alpha' \in A_2$. Then $\cap_{\alpha \in A_2} U_\alpha^c$ is a closed subset of $U_{\alpha'}^c$, and so is compact (theorem 3.5). Thus

$$\left(\bigcup_{\alpha \in A_2} U_\alpha \right)^c$$

is compact. Thus,

$$\left(\bigcup_{\alpha \in A} U_\alpha \right)^c = \left(\bigcap_{\alpha \in A_1} U_\alpha^c \right) \cap \left(\bigcap_{\alpha \in A_2} U_\alpha^c \right)$$

, which is a closed subset of the compact set $\cap_{\alpha \in A_2} U_\alpha^c$ and therefore is compact.

- Finite intersections: Now, let

$$\{U_\alpha\}_{\alpha \in A} \subseteq \beta'$$

be a finite subset of β' . As above, decompose A into $A_1 \cup A_2$. If $A_1 \neq \emptyset$, then

$$\infty \in \left(\bigcap_{\alpha \in A} U_\alpha \right)$$

And,

$$\left(\bigcap_{\alpha \in A} U_\alpha \right)^c = \left(\bigcap_{\alpha \in A_2} U_\alpha \right)^c = \bigcup_{\alpha \in A_2} U_\alpha^c$$

is a finite union of compact sets and therefore compact.

If $A_1 \neq \emptyset$, then

$$\infty \notin \left(\bigcap_{\alpha \in A} U_\alpha \right)$$

and

$$\left(\bigcap_{\alpha \in A} U_\alpha \right)^c = \left(\bigcup_{\alpha \in A_1} U_\alpha^c \right) \cup \left(\bigcup_{\alpha \in A_2} U_\alpha^c \right)$$

These are all finite unions, thus this last expression is the union of a closed and a compact and is therefore closed. Thus $(\bigcap_{\alpha \in A} U_\alpha)^c$ is closed. Thus $(\bigcap_{\alpha \in A} U_\alpha)$ is open.

$X \cup \{\infty\}$ is compact: Let $\{U_\alpha\}$ be an open cover of X . Then $\exists \alpha_0$ such that $\infty \in U_{\alpha_0}$. Since $U_{\alpha_0}^c$ is compact, and $\{U_\alpha\}$ is an open cover of $U_{\alpha_0}^c$, there are $U_{\alpha_1}, \dots, U_{\alpha_n}$ such that

$$U_{\alpha_0}^c \subseteq U_{\alpha_1} \cup \dots \cup U_{\alpha_n}$$

But, then

$$U_{\alpha_0}, U_{\alpha_1}, \dots, U_{\alpha_n}$$

is an open cover of $X \cup \{\infty\}$ and so we have found a finite subcover.

We now show $X \cup \{\infty\}$ is Hausdorff. Let $x, y \in X \cup \{\infty\}$. If $x, y \in X$ then since X itself is Hausdorff we know there are open sets that separate them. So suppose without loss of generality that $y = \infty$. Since X is locally compact there is an open set $U \subseteq X$ and a compact set $K \subseteq X$ with $x \in U \subseteq K$. Then $\infty \in K^c$ is an open set in $X \cup \{\infty\}$ and $K^c \cap U = \emptyset$.

It remains to show X is dense in $X \cup \{\infty\}$. The only way this is not true is if $\{\infty\}$ is an open set. But $\{\infty\}^c = X$ which was assumed to be *not* compact. \square

(14) We prove that $\mathbb{E}^n \cup \{\infty\}$ is homeomorphic to S^n :

Proof. As before ², there exists a homeomorphism

$$h : \mathbb{E}^n \rightarrow S^n - \{p\}, \quad p \in S^n$$

Clearly, the one point compactification of \mathbb{E}^n is $\mathbb{E}^n \cup \{\infty\}$. And further, the one point compactification of $S^n - \{p\}$ is S^n .

As both these spaces are T_2 , it follows by an extra lemma that

$$\mathbb{E}^n \cup \{\infty\} \simeq S^n$$

□

- (15) Let X , and Y be locally compact Hausdorff spaces, and let $f : X \rightarrow Y$ be an onto map. We show that f extends to a map from $X \cup \{\infty\}$ onto $Y \cup \{\infty\}$, if and only if, $f^{-1}[K]$ is compact for each compact subset K of Y . Further, we deduce that if X and Y are homeomorphic spaces, then so are their one-point compactifications and find two spaces which are not homeomorphic, but have homeomorphic one-point compactifications:

Proof. This problem was first shown as an extra lemma. Thus, we find two spaces which are not homeomorphic but have homeomorphic one point compactifications:

Consider

$$X_1 = [0, 1) \cup (1/2, 1] \quad X_2 = [0, 1)$$

and the result follows. □

3.4 Product Spaces

- (1) This is left to the reader as an exercise.
- (2) Suppose that A, B are compact in X, Y , respectively. We show that if W is a neighbourhood of $A \times B$ in $X \times Y$, that there exists a neighbourhood U of A in X , and a neighbourhood V of B in Y , such that

$$U \times V \subset W$$

Proof. Since A, B are compact, by theorem (3.15), $A \times B$ is compact. As this is the case, there exists a finite set of (a, b) , $a \in A$, $b \in B$, with neighbourhoods

$$U_{a_1} \times V_{b_1}, U_{a_2} \times V_{b_2}, \dots, U_{a_n} \times V_{b_n}$$

such that

$$A \times B \subset \bigcup_i U_{a_i} \times V_{b_i}$$

²reference this

Now, for each $a \in A$, let

$$E_a = \bigcap_{\substack{U_{a_i} \\ a \in U_{a_i}}} U_{a_i},$$

and

$$F_b = \bigcap_{\substack{V_{b_i} \\ b \in V_{b_i}}} V_{b_i}$$

for each $b \in B$. Then, as $\{U_{a_i} \times V_{b_i}\}_i$ is finite,

$$\{E_x \times F_y : x \in A, y \in B\}$$

is finite. Further, we see that

$$A \times B \subset \bigcup_{a,b} E_a \times F_b = U \times V$$

Consequently, $U \times V \subset W$, by construction. \square

(3) We prove that

- the product of two second-countable spaces is second-countable:

Proof. Let X, Y be second countable spaces. Then, as the finite union of countable sets is countable, it follows that the topology on $X \times Y$ has a countable base. \square

- the product of two separable spaces is separable:

Proof. Suppose that X, Y are separable. Then, they contain a countable dense subset, $A \subset X$, $B \subset Y$. From set theory, it follows that $A \times B$ is countable. Thus, we show that $\overline{A \times B} = X \times Y$:

Indeed; from exercise twenty on page fifty-five, we have

$$\overline{A \times B} = \overline{A} \times \overline{B} = X \times Y$$

\square

(4) We prove that $[0, 1) \times [0, 1) \simeq [0, 1] \times [0, 1)$:

Proof. As previously show, there exists some $h : [0, 1) \times [0, 1) \rightarrow D_1^2$,

$$D_1^2 = D^2 - \{x = (x_1, x_2) \in S^1 : x_1 > 0 \wedge x_2 > 0\},$$

a homeomorphism. Similarly, under the same map, $h : [0, 1] \times [0, 1) \rightarrow D_2^2$,

$$D_2^2 = D^2 - \{x = (x_1, x_2) \in S^1 : x_2 > \sqrt{2}/2\},$$

is a homeomorphism. But, $\{x = (x_1, x_2) \in S^1 : x_2 > \sqrt{2}/2\} = B$, $\{x = (x_1, x_2) \in S^1 : x_1 > 0 \wedge x_2 > 0\} = A$ are open in S^1 , and so homeomorphic via, say, h' .

Consequently,

$$S^1 - A \simeq_{h'} S^1 - B$$

But then,

$$D_1^2 = \text{Int } D^2 \cup (S^1 - A) \simeq_{h' \circ i} \text{Int } D^2 \cup (S^1 - B) = D_2^2$$

It follows that

$$[0, 1] \times [0, 1] \simeq_h D_1^2 \simeq D_2^2 \simeq_h [0, 1] \times [0, 1]$$

□

- (5) Let $x_0 \in X$ and $y_0 \in Y$. We prove that the functions $f : X \rightarrow X \times Y$, $g : Y \rightarrow X \times Y$ defined by $f(x) = (x, y_0)$, $g(y) = (x_0, y)$ are embeddings (as defined in problem 14):

Proof. By theorem (3.13), f is an embedding iff $p_1 \circ f$ and $p_2 \circ f$ are continuous. Thus, since $p_1 \circ f(x) = x$ (the identity function) and $p_2 \circ f(x) = y_0$ (a constant function), it follows that f is continuous. Now f is clearly one-to-one and onto its image

$$f(X) \subseteq X \times Y$$

Note that $p_1|_{f(X)}$ is the inverse of f on $f(X)$. Since p_1 is continuous, it follows that $p_1|_{f(X)}$ is continuous on $f(X)$. Thus f is a homeomorphism. The proof that g is a homeomorphism is basically identical. □

- (6) The first part is trivial. So, we show that X is Hausdorff if, and only if, $\Delta(\{X\})$ is closed in $X \times X$:

Proof. (\implies) Suppose that X is T_2 . We show that $\Delta(\{X\})^c$ is open in $X \times X$. Now, for $(x, y) \notin \Delta(\{X\})$, we have that there exists open sets, U_x, U_y , such that

$$x \in U_x, \quad y \in U_y, \quad U_x \cap U_y = \emptyset$$

Now, let $U_x \times U_y = W$, which is open in $X \times X$. If $p \in W \cap \Delta(\{X\})$, then $p \in W$, and so $U_x \cap U_y \neq \emptyset$. Thus,

$$W \subset \Delta(\{X\})^c$$

and so $\Delta(\{X\})$ is closed.

(\Leftarrow) Suppose that $\Delta(\{X\})$ is closed in $X \times X$. Now, if $(x, y) \notin \Delta(\{X\})$, then $(x, y) \in \Delta(\{X\})^c$, which is open. And so, by definition of the topology on $X \times X$, there exist open sets U, V , such that

$$(x, y) \in U \times V \subset \Delta(\{X\})^c$$

But, then

$$(U \times V) \cap \Delta(\{X\}) = \emptyset,$$

implies $U \cap V = \emptyset$, by set theory.

□

- (7) This is left to the reader as an exercise. Hint: Consider $f : \mathbb{R}^+ \rightarrow \mathbb{R}$, given by $f(x) = \frac{1}{x}$ and use the sequential criterion for closure.
- (8) Given a countable number of spaces X_1, X_2, \dots , a typical point of the product $\prod X_i$ will be written $x = (x_1, x_2, \dots)$. The *product topology* on $\prod X_i$ is the smallest topology for which all of the projections $p_i : \prod X_i \rightarrow X_i$, $p_i(x) = x_i$, are continuous. We construct a base for this topology from the open sets of the spaces X_1, X_2, \dots :

Proof. Define the base β for a topology on $\prod X_i$ by

$$\beta = \{\emptyset\} \cup \{U_1 \times U_2 \times \dots \mid U_i \in X_i\}$$

is open, and $U_i = X_i$ for all but finitely many i . Clearly,

$$\emptyset \in \beta \quad \prod X_i \in \beta$$

It is also obvious that β is closed under finite intersection since

$$\bigcap_{\alpha} (A_{\alpha} \times B) = \left(\bigcap_{\alpha} A_{\alpha} \right) \times B$$

Now let $U \subseteq X_i$. Then,

$$p_i^{-1}(U) = X_1 \times \dots \times X_{i-1} \times U \times X_{i+1} \times X_{i+2} \times \dots \in \beta$$

Thus, p_i is continuous.

Now, suppose B is the smallest topology for which p_i is continuous. Then for $U \subseteq X_i$,

$$p_i^{-1}(U) = X_1 \times \dots \times X_{i-1} \times U \times X_{i+1} \times X_{i+2} \times \dots$$

must be in B . Since any element of β is a finite intersection of such sets, it follows that $\beta \subseteq B$. Since B is the smallest topology for which p_i is continuous for all i , and the topology generated by β is contained in B , it must be that the topology generated by β equals B . □

(9) This is left to the reader as an exercise. Hint: Consider the function $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ given by $f(x) = \frac{x}{1+x}$.

(10) We show that

- the box topology contains the product topology:

Proof. As X_i is open in (X_i, τ_i) , the result follows. \square

- the box topology and the product topology are equal if, and only if, X_i is an indiscrete space for all but finitely many values of i :

Proof. If $U_i = \emptyset$, for any i , then

$$\prod X_i = \emptyset$$

And, so the only sets of the form

$$U_1 \times U_2 \times \dots$$

which are non-empty, are those which have $U_i = X_i$ for all but finitely many i . \square

3.5 Connectedness

(1) Let $X = \{(a, b) \in \mathbb{E}^2 : a \in \mathbb{Q} \vee b \in \mathbb{Q}\}$. We show that X with the induced topology is connected:

Proof. We seek to apply theorem (3.25), on page fifty-eight. Indeed, for each $q \in \mathbb{Q}$, let

$$X_q = \{(a, b) \in \mathbb{E}^2 : a = q \vee b = q\}$$

Then clearly

$$X_q = A_q \cup B_q,$$

where $A_q = \{(a, b) \in \mathbb{E}^2 : a = q, b \in \mathbb{E}\}$, $B_q = \{(a, b) \in \mathbb{E}^2 : a \in \mathbb{E} \vee b = q\}$. Further, for each $q \in \mathbb{Q}$,

$$A_q \cong \mathbb{E} \cong B_q$$

So, A_q, B_q are connected, and as $A_q \cap B_q = \{(q, q)\}$, X_q is connected by theorem (3.25).

For $t, q \in \mathbb{Q}$, we have

$$X_t \cap X_q = \{(q, t), (t, q)\} \neq \emptyset$$

and so, another application of 3.25 shows that

$$X = \bigcup_{q \in \mathbb{Q}} X_q$$

is connected. \square

- (2) Give the real numbers the finite-complement topology. We determine the connected components of the resulting space, and answer the same question for the half-open interval topology:

Proof.

- FC: The entire space is connected, since no set can be both open and closed.
- HO: The space is totally disconnected. Let $x < y$. Then by chapter 2, problem 11 every set $[a, b)$ is both open and closed. Thus, $[x, y)$ contains x and not y , and $[x, y)$ and $[x, y)^c$ are both open. Thus, x and y are not in the same connected component. Since x and y are arbitrary, connected components cannot contain more than one point.

□

- (3) If X has only a finite number of components, we show that each component is both open and closed and find a space none of whose components are open sets:

Proof. $X = C_1 \cup \cdots \cup C_n$ a disjoint union of connected components. By theorem (3.27), each C_i is closed. Since a finite union of closed sets is closed, $X - C_i = \cup_{j \neq i} C_j$ is closed. Thus,

$$C_i = \left(\bigcup_{j \neq i} C_j \right)^c$$

is open.

The space \mathbb{Q} with the natural metric topology is totally disconnected, but points in \mathbb{Q} are not open sets. □

- (4) (Intermediate Value Theorem) We show that if $f : [a, b] \rightarrow \mathbb{E}$ is a map such that $f(a) < 0$ and $f(b) > 0$, then there exists some $c \in [a, b]$ for which $f(c) = 0$:

Proof. By theorem (3.21), page fifty-eight, $f([a, b]) \subset \mathbb{E}$ is connected. By theorem (3.19), page fifty-seven, $f([a, b])$ is an interval. In particular, $f([a, b]) = [f(a'), f(b')]$, where $f(a')$, $f(b')$, are the minimum and maximum of f on $[a, b]$. Thus, $[f(a), f(b)] \subset [f(a'), f(b')]$, and $0 \in [f(a), f(b)]$. Thus, there exists $c \in [a, b]$, $f(c) = 0$. □

- (5) We show that

- \mathbb{E}^n is locally connected:

Proof. As has been shown,

$$\{I_1 \times I_2 \times \cdots \times I_n : I_i \in \tau_{\mathbb{E}}\}$$

is a base for the topology in \mathbb{E}^n . Thus, if $x \in \mathcal{O}_x$ for some open neighbourhood \mathcal{O}_x in \mathbb{E}^n , there exists

$$I = I_1 \times I_2 \times \cdots \times I_n \subset \mathcal{O}_x$$

And, as each I_i is connected, it follows by exercise 3.20, that I is connected. \square

- $X = \{0\} \cup \{1/n : n \in \mathbb{N}\}$ is *not* locally connected:

Proof. Let \mathcal{O} be an open neighbourhood of $0 \in X$, which is connected. But then, by theorem (3.19), \mathcal{O} is an interval. However, X clearly contains no intervals. \square

- (6) We show that local connectedness is preserved by a homeomorphism, but need not be preserved by a continuous function:

Proof. The first part of this proof is direct. For the second part, consider $Y = \{0\} \cup \{1/n : n \in \mathbb{N}\}$ as above, and $X = \mathbb{Z}^+$ as subspaces of \mathbb{E} . Let $f : X \hookrightarrow Y$ be the canonical map. Then, f is clearly continuous, but we have previously show that Y is not locally compact. \square

- (7) We show that (X, τ) is locally connected, if, and only if every component of each open subset of X is an open set:

Proof. (\implies): Suppose that X is connected, and let \mathcal{O} be an open set in X , and C a (maximal) connected component of \mathcal{O} . Then, for some $x \in C$, there exists a connected open set V_x , such that

$$x \in V_x \subset \mathcal{O}$$

But, since C is maximally connected, $V_x \subset C$. Then,

$$C = \bigcup_{x \in C} V_x$$

and so C is open.

(\impliedby): Now, let $x \in X$, and \mathcal{O} an open set containing x . Then, the components of \mathcal{O} are open. Thus, let V be a component such that $x \in V \subset \mathcal{O}$. But then, V is open, closed, and so connected. Thus, X is locally connected. \square

3.6 Joining Points by Paths

- (1) We show that the continuous image of a path-connected space is path-connected:

Proof. Without loss of generality, suppose that $f : X \rightarrow Y = f(X)$ is onto, and X is path-connected. Then, for

$$y_1 = f^{-1}(x_1), \quad y_2 = f^{-1}(x_2)$$

in Y , there exists a path h from x_1 to x_2 . Thus, $f \circ h$ is the desired path. \square

- (2) This is left to the reader as an exercise. Hint: Consider the case $x \neq -y$, and use the straight-line homotopy composed with the canonical map. In the case where $x = -y$, pick a third point which is not equal and apply the above technique.

- (3) We prove that the product of two path-connected spaces is path-connected:

Proof. This follows directly from the definition of the product topology and set theory. \square

- (4) If A and B are path-connected subsets of a space (X, τ) , and $A \cap B$ is non-empty, we show that $A \cup B$ is path-connected:

Proof. Suppose, without loss of generality, that

$$x, y \in A \cup B, \quad x \in A, \quad x \in B$$

Pick $c \in A \cap B$, and by assumption, there exists paths γ_1 in A , γ_2 in B , which connect x to c , and c to y , respectively.

It follows that $\gamma_1 \circ \gamma_2$ is the desired path. \square

- (5) We find a path-connected subset of a space whose closure is not path-connected:

Proof. As the comments on page sixty-two point out, letting

$$Z = \{(x, \sin(\pi/x)) : x \in \mathbb{R}^+\}$$

we see

$$\overline{Z} = Y \cup Z = X$$

\square

- (6) We show that any indiscrete space is path-connected:

Proof. As any function on an indiscrete space is continuous, the result follows. \square

- (7) We determine whether or not the space shown in fig. 3.4, page sixty-three, is locally path-connected, and convert $X = \{0\} \cup \{1/n : n = 1, 2, \dots\}$ into a subspace of \mathbb{E}^2 which is path-connected but not locally path-connected:

Proof. It is not the case that fig. 3.4 is locally path-connected. To see this, we note that any path at the origin must contain points from Y and Z . As the comments on page sixty-two point out, the result follows.

For the second part of the proof, let

$$X_0 = \{(0, y) : y \in [0, 1]\}, \quad X_n = \{(1/n, y) : y \in [0, 1]\}$$

Further, let $Y = \{(x, 0) : x \in [0, 1]\}$ and

$$X = Y \cup \left(\bigcup_n X_n \right)$$

Then, X is path connected as $Y \cap X_n \neq \emptyset$, for all n .

To see that X is not path connected, let $p \in (0, 1)$, and consider $\mathcal{O} \subset B(p, 1)$ be open. Then, $\mathcal{O} \cap Y = \emptyset$, so $\mathcal{O} \cap X$ is a collection of line segments separate from each other. So, $V \cap X$ is not path connected. \square

- (8) We prove that a space which is connected, and locally path-connected is path-connected:

Proof. Let $x \in X$. And let \mathcal{O}_x be the set of all $y \in X$ such that y is path-connected to x ; then, $\mathcal{O}_x \neq \emptyset$. We claim that $\mathcal{O}_x = X$.

Indeed; as \mathcal{O}_x is a maximally connected component it is open, by extra lemma³. Similarly, \mathcal{O}_x^c is open. Thus, \mathcal{O}_x is both open and closed. Therefore, as X is connected, $\mathcal{O}_x = X$. This completes the proof. \square

³cite this

Chapter 4

Identification Spaces

4.1 Constructing the Möbius Strip

There are no exercises listed for this section.

4.2 The Identification Topology

- (1) We check that the three descriptions of $\mathbb{R}P^n$ all lead to the same space:

Proof. We first note that the canonical construction of $\mathbb{R}P^n$ is formed via the relation

$$x \sim_{\mathcal{R}} \lambda x \quad \forall \lambda \neq 0, \quad \lambda \in \mathbb{R}$$

As λ can always be chosen so that $\|x \cdot \lambda\| = 1$, we consider such λ as equivalence class representatives.

To see that (a), (b) are equivalent, let $h : \mathbb{E}^{n+1} \rightarrow \mathbb{E} / \sim_{\mathcal{R}}$ be the aforementioned quotient map and $i : S^n \rightarrow \mathbb{E}^{n+1}$ be the natural embedding. Then, as S^n is compact, $\mathbb{E} / \sim_{\mathcal{R}}$ is Hausdorff, it follows by an extra lemma, and corollary (4.4), that $h \circ i$ is an identification map. So, (a), (b) are equivalent.

To see that (b) and (c) are equivalent, let $i : S^n \rightarrow B^n$ be the natural embedding, and g is quotient map given in (c). Then, again, by corollary (4.4), $g \circ i$ is an identification map. Thus, (b) and (c) are equivalent. \square

- (2) We determine which space do we obtain if we take a Möbius strip and identify its boundary circle to a point:

Proof. We obtain the projective plane $\mathbb{R}P^2$. To see this, let R be a rectangle as in Fig. 4.1 (page 66). Instead of first identifying the right and left edges to get a Möbius and then collapsing the boundary circle, we first collapse the top and bottom borders to two points a and b , then identify the (image of) the right and left edges as they would have been identified to form a Möbius

from R . This operation identifies the two points a and b . So after these two identifications we end up with the same space. So by this argument we can do the identifications in the opposite order. Proceeding in this way, we first collapse the top horizontal border to one point and the bottom border to another point. From this identification we get a space homeomorphic to a disc D . The identification map $f : R \rightarrow D$ can be seen as taking the left border of R to the left semi-circle of the boundary of D and the right border to the right semi-circle of the boundary of D . Then we follow f with the identification map from D to $\mathbb{R}P^2$ given in Problem 1 above. The result is an identification map from D to $\mathbb{R}P^2$ that (since it identifies antipodal points on the boundary) identifies the same two points on the two boundary semicircles that came from two points in R that would be identified if we formed the Möbius first. Thus the identification space is homeomorphic to the image $\mathbb{R}P^2$. \square

- (3) Let $f : X \rightarrow Y$ be an identification map, let A be a subspace of X and give $f(A)$ the induced topology from Y . We show that the restriction $f|_A : A \rightarrow f(A)$ need not be an identification map:

Proof. Let $X = [0, 1]$ and $Y = S^1 \subset \mathbb{C}$ and let $f : X \rightarrow Y$ be given by

$$x \mapsto e^{2\pi i x}$$

Then, by corollary (4.4), f is an identification map because X is compact and Y is Hausdorff.

Let $A = [0, 1)$. Then, $f|_A(A) = Y$, but $f|_A$ does not identify any points. Thus, if $f|_A$ were an identification map it would (by theorem (4.2)) induce a homeomorphism from A to Y . But we know f is not a homeomorphism¹. \square

- (4) With the terminology from the previous problem, we show that if A is open in X , and if f is an open map then $f|_A : A \rightarrow f(A)$ is an identification map:

Proof. This is theorem (4.3), page sixty-seven. \square

- (5) Let X denote the union of all circles of the form

$$\left(x - \frac{1}{n}\right)^2 + y^2 = \left(\frac{1}{n}\right)^2, \quad n \in \mathbb{N}$$

with the induced topology. Let Y denote the identification space obtained from the real line by identifying all the integers to a single point. We show that $X \not\cong Y$:

¹See section 1.4, example 3, page 14

Proof. We claim that X is compact, but Y is not.

Indeed; To show that X is compact, first let $\{\mathcal{O}_\alpha\}_\alpha$ be an open cover of X . Then, there exists some α_0 , such that

$$\{(0,0)\} \subset \mathcal{O}_{\alpha_0}$$

Thus, as \mathcal{O}_{α_0} is an open set containing $\{(0,0)\}$, there exists some $n_0 \in \mathbb{N}$ for which

$$\left(x - \frac{1}{n}\right)^2 + y^2 \subset B(\mathcal{O}_{\alpha_0}), \quad n \geq n_0$$

where $B(\mathcal{O}_{\alpha_0})$ denotes the solid ball with boundary \mathcal{O}_{α_0} . Consequently, only finitely many circles are outside $B(\mathcal{O}_{\alpha_0})$. Thus, compactness follows.

To show that Y is not compact, consider the open cover of \mathbb{R} given by

$$\{(n - 1/2, n + 1 + 1/2)\}_{n \in \mathbb{Z}}$$

There is clearly no finite subcover of \mathbb{R} . This completes the proof. \square

- (6) We give an example of an identification map which is neither open nor closed:

Proof. Let $\pi_1 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be projection onto the first coordinate and consider

$$A = \{(x, y) : x \geq 0 \vee y = 0\} \subset \mathbb{R} \times \mathbb{R}$$

We show that $\pi_1|_A$ is a quotient map which is neither open nor closed:

Indeed; We know that π_1 is a quotient map, as it sends open sets to open sets. In addition, $\pi_1|_A$ is obviously saturated and so a quotient map. To see that it is neither open nor closed, consider

$$C = \{(x, 1/x) : x \in \mathbb{R} - \{0\}\}$$

which we have previously shown is closed. Then, clearly $C \cap A$ is clearly closed in A , but

$$\pi_1|_A[C \cap A] = (0, \infty)$$

which is open. And,

$$V = A \cap (\mathbb{R} \times (-1, \infty))$$

is open in A , but

$$\pi_1|_A[V] = [0, \infty)$$

is closed in \mathbb{R} . This proves the result. \square

- (7) This is left to the reader as an exercise.

- (8) Suppose that X is a compact T_2 space. We show that

- the cone on X is homeomorphism to the one-point compactification of $X \times [0, 1)$:

Proof. Note that, as X is compact, $X \times [0, 1]$ is compact. Further,

$$X \times [0, 1] = X \times [0, 1) \cup X \times \{1\}$$

As the one point compactification of $X \times [0, 1)$ is $X \times [0, 1]$ and $X \times \{1\}$ is compact in $X \times [0, 1]$, the result follows. \square

- if A is closed in X , then X/A is homeomorphic to the one-point compactification of $X - A$:

Proof. Let $Y = X - A \cup \{\infty\}$ be the one point compactification of $X - A$. Further, let P be the image of A in X/A , under the identification map

$$i : X \rightarrow X/A$$

We show that the function $h : X/A \rightarrow Y$ given by

$$h(x) = \begin{cases} x & x \in X - A \\ \infty & x \in P \end{cases}$$

is a homeomorphism. We apply theorem (3.7) and claim that $h \circ i$ is continuous.

Indeed; If \mathcal{O} is open in Y and $\infty \notin \mathcal{O}$, then $\mathcal{O} \subset X - A$, and so,

$$(h \circ i)^{-1}(\mathcal{O}) = \mathcal{O} \subset X$$

is open. Now, if $\infty \in \mathcal{O}$, then \mathcal{O}^c is compact in $X - A$ by definition and is closed. Thus, $(h \circ i)^{-1}(\mathcal{O}^c)$ is closed in X . Consequently,

$$(h \circ i)^{-1}(\mathcal{O})^c = (h \circ i)^{-1}(\mathcal{O})$$

is open in X . So, h is continuous.

As $X - A \cup \{\infty\}$ is T_2 and $X - A$ compact (it is the continuous image under i of X), h is a homeomorphism by theorem (3.7), page forty-eight. This completes the proof. \square

- (9) Let $f : X \rightarrow X'$ be a continuous function and suppose that we have partitions $\mathcal{P}, \mathcal{P}'$ of X and X' respectively, such that if two points of X lie in the same member of \mathcal{P} , their images under f lie in the same member of \mathcal{P}' . We show that if Y, Y' are the identification spaces given by these partitions, that f induces a map $f' : Y \rightarrow Y'$, and that if f is an identification map then so is f' :

Proof. Let $\pi : X \rightarrow Y$, $\pi' : X' \rightarrow Y'$ be the given identification maps. We show that f induces a map $f' : Y \rightarrow Y'$:

Define $f' : Y \rightarrow Y'$ by $f'(\pi(\mathcal{P})) = \pi'(f(\mathcal{P}))$, for all $\mathcal{P} \in \mathcal{P}$. To show that f' is a map, we first show that it is well-defined. For all $\mathcal{P} \in \mathcal{P}$, there exists $\mathcal{P}' \in \mathcal{P}'$, such that $f(\mathcal{P}) \subset \mathcal{P}'$, by assumption. The fact that f' is well-defined follows and everywhere defined is clear.

To show that f' is a map, let \mathcal{O} be open in Y' . Then, we have

$$f'^{-1}(\mathcal{O}) = (\pi' \circ f)^{-1}(\mathcal{O}) = f^{-1}(\pi'^{-1}(\mathcal{O})),$$

which is open as f is continuous. Thus, f' is a map.

To conclude the proof, we show that if f is an identification map, then so is f' : Indeed; Let \mathcal{O} be open in Y . We claim that $f'(\mathcal{O})$ is open in Y' . By definition, if \mathcal{O} is open in Y , then as π is an identification map, $\pi^{-1}(\mathcal{O})$ is open in X . Further,

$$(\pi' \circ f \circ \pi^{-1})(\mathcal{O}) = \pi'(f(\pi^{-1}(\mathcal{O}))) = f'(\pi(\pi^{-1}(\mathcal{O}))) = f'(\mathcal{O})$$

and,

$$f^{-1}(\pi'^{-1}(f'(\mathcal{O}))) = f^{-1}(\pi'^{-1}(\pi'(f(\pi^{-1}(\mathcal{O})))))) = \pi^{-1}(\mathcal{O})$$

Thus, by definition of identification map, $f'(\mathcal{O})$ is open in Y' . This concludes the proof. \square

- (10) Let S^2 be the unit sphere in \mathbb{E}^3 and define $f : S^2 \rightarrow \mathbb{E}^4$ by $f(x, y, z) = (x^2 - y^2, xy, xz, yz)$. We show that f induces an embedding of the projective plane in \mathbb{E}^4 :

Proof. Let $X = \text{im}(f) \subseteq \mathbb{E}^4$. Since \mathbb{E}^4 is Hausdorff and a subspace of a Hausdorff space is Hausdorff, X is Hausdorff. And S^2 is compact. Thus by Corollary 4.4 $f : S^2 \rightarrow X$ is an identification map. By the example “Projective spaces (a)” on page 71, \mathbb{P}^2 is S^2 with antipodal points identified. Thus by Theorem 4.2 (a) we will be done if we show f identifies antipodal points (and no others). It clearly does identify antipodal points, $f(x, y, z) = f(-x, -y, -z)$. So suppose $f(x_1, y_1, z_1) = f(x_2, y_2, z_2)$. We must show $(x_1, y_1, z_1) = (x_2, y_2, z_2)$ or $(x_1, y_1, z_1) = (-x_2, -y_2, -z_2)$. We have

$$x_1^2 - y_1^2 = x_2^2 - y_2^2 \tag{4.1}$$

$$x_1 y_1 = x_2 y_2 \tag{4.2}$$

$$x_1 z_1 = x_2 z_2 \tag{4.3}$$

$$y_1 z_1 = y_2 z_2 \tag{4.4}$$

From (3) and (4) it follows that

$$(x_1^2 - y_1^2)z_1^2 = (x_2^2 - y_2^2)z_2^2 \tag{4.5}$$

case 1: $x_1^2 - y_1^2 \neq 0$.

Combining (1) and (5) we get $z_1^2 = z_2^2$. Thus $z_1 = \pm z_2$. If $z_1 = z_2$, then

case a: $z_1 \neq 0$. Then (3) and (4) imply $x_1 = x_2$ and $y_1 = y_2$. If $z_1 = -z_2$ then (3) and (4) imply $x_1 = -x_2$ and $y_1 = -y_2$. Thus either $(x_1, y_1, z_1) = (x_2, y_2, z_2)$ or $(x_1, y_1, z_1) = (-x_2, -y_2, -z_2)$.

case b: $z_1 = 0$. Then $x_1^2 + y_1^2 = 1$. So one of x_1 or y_1 must be different from zero. Assume wlog that $x_1 \neq 0$. Now $x_1^2 + y_1^2 = 1 = x_2^2 + y_2^2$ together with (1) implies $2x_1^2 = 2x_2^2$. Thus $x_1 = \pm x_2$. If $x_1 = x_2$ then (2) implies $y_1 = y_2$, thus $(x_1, y_1, z_1) = (x_2, y_2, z_2)$. If $x_1 = -x_2$ then (2) implies $y_1 = -y_2$, thus $(x_1, y_1, z_1) = (-x_2, -y_2, -z_2)$.

case 2: $x_1^2 - y_1^2 = 0$.

Suppose $x_1 = 0$. Then $y_1 = 0$, and combining (1) and (2) it follows that $x_2 = 0$ and $y_2 = 0$. Now if $x_1 = y_1 = 0$ then necessarily $z_1 = \pm 1$. Likewise $z_2 = \pm 1$. Thus in this case either $(x_1, y_1, z_1) = (x_2, y_2, z_2)$ or $(x_1, y_1, z_1) = (-x_2, -y_2, -z_2)$. By symmetry, the same thing happens if $y_1 = 0$, $x_2 = 0$ or $y_2 = 0$.

Therefore we have reduced to the case that none of x_1, x_2, y_1, y_2 are zero. By assumption $x_1 = \pm y_1$ and from (1) it follows that $x_2 = \pm y_2$. It then follows from (2) that $x_1 = \pm x_2$.

case a: $x_1 = x_2$. Then (2) implies $y_1 = y_2$ and (4) implies $z_1 = z_2$. Thus $(x_1, y_1, z_1) = (x_2, y_2, z_2)$.

case b: $x_1 = -x_2$. Then (2) implies $y_1 = -y_2$ and (4) implies $z_1 = -z_2$. Thus $(x_1, y_1, z_1) = (-x_2, -y_2, -z_2)$. \square

- (11) We show that the function $f : [0, 2\pi] \times [0, \pi] \rightarrow \mathbb{E}^5$ defined by $f(x, y) = (\cos x, \cos 2y, \sin 2y, \sin x \cos y, \sin x \sin y)$ induces an embedding of the Klein bottle in \mathbb{E}^5 :

Proof. Let $Y = [0, 2\pi] \times [0, \pi]$. Let $X = \text{im}(f) \subseteq \mathbb{E}^5$. Since \mathbb{E}^5 is Hausdorff and a subspace of a Hausdorff space is Hausdorff, X is Hausdorff. And $[0, 2\pi] \times [0, \pi]$ is compact. Thus by Corollary 4.4 $f : Y \rightarrow X$ is an identification map. We know the Klein bottle can be obtained from Y by identifying two opposite edges in the same orientation and the other two in the opposite orientation (see Figure 1.12, page 10).

First note that f identifies all four corners of Y together. So in what follows we will examine what happens to all of the other points.

Suppose $f(x_1, y_1) = f(x_2, y_2)$. First suppose $y_1 \neq y_2$ and assume $y_1, y_2 \in (0, \pi)$. Inspection of the graph of $\sin(x)$ tells us that if $\sin 2y_1 = \sin 2y_2$ then one of two cases hold: (i) $0 < 2y_1, 2y_2 < \pi$ and $2y_1 = \pi - 2y_2$ or (ii) $\pi < 2y_1, 2y_2 < 2\pi$ and $2y_1 = 3\pi - 2y_2$. Now $\cos(2y_1) = \cos(2y_2)$ implies $2y_1 = 2\pi - 2y_2$, or equivalently $y_1 = \pi - y_2$. But this is incompatible with both (i) and (ii). Thus if we continue to assume $y_1 \neq y_2$, it must be that at least one of y_1 or y_2 equals zero or π . Since $y_1 = \pi - y_2$, if $y_1 = 0$ then $y_2 = \pi$ and conversely. Thus we have shown that if $f(x_1, y_1) = f(x_2, y_2)$ and $y_1 \neq y_2$ then one of (x_1, y_1) or (x_2, y_2) must be on the top side and the other on the bottom side. In other words $\{y_1, y_2\}$ equals $\{0, \pi\}$. We need to examine what happens to the x -coordinate of such points.

So suppose $f(a, 0) = f(b, \pi)$. And suppose $0 < a, b < 2\pi$. We have $\cos a = \cos b$ and $\sin a = -\sin b$ (from the first and fourth coordinates of f). From $\cos a = \cos b$ we know that either $a = b$ or $a = 2\pi - b$. From $\sin a = -\sin b$ we know that the only one of these that is possible is $a = 2\pi - b$ (keep in mind we are assuming here that a and b are strictly between 0 and 2π).

Thus the points where $f(x_1, y_1) = f(x_2, y_2)$ with $y_1 \neq y_2$ and $x_1, x_2 \in (0, 2\pi)$ are exactly the points of the form $((x_1, y_1), (x_2, y_2)) = ((x, 0), (2\pi - x, \pi))$ for some $x \in (0, 2\pi)$. The only other points on the top or bottom sides are the corner points which we have already handled.

Suppose now that $y_1 = y_2$, and assume $x_1 \neq x_2$. Assume $y_1, y_2 \in (0, \pi)$. We know from the first coordinate of $f(x_1, y_1) = f(x_2, y_2)$ that $\cos x_1 = \cos x_2$. Since $x_1 \neq x_2$, this holds only if $x_1 = 2\pi - x_2$. Now from the fifth coordinate of $f(x_1, y_1) = f(x_2, y_2)$ we also know $\sin x_1 = \sin x_2$ (substitute $y_2 = y_1$ and then we can cancel $\sin y_1$ from both sides since we are assuming $y_1 \in (0, \pi)$), which together with $x_1 = 2\pi - x_2$ implies $-\sin x_2 = \sin x_2$, which implies $\sin x_2 = 0$. Thus $x_2 = 0$ or $x_2 = \pi$. We do not need to further evaluate the y -coordinates of such points since in this case we have assumed $y_1 = y_2$. Thus f identifies points on the opposite vertical sides at the same vertical height to each other. The only other points on the left or right sides are the corner points which we have already handled.

In summary we have shown that any points that f identifies to each other must lie on the boundary of the square; and that f identifies the four corners of the square to one point; and f identifies points on the top and bottom sides in pairs, where the point with x -coordinate x on the top is identified with the point with x -coordinate $2\pi - x$ on the bottom; and finally f identifies points on the left and right sides in pairs, where points with the same y -coordinate are identified to each other.

It follows that the identification space is the Klein bottle. \square

- (12) With the notation of problem 11, show that if $(2 + \cos x) \cos 2y = (2 + \cos x') \cos 2y'$ and $(2 + \cos x) \sin 2y = (2 + \cos x') \sin 2y'$, then $\cos x = \cos x'$, $\cos 2y = \cos 2y'$, and $\sin 2y = \sin 2y'$. Deduce that the function $g : [0, 2\pi] \times$

$[0, \pi] \rightarrow \mathbb{E}^4$ given by $g(x, y) = ((2 + \cos x) \cos 2y, (2 + \cos x) \sin 2y, \sin x \cos y, \sin x \sin y)$ induces an embedding of the Klein bottle in \mathbb{E}^4 :

Proof. Suppose that

$$(2 + \cos x) \cos 2y = (2 + \cos x') \cos 2y' \quad (4.1)$$

$$(2 + \cos x) \sin 2y = (2 + \cos x') \sin 2y' \quad (4.2)$$

We want to show

$$\cos x = \cos x' \quad (4.3)$$

$$\cos 2y = \cos 2y' \quad (4.4)$$

$$\sin 2y = \sin 2y' \quad (4.5)$$

case 1: $\{y, y'\} \cap \{\pi/4, 3\pi/4\} = \emptyset$. Then $\cos 2y \neq 0$ and $\cos 2y' \neq 0$. Thus we can divide (2) by (1) to get $\tan 2y = \tan 2y'$. Since $y, y' \in [0, \pi]$, the only way this is possible is if $y = y'$ or (wlog) $2y = 2y' + \pi$. If $y = y' + \pi/2$ then $\cos(2y) = -\cos(2y')$. But then (1) cannot hold (remember $\{y, y'\} \cap \{\pi/4, 3\pi/4\} = \emptyset$). Thus $y = y'$, from which (4) and (5) obviously hold and it follows immediately from (1) or (2) that (3) holds.

case 2: $\{y, y'\} \cap \{\pi/4, 3\pi/4\} \neq \emptyset$. If $y = \pi/4$, then (1) implies $y' = \pi/4$ or $y' = 3\pi/4$. But if $y' = 3\pi/4$ then (2) does not hold. So it must be $y = y' = \pi/4$. Then (4) and (5) hold and (2) implies (3). If on the other hand $y = 3\pi/4$, then as before (1) implies $y' = \pi/4$ or $y' = 3\pi/4$. But if $y' = \pi/4$ then (2) does not hold. So it must be that $y = y' = 3\pi/4$. Then (4) and (5) hold and (2) implies (3).

Now let f be the function defined in Problem 11. Then g is an identification map for the same reasons f is. And it follows from what we just proved above that $g(x_1, y_1) = g(x_2, y_2) \Leftrightarrow f(x_1, y_1) = f(x_2, y_2)$. Thus the identification space for g is the same as the identification space for f . Thus the image of g is homeomorphic to the Klein bottle. \square

4.3 Topological Groups

Throughout this section, and the rest of the book, the author assumes that such groups are T_2 . Unless otherwise stated, this will be the case throughout the solutions.

(1) This is left to the reader as an exercise.

(2) Suppose that (G, m, τ) is a topological group. We show that

- if H is a subgroup of G , then its closure \overline{H} is also a subgroup:

Proof. Let $a, b \in \overline{H}$. We claim that $a + b \in \overline{H}$. Now, as $H \subset \overline{H}$, $m^{-1}(H) \subset m^{-1}(\overline{H})$. It follows that as

$$m^{-1}(H) = \{(a, b) : a, b \in H\} = H \times H$$

we have $H \times H \subset \overline{H} \times \overline{H} \subset m^{-1}(\overline{H})$. But, as \overline{H} is closed, and m is continuous, we have

$$m(\overline{H} \times \overline{H}) \subset \overline{H}$$

Similarly we can show this for the inverse function. \square

- if H is normal, then so is \overline{H} :

Proof. We use the characterization for normality give by left and right co-set equality. As previously show, $\overline{H} \leq G$. As $gH = Hg$, we have that

- $gH = Hg \subset g\overline{H}$, and so $\overline{H}g \subset g\overline{H}$.
- Similarly, $g\overline{H} \subset \overline{H}g$.

\square

- (3) Let G be a compact Hausdorff space which has the structure of a group. We show that G is a topological group if the multiplication $m : G \times G \rightarrow G$ is continuous:

Proof. Let π be the canonical homeomorphism between G and $\{e\} \times G$, and p_2 projection onto the second coordinate. From the fact that compositions of homeomorphisms are homeomorphisms, and the comments on page seventy-five, it follows that $L_{g^{-2}}$ is a homeomorphism and

$$p_2 \circ (\pi \circ L_{g^{-2}}) \equiv i : G \rightarrow G^{-1}$$

is continuous. \square

- (4) We prove that $O(n)$ is homeomorphic to $SO(n)$ and that they are isomorphic as topological groups:

Proof. Consider the determinate function restricted to $O(n)$. It follows that the function $g : O(n) \rightarrow SO(n) \times \mathbb{Z}_2$ given by

$$g(X) = (\det(X)X, \det X)$$

is 1-1 and onto. We show that g is continuous. Let \mathcal{O} be open in $SO(n) \times \mathbb{Z}_2$. Then, $\mathcal{O} = U \times \{1\}$, or $\mathcal{O} = U \times \{0\}$ for some open U in $SO(n)$. But, as $SO(n)$ is a subgroup of $O(n)$, U is open in $O(n)$. Thus, as $U, -U$ are open by extra lemma, it follows that

$$g^{-1}(U \times \{1\}) = -U, \quad g^{-1}(U \times \{0\}) = U$$

are open. Thus, g is continuous. As $O(n)$ is compact, and $SO(n) \times \mathbb{Z}_2$ is Hausdorff, it follows from theorem (3.7), page forty-eight, that g is a homeomorphism.

To conclude the proof, we note that $\det XY = \det X \det Y$. \square

- (5) Let A, B be compact subsets of a topological group. We show that the product set AB is compact:

Proof. As $m : G \times G \rightarrow G$ is continuous, and $m(A \times B) = AB$, compactness follows from theorem (3.4), page forty-seven. \square

- (6) We show that if U is a neighbourhood of e in a topological group, there is a neighbourhood V of e for which $VV^{-1} \subset U$:

Proof. Consider $m^{-1}(U)$. Then, $m^{-1}(U)$ is an open neighbourhood of (e, e) in $G \times G$. It follows from the definition of product topology that $m^{-1}(U) = V \times V$ for some open set V in G . Via $L_{v^{-2}}$, we have

$$V \cong_{L_{v^{-2}}} V^{-1}$$

and so,

$$V \times V \cong V \times V^{-1}$$

And, as m is continuous, we have

$$m(V \times V^{-1}) = VV^{-1} \subset U = m(m^{-1}(U))$$

\square

- (7) Let H be a discrete subgroup of a topological group (G, m, τ) . We find a neighbourhood N of e in G such that the translates $hN = L_h(N)$, $h \in H$ are all disjoint:

Proof. As H is discrete, for some open cover $\{\mathcal{O}_\alpha\}_\alpha$, each \mathcal{O}_α contains one element of H . In particular, there exists some \mathcal{O}_{α_0} , such that

$$\{e\} \subset \mathcal{O}_{\alpha_0}$$

Thus, $\mathcal{O}_{\alpha_0} \cap H \neq \{e\}$. By an exercise above, there exists some open N in G , for which $NN^{-1} \subset \mathcal{O}_{\alpha_0}$. Now, suppose to the contrary that for some $h, g \in H$, $h \neq g$, $hN \cap gN = \emptyset$.

Then, there exists some $n_1, n_2 \in N$, for which $hn_1 = gn_2$, implying that

$$g^{-1}h = n_2n_1^{-1} \in NN^{-1} \subset \mathcal{O}_{\alpha_0}$$

But then, $g^{-1}h = n_2n_1^{-1} \in \mathcal{O}_{\alpha_0}$. This implies that

$$g^{-1}h \in \mathcal{O}_{\alpha_0} \cap H = \{e\};$$

a contradiction. \square

- (8) We show that if C is a compact subset of a topological group (G, m, τ) , and if H is a discrete subgroup of G that $H \cap C$ is finite:

Proof. By extra lemma, $\overline{H} = H$, and as C is compact in a T_2 space, C is closed. This implies that $H \cap C \subset C$ is closed, and compact by theorem (3.5), page forty-seven.

Suppose, to the contrary, that $H \cap C$ was not finite. As

$$\bigcup_{x \in C} \{x\}$$

is an open cover of $H \cap C$, no finite set of $\{x : x \in C\}$ could cover $H \cap C$, contradicting the compactness of $H \cap C$. \square

- (9) We prove that every nontrivial discrete subgroup of \mathbb{R} is infinite cyclic:

Proof. Let G be a non-trivial discrete subgroup of \mathbb{R} . Let $g \in G$, $g \neq 0$. Then, as (G, m, τ) is discrete, $\{0, g\}$ is a neighbourhood of 0, and so, g generates G (theorem (4.11), page seventy-five). The proof that $G = \langle g \rangle$ is clear from the division algorithm. To conclude, we show that $|G| = \infty$:

Suppose, to the contrary, that $|G| = p$, for some $p \in \mathbb{N}$. Then, as $\overline{G} = G$, $\inf G \in G$. But then, by the Archimedean Principle, there exists some $n \in \mathbb{N}$, for which $ng > \inf G$, $hg \in \langle g \rangle$. Thus, $|G| = \infty$. \square

- (10) We prove that every non-trivial discrete subgroup of the circle is finite and cyclic:

Proof. Let G be a discrete subgroup of S^1 . Define $f : [0, 1] \rightarrow S^1$ by $f(t) = e^{2\pi it}$. Then, G is finite, since it is discrete, and compact. It is easy to see that $f^{-1}(G)$ is a discrete subgroup of \mathbb{R} , so is cyclic. Thus, $G = f(f^{-1}(G))$ is cyclic. \square

- (11) Suppose that $A, B \in O(2)$, such that $\det A = 1$, $\det B = -1$. We show that $B^2 = I$, and $BAB^{-1} = A^{-1}$, and deduce that every discrete subgroup of $O(2)$ is either cyclic or dihedral:

Proof. By exhaustion,

$$B = \begin{bmatrix} \pm 1 & 0 \\ 0 & \mp 1 \end{bmatrix}$$

Thus, either way, $B^2 = I$. The fact that $BAB^{-1} = A^{-1}$ is routine. To complete the proof, we show that every discrete subgroup of $O(2)$ is either cyclic or dihedral:

From the comments on page seventy-seven, and exercise sixteen, we have that

$$O(2) \cong SO(2) \times \mathbb{Z}_2 \cong S^1 \times \mathbb{Z}_2$$

Now, let G be a discrete subgroup of $S^1 \times \mathbb{Z}_2$. Then, for every $X \in O(2)$, $\det X \neq \pm 1$, we have

$$f(X) \cong SO(2) \times \{1\} \cong S^1 \times \{1\} \cong S^1$$

as in the previous exercise. By the exercise a above, we have that $f(X)$ must be finite cyclic. In other-words, $f^{-1}(f(X)) = X$ is finite cyclic.

To finish the proof, suppose that $X \in G$ was such that $\det X = -1$, without loss of generality. Now, let $K = G \cap SO(2)$. Then, K is cyclic and K is the set of $Z \in G$, such that $\det Z = 1$. Let M be the generator of K . Likewise, let N be the generator of $G - K$. Then, as $N^2 = I$, $NMN^{-1} = N^{-1}$, $\langle M, N \rangle$ is dihedral.

We claim that $G = \langle M, N \rangle$; Let $L \in G - \langle M, N \rangle$. Then, $L \notin K$, and so $\det L = -1$. Consequently, $\det LMN = 1$, and so, $LMN \in K$. But then, $L \in G$; a contradiction. This completes the proof. \square

(12) We show that

- if T is an automorphism of the topological group \mathbb{R} , that $T(r) = rT(1)$, for any rational r :

Proof. If $n \in \mathbb{N}$, then we must have $T(n+x) = T(n) + T(x)$. Further, as T sends generators to generators, we must have

$$T(n) = n, \quad T(n+x) = n + T(x)$$

It follows from the equality

$$T(1) = \sum_{i=1}^n T(1/n) = n + T(1/n)$$

that

$$T(m/n) = mT(1/n) = \frac{m}{n}T(1)$$

Note the change in notation. Thus, as \mathbb{Q} is dense in \mathbb{R} , it follows from the sequential criterion for continuity, that $xT(1) = T(x)$ for all $x \in \mathbb{R}$. \square

- the automorphism group of \mathbb{R} is isomorphic to $\mathbb{R} \times \mathbb{Z}_2$:

Proof. We first show that $\text{Aut } \mathbb{R} \cong \mathbb{R} \times \mathbb{Z}_2$; The map $T_x : \mathbb{R} \rightarrow \mathbb{R}$, given by $T_x(y) = xy$, $x \neq 0$ is an automorphism of \mathbb{R} , such that $T_x(1) = x$. Thus, consider $f : \mathbb{R} - \{0\} \rightarrow \text{Aut } \mathbb{R}$ given by $f(x) = x = T_x(1)$, for each $T_x \in \text{Aut } \mathbb{R}$, as described above. Then, clearly, f is 1-1 and onto. Further,

$$f(xy) = T_x(1)T_y(1) = xy = f(x)f(y)$$

To conclude, we show that $\mathbb{R} - \{0\} \cong \mathbb{R} \times \mathbb{Z}_2$; Consider the map

$$g : \mathbb{R} \times \mathbb{Z}_2 \rightarrow \mathbb{R}, \quad g(x, y) = \begin{cases} e^x & y = 0 \\ -e^x & y = 1 \end{cases}$$

We claim that g is a homeomorphic isomorphism. As $e^x : \mathbb{R} \rightarrow \mathbb{R}^+$ is a homeomorphism from \mathbb{R} to \mathbb{R}^+ , it follows that g is a homeomorphism. To show that it is an isomorphism, we show the homomorphism property: Without loss of generality, suppose that $(x, y), (x', y') \in \mathbb{R} \times \mathbb{Z}_2$ is such that $y = 0, y' = 1$; then,

$$g(x, y)g(x', y') = e^x(-e^{x'}) = -e^{x+x'} = g(x + x', y + y')$$

This completes the proof. \square

- (13) We show that the automorphism group of the circle group is isomorphic to \mathbb{Z}_2 :

Proof. We first note that the identity map $i : S^1 \rightarrow S^1$ is the trivial automorphism. In addition, we have shown that the conjugate map given by $\bar{i}(x) = \bar{x}$ is an automorphism. We show that $|\text{Aut } S^1| = 2$:

We show that if f is any other automorphism, the $f \equiv i$, or $f \equiv \bar{i}$. Indeed; first note that, as

$$S^1 \cong U(1) = \bigcup_n U_n, \quad U_n = \{c \in \mathbb{C} : c^n = 1\},$$

if $f \in \text{Aut } S^1$, then f must send generators to generators. So, consider the generators of $U(1)$, given by U_2 , and U_3 . Then,

$$U_2 \cong \mathbb{Z}_2, \quad U_3 \cong \mathbb{Z}_3$$

Further, for $f \in \text{Aut } S^1$, we must have $f(-1) = -1$, and thusly,

$$f\left(\frac{-1 + i\sqrt{3}}{2}\right) = \frac{-1 \pm i\sqrt{3}}{2}$$

In which case, $f \cong i$ and $f \cong \bar{i}$, respectively. Thus, by extra lemma, we have that $|\text{Aut } S^1| = 2$ implies $\text{Aut } U(1) \cong \mathbb{Z}_2$. \square

4.4 Orbit Spaces

- (1) We give an action of \mathbb{Z} on $\mathbb{E} \times [0, 1]$ which has the Möbius Strip as an orbit space:

Proof. Define $\pi : \mathbb{Z} \times (\mathbb{E} \times [0, 1]) \rightarrow \mathbb{E}^2$, by

$$\pi(z, (x, y)) = \begin{cases} (x + z, y) & z \in 2\mathbb{Z} \\ (x + z, 1 - y) & z \in 2\mathbb{Z} + 1 \end{cases}$$

We show that π is a topological group action:

- Clear by construction.
- As, $\pi(0, (x, y)) = (x + 0, y) = (x, y)$, the result follows.
- As the components are continuous, π is continuous.

□

- (2) We find an action of \mathbb{Z}_2 on the torus with orbit space the cylinder:

Proof. Consider the cylinder on page eighty. Define $\pi : \mathbb{Z}_2 \times T \rightarrow T$, by

$$\pi(g, (x, y, z)) = \begin{cases} (x, -y, z) & g = 1 \\ (x, y, z) & g = 0 \end{cases}$$

We check that π is a T group action:

- By definition.
- As $0 \equiv e \in \mathbb{Z}_2$, clearly $\pi(0, (x, y, z)) = (x, y, z)$.
- As the components are continuous, π is continuous.

To show that $\pi(\mathbb{Z}_2 \times T) \cong C$, we note that $\{(x, y, z) : y \in \mathbb{R}\}$ is the canonical cylinder. □

- (3) We describe the orbits of the natural action of $SO(n)$ on \mathbb{E}^n as a group of linear transformation and identify the orbit space:

Proof. Let $r \in \mathbb{R}$, $r \geq 0$, let

$$S_r = \{p \in \mathbb{E}^n \mid \|p\| = r\}$$

Then since $SO(n)$ preserves distances, $SO(n)$ must take S_r to itself. Furthermore, the action on S_r is transitive, because it is transitive on $S^{n-1} \subset \mathbb{E}^n$ and $S_r = r \cdot S^{n-1}$. To see the action on S^{n-1} is transitive, for any vector $\mathbf{v} \in S^{n-1}$, it can be put into an orthonormal basis \mathbf{B} . Then there is a change of coordinates matrix M from the standard basis $\mathbf{e}_1, \dots, \mathbf{e}_n$ to \mathbf{B} that takes \mathbf{e}_1 to \mathbf{v} . Since both bases are orthonormal, $M \in O(n)$. Clearly M can be chosen to be in $SO(n)$ such that $M(\mathbf{e}_1) = \mathbf{v}$ (if it's not already in $SO(n)$, just multiply one of the other \mathbf{e}_i 's by -1). Since any element of S^{n-1} can be taken to \mathbf{e}_1 , the action must be transitive. Now if $\mathbf{v} \in \mathbb{E}^n$ is arbitrary ($\mathbf{v} \neq 0$) then just scale \mathbf{v} to be in S^{n-1} , transform within S^{n-1} and scale

back. The two scaling operations assure that the resulting transformation has determinant equal to one, and therefore is in $SO(n)$.

Now let $r, r' \in \mathbb{R}$, $r, r' > 0$, $r \neq r'$. Since things in $SO(n)$ are length preserving, an element of $SO(n)$ cannot take an element of S_r to an element of $S_{r'}$. Thus each S_r is exactly one orbit (true also if $r = 0$ since S_0 consists of one point).

Let $f : \mathbb{E}^n \rightarrow [0, \infty)$ be given by $f(\mathbf{v}) = \|\mathbf{v}\|$. Then f is a continuous function that identifies each orbit of the action to a single point. Let B be an open ball in \mathbb{E}^n . Then clearly $f(B)$ is an interval, open in $[0, \infty)$. Since functions respect unions, it follows that f is an open map. By Corollary 4.4 f is an identification map. Thus the identification space is homeomorphic to the image of f , which is $[0, \infty)$. Thus the orbit space of $SO(n)$ on \mathbb{E}^n is homeomorphic to $[0, \infty)$. \square

(4) Suppose that $\pi : X \rightarrow X/G$ is the natural identification map. We show that

- if \mathcal{O} is open in X , then $\pi^{-1}(\pi(\mathcal{O}))$ is the union of the sets $g(\mathcal{O})$, $g \in G$:

Proof. Let $a \in \mathcal{O}$ and $g \in G$. Since ga is in the same orbit as a , $\pi(ga) = \pi(a)$. Thus $\pi(g(\mathcal{O})) = \pi(\mathcal{O})$, for all $g \in G$,

$$g(\mathcal{O}) \subset \pi^{-1}(\pi(\mathcal{O})) \quad \forall g \in G$$

and,

$$\bigcup_{g \in G} g(\mathcal{O}) \subset \pi^{-1}(\pi(\mathcal{O}))$$

Now suppose $x \in \pi^{-1}(\pi(\mathcal{O}))$; Then $\pi(x) \in \pi(\mathcal{O})$. Thus x is in the same orbit as some element of \mathcal{O} , i.e. $x = ga$ for some $g \in G$ and $a \in \mathcal{O}$. Consequently, $x \in g(\mathcal{O})$ implies

$$\pi^{-1}(\pi(\mathcal{O})) \subset \bigcup_{g \in G} g(\mathcal{O})$$

Since we have containment in both directions, we can conclude that

$$\pi^{-1}(\pi(\mathcal{O})) = \bigcup_{g \in G} g(\mathcal{O})$$

\square

- π is an open map:

Proof. Now suppose \mathcal{O} is open in X . Recall a set $U \subset X/G$ is open in X/G if, and only if, $\pi^{-1}(U)$ is open in X . Now, $\pi^{-1}(\pi(\mathcal{O}))$ is a union of sets of the form $g(\mathcal{O})$ and (since each g induces a homeomorphism of X) $g(\mathcal{O})$ is open in X , for all $g \in G$. Thus $\pi^{-1}(\pi(\mathcal{O}))$ is open. Consequently, $\pi(\mathcal{O})$ is open in X/G . \square

- π is not necessarily a closed map:

Proof. We will show by counter-example that π is not a closed map. Let \mathbb{Z} act on \mathbb{R} by translation: $x \mapsto x + z$. This orbit space is S^1 . For each $n = 0, 1, 2, \dots$, let

$$A_n = [n + \frac{1}{n+3}, n + \frac{1}{n+2}]$$

So,

$$A_0 = [1/3, 1/2]$$

$$A_1 = [1 + \frac{1}{4}, 1 + \frac{1}{3}]$$

$$A_2 = [2 + \frac{1}{5}, 2 + \frac{1}{4}]$$

$$\vdots$$

Let $A = \bigcup_n A_n$. Then A is closed in \mathbb{E} , but $\pi(A) = \pi((0, 1/2])$ which is not a closed subset of S^1 . \square

(5) We show that

- if X is Hausdorff, it is not necessarily the case that X/G be Hausdorff:

Proof. Consider $X = S^1 \times S^1$. We claim that

$$X/\mathbb{R}$$

given in example seven, page eighty-three is not T_2 . Clearly, $S^1 \times S^1 = T$ is T_2 . Suppose to the contrary, that for some $x, y \in X/\mathbb{R}$, there exists disjoint neighbourhoods $\mathcal{O}_x, \mathcal{O}_y$. But then, as the comments on page eighty-three point out, $\text{Orb } x$ is dense in T . But then,

$$\mathcal{O}_x \cap \mathcal{O}_y \neq \emptyset$$

\square

- if (X, m, τ) is a topological group and G is a closed subgroup acting on X by left translation, that X/G is Hausdorff:

Proof. Let

$$C = \{(x, y) \in X \times X \mid x^{-1}y \in G\}$$

and $h : X \times X \rightarrow X$ be the map $h(x, y) = x^{-1}y$. Then $h^{-1}(G) = C$. Since G is closed and h is continuous, it follows that C is closed. Let $f : X \rightarrow X/G$ be the identification map. Let $g : X \times X \rightarrow X/G \times X/G$ be the map $g(x, y) = (f(x), f(y))$. By Problem 29 f is an open

map. It follows that g is an open map. Thus by Theorem 4.3 g is an identification map. Let Δ be the diagonal in $X/G \times X/G$. Then $g^{-1}(\Delta) = C$. Since g is an identification map and C is closed in $X \times X$, it follows that Δ is closed in $X/G \times X/G$. By Chapter 3, Problem 25 (page 55) it follows that X/G is Hausdorff. \square

- (6) We show that the stabilizer of any point is closed subgroup of G when X is Hausdorff, and that points in the same orbit have conjugate stabilizers for any X :

Proof. If g and g' are in the stabilizer of x , then $gg'x = gx = x$ so gg' is in the stabilizer of x . And $g^{-1}gx = 1 \cdot x = x$, but also $g^{-1}gx = g^{-1}x$. Thus $g^{-1}x = x$ so g^{-1} is in the stabilizer of x . It follows that the stabilizer of x is a subgroup of G . Now, let $f : G \rightarrow X$ be given by $f(g) = gx$. Then f is continuous. Since X is Hausdorff, by Theorem 3.6 points are closed (finite sets are always compact). Thus $f^{-1}(x)$ is closed in G . But $f^{-1}(x)$ is exactly the stabilizer of x . Thus the stabilizer of x is closed in G .

It remains to show points in the same orbit have conjugate stabilizers. Let x, y be in the same orbit, so $x = gy$ for some $g \in G$ and, let $a \in \text{stab } x$. Then

$$\begin{aligned} g^{-1}agy &= g^{-1}ax = g^{-1}x = y \\ \implies g^{-1}ag &\in \text{stab } y \\ \implies g^{-1}(\text{stab } x)g &\subset \text{stab } y \end{aligned}$$

Now let $a \in \text{stab } y$. Then,

$$gag^{-1}x = gay = gy = x$$

So,

$$gag^{-1} \in \text{stab } x \implies \text{stab } y \subset g^{-1}(\text{stab } x)g$$

Since we have set containment in both directions it follows that

$$g^{-1}(\text{stab } x)g = \text{stab } y$$

\square

- (7) Suppose that G is compact, X is Hausdorff and that G acts transitively on X . We show that X is homeomorphic to the orbit space $G/(\text{stabilizer of } x)$ for any $x \in X$:

Proof. Let $x \in X$, $f : G \rightarrow X$ be given by $f(g) = gx$. Since G acts transitively, f is onto. Since G is compact and X is Hausdorff, f is an identification map by corollary (4.4). So G^* , the identification space associated to f , is homeomorphic to X .

Now, suppose $f(g_1) = f(g_2)$. Then $g_2^{-1}g_1 \in \text{stab } x$. Consequently, g_1 is in the same coset as g_2 with respect to the subgroup (stabilizer of x). Thus G^* is exactly $G/(\text{stabilizer of } x)$. \square

- (8) We prove that the resulting space is homeomorphic to the Lens space $L(p, q)$:

Proof. \square

- (9) We show that $L(2, 1)$ is homeomorphic to $\mathbb{R}P^3$ and that if p divides $q - q'$, that $L(p, q)$ is homeomorphic to $L(p, q')$:

Proof. Using the definition of $L(2, 1)$ on page eighty-two, $L(2, 1)$ is an identification space of S^3 , the orbit space under the action of \mathbb{Z}_2 . The action of the generator of \mathbb{Z}_2 is the homeomorphism $(z_0, z_1) \mapsto (e^{\pi i} z_0, e^{\pi i} z_1) = (-z_0, -z_1)$. And $(-z_0, -z_1)$ is the antipodal point to (z_0, z_1) . Thus, by example 2 on page eighty, the identification space is exactly \mathbb{P}^3 .

Now suppose $p|(q - q')$. Then $q = q' + np$ for some $n \in \mathbb{N}$. Thus,

$$e^{2\pi q i/p} = e^{2\pi(q' + np)i/p} = e^{2\pi q' i/p + 2\pi n p i/p} = e^{2\pi q' i/p} e^{2\pi n i} = e^{2\pi q' i/p}$$

And, so, the action of \mathbb{Z}_p is identical in both cases; the resulting orbit spaces are identical (and therefore homeomorphic). \square

Chapter 5

The Fundamental Group

5.1 Homotopic Maps

Throughout this section i will typically denote the identity map, unless otherwise stated.

- (1) Let C denote the unit circle in the plane. Suppose that $f : C \rightarrow C$ is a map which is not homotopic to the identity. We show that $f(x) = -x$ for some $x \in C$:

Proof. Suppose, to the contrary, that for any $x \in C$, $f(x) \neq -x$. Then, we claim that $f \simeq i$. Indeed; as the example on page eighty-nine points out,

$$F(x, t) = \frac{(1-t)f(x) + tg(x)}{\|(1-t)f(x) + tg(x)\|}$$

is such that

$$f \simeq_F i \equiv g;$$

a contradiction. □

- (2) With C as above, we show that the map which takes each point of C to its antipodal is homotopic to the identity:

Proof. Consider the matrix

$$A_\theta = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

Then, $A_\pi \equiv f(x)$. Thus, define $F : C \times I \rightarrow C$ by

$$F(x, t) = \begin{bmatrix} \cos(\pi(1-t)) & -\sin(\pi(1-t)) \\ \sin(\pi(1-t)) & \cos(\pi(1-t)) \end{bmatrix} x$$

We show that F is homotopy from f to i :

Clearly,

$$\begin{aligned} F(x, 0) &= A_\pi x \equiv f(x) \\ F(x, 1) &= A_0 x = Ix \equiv i \end{aligned}$$

Further, as F is the sum of continuous functions, it is continuous. Thus,

$$f \simeq_F i$$

□

- (3) Let D be the disc bounded by C , and parametrize D using polar coordinates. Let $h : D \rightarrow D$ be the homeomorphism defined by $h(0) = 0$, $h(r, \theta) = (r, \theta + 2\pi r)$. We find a homotopy F , from h to the identity, such that the functions

$$F|_{D \times \{t\}} : D \times \{t\} \rightarrow D, \quad t \in [0, 1]$$

are all homeomorphisms:

Proof. Let $i(r, \theta) = (r, \theta)$ the identity map. Define $F : D \times I \rightarrow D$ by $F((r, \theta), t) = (r, \theta + 2\pi r(1 - t))$. F is given by polynomials in r, θ and t , so F is continuous. Further,

$$\begin{aligned} F((r, \theta), 0) &= h(r, \theta) \\ F((r, \theta), 1) &= (r, \theta) = i(r, \theta) \end{aligned}$$

Thus,

$$h \simeq_F i$$

Since $F|_{D \times \{t\}} \rightarrow D$ is a one-to-one continuous map from a compact space to a Hausdorff space, theorem (3.7) implies that $F|_{D \times \{t\}} \rightarrow D$ is a homeomorphism. □

- (4) With the above terminology, we show that h is homotopic to the identity relative to C :

Proof. As we are consider h relative to C , $r = 1$ for all $x \in C$. Thus,

$$h = h' = (1, \theta + 2\pi)$$

Define $F : \partial D \times I \rightarrow \partial D$ by

$$F(x, t) = (1, \theta + 2\pi(1 - t))$$

Then,

$$\begin{aligned} F(x, 0) &= (1, \theta + 2\pi) = h'(x) \\ F(x, 1) &= (1, \theta) = i(x), \end{aligned}$$

where i is the identity on C . Likewise, F is continuous as it is polynomial in θ , t . Consequently,

$$h' \equiv h|_{\partial D} \simeq_F i$$

□

- (5) Let $f : X \rightarrow S^n$ be a map that is not onto. We show that f is null-homotopic:

Proof. Let p be a point in S^n such that its antipodal point $-p$ is not in the image of f . Now let $g : X \rightarrow S^n$ be the constant function $g(x) = p$. Then $g(x)$ and $f(x)$ never give a pair of antipodal points for any $x \in X$. By example two on page eighty-nine, f and g are homotopic. □

- (6) Let CY denote the cone on Y . We show that any two maps $f, g : X \rightarrow CY$ are homotopic:

Proof. By extra lemma ¹ f, g are null-homotopic. □

- (7) We show that a map from X to Y is null homotopic if, and only, if it extends to a map from the cone on X to Y :

Proof. (\implies): Suppose that $f : X \rightarrow Y$ is null-homotopic, via F .

Thus,

$$F : CX \rightarrow Y, \quad \text{as } CX = X/X \times \{1\}$$

Thus, we need to show that F is an extension of f .

Indeed, $X \times \{0\} \subset X \times I$, and

$$\{f(x) : x \in X\} \cup \{0\} \subset Y$$

So,

$$F|_{X \times \{0\}}(x, t) = F(x, 0) = f(x), \quad \forall x \in X \times \{0\} \cong X.$$

As f is continuous, it follows that $F|_{X \times \{0\}}$ is.

(\impliedby): Suppose that $f : X \rightarrow Y$ is a map and that $F : CX \rightarrow Y$ is an extension of f . Then, consider the map $H(x, t) = F(x, t)$. As

$$F(x, 1) \equiv X \times \{1\},$$

we have

$$F(x, 1) = H(x, 1) = c, \quad \text{for some } c \in Y.$$

In addition,

$$H(x, 0) \equiv F|_{X \times \{0\}} \equiv f$$

and we have

$$f \simeq_H i$$

□

¹reference this

- (8) Let A denote the annulus $\{(r, \theta) : r \in [1, 2], \theta \in [0, 2\pi]\}$, and let h be a homeomorphism of A defined by $h(r, \theta) = (r, \theta + 2\pi(r - 1))$. We show that h is homotopic to the identity map:

Proof. Consider the function $F : A \times I \rightarrow A$ given by

$$F((r, \theta), t) = (r, \theta + 2\pi t(r - 1))$$

Then,

$$\begin{aligned} F((r, \theta), 0) &= (r, \theta) = i(r, \theta) \\ F((r, \theta), 1) &= (r, \theta + 2\pi(r - 1)) \end{aligned}$$

As F is polynomial in θ, t, r , it follows that F is continuous. Thus,

$$h \simeq_F i$$

□

5.2 Construction of the Fundamental Group

- (1) Let α, β, γ be loops in a space X , all based at p . We write out the formulae for $(\alpha.\beta).\gamma$, and $\alpha.(\beta.\gamma)$ and show that they are homotopic loops relative $\{0, 1\}$:

Proof. By definition, we have

$$((\alpha.\beta).\gamma)(t) = \begin{cases} \alpha(4t) & t \in [0, 1/4] \\ \beta(4t - 1) & t \in [1/4, 1/2] \\ \gamma(2t - 1) & t \in [1/2, 1] \end{cases}$$

and,

$$(\alpha.(\beta.\gamma))(t) = \begin{cases} \alpha(2t) & t \in [0, 1/2] \\ \beta(4t - 2) & t \in [1/2, 3/4] \\ \gamma(4t - 3) & t \in [3/4, 1] \end{cases}$$

Considering the canonical diagram, we have claim that $H : X \times I \rightarrow X$, defined by

$$H(t, s) = \begin{cases} \alpha(4t/s+1) & t \in [0, \frac{s+1}{4}] \\ \beta(4t - 1 - s) & t \in [\frac{s+1}{4}, \frac{s+2}{4}] \\ \gamma(4t-2-s/2-s) & t \in [\frac{s+2}{4}, 1] \end{cases}$$

is a homotopy. Indeed;

$$H(t, 0) = \begin{cases} \alpha(4t) & t \in [0, 1/4] \\ \beta(4t - 1) & t \in [1/4, 1/2] \\ \gamma(2t - 1) & t \in [1/2, 1] \end{cases}$$

and, similarly,

$$(H(t, 1) = \alpha.(\beta.\gamma))(t)$$

By two applications of the Gluing Lemma, H is continuous. So,

$$(\alpha.\beta).\gamma \simeq_H \alpha.(\beta.\gamma) \text{ rel}\{0, 1\}$$

□

- (2) Let γ, σ be two paths in the space X which start at p and end at q . We show that σ_* is the composition of γ_* and the inner automorphism of $\pi_1(X, q)$ induced by the element $\langle \sigma^{-1}\gamma \rangle$:

Proof. The isomorphism defined by σ_* and γ_* are

$$\sigma_*(\langle \alpha \rangle) = \sigma^{-1}.\alpha.\sigma$$

and,

$$\gamma_*(\langle \alpha \rangle) = \gamma^{-1}.\alpha.\gamma$$

where α is a loop based at p . The claim is that

$$\sigma_*(\langle \alpha \rangle) = \gamma_*(\langle \gamma.\sigma^{-1} \rangle \langle \alpha \rangle \langle \sigma.\gamma^{-1} \rangle)$$

Indeed;

$$\begin{aligned} \gamma_*(\langle \gamma.\sigma^{-1} \rangle \langle \alpha \rangle \langle \sigma.\gamma^{-1} \rangle) &= \gamma_*(\langle \gamma.\sigma^{-1} \rangle) \gamma_*(\langle \alpha \rangle) \gamma_*(\langle \sigma.\gamma^{-1} \rangle) \\ &= (\gamma^{-1}.\gamma.\sigma^{-1}.\gamma).(\gamma^{-1}.\sigma.\gamma).(\gamma^{-1}.\sigma.\gamma^{-1}.\gamma) \\ &= \sigma^{-1}.\alpha.\sigma \\ &= \sigma_*(\langle \alpha \rangle) \end{aligned}$$

□

- (3) Let X be a path connected space. We describe when it is true that for any two points $p, q \in X$, all paths from p, q induce the same isomorphism between $\pi_1(X, p)$ and $\pi_1(X, q)$:

Proof. As the above proof shows, if γ, σ are paths from p to q , then

$$\sigma_*(\langle \alpha \rangle) = \sigma^{-1}.\alpha.\sigma$$

Thus, $\sigma_* = \gamma_*$ if, and only, if

$$\gamma_*(\langle \gamma.\sigma^{-1} \rangle \langle \alpha \rangle \langle \sigma.\gamma^{-1} \rangle) = \gamma_*(\langle \alpha \rangle),$$

i.e. $\langle \alpha \rangle.\langle \beta \rangle = \langle \beta \rangle.\langle \alpha \rangle$. Thus, $\pi_1(X)$ must be abelian.

□

- (4) We show that any indiscreet space has trivial fundamental group:

Proof. By a previous problem, if (X, τ) is indiscrete, then X is path connected as $\tau = \{X, \emptyset\}$. The idea here is that there are no open sets (holes) for a path to get caught on. We claim that X is contractible:

Consider $\pi_1(X, x_0)$, for some fixed $x_0 \in X$. Define $i \in \pi_1(X, x_0)$ by

$$i(x) = x_0, \quad \forall x \in X$$

Now, for an arbitrary $f \in \pi_1(X, x_0)$, define $H : X \times I \rightarrow X$ by

$$H(x, t) = \begin{cases} i(x) & t \in [0, 1/2] \\ f(x) & t \in [1/2, 1] \end{cases}$$

Then, H is the desired homotopy as

$$\begin{aligned} H(x, 0) &= i(x) \\ H(x, 1) &= f(x) \end{aligned}$$

and it is continuous as the only non-trivial open sets in τ is X and $H^{-1}(X) = X \times I$, which is open in $X \times I$. Therefore,

$$f \simeq_H i$$

□

- (5) Let (G, m, τ) be a path-connected topological group. Given two loops α, β based at e in G , define a map $F : [0, 1]^2 \rightarrow G$ by

$$F(s, t) = \alpha(s) \cdot \beta(t)$$

We show the effect of this map on the square, and prove that the fundamental group of G is abelian:

Proof. We want to show that for any two loops α, β that

$$\langle \alpha \rangle \langle \beta \rangle = \langle \beta \rangle \langle \alpha \rangle$$

As

$$F(s, 0) = F(s, 1) = \alpha(s), \quad F(0, t) = F(1, t) = \beta(t),$$

we have that

$$\alpha \simeq_F \alpha(s), \quad \beta \simeq_F \beta(t).$$

We first show that $\alpha \cdot \beta \simeq m(\alpha(t), \beta(t))$:

Define $P : [0, 1]^2 \rightarrow G$ by

$$P(s, t) = \begin{cases} \alpha(\frac{2t}{1+s}) & t \in [0, \frac{1+s}{2}] \\ e & t \in (\frac{1+s}{2}, 1] \end{cases}$$

and, $Q : [0, 1]^2 \rightarrow G$ by

$$Q(s, t) = \begin{cases} e & t \in [0, \frac{1-s}{2}] \\ \beta(\frac{2t-1+s}{1+s}) & t \in (\frac{1-s}{2}, 1] \end{cases}$$

Then, let $H(s, t) = m(P(s, t), Q(s, t))$. Note that H is a homotopy between $\alpha.\beta$ and $m(\alpha(t), \beta(t))$. Similarly, we can show that

$$m(\alpha(t), \beta(t)) \simeq_{H'} \beta.\alpha,$$

where $H' = QP$. Therefore,

$$\alpha.\beta \text{ rel}\{0, 1\} \simeq_H m(\alpha(t), \beta(t)) \simeq_{H'} \beta.\alpha \text{ rel}\{0, 1\}$$

I.e. $\pi_1(G)$ is abelian. □

- (6) We show that the space $\mathbb{E}^n - B$, where $B = \{(x, y, z) : y = 0 \wedge 0 \leq z \leq 1\}$ has trivial fundamental group:

Proof. As B is homeomorphic to a point, it follows by extra lemma that $\mathbb{E}^n - B$ has trivial fundamental group. □

5.3 Calculations

- (1) We use theorem (5.13) to show that the Möbius strip and the cylinder both have fundamental group \mathbb{Z} :

Proof. Letting $X = \mathbb{E}^1 \times I$, it is clear that as \mathbb{E} and $[0, 1]$ are convex, that X is simply connected. Consider the action of \mathbb{Z} on the identification square which yields the Möbius loop. It follows that

$$\pi_1(\mathbb{E} \times [0, 1] / \mathbb{Z}) = \pi_1(M) \cong \mathbb{Z}$$

This is similar for the torus. □

- (2) Consider $S^n \subset \mathbb{E}^{n+1}$. Given a loop α in S^n , we find a loop β in \mathbb{E}^{n+1} which is based at the same point as α , and is made up of a finite number of straight line segments, and satisfy

$$\|\alpha(s) - \beta(s)\| < 1, \quad s \in [0, 1]$$

And, we use this to deduce that S^n is simply connected when $n \geq 2$:

Proof. Note that for each n , S^n is compact. Likewise, as $[0, 1]$ is compact, $\alpha([0, 1])$ is compact. Now, either α contains antipodal points, or it doesn't. Let

$$\{\mathcal{O}_b\}_b$$

be an open cover of $\alpha([0, 1])$. Then, it has a finite subcover; say,

$$\{\mathcal{O}_{b_i}\}_{i=1}^N$$

If α has no antipodal points, then we are done as the path β which connects one point from each \mathcal{O}_{b_i} at the same value is the desired path.

Now, if α has antipodal points for some $s \in [0, 1]$ we can disjointize neighbourhoods so that $\alpha(s) \in V_i$, and $-\alpha(s) \in V_j$, such that $\alpha(s) \notin V_j$ and $-\alpha(s) \in V_i$, such that

$$\bigcup V_i = \bigcup \mathcal{O}_{b_i}$$

and the construction of β follows, as above.

To conclude the proof, we show that S^n is simply connected: Indeed; by construction, no line adjoining $\alpha(s)$ and $\beta(s)$ goes through the origin. Thus, the projective straight line homotopy shows that S^n is simply connected, as \mathbb{E}^{n+1} is for $n \geq 2$.

Note, this breaks down in the case where the stereographic projective homotopy fails. This completes the proof. \square

- (3) Considering the 'proof' of theorem (5.13), we show that for $g_1, g_2 \in G$, $\gamma_1 \cdot (g_1 \circ \gamma_2)$ joins x_0 to $g_1 g_2(x_0)$, where γ_1, γ_2 are paths from x_0 to $g_1(x_0), g_2(x_0)$, respectively. In addition, we use this to deduce that ϕ is a homomorphism:

Proof. As the statement on page one-hundred-two points out,

$$\gamma_1 \cdot (g_1 \circ \gamma_2)$$

is a path from x_0 to $\gamma_1 g_2(x_0)$. Thus, as in the proof of (5.13),

$$\begin{aligned} \phi(g_1 g_2) &= \langle \pi \circ (\gamma_1 \cdot g_1 \circ \gamma_2) \rangle \\ &= \langle \pi \circ \gamma_1 \cdot \pi \circ \gamma_2 \rangle \\ &= \langle \pi \circ \gamma_1 \rangle \langle \pi \circ \gamma_2 \rangle \\ &= \phi(g_1) \phi(g_2) \end{aligned}$$

As $\pi \circ \gamma_1 \cdot \pi \circ \gamma_2$ is a loop at $\pi(x_0)$ (it follows that they are in the same homotopy class). \square

- (4) Let $\pi : X \rightarrow Y$ be a covering map and α a path in Y . We show that α lifts to a (unique) path in X which begins at any preassigned point of $\pi^{-1}(\alpha(0))$:

Proof. As $X = \bigcup_{y \in Y} \{y\}$, we can form an open cover of Y by evenly covered, i.e. canonical, neighbourhoods. Let \mathcal{O} be the aforementioned open cover, and set

$$\alpha^{-1}(\mathcal{O})$$

to be the family of open sets which cover $[0, 1]$. As $[0, 1] \subset \mathbb{R}$, is compact, and

$$\bigcup \alpha^{-1}(\mathcal{O})$$

is an open cover of $[0, 1]$, $\bigcup \alpha^{-1}(\mathcal{O})$ has a Lebesgue number, δ .

Now, choose $n \in \mathbb{N}$, so that $1/n < \delta$. Then, consider the partition of $[0, 1]$, given by

$$\{0, 1/n, 2/n, \dots, n-1/n\}$$

Then, for each $i = 1, 2, \dots, n$,

$$\alpha\left(\left[\frac{i-1}{n}, \frac{i}{n}\right]\right)$$

lies inside a canonical neighbourhood of Y . Thus, by setting

$$t_i = \frac{i-1}{n},$$

the result follows. \square

- (5) Let $\pi : X \rightarrow Y$ be a covering map, $p \in Y$, $q \in \pi^{-1}(p)$, and $F : I \times I \rightarrow Y$ a map such that

$$F(0, t) = F(1, t) = p, \quad t \in [0, 1]$$

We use the argument of lemma (5.11) to find a map $F' : I \times I \rightarrow X$ which satisfies

$$\pi \circ F' = F, \quad F'(0, t) = q, \quad t \in [0, 1]$$

and is unique:

Proof. As the argument of lemma (5.11) is really quite bad, we proceed with a more technical proof via problem twenty. \square

- (6) With the terminology above, we note that for each $t \in [0, 1]$, we have a path $F_t(s) = F(s, t)$ in Y which begins at p . Let F'_t be its unique lift to a path X which begins at q , and set $F'(s, t) = F'_t(s)$. We show that F' is continuous and lifts F :

Proof. By the homotopy lifting lemma, there exists a unique homotopy lift, B of F . By the proof so mentioned, B is continuous and satisfies

$$\pi \circ B \equiv F$$

But, by the path lifting lemma, as $F'(s, t_0)$, $t_0 \in [0, 1]$ is a lift of a path $F(s, t)$ with initial point q ,

$$B = F'$$

\square

- (7) We describe the homomorphism $f_* : \pi_1(S^1, 1) \rightarrow \pi_1(S^1, f(1))$ induced by each of the following maps:

- $f(e^{i\theta}) = e^{i(\theta+\pi)}, \theta \in [0, 2\pi]$:

Proof. Note, $f_*(\langle \alpha \rangle) = \langle f(\alpha) \rangle = \langle -\alpha \rangle$. And, as $f(1) = -1$, $f_* \equiv id$. \square

- $f(e^{i\theta}) = e^{in\theta}, \theta \in [0, 2\pi], n \in \mathbb{Z}$:

Proof. Note, $f(1) = 1$. Thus, for each loop, we have $f_*(\langle \alpha \rangle) = \langle n\alpha \rangle$. So, f_* sends the loop component to the n -th degree of α . \square

- $f(e^{i\theta}) = \begin{cases} e^{i\theta} & \theta \in [0, \pi] \\ e^{i(2\pi-\theta)} & \theta \in [\pi, 2\pi] \end{cases}$:

Proof. Note, $f(1) = 1$. f_* means that any loop part on the upper half of S^1 is identified to the lower part of S^1 . Thus, it returns the straight line for each loop on S^1 . \square

- (8) For each of the three different action of \mathbb{Z}_2 on the torus, in section 4.4, we describe the homomorphism from the fundamental group of the torus to that of the orbit space induced by the natural identification map:

Proof. \square

- (9) As in problem eight, page ninety-one, we show that it is impossible to find a homotopy from h to the identity which is relative to the two boundary circles of A :

Proof. If $h \simeq_F i \text{ rel } \{c_1, c_2\}$, then

$$\alpha^{-1}\beta \simeq_F c \text{ rel } \{0, 1\}$$

However, $\alpha^{-1}\beta$ is homotopic to the loop of constant radius which we know is non-trivial via a deformation retract. \square

5.4 Homotopy Type

Throughout this section, \simeq will typically denote homotopy equivalence.

- (1) If $X \simeq Y$ and $X' \simeq Y'$, we show that $X \times X' \simeq Y \times Y'$:

Proof. We are given maps $f : X \rightarrow Y$, $g : Y \rightarrow X$, $f' : X' \rightarrow Y'$ and $g' : Y' \rightarrow X'$, such that $g \circ f \simeq_{\gamma'} 1_X$, $f \circ g \simeq_{\gamma} 1_Y$, $g' \circ f' \simeq_{\delta'} 1_{X'}$ and $f' \circ g' \simeq_{\delta} 1_{Y'}$.

From the component-wise definition of the product topology, $F : X \times X' \rightarrow Y \times Y'$, defined by $F(x, x') = (f(x), f'(x'))$, and $G : Y \times Y' \rightarrow X \times X'$, defined by $G(y, y') = (g(y), g'(y'))$ are continuous. Further, $F \circ G \simeq 1_{Y \times Y'}$, and $G \circ F \simeq 1_{X \times X'}$, via the canonical maps, component-wise defined: $\mathcal{F} : (X \times X') \times I \rightarrow X \times X'$, $\mathcal{F} \equiv (\gamma', \delta')$, and $\mathcal{G} : (Y \times Y') \times I \rightarrow Y \times Y'$, $\mathcal{G} \equiv (\gamma, \delta)$, respectively.

Thus, $X \times X' \simeq Y \times Y'$. \square

- (2) We show that the cone, CX , is contractible for any space (X, τ_X) :²

Proof. By definition, we have $CX = X \times I / X \times \{1\}$. Define $H : CX \times I \rightarrow CX$ by

$$H((x, t), s) = (x, t(1 - s))$$

Then, we have

$$H((x, t), 0) = (x, t) = 1_{CX}$$

$$H((x, t), 1) = (x, 0)$$

Now, if $\mathcal{O}_{CX} \in \tau_{CX}$, then $H^{-1}(\mathcal{O}_{CX}) = (\mathcal{O}_{CX}, I) \in CX \times I$. Thus, H is continuous. This proves the result. \square

- (3) We show that the punctured torus deformation retracts onto the one-point union of two circles.

Proof. We consider the torus as the identification space of a square, X , bounded by the box whose edge points are $(-1, -1), (-1, 1), (1, 1), (1, -1)$. Assume, without loss of generality, that the point $(0, 0)$ is removed.

We show that $F : T \times I \rightarrow T$ defined by

$$F(x, t) = (1 - t)x + t \frac{x}{\|x\|}$$

is a deformation retract onto the one point union of two circles:

As

$$F(x, 0) = x$$

$$F(x, 1) = \frac{x}{\|x\|}$$

²We proved another proof for this. The first of which is in the extra lemmas section.

F is a deformation retract onto ∂X . The function $f : \partial X \rightarrow S_1^1 \vee S_2^1$, given by

$$f(x) = \begin{cases} g(x) & x \in (-1, \pm 1)t + (1, \pm 1)(1-t) \\ s(x) & x \in (\pm 1, 1)t + (\pm, -1)(1-t) \end{cases}$$

for all $t \in I$, where g, s are the guaranteed homeomorphisms from $[a, b]$ to S^1 , is a homeomorphism by the Gluing Lemma. \square

- (4) For each of the following cases, we choose as base point in C and describe the generators for the fundamental groups of C and S . Further, we write down the homomorphism, in terms of these generators, the fundamental groups induced by the inclusion of C in S .

Consider the following examples of a circle C embedded in a surface S :

- (a) $S = \text{Möbius Strip}$ and $C = \partial S$:
 - (b) $S = S^1 \times S^1 = T^1$ and $C = \{(x, y) \in T^1 : x = y\}$:
 - (c) $S = S^1 \times I$ and $C = S^1 \times 1$:
- (5) Suppose that $f, g : S^1 \rightarrow X$ are homotopic maps. We prove that the spaces formed from X by attaching a disc, using f and using g are homotopy equivalent; in other words, we prove that $X \cup_f D \simeq X \cup_g D$:

Proof. We show that this is true for S^{n-1} . To show this, we claim that $X \cup_f D^n$ and $X \cup_g D^n$ are deformation retracts of the same space, $X \cup_F (D^n \times I)$, where

$$f \simeq_F g$$

As $F : S^1 \times I \rightarrow S^1$ is a homotopy between f and g , we have that

$$\begin{aligned} F(x, 0) &= f(x) \\ F(x, 1) &= g(x) \end{aligned}$$

We claim that the function $r : D^n \times I \rightarrow D^n \times \{0\} \cup S^{n-1} \times I$ given by

$$r(x, t) = \begin{cases} (\frac{2x}{2-t}, 0) & t \in [0, 2(1 - \|x\|)] \\ (\frac{x}{\|x\|}, 2 - \frac{2-t}{\|x\|}) & t \in [2(1 - \|x\|), 1] \end{cases}$$

is a retraction.

Lemma. r , as defined above is a retraction:

In the first case, suppose that

$$(x, t) \in D^n \times \{0\} \cup S^{n-1} \times I = A$$

is such that $x \in D^n$, and $t = 0$. Then, we have that

$$r(x, t) = r(x, 0) = (x, 0)$$

In the second case, if $t > 0$, and $x \in S^{n-1}$, then $\|x\| = 1$ and so $t \in [0, 1]$ and

$$r(x, t) = \left(\frac{x}{1}, 2 - \frac{2-t}{1} \right) = (x, t)$$

Thus, $r|_A = 1_A$. A similar argument shows that $r(D^n \times I) = A$.

To complete the proof of the lemma, we note that as r is polynomial in x and t , it follows that r is continuous. Consequently, r is a retraction. \square

We further claim that r is a deformation retraction. To show this, consider the function

$$\begin{aligned} d : (D^n \times I) \times I &\rightarrow D^n \times I \\ d((x, t), s) &= s(r(x, t)) + (1 - s)(x, t) \end{aligned}$$

As

$$\begin{aligned} d((x, t), 0) &= (x, t) \\ d((x, t), 1) &= r(x, t) \end{aligned}$$

it follows that d is a homotopy between r and the identity. Consequently, r is a deformation retraction.

Using the fact that r is a deformation retraction we have that $X \cup_F (D^n \times I)$ deformation retracts onto

$$X \cup_F (D^n \times \{0\} \cup S^{n-1} \times I) = X \cup_f (D^n \times \{0\}) \cong D^n$$

(where $X \cup_F (D^n \times \{0\} \cup S^{n-1} \times I) \cong (D^n \times \{0\} \cup_f X)$ via the map sending $D^n \times \{0\}$ and X to itself, and $S^{n-1} \times I$ to $F(S^{n-1} \times I)$.)

A similar argument show a deformation retract of $D^n \times I$ onto

$$D^n \times \{1\} \cup S^{n-1} \times I$$

which gives the identification space with g as the attaching map. And, the result follows. \square

- (6) We use the previous problem, and the third example of homotopy given in section 5.1, to show that the 'dunce hat' has the homotopy type of a disc, and is therefore contractible:

Proof. This is so horrendously explained by Armstrong. By "the third example of homotopy given in section 5.1," we assume that Armstrong really means "(to take as the canonical definition of 'dunce cap') that the dunce cap is constructed by gluing D^2 to S^1 via the map $g : S^1 \rightarrow S^1$ given by

$$g(e^{i\theta}) = \begin{cases} e^{4i\theta} & 0 \leq \theta \leq \pi/2 \\ e^{4i(2\theta-\pi)} & \pi/2 \leq \theta \leq 3\pi/2 \\ e^{8i(\pi-\theta)} & 3\pi/2 \leq \theta \leq 2\pi \end{cases}$$

That is, $S^1 \cup_g D^2$ is the dunce hat. With this assumption, we continue with the proof:

Consider the adjunct space given by $i : S^1 \rightarrow D^2$; $X \cup_i D^2$. To conclude, we show that g is homotopic to the identity map:

Consider $F : S^1 \times I \rightarrow S^1$ given by

$$F(e^{i\theta}, t) = tg(e^{i\theta}) + (1 - t)e^{i\theta}$$

This is the straight line homotopy;

$$g \simeq_F i$$

Thus, by exercise twenty-seven, page one-hundred-nine, $i, g : S^1 \rightarrow D^2$ are homotopic, so that

$$S^1 \cup_g D^2 \simeq S^1 \cup_i D^2 \cong D^2$$

and the result follows. \square

- (7) We show that the 'house with two rooms' is contractible:

Proof. A rigorous proof of this theorem is probably beyond the scope of this section. However, a sketch of it is given in Allen Hatcher's *Algebraic Topology*, Chapter 0. text³. \square

- (8) We give a detailed proof to show that the cylinder and the Möbius strip have the homotopy type of the circle:

Proof. We have previously shown that both the cylinder and the Möbius strip are homotopic to S^1 . For a more general argument:

A deformation retract $F : X \times I \rightarrow X$ of X onto $A \subset X$ induces a homotopy equivalence by taking $F(-, 1) : X \rightarrow A$ and the inclusion $\iota : A \rightarrow X$. Thus, the composition of the homotopy equivalences $M \rightarrow S^1 \rightarrow S^1 \times I$ prove the result. \square

- (9) Let X be the comb space. We prove that the identity map of X is not homotopic rel $\{p\}$, to the constant map, $p = (0, 1/2)$:

Proof. This is a problem in Munkers's *Topology*. In his text, the point p is $(0, 1)$. We proceed with this assumption:

Let $p = x_0 = (0, 1)$. Consider a neighborhood U of x_0 which is disjoint from $I \times \{0\}$. Suppose $H : X \times I \rightarrow X$ is the homotopy starting with the identity on X and ending with the constant map $X \rightarrow \{x_0\}$ such that $H(x_0, t) = x_0$ for all times t . That means $\{x_0\} \times I$ has a neighborhood $H^{-1}(U)$. By the tube lemma (this is the statement that in the product topology, there is an

³cite this

open set containing a compact product), there is an open set V such that $\{x_0\} \times I \subset V \times I \subset H^{-1}(U)$. That means every point in V stays in U during the entire deformation. However a point $y = (a, 1)$ must traverse a path to x_0 and no such path exists within U . \square

(10) *FTA*: We prove the fundamental theorem of algebra:

Proof. This is shown in Munkers's *Topology*, page three-hundred-fifty. \square

5.5 Brouwer Fixed-Point Theorem

Throughout, we say a topological space (X, τ_X) has the fixed-point property if every continuous function from X to itself has a fixed point.

(1) We determine which of the following have the fixed point property:

- The 2-Sphere: S^2 does not exhibit the fixed point property.

Proof. We have previously shown that the map $f : S^n \rightarrow S^n$, given by $f(x) = -x$ is a homeomorphism. \square

- The Torus: $T^1 = S^1 \times S^1$ does not exhibit the fixed point property.

Proof. As $T^1 = S^1 \times S^1$, and the antipodal map, f , on S^1 is a homeomorphism, we have that $F : T^1 \rightarrow T^1$, defined by $F((x, y)) = (f(x), f(y))$, is continuous and doesn't exhibit the fixed point property. \square

- The interior of the unit disc: $\text{Int}D^1$ does not exhibit the fixed point property:

Proof. Note that the interior of the unit disc is homeomorphic to the euclidean plane by a homeomorphism $h : \text{Int}D^1 \rightarrow \mathbb{E}^2$. We define a function g from the euclidean plane to the euclidean plane by $g((x, y)) = (x + 1, y)$. Then, $h^{-1} \circ g \circ h$ is a continuous function from the disc to itself that has no fixed point. \square

- The one point union of two circles: $X \vee Y = X \sqcup Y / \{p\}$ does not exhibit the fixed point property:

Proof. Define a function f as follows: If $x \in X$ then we map e^{ix} to $e^{i(x+\pi)}$. Also if $y \in Y$, we map y to $f(p) = e^{i\pi/2}$. This function f , as defined on the one point union, leaves no points fixed. \square

(2) Suppose X and Y are of the same homotopy type and X has the fixed-point property. We prove that Y does not necessarily have the fixed point property:

Proof. Let Y be the subspace $(0, 1) \subset \mathbb{E}^n$, and let $X = \{1/2\}$. Note that Y is homotopic to X by the straight line homotopy, and every map from X to itself has a fixed point. Yet, the function $f : Y \rightarrow Y$, defined by $f(y) = y^2$ has no fixed point. \square

- (3) Suppose that X retracts onto the subspace $A \subset X$, and that A has the fixed point property. We show that X may not exhibit the fixed point property:

Proof. Take X and Y , as above. The straight-line homotopy proves the assertion. \square

- (4) We show that if X retracts onto the subspace A , and X has the fixed-point property, then A also has it:

Proof. Let $f : A \rightarrow X$ be a continuous function. Since X retracts onto A , there exists a map $g : X \rightarrow A$ such that $g|_A \equiv 1_A$. Then, $f \circ g$ is a continuous function from X to X , and so has a fixed point. Hence, there exists an $a \in A$ such that $f(g(a)) = f(a) = a$. This completes the proof. \square

- (5) We deduce that the fixed-point property holds for the 'house with two rooms', X :

Proof. As was previously show, X is contractible to some x_0 . Thus, there exists some map $F : X \times I \rightarrow X$ such that $F(x, 0) = x$, and $F(x, 1) = x_0$. Thus, by extra lemma ⁴, every map $f : X \rightarrow X$ is null-homotopic. In particular, this includes that maps $g : X \rightarrow X$. Thus, X has the fixed point property⁵.

To use the previous problems hints, we could think about starting with the unit cylinder, and pushing in the areas from the top and bottom. However, this method is not rigorous. \square

- (6) Let f be a fixed-point-free map from a compact metric space (X, d) to itself. We prove there is a positive number ϵ such that $d(x, f(x)) > \epsilon$, $\forall x \in X$:

Proof. We show the contrapositive; Suppose that for all $\epsilon > 0$ there exists an element $x \in X$ such that $d(x, f(x)) \leq \epsilon$. Pick $x_1 \in X$ such that $d(x_1, f(x_1)) \leq \frac{1}{2} < 1$.

It follows that there exists a set, $\{x_1, \dots, x_n\} \subset X$ such that $d(x_i, f(x_i)) < d(x_k, f(x_k))$, where $i < k$ and $d(x_k, f(x_k)) < \frac{1}{k}$, for all $1 \leq k \leq n$.

Now, let

$$\epsilon_0 = \frac{1}{2} \min \left\{ d(x_1, f(x_1)), \dots, d(x_n, f(x_n)), \frac{1}{n+1} \right\}$$

⁴include this

⁵If this is not totally clear, see extra lemma.

Then there exists an element $x \in X$ such that $d(x, f(x)) \leq \epsilon_0$. Pick $x \in X$ that satisfies this property, and call this x_{n+1} . Note here that $d(x_n, f(x_n)) < d(x_{n+1}, f(x_{n+1}))$ by construction. So that for all $i < n + 1$, $d(x_i, f(x_i)) \leq d(x_n, f(x_n)) < d(x_{n+1}, f(x_{n+1}))$. Thus by induction, we have created an infinite sequence $\{x_n\}_{n=1}^\infty$ of distinct points such that $d(x_k, f(x_k)) < 1/k$ for each positive integer k . And, $\{d(x_n, f(x_n))\}_n$ is a monotone decreasing sequence.

Since X is compact, every infinite subset has a limit point. Therefore $\{x_n\}_{n=1}^\infty$ has a limit point $x \in X$.

Let $\epsilon > 0$ be given. Since f is continuous at x , then there exists a $\delta > 0$ such that $d(x, a) < \delta \implies d(f(x), f(a)) < \epsilon/3$ for all $a \in X$. By the Archimedian Property, we can find a positive integer k , such that

$$\frac{1}{k} < \frac{\epsilon}{3}$$

Set $r = \min\{\frac{\epsilon}{3}, \delta\}$. Then, since x is a limit point of $\{x_n\}_{n=1}^\infty$, there are infinitely many points of the sequence, such that $d(x_N, x) < r$. Thus, there exists a point x_N such that $N > k$ and $d(x_N, x) < r$. Since $d(x_N, x) < \delta$, then $d(f(x_N), f(x)) < \epsilon/3$. Since $N > k$,

$$d(x_N, f(x_N)) < d(x_k, f(x_k)) < 1/k < \frac{\epsilon}{3}$$

Therefore

$$d(x, f(x)) \leq d(x, x_N) + d(x_N, f(x_N)) + d(f(x_N), f(x)) < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon$$

Thus, for all $\epsilon > 0$, we have that $d(x, f(x)) < \epsilon$. Thus, $d(x, f(x)) \leq 0$, and hence $d(x, f(x)) = 0$. Consequently, $f(x) = x$, and so f is not a fixed-point-free map. \square

- (7) We show that the unit ball, B^n , in \mathbb{E}^n , with $(1, 0, \dots, 0)$ removed does not exhibit the fixed point property:

Proof. Let $p = (1, \dots, 0)$. Consider the function $f : B^n \setminus \{p\} \rightarrow B^n \setminus \{p\}$, defined by $f(x) = \frac{x+p}{2}$. We show that f is continuous:

Let \mathcal{O} be an open set in $B^n \setminus \{p\}$. Let $x \in f^{-1}(\mathcal{O})$. Then $f(x) \in \mathcal{O}$, and since \mathcal{O} is open there exists an $r > 0$ such that $B_r(f(x)) \subset \mathcal{O}$. Let $y \in B_{2r}(x)$. Then $|x - y| < 2r$. Then,

$$|f(x) - f(y)| = \left| \frac{x+p}{2} - \frac{y+p}{2} \right| = \frac{1}{2}|x - y| < r$$

Thus, $y \in B_r(f(x))$. Hence, $y \in f^{-1}(\mathcal{O})$. Thus, $B_{2r}(x) \subset f^{-1}(\mathcal{O})$. Therefore, $f^{-1}(\mathcal{O})$ is open and f is continuous.

Further, f does not have the fixed-point property; If $f(x) = x = (x_1, \dots, x_n)$, then we have $2x = x + p$. Implying that $x_i = 2x_i$, $2 \leq i \leq n$. Thus, $x_i = 0$. However, then the only solution to $x_1 + 1 = 2x_1$ is $x_1 = 1$. This is a contradiction since we must have $x = p \notin B^n \setminus \{p\}$. \square

- (8) We show that the one-point union of X and Y , $X \vee Y$, has the fixed-point property if, and only if, both X and Y have it:

Proof.

(\implies): Suppose that the one point union $X \vee Y$ has the fixed-point property. Let $f : X \rightarrow X$ be a map, and let p be the point glued together in the one point union. Then, define $g : X \vee Y \rightarrow X \vee Y$, as $g(x) = f(x)$, if $x \in X$, and $g(x) = f(p)$, if $x \in Y$. By the gluing lemma, g is a continuous map and so, by hypothesis, has a fixed point x_0 . Note that g is a map into X . Thus the fixed point must be in X . Hence, by construction, $x_0 = g(x_0) = f(x_0)$, so that f has a fixed point. Similarly, any continuous function from Y to itself has a fixed point.

(\impliedby): Suppose X and Y have the fixed point property and let $f : X \vee Y \rightarrow X \vee Y$ be a map. Then, suppose $f(p) \in X$, and define a map $g : X \rightarrow X$ such that $g(x) = f(x)$, if $f(x) \in X$, and $g(x) = p$, if $f(x) \notin X$. Then, since g is continuous, it has a fixed point. By construction, the fixed point must be one $x = g(x) = f(x)$. Thus, f has a fixed point. Next, suppose $f(p) \in Y$, and define a map $g : Y \rightarrow Y$ such that $g(y) = f(y)$, if $f(y) \in Y$, and $g(y) = p$, if $f(y) \notin Y$. Then, since g is continuous, it has a fixed point. By construction, the fixed point must be one $y = g(y) = f(y)$. Consequently, f has a fixed point. \square

- (9) How does changing 'continuous function' to 'homeomorphism' in the definition of the fixed-point property affect problem 33, 37?

Proof. We first examine problem 33:

S^2 : This topological space would not exhibit the fixed point property. We know that the antipodal map is a homeomorphism which leaves no points fixed.

$S^1 \times S^1$: Likewise, the antipodal map on each component, S^1 , provides another counter example.

Int D : Neither does this space have the homeomorphic fixed point property. To see this, we note that $\text{Int } D \simeq \mathbb{E}$, and the map $g : \mathbb{E} \rightarrow \mathbb{E}$, given by $g(x) = x + 1$ is a homeomorphism which leaves no points fixed.

$X \vee Y$: First, consider the map $E_1 : X \vee Y \rightarrow X \vee Y$, given by $E_1(x) = x$, if $x \in X - \{p\}$. And, $E_2(y) = e^{yi\pi/2}$. Then, E_1 is a homeomorphism, which leaves X fixed, except for p . Similarly, define E_2 for Y . Then, it follows that $E_2 \circ E_1 : X \vee Y \rightarrow X \vee Y$ is a homeomorphism that leaves no points of $X \vee Y$ fixed. Thus, $X \vee Y$ does not exhibit the fixed point property⁶.

Now, we examine problem 37:

This shape does not exhibit the homeomorphic fixed point property. Consider $\partial B^n = S^{n-1}$, and let $p = (1, 0, \dots, 0)$. Then, $B^n - \{p\} = \text{Int } B^n \cup S^{n-1} - \{p\}$. Furthermore, without loss of generality, $S^{n-1} - \{p\} \simeq \{-p\}$.⁷ It follows that $\text{Int } B^n \cup S^{n-1} - \{p\} \simeq \text{Int } B^n \cup \{-p\}$. To conclude, we exhibit a homeomorphism on $\text{Int } B^n \cup \{-p\}$ which does not have a fixed point:

Consider $f : \text{Int } B^n \cup \{-p\} \rightarrow \text{Int } B^n \cup \{-p\}$ defined by

$$f(x) = \begin{cases} -x & x \notin \{-p, 0\} \\ 0 & x = -p \\ -p & x = 0 \end{cases}$$

The fact that f is 1-1 and onto is clear by construction. Further, f is continuous, by the Gluing Lemma⁸. It follows that f is a homeomorphism, as $f^{-1} \equiv -f$. However, f clearly leaves no points fixed. \square

5.6 Separation of the Plane

Most of the exercises in the section are too advanced for an undergraduate course in algebraic topology. As a result, we refer to well-known sources, which prove the results.

- (1) Let A be a compact subset of \mathbb{E}^n . We show that $\mathbb{E}^n - A$ has exactly one unbounded component:

Proof. As A is compact, it is closed and bounded. Further, we identify \mathbb{E}^n with $S^n - \{p\}$ under stereographic projection, π . Then, A is a compact set in S^n such that $\{p\} \notin A \cap S^n$. The remainder of this proof is an extension of lemma (61.1), Munkers' *Topology*⁹. \square

- (2) Let J be a polygonal Jordan curve in the plane. Let p be a point in the unbounded component of $\mathbb{E}^2 - J$ which does not lie on any of the lines produced by extending each of the segments of J in both directions. Given

⁶We make use of the previous proof; that is $X \vee Y$ has the fixed point property iff X and Y both have it.

⁷See previous exercises.

⁸Armstrong, pg. 69.

⁹cite this

a point x of $\mathbb{E}^2 - J$, say that x is inside (outside) J if the straight line joining p to x cuts across J an odd (even) number of times. We show that the complement of J has exactly two components, namely the set of inside points and the set of outside points:

Proof. See *The Jordan-Schönflies Theorem and the Classification of Surfaces*, by Carsten Thomassen. \square

- (3) Let J be a polygonal Jordan curve in the plane, and let X denote the closure of the bounded component of J . We show that X is homeomorphic to a disc:

Proof. See *The Jordan-Schönflies Theorem and the Classification of Surfaces*, by Carsten Thomassen. \square

- (4) We prove Schönflies theorem for polygonal Jordan curves:

Proof. See *The Jordan-Schönflies Theorem and the Classification of Surfaces*, by Carsten Thomassen. \square

- (5) If J is a Jordan curve in the plane, we use theorem (5.12) to show that the frontier of any component of $\mathbb{E}^2 - J$ is J :

Proof. See *The Jordan-Schönflies Theorem and the Classification of Surfaces*, by Carsten Thomassen. \square

- (6) We give an example of a subspace of the plane which has the homotopy type of a circle, which separates the plane into two components, but which is not the frontier of both these components:

Proof. Consider the standard annulus in the plane,

$$A = \{(x, y) : x^2 + y^2 = r \wedge 1 \leq r \leq 2\}$$

Then, as we have previously shown, A deformation retracts onto $S^1 \subset A$, and so A has the homotopy type of a circle. Further,

$$\text{Int } A = \{(x, y) : x^2 + y^2 < 1\},$$

with $\partial \text{Int } A = S^1 \neq A$. In addition,

$$\text{Out } A = \{(x, y) : x^2 + y^2 > 2\},$$

and $\partial \text{Out } A = 2S^1 \neq A$. \square

- (7) We give example of simple closed curves which separate, and fail to separate
- the torus:

Proof. Considering the torus as the identification space of $[0, 1]^2$, the circle of radius $1/4$ centered at $(1/2, 1/2)$, separates T , while the line $x = 1/2$ does not. \square

• $\mathbb{R}P^2$:

Proof. Note that $\mathbb{R}P^2$ is given by the equivalence relation $x \sim \lambda x$, $\lambda \neq 0$. Thus, a curve in $\mathbb{R}P^2$ is a set of equivalence classes. Thus, a circle of radius $1/4$ centered at $(1/2, 1/2)$, which is equivalent to an arc on the first quadrant of S^1 separates $\mathbb{R}P^2$. And, S^1 does not separate $\mathbb{R}P^2$. \square

- (8) Let X be the subspace of the plane which is homeomorphic to a disc. We generalize the argument of theorem (5.21) to show that X cannot separate the plane:

Proof. See *The Jordan-Schönflies Theorem and the Classification of Surfaces*, by Carsten Thomassen. \square

- (9) Suppose that X is both connected and locally path-connected. We show that a map $f : X \rightarrow S^1$ lifts to a map $f' : X \rightarrow \mathbb{R}$ if, and only, if the induced homomorphism $f_* : \pi_1(X) \rightarrow \pi_1(S^1)$ is the zero homomorphism:

Proof. See the accompanying solution¹⁰. \square

5.7 The Boundary of a Surface

- (1) We use an argument similar to that of theorem (5.23) to prove that \mathbb{E}^2 and \mathbb{E}^3 are not homeomorphic:

Proof. We have previously show that $\pi_1(\mathbb{E}^m)$ is trivial for $m \geq 3$, while $\pi_1(\mathbb{E}^2)$ is not. The result follows. \square

- (2) We use the material of this section to show that the spaces X , and Y illustrated in problem twenty-four of chapter one are not homeomorphic:

Proof. Suppose, to the contrary, that X and Y were homeomorphic. Then, by corollary (5.25), they have homeomorphic boundaries. Further, it follows from the proof of theorem (5.24), that the outer boundaries, $\text{Out } X$, of X and $\text{Out } Y$ of Y must be homeomorphic, as well as the inner boundaries. However, we have previously shown that they are not homeomorphic, as one is separable by removing a point, and the other is not. \square

¹⁰<http://stanford.edu/class/math215b/Sol3.pdf>

Chapter 6

Triangulations

Unfortunately, this is the point in Armstrong when the definitions presented are even less clear. For this reason, the reader is referred to this short paper which provides an excellent introduction to simplicial complexes and triangulations; the definitions presented are inline with those in Armstrong.

6.1 Triangulating Spaces

(1) We construct triangulations for

- the cylinder:

Proof. Please see figure 6.1, with the usual identifications. □

- the Klein Bottle:

Proof. Please see figure 6.2, with the usual identifications. □

- the double Torus:

Proof. Please see figure 6.3, with the usual identifications. □

(2) We finish off the proof of lemma (6.3):

Proof. To do this, we mimic the proof theorem (3.30) and show that $|K|$ is path-connected;

Let

$$Z = \bigcup_{0 < \delta < \epsilon} (B(x, \delta) \cap |K|) = \left(\bigcup_{0 < \delta < \epsilon} \right) \cap |K|$$

We claim that $|K| = Z$. Since $B(x, \delta)$ is open in \mathbb{E}^n for each δ , and $|K|$ has the subspace topology, Z is open. Further,

$$|K| - Z = |L| = \bigcup_{y \in |L|} B(y, \delta_y)$$




Figure 6.1: Triangulation of the cylinder.




Figure 6.2: Triangulation of Klein bottle.




Figure 6.3: Triangulation of the double torus.

where $\delta_y = d(x, y) - \epsilon$. So, $|K| - Z$ is open. And, since $|K|$ is connected, $Z \neq \emptyset$, theorem (3.20) (page fifty-seven), implies $Z = |K|$. \square

- (3) If $|K|$ is a connected space, we show that any two vertexes of K can be connected by a path whose image is a collection of vertexes and edges of K :

Proof. Let v be a 0-simplex in K and let \mathcal{P}_v be the subcomplex of K consisting of all edge path starting with v , along with the edge paths they span.

Now, if σ is a simplex of K that has a vertex, call it w in \mathcal{P}_v , then every vertex w' of σ is in \mathcal{P}_v (by definition). Thus, $\sigma \in \mathcal{P}_v$.

To conclude the proof, we show that $|\mathcal{P}_v|$ is clopen in $|K|$: The fact that $|\mathcal{P}_v|$ is closed follows from the fact that each 1-simplex is closed and there are only finitely many. Likewise, $|\mathcal{P}_v|$ is open, as each 1-simplex is closed.

Thus, as $v \in \mathcal{P}_v \neq \emptyset$, we have by theorem (3.20) that $\mathcal{P}_v = K$ and the result follows. \square

- (4) We check that $|CK|$ and $C|K|$ are homeomorphic spaces:

Proof. Let K be a complex in \mathbb{E}^n . Then, CK consists of the simplices of K and the join of each of them to some vertex, v via 1-simplices and the 0-simplex v itself. So, $|CK|$ is the geometric cone on $|K|$. By lemma (4.5), page sixty-eight,

$$|CK| \simeq C|K|$$

\square

- (5) We show that if X and Y are triangulable spaces, then $X \times Y$ is triangulable:

Proof. Let $\kappa : |K| \rightarrow X$, and $\lambda : |L| \rightarrow Y$ be the triangulations of X , and Y . We claim that the function

$$\phi : |K| \times |L| \rightarrow X \times Y, \quad \phi(k, l) = (\kappa(k), \lambda(l))$$

is the desired triangulation.

As κ, λ are homeomorphisms, the result follows. \square

- (6) We show that if K and L are complexes in \mathbb{E}^n , that $|K| \cap |L|$ is a polyhedron:

Proof. Unfortunately, this is another ill-posed question; what Armstrong neglects to say is that X is a polyhedron iff X can be triangulated (this is the canonical definition given elsewhere). Further, this question is really beyond the scope of this chapter. As such, we give a proof which assumes that every convex, compact, connected subset of \mathbb{E}^n can be triangulated; we actually show that $|K| \cap |L|$ is the union of such subsets of \mathbb{E}^n , where

connected implies a maximally connected component (we note that it need not be the case that $|K| \cap |L|$ is connected. Consider figure):

Let

$$|K| = \bigcup_{\sigma \in |K|} \sigma, \quad |L| = \bigcup_{\sigma \in |L|} \sigma$$

be the simplicial decompositions of $|K|$ and $|L|$. Then, as the intersection of convex sets is convex, we have

$$|K| \cap |L| = \bigcup_{\substack{\sigma \in |K| \\ \sigma' \in |L|}} (\sigma \cap \sigma')$$

is convex and each $\sigma \cap \sigma'$ is convex.

Now, as $\dim K, \dim L < N < \infty$, by the well-ordering principle, we start with the smallest dimensional intersection of simplices, say $n \in \mathbb{N}$. Pick any such (non-empty) intersection and label it κ_1^n . Problem 3, shows that κ_1^n is edge connected and further, as each simplex is completely determined by its vertexes, κ_1^n determines some maximally connected, convex, compact subset of \mathbb{E}^n .

To conclude the proof, we extend this process to some other intersections

$$\kappa_1^n, \kappa_2^n, \kappa_3^n, \dots, \kappa_q^n, \quad \dim \kappa_i^n = \dim \kappa_j^n = n$$

This method can be extended to the next higher' dimension $n + 1$ to show that

$$|K| \cap |L| = \bigcup_{\substack{n \leq i \leq N \\ j}} |\kappa_j^i|$$

which is indeed triangulable, and so a polyhedron. \square

(7) We show that S^n and P^n are both triangulable:

Proof. We give a triangulation of S^n and use the antipodal map to establish a triangulation for P^n :

Let $(S^n)^+$ denote the northern hemisphere of S^n and consider $S^n \subset \mathbb{E}^n$ and the canonical basis

$$\{e_1, e_2, \dots, e_n\}, \quad e_i = (0, \dots, 0, 1_i, 0, \dots, 0)$$

We claim that the set X of 1-simplices given by

$$\sigma_i = e_i e_{i+1}, \quad 2 \leq i \leq n, \quad n+1 \sim 2,$$

along with the set of 1-simplices defined by

$$\sigma_{Ni} = N e_i, \quad N = (1, \dots, 0), \quad 2 \leq i \leq n$$

along with the 2-simplexes σ_i, σ_{N_i} bound, and all their vertexes is a triangulation of the norther hemisphere of S^n :

Indeed; The fact that X is a simplicial complex is clear from construction and the definition of S^n and by radial projection, π , we have

$$\pi : |X| \rightarrow (S^n)^+$$

is a homeomorphism, and so π is a triangulation of $(S^n)^+$. Further by considering the simplexes

$$\sigma_{S_i} = Se_i, \quad S = (0, \dots, 1), \quad 2 \leq i \leq n$$

we extend π to a map

$$\Pi : |X \cup Y| \hookrightarrow S^n$$

It follows that S^n is triangulated by Π .

To show that $\mathbb{R}P^n$ is triangulable, we note that the antipodal map

$$i_{-1} : S^n \rightarrow S^n$$

is a homeomorphism and that $\mathbb{R}P^n$ is defined as the identification map induced by i_{-1} , call it ϕ .

Consequently,

$$\phi\Pi : |X \cup Y| \rightarrow \mathbb{R}P^n$$

is a homeomorphism, and so a triangulation. \square

- (8) We show that the 'dunce hat' is triangulable, but the 'comb space' is not:

Proof. Please see figure 6.4, with the usual identifications.

To show that the comb space X is not triangulable, we first prove that it is not locally connected:

Lemma. If X were locally connected we could find a connected open neighbourhood \mathcal{O} for each neighbourhood of p . But, each open neighbourhood \mathcal{O} contains infinitely many terms of the sequence

$$\left\{ \left(\frac{1}{n}, \frac{1}{2} \right) \right\}_{n=1}^{\infty}$$

As

$$\left\{ \left(\frac{1}{n}, \frac{1}{2} \right) : n \geq n_0 \right\}$$

is clearly disconnected, it follows that X is not locally connected. \square

From lemma (6.3), it follows that $|K|$ is locally connected (this is shown in [?]). Now, since local connectedness is a topological property, if X were triangulable then it must be locally connected. However, we have just shown that X is not locally connected. \square




Figure 6.4: The second barycentric subdivision of the 'dunce cap' yields a triangulation.

6.2 Barycentric Subdivision

- (1) We present a visualization of the first barycentric subdivision of a 3-Simplex:

Proof.

□

- (2) Let \mathcal{F} be an open cover of $|K|$, we show the existence of a barycentric subdivision K^r , with the property that given a vertex v of K^r , there is an open set $U \in \mathcal{F}$ which contains all the simplexes of K^r that have v as a vertex:

Proof.

□

- (3) Let L be a subcomplex of K , and let N be the following collection of simplexes of K^2 : a simplex B lies in N if we can find a simplex C in L^2 such that the vertexes of B and C together determine a simplex of K^2 . We show that

- N is a subcomplex of K^2 :

Proof.

□

- $|N|$ is a neighbourhood of $|L|$ in $|K|$:

Proof.

□

- (4) We use the construction of problem 11 to prove that if X is a triangulable space, and Y is a subspace of X which is triangulated by a subcomplex of some triangulation of X , then the space obtained from X by shrinking Y to a point is triangulable:

Proof.

□

6.3 Simplicial Approximation

- (1) We use the simplicial approximation theorem to show that the n -sphere is simply connected for $n \geq 2$:

Proof.

□

- (2) We show that if $k < m, n$ that any map from S^k to S^m is null homotopic, and that the same is true of any map from S^k to $S^m \times S^n$:

Proof.

□

- (3) We show that a simplicial map from $|K|$ to $|L|$ induces a simplicial map from $|K^m|$ to $|L^m|$ for any m :

Proof. □

- (4) Suppose that $s : |K^m| \rightarrow |L|$ simplicially approximates $f : |K^m| \rightarrow |L|$, and $t : |L^n| \rightarrow |M|$ simplicially approximates $g : |L^n| \rightarrow M$. We determine if $ts : |K^{m+n}| \rightarrow |M|$ is always a simplicial approximation for $gf : |K^{m+n}| \rightarrow |M|$:

Proof. □

- (5) We use the simplicial approximation theorem to show that the set of homotopy classes of maps from one polyhedron to another is always countable:

Proof. □

- (6) This is left to the reader as an exercise.

6.4 The Edge Group of a Complex

- (1) We use van Kampen's theorem to calculate the fundamental group of the double torus by dividing the surface into two halves (each of which is a punctured torus). And then, we do the calculation again, but this time splitting the surface into a disc and the closure of the complement of the disc:

Proof. □

- (2) We show that the edge paths E_1 and E_2 introduced in the proof of theorem (6.10) are equivalent:

Proof. □

- (3) We prove that the 'dunce hat' is simply connected, using van Kampen's theorem:

Proof. □

- (4) Let X be a path-connected triangulable space. We examine how attaching a disc to X affects the fundamental group of X :

Proof. □

- (5) Let G be a finitely presented group. We construct a compact triangulable space which has fundamental group G :

Proof. □

6.5 Triangulating Orbit Spaces

- (1) Suppose $\{V, S\}$ satisfies the hypotheses of the realization theorem, and label the elements of V as v_1, \dots, v_k . Suppose that x_i denotes the point

$$(i, i^2, \dots, i^{2m+1}) \in \mathbb{E}^{2m+1}$$

We show that any $2m + 2$ of the points x_1, \dots, x_k are in general position, and so that the correspondence $v_i \leftrightarrow x_i$ can be used to realize $\{V, S\}$ in \mathbb{E}^{2m+1} .

Proof.

□

- (2) By the previous problem the vertex scheme of any one-dimensional complex can be realized in \mathbb{E}^3 . We find a one-dimensional complex whose vertex scheme cannot be realized in \mathbb{E}^2 :

Proof.

□

- (3) Consider the antipodal action on S^2 and the triangulation shown in fig. 6.17. We show that the map $\psi : |K|/G \rightarrow |K^1/G|$ is a homeomorphism, and draw the resulting triangulation of the projective plane:

Proof.

□

- (4) We show that the map $\psi : |K|/G \rightarrow |K/G|$ is a homeomorphism iff the action of G on $|K|$ satisfies:

- The vertexes of a 1-simplex of K never lie in the same orbit:

Proof.

□

- If the sets of vertexes of v_0, \dots, v_k, a and v_0, \dots, v_k, b span simplexes of K , and if a, b lie in the same orbit, then there exists $g \in G$ such that $g(v_i) = v_i$ for $0 \leq i \leq k$ and $g(a) = b$.

Proof.

□

- (5) We check that the previous conditions are always satisfied if we replace K by its second barycentric subdivision:

Proof.

□

6.6 Infinite Complexes

- (1) We find the triangulations of the following spaces from the crystallographic groups shown in fig. 4.5, page eighty-four;

- the sphere:

Proof. □

- the torus:

Proof. □

- the Klein bottle:

Proof. □

- (2) We check that the construction illustrated in fig. 6.21 really can be carried out to produce a realization of T in \mathbb{E}^2 :

Proof. □

- (3) We show that the following collection of simplexes in \mathbb{E}^2 is not a simplicial complex: For each $n \in \mathbb{N}$, we have a vertical 1-simplex joining $(1/n, 0)$ to $(1/n, 1)$ and a sloping 1-simplex with vertexes $(1/n, 0)$, $(1/n + 1, 1)$. In addition, we have a 1-simplex on the y axis joining $(0, 0)$ to $(1, 1)$.

Proof. We now consider the case of removing the 1-simplex on the y axis: □

- (4) We determine whether the comb space, or the space illustrated in figure 3.4, page sixty-three, can be triangulated by an infinite simplicial complex:

Proof. □

- (5) We show that the free group on a countable number of generators is a subgroup of $\mathbb{Z} \otimes \mathbb{Z}$, and deduce that any subgroup of this group must be free:

Proof. □

- (6) We show

- if X is the polyhedron of a finite complex, and if h is simplicial, the pointwise periodic implies periodic:

Proof. □

- a connected infinite complex K and a simplicial homeomorphism of $|K|$ which is pointwise periodic but not periodic:

Proof. □

- (7) We determine whether or not a pointwise periodic homeomorphism of a compact space need be periodic:

Proof. □

- (8) Let G be a group of homeomorphisms of the space X . We show

- that if $N \trianglelefteq G$, G/N acts in a natural way on X/N and that X/G is homeomorphic to $(X/N)/(G/N)$:

Proof. □

- if F is the smallest normal subgroup of G which contains all the elements that have fixed points, then G/F acts freely on X/F in the sense that only the identity element has fixed points:

Proof. □

- (9) Suppose that in addition to the conditions in the previous problem that X is a simply connected polyhedron, G acts simplicially, and X/G is triangulated so that the projection $p : X \rightarrow X/G$ is simplicial. Choose a vertex v of X as a base point and define $\phi : G \rightarrow \pi_1(X/G, p(v))$ as follows: given $g \in G$, join v to $g(v)$ by an edge path E in X ; then $\phi(g)$ is the homotopy class of the edge loop $p(E)$.

We show that ϕ is a homomorphism, and that each element of F goes to the identity under ϕ , and ϕ is onto:

Proof. □

- (10) Under the assumptions of the previous problem, we show that X/F is simply connected, and that the action of G/F on X/F satisfies the hypothesis of theorem (5.13):

Proof. □

- (11) We deduce theorem (6.18):

Proof. □

Chapter 7

Surfaces

7.1 Classification

- (1) We show that punching a disc out of the sphere and adding a cross cap in its place gives a representation of $\mathbb{R}P^2$ as a surface in \mathbb{E}^3 with self-intersections:

Proof.

□

- (2) Let X consist of S^2 plus one extra point p . The neighbourhoods of the points of S^2 are the usual ones, and those of p are sets of the form $[U - \{(0, 0, 1)\}] \cup \{p\}$ where U is a neighbourhood of $(0, 0, 1)$ in S^2 . We show that X is not Hausdorff, but is locally homeomorphic to the plane:

Proof. We now consider whether or not it seems reasonable to call X a surface:

□

- (3) We show that the connected sum of a torus with itself is a sphere with two handles, and the connect sum of a projective plane with itself is a Klein bottle:

Proof.

□

- (4) We determine what the connect sum of a torus and a projective plane is:

Proof.

□

7.2 Triangulation and Orientation

- (1) Suppose we want to triangulate a surface which has a boundary. We examine how the definition of a combinatorial surface needs to be adjusted:

Proof. □

- (2) Let K be a combinatorial surface. We

- show that the triangles of K can be labelled T_1, \dots, T_s in such a way that T_i always has an edge in common with at least one of T_1, \dots, T_{i-1} :

Proof. □

- build a model for the surface of $|K|$ in the form of a regular polygon in the plane, which has an even number of sides, and whose sides have to be identified in pairs in some way:

Proof. □

- (3) We show that if $h : |K| \rightarrow S$ is a triangulation of a closed surface, then K must be a combinatorial surface:

Proof. □

- (4) Let G be a finite group which acts as a group of homeomorphisms of a closed surface S in such a way that the only element with any fixed points is the identity. We show

- that the orbit space S/G is a closed surface:

Proof. □

- S may be orientable, yet S/G non-orientable:

Proof. □

- that if S/G is orientable, then S does not have to be:

Proof. □

- (5) Let K be an orientable combinatorial surface, orient all its triangles in a compatible manner, and suppose that $h : |K| \rightarrow |K|$ is a simplicial homeomorphism. Suppose, further, that there is a triangle A , oriented by the ordering (uvw) whose image $h(A)$ occurs with the orientation $(h(u)h(v)h(w))$ induced by h . We prove that the same must hold for any other triangle of K :

Proof. We give an example of an orientable combinatorial surface and a simplicial homeomorphism which is not orientation-preserving: \square

- (6) Let K be an orientable combinatorial surface. Suppose that G acts simplicially on $|K|$. We show that if the action is fixed-point free and each element of G is orientation-preserving that the complex K^2/G is an orientable combinatorial surface:

Proof. \square

7.3 Euler Characteristics

- (1) We prove lemma (7.8):

Proof. \square

- (2) We prove lemma (7.9):

Proof. \square

- (3) We prove from the previous problem that the Euler characteristic of a graph Γ is a topological invariant of $|\Gamma|$:

Proof. \square

- (4) Let K be a finite complex. Suppose that G acts simplicially on $|K|$, and that the action is fixed-point free. We show that

$$\chi(K) = |G|\chi(K^2/G)$$

where $|G| = \text{card } G$:

Proof. \square

- (5) Let K be a combinatorial surface such that $|K|$ is in the plane as in problem 6, and let J denote the boundary curve of the resulting regular polygon. We show that

$$\chi(K) = \chi(\Gamma) + 1$$

and deduce lemma(.6) from lemma (7.5):

Proof. \square

- (6) Continuing from the previous problem, suppose that Γ has an edge, one end of which is not joined to any other edges. We show that $\chi(K) = 2$ implies

$$|K| \cong S^2 :$$

Proof. \square

7.4 Surgery

- (1) We thicken each curve in fig. 7.16 and decide whether the result is a cylinder or a Möbius strip and describe the effect of doing surgery along the curve:

Proof. □

- (2) We show that the surface illustrated in fig. 7.17 is homeomorphic to one of the standard ones using the procedure of theorem (7.14):

Proof. □

- (3) Let $Z \subset Y \subset X$ be three concentric discs in the plane. We find a homeomorphism from X to itself which is the identity on the boundary circle of X and which throws Y onto Z :

Proof. □

- (4) Suppose that we have two discs in the plane, both of which are bounded by polygonal curves, and one which lies in the interior of the other. We prove that the region between them is homeomorphic to an annulus:

Proof. □

- (5) with the notation of lemma (7.13) and problem 19, we find a homeomorphism $h : D^1 \rightarrow X$ such that $h(D) = Y$ and

$$h(\overline{\text{star}(\hat{A}, K^2)}) = Z$$

and prove lemma (7.13):

Proof. □

7.5 Surface Symbols

- (1) We show that the two surfaces shown in fig. 7.22 are not homeomorphic:

Proof. □

- (2) We determine what happens if we remove the interiors of two disjoint discs from a closed surface, then glue the two resulting boundary circles together:

Proof. □

- (3) We use the classification theorem to show that the operation of connected sum is well-defined:

Proof. □

- (4) Assume that every compact surface can be triangulated. We show that if the boundary component of a surface is non-empty, then it consists of a finite number of disjoint circles:

Proof. □

- (5) We show that any compact connected surface is homeomorphic to a closed surface from which the interiors of a finite number of disjoint discs have been removed:

Proof. □

- (6) We determine the fundamental group of the space obtained by punching k holes in the sphere:

Proof. □

- (7) Write $H(p, r)$ for $H(p)$ with the interiors of r disjoint discs removed, and write $M(q, s)$ for $M(q)$ with s discs similarly removed. We show that $H(p, r)$ can be obtained from a $(4p + 3r)$ -sided polygonal region in the plane by gluing up its edges according to the surface symbol

$$a_1 b_1 a_1^{-1} b_1^{-1} \dots a_p b_p a_p^{-1} b_p^{-1} x_1 y_1 x_1^{-1} \dots x_r y_r x_r^{-1} :$$

Proof. □

- (8) We find a surface symbol for $M(q, s)$, as defined in the previous problem:

Proof. □

- (9) We calculate the fundamental groups of $H(p, r)$ and $M(q, s)$:

Proof. □

- (10) We show

- that $H(p, r) \cong H(p', r')$ implies $p = p'$ and $r = r'$:

Proof. □

- that $M(q, s) \cong M(q', s')$ implies $q = q'$ and $s = s'$:

Proof. □

- that there are no values of p, q, r, s for which $M(p, r) \cong H(q, s)$:

Proof.

□

- (11) Define the genus of a compact connect surface to be the genus of the closed surface obtained by capping off each boundary circle with a disc. We show that a compact connect surface is completely determined by wheter or not it is orientable, together with its genus and its number of boundary circles:

Proof.

□

- (12) We posit a general result from the two picture in fig. 7.24:

Proof.

□

Chapter 8

Simplicial Homology

8.1 Cycles and Boundaries

(1)

8.2 Homology Groups

(1)

8.3 Examples

(1)

8.4 Simplicial Maps

(1)

8.5 Stellar Subdivision

(1)

8.6 Invariance

(1)