

# Microstructure-derived flexibility in the acorn weevil exoskeleton

M. Andrew Jansen,<sup>1,\*</sup> Nikhilesh Chawla,<sup>2</sup> and Nico M. Franz<sup>1</sup>

<sup>1</sup>*School of Life Sciences, Arizona State University, Tempe, AZ 85287, USA*

<sup>2</sup>*School for Engineering of Matter, Energy, and Transport,  
Arizona State University, Tempe, AZ 85287, USA*

(Dated: June 16, 2018)

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

We report novel modifications to the composite microstructure of the exoskeleton in the snout of acorn weevils (Coleoptera: Curculionidae) belonging to the genus *Curculio* Linnaeus, 1756.

As a weevil (snout beetle), members of the genus *Curculio* are typified by the presence of a highly elongate structure on the head, called the rostrum (snout). This structure is a hollow, cylindrical extension of the exoskeleton of the otherwise nearly-spherical head, which bears at its apex the terminal chewing mouthparts. The space inside of the rostrum contains the esophagus, various muscles and tendons used for feeding, and hemolymph that serves as a rough equivalent to blood in insects. By contrast, the solid shell of the rostrum is comprised entirely of cuticle, which can be considered a laminate composite consisting of various arrangements of chitin fibers embedded in a protein matrix (see ). We have found that the composite profile of the rostrum is strongly differentiated from the head capsule and other body parts, with modification of both the relative layer thicknesses and fiber orientation angles of cuticle regions (viz. exocuticle and endocuticle), which we describe in detail below.

These modifications enable the snout to be flexed until straight, remaining within the elastic limits of the material, and without evident alteration of the mechanical properties of the individual components of the cuticle across the structure. Thus, the flexibility of the rostrum appears to be derived exclusively through modification of the composite architecture and fiber arrangement in the exoskeleton. Support for this hypothesis has come from three lines of evidence: first, examination of the cuticle microstructure across the length of the snout has revealed modifications to the composite structure of the rostrum; second, tensile testing of the structure has demonstrated that the mechanical strength of the cuticle components are consistent along the length of the structure among species; and, third, fatigue testing has shown that a highly curved rostrum is capable of flex-

ing hundreds of thousands of times without damage to the structure. We additionally report on the fracture mechanics of the snout, as pertains to both cuticle composite structure and tensile behavior, and consider how modification of the cuticle may reduce the risk of rostral fracture during oviposition. To our knowledge, this is the first time such modifications have been reported for enhancing structural elasticity in the insect exoskeleton.

## MICROSTRUCTURE

Display 1: Pictures of heads, macrofiber arrangement, and exo-endo ratio at base and apex, across species Explain how everything is laid out across the length of the rostrum, emphasizing that this is key to predicting and understanding the mechanical behavior of the snout during bending.

## TENSILE TESTING AND FRACTURE MECHANICS

The predicted behavior of the snout is borne out by the data, where UTS is strongly correlated with the cross-sectional area of the endocuticle across species.

## FATIGUE TESTING OF *CURCULIO CARYAE*

## CONCLUSION

## METHODS

Methods, including statements of data availability and any associated accession codes and references, are available in the online version of this paper.

\* corresponding author, email: majansel@asu.edu

---

REFERENCES

[1] Leslie Lamport, *LaTeX: a document preparation system*, Addison Wesley, Massachusetts, 2nd edition, 1994.

ACKNOWLEDGMENTS

AUTHOR CONTRIBUTIONS

ADDITIONAL INFORMATION

Supplementary information is available in the online version of the paper. Reprints and permissions information is available online at [www.nature.com/reprints](http://www.nature.com/reprints). Correspondence and requests for materials should be addressed to M.A.J.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

METHODS

Method 1

Method 2

Method 3

Statistical analysis

Code availability

Data availability

---

REFERENCES

[1] Leslie Lamport, *LaTeX: a document preparation system*, Addison Wesley, Massachusetts, 2nd edition, 1994.