

# kreyszig-10-1

March 23, 2025

## 0.1 Erwin Kresyzig 10.1 Problems

### 0.1.1 2-11 LINE INTEGRAL. WORK.

Calculate

$$\int_C \mathbf{F}(\mathbf{r}) \cdot d\mathbf{r}$$

for the given data. If  $\mathbf{F}$  is a force, this gives the work done by the force in the displacement along  $C$ . Show the details.

[ ]:

We repeat for clarity the definition of the “line” integral  $\mathbf{F} \cdot d\mathbf{r}$  along the curve  $C$  from first principles.

The integral along  $C$  is defined as:

$$\int_C \mathbf{F} \cdot d\mathbf{r}$$

where  $\mathbf{F}$  and  $\mathbf{r}$  are vector functions to be expanded in vector components as:

$$\mathbf{F} = [F_1, F_2, F_3] = F_1 \mathbf{i} + F_2 \mathbf{j} + F_3 \mathbf{k},$$

$$\mathbf{r}(t) = [x(t), y(t), z(t)] = x(t) \mathbf{i} + y(t) \mathbf{j} + z(t) \mathbf{k},$$

$$\text{and the differential } d\mathbf{r} = [dx, dy, dz].$$

$\mathbf{F} = [F_1, F_2, F_3]$  is a vector function in the 3-dimensional space to be evaluated along the path  $C$  defined by  $\mathbf{r}(t)$ , i. e.,  $\mathbf{F}$  is  $([F_1, F_2, F_3])(\mathbf{r}(t))$ .

The differential  $d\mathbf{r}$  may be calculated by chain rule:

$$d\mathbf{r} = \frac{d\mathbf{r}(t)}{dt} dt = \mathbf{r}'(t) dt.$$

The components are,

$$\frac{d}{dt} \mathbf{r}(t) = \frac{d}{dt} [x(t), y(t), z(t)] = \left[ \frac{d}{dt} x(t), \frac{d}{dt} y(t), \frac{d}{dt} z(t) \right] = [x'(t), y'(t), z'(t)]$$

or more concisely:

$$d\mathbf{r} = [dx, dy, dz] = [x'(t), y'(t), z'(t)] dt.$$

Thus the dot product in the integrand is, along  $C$ ,

$$\begin{aligned}\mathbf{F} \cdot d\mathbf{r} &= [F_1, F_2, F_3] \cdot [dx, dy, dz] \\ &= ([F_1, F_2, F_3](\mathbf{r}(t))) \cdot [x'(t), y'(t), z'(t)] dt \\ &= ((F_1(\mathbf{r}(t)) \cdot x'(t) + (F_2(\mathbf{r}(t))) \cdot y'(t) + (F_3(\mathbf{r}(t))) \cdot z'(t)) dt\end{aligned}$$

Thus the line integral in the 3-dimensional space is reduced to an integral in one variable along a closed interval in the real line, say,  $[a, b]$ .

[ ]:

$$2. \quad F = [y^2, -x^2], \quad C : y = 4x^2 \text{ from } (0, 0) \text{ to } (1, 4)$$

Given  $\mathbf{F} = [y^2, -x^2]$  and  $C : y = 4x^2$  from  $(0, 0)$  to  $(1, 4)$ , we have to compute the line integral:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C (y^2 dx - x^2 dy).$$

Here  $\mathbf{F} = y^2 \mathbf{i} - x^2 \mathbf{j}$ .

So  $F_1 = y^2$  and  $F_2 = -x^2$ .

Parameterize  $C$  using  $x = t$ , so  $y = 4t^2$ , and  $t$  goes from 0 to 1.

For differentials,  $dx = dt$  and  $dy = 8t dt$ .

Substituting into the integrand:  $\mathbf{F} = y^2 \mathbf{i} - x^2 \mathbf{j}$  and expanding dot product:

$$\begin{aligned}\mathbf{F} \cdot d\mathbf{r} &= F_1 dx + F_2 dy = y^2 dx - x^2 dy \\ &= (4t^2)^2 dt - t^2 (8t dt) \\ &= 16t^4 dt - 8t^3 dt \\ &= (16t^4 - 8t^3) dt\end{aligned}$$

Integrating:

$$\begin{aligned}\int_0^1 (16t^4 - 8t^3) dt &= \left[ \frac{16t^5}{5} - \frac{8t^4}{4} \right]_0^1 \\ &= \frac{16}{5} - \frac{8}{4} \\ &= \frac{16}{5} - 2 = \frac{6}{5}.\end{aligned}$$

[ ]:

3  $F$  as in Prob. 2,  $C$  from  $(0, 0)$  straight to  $(1, 4)$ . Compare.

---

Using same  $\mathbf{F} = [y^2, -x^2]$  from previous problem. Now  $C$  is the straight line from  $(0, 0)$  to  $(1, 4)$ , i.e.,  $y = 4x$ .

Parameterize:  $x = t$ ,  $y = 4t$ ,  $t$  from 0 to 1.

Differentials:  $dx = dt$ ,  $dy = 4 dt$ .

Substituting into integrand:

$$\begin{aligned}\mathbf{F} \cdot d\mathbf{r} &= (4t)^2 dt - t^2(4 dt) \\ &= 16t^2 dt - 4t^2 dt \\ &= 12t^2 dt\end{aligned}$$

Integrating:

$$\int_0^1 12t^2 dt = \left[ 12 \frac{t^3}{3} \right]_0^1 = 12 \cdot \frac{1}{3} = 4.$$

**Comparison:** The work done differs from problem 2 for same function along the parabolic path, where it was  $\frac{6}{5} = 1.2$ . Here, it is 4. Thus integral of  $\mathbf{F}$  is not path-independent.

[ ]:

4.  $F = [xy, x^2y^2]$ ,  $C$  from  $(2, 0)$  straight to  $(0, 2)$

---

$C$  is the straight line from  $(2, 0)$  to  $(0, 2)$ , i.e.,  $y = -x + 2$ .

Parameterize:  $x = 2 - t$ ,  $y = t$ ,  $t$  from 0 to 2.

Differentials:  $dx = -dt$ ,  $dy = dt$ .

Substitute:

$$\begin{aligned}\mathbf{F} \cdot d\mathbf{r} &= (2-t)t(-dt) + (2-t)^2t^2(dt) \\ &= (t^4 - 4t^3 + 5t^2 - 2t) dt\end{aligned}$$

Integrate:

$$\begin{aligned}\int_0^2 (t^4 - 4t^3 + 5t^2 - 2t) dt &= \left[ \frac{t^5}{5} - \frac{4t^4}{4} + \frac{5t^3}{3} - \frac{2t^2}{2} \right]_0^2 \\ &= \frac{32}{5} - 16 + \frac{40}{3} - 4 \\ &= -\frac{4}{15}.\end{aligned}$$

[ ]:

**5**  $F$  as in Prob. 4,  $C$  the quarter-circle from  $(2, 0)$  to  $(0, 2)$  with center  $(0, 0)$

---

Using  $\mathbf{F} = [xy, x^2y^2]$ ,  $C$  is the quarter-circle from  $(2, 0)$  to  $(0, 2)$ , center  $(0, 0)$ , radius 2.

Parameterize:  $\mathbf{r} = [2 \cos t, 2 \sin t]$ ,  $t$  from 0 to  $\frac{\pi}{2}$ .

Differentials:  $dx = -2 \sin t \, dt$ ,  $dy = 2 \cos t \, dt$ .

Thus

$$\mathbf{F} = [4 \cos t \sin t, 16 \cos^2 t \sin^2 t].$$

$$\mathbf{F} \cdot d\mathbf{r} = 8 \cos t \sin^2 t (3 - 4 \sin^2 t) \, dt$$

Substitute  $u = \sin t$ ,  $du = \cos t \, dt$ ,  $u$  from 0 to 1:

$$\int_0^1 8(3u^2 - 4u^4) \, du = 8 \left( 1 - \frac{4}{5} \right) = \frac{8}{5}.$$

[ ]:

**6**  $F = [x - y, y - z, z - x]$ ,  $C : \mathbf{r} = [2 \cos t, t, 2 \sin t]$  from  $(2, 0, 0)$  to  $(2, 2\pi, 0)$

---

$C : \mathbf{r} = [2 \cos t, 2 \sin t, 0]$ ,  $t$  from 0 to  $2\pi$  (a closed loop).

Differentials:  $dx = -2 \sin t \, dt$ ,  $dy = 2 \cos t \, dt$ ,  $dz = 0$ .

Thus

$$\mathbf{F} = [2 \cos t - 2 \sin t, 2 \sin t, -2 \cos t].$$

$$\mathbf{F} \cdot d\mathbf{r} = 4 \sin^2 t \, dt.$$

Integrate using  $\sin^2 t = \frac{1 - \cos 2t}{2}$ :

$$\int_0^{2\pi} 4 \sin^2 t \, dt = 2 \left[ t - \frac{\sin 2t}{2} \right]_0^{2\pi} = 4\pi.$$

[ ]:

7  $F = [x^2, y^2, z^2]$ ,  $C : \mathbf{r} = [\cos t, \sin t, e^t]$  from  $(1, 0, 1)$  to  $(1, 0, e^{2\pi})$ . Sketch  $C$ .

---

Given  $\mathbf{F} = [x^2, y^2, z^2]$ ,  $C : \mathbf{r} = [\cos t, \sin t, e^t]$ ,  $t$  from 0 to  $2\pi$ .

Differentials:  $dx = -\sin t \, dt$ ,  $dy = \cos t \, dt$ ,  $dz = e^t \, dt$ .

$\mathbf{F} = [\cos^2 t, \sin^2 t, e^{2t}]$ .  $\mathbf{F} \cdot d\mathbf{r} = \sin t \cos t (\sin t - \cos t) \, dt + e^{3t} \, dt$ .

Integrate: First term is 0 (closed loop in  $u = \sin t$ ). Second term:

$$\int_0^{2\pi} e^{3t} \, dt = \frac{e^{6\pi} - 1}{3}.$$

**Sketch:** The curve is a spiral on the cylinder  $x^2 + y^2 = 1$ , with  $z$  increasing from 1 to  $e^{2\pi}$  at the speed of exponential function.

```
[3]: import numpy as np
import matplotlib.pyplot as plt
from mpl_toolkits.mplot3d import Axes3D

# Create parameter t from 0 to 2
t = np.linspace(0, 2*np.pi, 100)

# Calculate x, y, z coordinates
x = np.cos(t)
y = np.sin(t)
z = np.exp(t)

# Create 3D plot
fig = plt.figure(figsize=(10, 8))
ax = fig.add_subplot(111, projection='3d')

# Plot the curve
ax.plot(x, y, z, 'b-', label='r(t) = [cos t, sin t, e^t]')

# Plot start and end points
ax.scatter([1], [0], [1], color='green', s=100, label='Start (1,0,1)')
ax.scatter([1], [0], [np.exp(2*np.pi)], color='red', s=100, label='End ↴
↵(1,0,e^(2))')


# Set labels
ax.set_xlabel('X axis')
ax.set_ylabel('Y axis')
ax.set_zlabel('Z axis')
ax.set_title('Parametric Curve C: r(t) = [cos t, sin t, e^t]')

# Add legend
ax.legend()
```

```
# Adjust the view angle
ax.view_init(elev=20, azim=45)

plt.show()
```

Parametric Curve C:  $\mathbf{r}(t) = [\cos t, \sin t, e^t]$



[ ]:

8 Let  $F = [e^x, \cosh y, \sinh z]$ ,  $C : \mathbf{r} = [t, t^2, t^3]$  from  $(0, 0, 0)$  to  $(\frac{1}{2}, \frac{1}{4}, \frac{1}{8})$ . Sketch  $C$ .

Let  $\mathbf{F} = [e^x, \cosh y, \sinh z]$ ,  $C : \mathbf{r}(t) = [t, t^2, t^3]$  from  $(0, 0, 0)$  to  $(\frac{1}{2}, \frac{1}{4}, \frac{1}{8})$ , so  $t$  ranges from 0 to  $\frac{1}{2}$ .

Differentials:  $dx = dt$ ,  $dy = 2t dt$ ,  $dz = 3t^2 dt$ . (Or,  $d\mathbf{r} = [1, 2t, 3t^2]dt$ .)

Then,  $d\mathbf{r} = [1, 2t, 3t^2] dt$ , and along  $C$ ,  $\mathbf{F} = [e^t, \cosh(t^2), \sinh(t^3)]$ , so:

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &= \int_0^{\frac{1}{2}} [e^t + 2t \cosh(t^2) + 3t^2 \sinh(t^3)] dt \\ &= \int_0^{\frac{1}{2}} e^t dt + \int_0^{\frac{1}{4}} \cosh u du + \int_0^{\frac{1}{8}} \sinh v dv \\ &= [e^t]_0^{\frac{1}{2}} + [\sinh u]_0^{\frac{1}{4}} + [\cosh v]_0^{\frac{1}{8}} \\ &= (e^{\frac{1}{2}} - 1) + \sinh \frac{1}{4} + (\cosh \frac{1}{8} - 1) \\ &= e^{\frac{1}{2}} + \sinh \frac{1}{4} + \cosh \frac{1}{8} - 2.\end{aligned}$$

(Substitutions:  $u = t^2$ ,  $v = t^3$ .)

**Sketch:** The curve follows  $y = x^2$ ,  $z = x^3$ , a winding 3D curve from  $(0, 0, 0)$  to  $(\frac{1}{2}, \frac{1}{4}, \frac{1}{8})$ . The projections to  $xy$  and  $xz$  axes give  $y = x^2$  and  $z = x^3$ .

```
[1]: # Import required libraries
import numpy as np
import matplotlib.pyplot as plt
from mpl_toolkits.mplot3d import Axes3D

# Define the parameter t
t = np.linspace(0, 0.5, 100) # 100 points from t=0 to t=0.5

# Parametric equations
x = t
y = t**2
z = t**3

# Create a 3D plot
fig = plt.figure(figsize=(8, 6))
ax = fig.add_subplot(111, projection='3d')

# Plot the curve
ax.plot(x, y, z, label=r'\mathbf{r}(t) = [t, t^2, t^3]', color='blue')

# Mark the start and end points
ax.scatter([0], [0], [0], color='green', s=100, label='Start (0, 0, 0)')
ax.scatter([0.5], [0.25], [0.125], color='red', s=100, label=r'End_
↪\left(\frac{1}{2}, \frac{1}{4}, \frac{1}{8}\right)')

# Set labels and title
ax.set_xlabel('X')
ax.set_ylabel('Y')
```


```

ax.set_zlabel('Z')
ax.set_title('Sketch of the Curve C')
ax.legend()

# Set axis limits for better visualization
ax.set_xlim(0, 0.6)
ax.set_ylim(0, 0.3)
ax.set_zlim(0, 0.15)

# Show the plot
plt.show()

```



[ ]:

9 Let  $F = [x + y, y + z, z + x]$ ,  $C : \mathbf{r} = [2t, 5t, t]$  from  $t = 0$  to 1. Also from  $t = -1$  to 1.



Given  $\mathbf{F} = [x + y, y + z, z + x]$ ,  $C : \mathbf{r} = [2t, 5t, t]$ .

Differentials:  $dx = 2 dt$ ,  $dy = 5 dt$ ,  $dz = dt$ .

$$\mathbf{F} = [7t, 6t, 3t].$$

$$d\mathbf{r} = [2, 5, 1]dt.$$

$$\mathbf{F} \cdot d\mathbf{r} = 47t dt.$$

Integrate from  $t = 0$  to  $1$ :

$$\int_0^1 47t dt = \frac{47}{2}.$$

From  $t = -1$  to  $1$ :

$$\int_{-1}^1 47t dt = 0.$$

[ ]:

**10** Let  $F = [x, -z, 2y]$  from  $(0, 0, 0)$  straight to  $(1, 1, 0)$ , then to  $(1, 1, 1)$ , back to  $(0, 0, 0)$ .

---

$C$  is a closed triangular loop:  $(0, 0, 0)$  to  $(1, 1, 0)$ , to  $(1, 1, 1)$ , to  $(0, 0, 0)$ .

Side 1:

$\mathbf{r}(t) = [t, t, 0]$ ,  $t$  from  $0$  to  $1$ ,  $d\mathbf{r} = [1, 1, 0] dt$ ,  $\mathbf{F} = [t, 0, 2t]$ , so  $\mathbf{F} \cdot d\mathbf{r} = t dt$ . Then,

$$\int_0^1 t dt = \frac{1}{2}.$$

Side 2:

$\mathbf{r}(t) = [1, 1, t]$ ,  $t$  from  $0$  to  $1$ ,  $d\mathbf{r} = [0, 0, 1] dt$ ,  $\mathbf{F} = [1, -t, 2]$ , so  $\mathbf{F} \cdot d\mathbf{r} = -t dt$ . Then,

$$\int_0^1 -t dt = -\frac{1}{2}.$$

Side 3:

$\mathbf{r}(t) = [1 - t, 1 - t, 1 - t]$ ,  $t$  from  $0$  to  $1$ ,  $d\mathbf{r} = [-1, -1, -1] dt$ ,  $\mathbf{F} = [1 - t, -(1 - t), 2(1 - t)]$ , so:

$$\begin{aligned}\mathbf{F} \cdot d\mathbf{r} &= (1 - t)(-1) + (t - 1)(-1) + 2(1 - t)(-1) \\ &= (4t - 4) dt,\end{aligned}$$

and

$$\int_0^1 (4t - 4) dt = [2t^2 - 4t]_0^1 = -2.$$

Total:  $\frac{1}{2} - \frac{1}{2} - 2 = -2$ .

[ ]:

11 Let  $F = [e^{-x}, e^{-y}, e^{-z}]$ ,  $C : \mathbf{r} = [t, t^2, t]$  from  $(0, 0, 0)$  to  $(2, 4, 2)$ . Sketch  $C$ .

For the given endpoints,  $t$  is from 0 to 2.

Differentials:  $dx = dt$ ,  $dy = 2t dt$ ,  $dz = dt$ .

Integrand:

$$\mathbf{F} \cdot d\mathbf{r} = 2e^{-t} + 2te^{-t^2} dt.$$

To integrate:

$$2 \int_0^2 e^{-t} dt + 2 \int_0^2 te^{-t^2} dt = 2(1 - e^{-2}) + (1 - e^{-4}) = 3 - 2e^{-2} - e^{-4}.$$

**Sketch:** The curve follows  $y = x^2$ ,  $z = x$ , a parabolic path in 3D from  $(0, 0, 0)$  to  $(2, 4, 2)$ .

```
[2]: # Import required libraries
import numpy as np
import matplotlib.pyplot as plt
from mpl_toolkits.mplot3d import Axes3D

# Define the parameter t
t = np.linspace(0, 2, 100) # 100 points from t=0 to t=2

# Parametric equations
x = t
y = t**2
z = t

# Create a 3D plot
fig = plt.figure(figsize=(8, 6))
ax = fig.add_subplot(111, projection='3d')

# Plot the curve
ax.plot(x, y, z, label=r'\mathbf{r}(t) = [t, t^2, t]', color='blue')

# Mark the start and end points
ax.scatter([0], [0], [0], color='green', s=100, label='Start (0, 0, 0)')
ax.scatter([2], [4], [2], color='red', s=100, label='End (2, 4, 2)')

# Set labels and title
ax.set_xlabel('X')
ax.set_ylabel('Y')
ax.set_zlabel('Z')
ax.set_title('Sketch of the Curve C')
```

```

ax.legend()

# Set axis limits for better visualization
ax.set_xlim(0, 2.5)
ax.set_ylim(0, 4.5)
ax.set_zlim(0, 2.5)

# Show the plot
plt.show()

```

Sketch of the Curve C

