



PROFOUND HYPERTRIGLYCERIDAEMIA (51 MMOL/L) FOLLOWING CHEMOTHERAPY IN POST-TRANSPLANT LYMPHOPROLIFERATIVE DISEASE.



C BURNETT^{1,2}, B DOUCET^{1,2}, K MORRIS³, G JOHN^{1,2}

¹Kidney Health Service, Royal Brisbane and Women's Hospital, HERSTON, QLD, Australia

²School of Clinical Medicine, The University of Queensland, QLD, Australia

³Cancer Care Services, Royal Brisbane and Women's Hospital, HERSTON, QLD, Australia

Background

- Kidney transplant recipients are at increased risk of malignancy, hypercholesterolaemia, cardiovascular disease. Important to monitor and control cholesterol to avoid increased CV risk.
- Predisposing factors to high cholesterol:
 - diabetes
 - weight gain / obesity
 - steroids
 - mTOR inhibitors, calcineurin inhibitors
 - platinum chemotherapy (1)
 - Lymphoma lipid mechanic alterations
- Post transplant lymphoproliferative disease (PTLD) is a life-threatening complication with an incidence of 2-20%.
- Hypertriglyceridemia is increasingly recognised as a hallmark of lymphoma yet the mechanism is not well understood.

Cholesterol Management

- Baseline triglycerides = 4-6 mmol/L (<2.1) Profound hypercholesterolaemia (51 mmol/L) after platinum-based chemotherapy with dexamethasone.
- No pancreatitis symptoms or eruptive xanthomas or xanthelasma.
- Hba1c = 6.2%. TFT=N. Lipase=N. ApoB=1.51H (<1.2), apoA1=0.97L, Lipid EPP = Type V hyperlipoproteinaemia.
- Treated with insulin infusion, initiation of fenofibrate 145mg once daily, fish oil 2g daily, rosuvastatin 40mg Partial recovery to 10 mmol/L after cessation of Tx.


Case Report

41-year-old teacher, 13 years post kidney transplant

- Kidney Transplant: 27/9/11, 5MM, cPRA null, EBV +/+, no rejection.
- Complications: New onset diabetes, AZA conversion due to low WCC.
- Underlying kidney disease: MPGN.
- PMHx: NAFLD, depression, OSA, distant ex-smoker. ECOG1.
- Meds: CyA 25mgBD, Pred 5mg, metformin, insulin, bactrim, apixaban


PTLD: Monomorphic PTLD (Diffuse large B cell variant, EBV negative)

- Initial disease (abdomen, 29/4/21):** Rituximab, cyclophosphamide, doxorubicin, vincristine and prednisolone.
 - No substantial change in cholesterol. Remission.
- Relapse disease:** Rituximab, dexamethasone, **oxaliplatin**, cytarabine.
 - Deterioration of lipid levels without evidence of pancreatitis.
 - Partial response, complicated by hearing loss / VTE.
- Outcome:** Due to intolerable side effects and pt wishes – declined further salvage chemotherapy or CAR-T / experimental therapy. Local abdominal radiation (15#) given in Dec 2023.



Possible mechanisms of hypertriglyceridemia

- Type V hyperlipoproteinemia: Resistance to treatment, predisposition to high levels
- Platinum chemo and dexamethasone are associated apolipoprotein dysfunction (1, 2).
- Cholesterol regulation is altered in lymphomas, triglycerides can rise with tumour lysis, and burden / relapse (4).



Take home messages

- Mechanism and treatment of hypercholesterolaemia in this population is not well understood, persisting hypercholesterolaemia should prompt lipid electrophoresis.
- In this case, multiple factors contributed to severity of triglycerides: type V dyslipidaemia, calcineurin inhibition, steroids, platinum chemotherapy and lymphoma
- Regular lipid testing is important during treatment of PTLD and may be of use in the monitoring of patients during cancer survivorship.

References

- Wang, G, Su, C, Yin, T 2017, 'Paclitaxel and platinum-based chemotherapy results in transient dyslipidaemia in cancer patients', *Mol Clin Oncol*, 6(2): 261-265. doi: 10.3892/mco.2016.1107
- Koneshamoorthy et al. 2022, 'Massive hypertriglyceridemia associated with paclitaxel; a case report', *Gynecol Oncol Rep*, 42: 101030. doi: 10.1016/j.gore.2022.101030
- Dai et al. 2021, 'Excipient of paclitaxel induces metabolic dysregulation and unfolded protein response', *iScience*, 24(10): 103170. doi: 10.1016/j.isci.2021.103170
- Spiegel, et al. 1982, 'Plasma lipid alterations in leukemia and lymphoma', *American Journal of Medicine*, 72(5):775-782.