

How does excitation synchronize two PING networks?

exp085 · 19 August 2026 · Draft

Abstract

Two frequency-detuned PING networks are connected by reciprocal excitatory projections. This experiment asks whether phase locking is caused mainly by direct excitation of pyramidal cells, recruitment of local inhibition, or interaction between both pathways.

Prior art

Lowet et al. used two 80-excitatory, 20-inhibitory PING networks connected through E-to-E and E-to-I projections. They found that locking depends mainly on frequency detuning and coupling strength. Their topology provides the starting point for this experiment: Lowet et al. (2015).

Phase-response analysis provides a way to connect individual coupling events to network-level locking. It also warns that a population rhythm can respond differently from any one neuron: Lowet et al. (2017) and Xu and Rieke (2021).

Methods

Methods 1–3 are complete. Methods 4–5 are not yet run.

1. **Define the networks.** Build two matched PING networks. Each contains 80 excitatory and 20 inhibitory neurons connected through local E-to-I and I-to-E synapses. Drive Network A at 300 Hz and Network B at 240 Hz; these values place both uncoupled rhythms in the gamma band. Use a 1 ms local E-to-I AMPA decay so each inhibitory neuron fires once, rather than twice, after an excitatory volley. Connect the networks reciprocally through E-to-E and E-to-I projections. Each target receives exactly 8 randomly selected excitatory afferents from the other network. Give the pathways separate weights K_{EE} and K_{EI} and a shared delay d .




Figure 1: SNNLANG circuit export. Each network contains a local E-to-I-to-E PING loop. Reciprocal cross-network excitation targets both populations with exact fan-in 8, separate K_{EE} and K_{EI} controls, and shared delay d .

Generates Result 1.

2. **Confirm uncoupled phase drift.** Set $K_{EE} = K_{EI} = 0$ and run both networks for 2 s with independent input streams. Discard the first 300 ms. Detect excitatory population volleys, interpolate phase between successive volleys, and calculate the wrapped phase difference. Require both rhythms to remain in the gamma band with an inter-volley-interval coefficient of variation below 0.2. Generates Result 2.
3. **Measure each pathway's phase response.** Use Network A and one baseline cycle. Deliver a synchronous probe volley at nine phases from 0.1 to 0.9 of that cycle. Send it separately through coupling-matched E-to-E and E-to-I projections. Each target receives exactly 8 probe afferents with the corresponding coupling strength and delay. Repeat the fixed input without a probe and measure the shift in the next excitatory volley. Positive values mean an advance; negative values mean a delay. Generates Result 3.
4. **Test the coupling pathways.** Continue from the same saved state and input in four conditions: no coupling, E-to-E only, E-to-I only, and both pathways. Track the relative phase in each condition. Generates Result 4.
5. **Trace the mechanism.** Align activity on incoming excitatory volleys. Measure target excitatory activity, target inhibitory activity, inhibition received by the target excitatory population, and the phase correction in the following cycle. Generates Result 5.

Results

Result 1: Network definition

The authored graph contains two matched 80-excitatory, 20-inhibitory PING circuits. It keeps inhibition local and uses reciprocal long-range E-to-E and E-to-I projections. The two

pathways have separate parameters, $K_{EE} = 0.08$ and $K_{EI} = 0.08$, with delay $d = 0.5$ ms. SNNLANG represents the sparse projections through exact-fan-in initialization.

Result 2: Uncoupled phase drift




Figure 2: Uncoupled activity. The upper panels show normalized excitatory and inhibitory population rates over the same 250 ms interval. The lower panel shows the wrapped phase difference across the full post-burn recording. Vertical jumps are phase wrapping.

Network A oscillates at 50.7 Hz with inter-volley-interval CV 0.037. Network B oscillates at 44.5 Hz with CV 0.04. Every inhibitory neuron fires exactly once in every measured cycle. Both networks therefore produce clean, regular gamma rhythms without baseline inhibitory doublets. Their relative phase wraps 10 times, establishing sustained phase drift before coupling.

Result 3: Phase-response curve




Figure 3: Next-volley response to coupling-matched probe volleys. Positive values indicate an advance; negative values indicate a delay.

Direct excitation of the excitatory population has little effect early in the cycle and advances the next volley later in the cycle. Excitation of the inhibitory population has a narrow early delay window. The earliest I-targeted probe produces a second 20-spike inhibitory volley 3.34 ms after the normal volley. This extra inhibition delays the next excitatory volley by 5.9 ms. The strong E-to-I response is therefore a narrow doublet mechanism rather than a broad inhibitory effect.

Result 4: Pathway comparison

Axes. Time against wrapped relative phase.

Traces. No coupling, E-to-E only, E-to-I only, and both pathways.

Why. Identify which pathway is sufficient for locking.

Expectation. The conditions separate direct excitation, feedforward inhibition, and any interaction between them.

Result 5: Event-aligned mechanism

Axes. Time from a source excitatory volley against normalized activity or conductance.

Traces. Arriving excitation, target inhibitory activity, target inhibition, and target excitatory activity.

Why. Establish the order of events that produces each phase correction.

Expectation. The pathway responsible for locking changes before the target network's next corrected volley.