

ABSTRACT

SEARCH FOR CHARGED HIGGS BOSONS IN THE $\tau + \ell$ FINAL STATE WITH 36.1 fb⁻¹ OF pp COLLISION DATA AT $\sqrt{s} = 13$ WITH THE ATLAS EXPERIMENT

Elliot Wesley Parrish, Ph.D.
Department of Physics
Northern Illinois University, 2022
Dhiman Chakraborty and Jahred Adelman, Director

This dissertation uses 139 fb⁻¹ of pp collision data collected at a center of mass energy of $\sqrt{s} = 13$ by the ATLAS detector to search for charged Higgs bosons decaying to a tau lepton and a neutrino ($H^\pm \rightarrow \tau^\pm \nu_\tau$) in association with a leptonically decaying top quark. No significant excess was found, therefore limits are set at the 95% confidence level on the charged Higgs production cross section times the branching fraction into the $\tau^\pm \nu_\tau$ ranging from XX pb to XX fb. These limits are interpreted in the hMSSM benchmark scenario as an exclusion at 95% confidence on $\tan\beta$ as a function of m_{H^\pm} . In this scenario, for $\tan\beta = 60$, the H^\pm mass range up to XXXX GeV is excluded, with all values of $\tan\beta$ excluded for $m_{H^\pm} \leq XXX \text{ GeV}$.

NORTHERN ILLINOIS UNIVERSITY
DE KALB, ILLINOIS

DECEMBER 2022

**SEARCH FOR CHARGED HIGGS BOSONS IN THE $\tau + \ell$ FINAL STATE
WITH 36.1 fb⁻¹ OF pp COLLISION DATA AT $\sqrt{s} = 13$ WITH THE ATLAS
EXPERIMENT**

BY

ELLIOT WESLEY PARRISH
© 2022 Elliot Wesley Parrish

A DISSERTATION SUBMITTED TO THE GRADUATE SCHOOL
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE
DOCTOR OF PHILOSOPHY

DEPARTMENT OF PHYSICS

Dissertation Director:
Dhiman Chakraborty and Jahred Adelman

ACKNOWLEDGEMENTS

DEDICATION

To Dr. Dhiman Chakraborty. Thank you for everything.

TABLE OF CONTENTS

Chapter	Page
1 Theory	1
1.1 The Standard Model	1
1.1.1 Particles	2
1.1.1.1 Fermions	2
1.1.1.2 Bosons	2
1.1.2 Interactions	3
1.1.2.1 Electromagnetic Interaction	3
1.1.2.2 Weak Interaction	3
1.1.2.3 Strong Interaction	4
1.1.3 The Higgs Mechanism	4
1.2 Supersymmetry	4
1.2.1 MSMM Particles	4
1.2.2 R-Parity	4
1.2.3 The MSSM Higgs Sector	4
1.3 Charged Higgs Bosons	5
1.3.1 Previous Result	5
2 The LHC and ATLAS Experiment	6
2.1 The Large Hadron Collider	6

Chapter	Page
2.2 The ATLAS Detector	6
2.2.1 Inner Detector	6
2.2.1.1 Pixel	6
2.2.1.2 Semiconductor Tracker.	7
2.2.1.3 Transition Radiation Tracker	7
2.2.2 Calorimeters.	7
2.2.2.1 Liquid Argon Electromagnetic Calorimeter	7
2.2.2.2 Tile Hadronic Calorimeter	7
2.2.3 Muon System	7
2.2.3.1 Monitored Drift Tubes.	8
2.2.3.2 Cathode Strip Chambers	8
2.2.3.3 Resistive Plate Chambers.	8
2.2.3.4 Thin Gap Chambers	8
2.2.4 Magnet Systems	8
2.2.4.1 Solenoid Magnet	8
2.2.4.2 Toroid Magnet	9
3 Event Reconstruction	10
3.1 Trigger	10
3.2 Inner Detector	10
3.3 Calorimeters	10
3.4 Muon	10
3.5 E Gamma.	10
3.6 Jets	10
3.6.1 Flavor Tagging	10

Chapter	Page
3.6.2 Tau	10
3.7 E_T^{miss}	10
4 Search for Charged Higgs Bosons	11
4.1 Signature and Event Selection	11
4.1.1 Object Definitions.	11
4.1.2 Event Selections	11
4.2 Datasets.	11
4.2.1 Signal Modeling	11
4.3 Background Modeling	12
4.4 Analysis Strategy	12
4.4.1 Multivariate Analysis Techniques	12
4.4.2 Training.	12
4.4.3 Feature Selection	12
4.4.4 Hyperparameter Optimization	12
4.5 Systematic Uncertainties	13
4.6 Results.	13
5 Conclusion	14

CHAPTER 1

THEORY

In this chapter, the theoretical motivation of a search for $H^\pm \rightarrow \tau^\pm \nu_\tau$ is described. Firstly, a review of the Standard Model of particle physics (SM) is laid out, then a brief overview of Supersymmetry focusing on the Minimal Supersymmetric Standard Model (MSSM) Higgs sector is detailed. Finally, the Type II 2-Higgs Doublet Model's (2HDM) relation to the H^\pm production cross section and subsequent branching ratio into SM particles is described as motivation for the choice of studying $H^\pm \rightarrow \tau^\pm \nu_\tau$.

1.1 The Standard Model

The Standard Model of particle physics is a mathematical model that describes all known matter and forces. The Standard Model is built upon a gauge group of type $SU(3)_C \times SU(2)_L \times U(1)_Y$. The $SU(3)_C$ term dictates the strong interaction while the $SU(2)_L \times U(1)_Y$ term describes the electroweak interaction. These interactions occur between fundamental particles called fermions that comprise the known matter of the universe. The interactions, or forces, are mediated by fundamental particles called bosons.

1.1.1 Particles

The particles that make up the Standard Model are separated into two groups according to their intrinsic angular momentum charge, or spin. Fermions are those that carry half-integer spin while Bosons carry full integer spin values.

Include standard model diagram

1.1.1.1 Fermions

Fermions are even further divided into two groups, quarks and leptons. The quarks participate in the strong interaction via their color charge. Quarks cannot exist as a singular particle and thus combine into hadrons. Leptons carry no color charge and therefore do not participate in strong force interactions. The fermions in the standard model all participate in the electroweak interaction. However, the electromagnetic interaction is limited to those fermions that carry an electromagnetic charge.

Fermions can then be split further into three generations, each lepton has a weak force partner in the form of a neutrino.

1.1.1.2 Bosons

NEEDS TO BE DONE

1.1.2 Interactions

At its core, the SM relies upon symmetries. From these symmetries, conservation laws follow. It is these laws of conservation, and the breaking of said symmetries, that dictate the allowed interactions of matter. The first, being a symmetry under charge conjugation, mirror reflection, and time reversal is known as CPT symmetry. The symmetry between charge conjugation and mirror reflection (CP) can be broken in certain circumstances, but holds in strong and electromagnetic interactions. This breaking of CP symmetry occurs in the weak interaction and implies a non-symmetry between matter and antimatter. Since this symmetry holds for strong and electromagnetic interactions, baryon number ($B = \frac{1}{3}(n_q - n_{\bar{q}})$) and lepton number are conserved in SM interactions. Lepton generation number ¹, electric charge, color charge, 4-momentum ($p = (E, \vec{p})$), and angular momentum are all conserved in the SM.

1.1.2.1 Electromagnetic Interaction

NEEDS TO BE DONE

1.1.2.2 Weak Interaction

NEEDS TO BE DONE

¹Ignoring neutrino oscillations

1.1.2.3 Strong Interaction

NEEDS TO BE DONE

1.1.3 The Higgs Mechanism

NEEDS TO BE DONE

1.2 Supersymmetry

NEEDS TO BE DONE

1.2.1 MSMM Particles

NEEDS TO BE DONE

1.2.2 R-Parity

NEEDS TO BE DONE

1.2.3 The MSSM Higgs Sector

NEEDS TO BE DONE

1.3 Charged Higgs Bosons

NEEDS TO BE DONE

1.3.1 Previous Result

NEEDS TO BE DONE

CHAPTER 2

THE LHC AND ATLAS EXPERIMENT

2.1 The Large Hadron Collider

In order to study the Standard Model, the Higgs boson, and hints of new physics, the Large Hadron Collider (LHC) was built outside of Geneva, Switzerland. At 27 km in circumference with a center of mass energy of 13.6 TeV, the LHC is the largest and highest energy particle accelerator ever built. It consists of **NUM SECTORS** magnet sectors split between dipole and quadrupole magnets.

2.2 The ATLAS Detector

NEEDS TO BE DONE [1]

2.2.1 Inner Detector

NEEDS TO BE DONE

2.2.1.1 Pixel

NEEDS TO BE DONE

2.2.1.2 Semiconductor Tracker

NEEDS TO BE DONE

2.2.1.3 Transition Radiation Tracker

NEEDS TO BE DONE

2.2.2 Calorimeters

NEEDS TO BE DONE

2.2.2.1 Liquid Argon Electromagnetic Calorimeter

NEEDS TO BE DONE

2.2.2.2 Tile Hadronic Calorimeter

NEEDS TO BE DONE

2.2.3 Muon System

NEEDS TO BE DONE

2.2.3.1 Monitored Drift Tubes

NEEDS TO BE DONE

2.2.3.2 Cathode Strip Chambers

NEEDS TO BE DONE

2.2.3.3 Resistive Plate Chambers

NEEDS TO BE DONE

2.2.3.4 Thin Gap Chambers

NEEDS TO BE DONE

2.2.4 Magnet Systems

NEEDS TO BE DONE

2.2.4.1 Solenoid Magnet

NEEDS TO BE DONE

2.2.4.2 Toroid Magnet

NEEDS TO BE DONE

CHAPTER 3

EVENT RECONSTRUCTION

3.1 Trigger

3.2 Inner Detector

3.3 Calorimeters

3.4 Muon

3.5 E Gamma

3.6 Jets

3.6.1 Flavor Tagging

3.6.2 Tau

3.7 $E_{\text{T}}^{\text{miss}}$

CHAPTER 4

SEARCH FOR CHARGED HIGGS BOSONS

4.1 Signature and Event Selection

NEEDS TO BE DONE

4.1.1 Object Definitions

NEEDS TO BE DONE

4.1.2 Event Selections

NEEDS TO BE DONE

4.2 Datasets

NEEDS TO BE DONE

4.2.1 Signal Modeling

NEEDS TO BE DONE

4.3 Background Modeling

NEEDS TO BE DONE

4.4 Analysis Strategy

NEEDS TO BE DONE

4.4.1 Multivariate Analysis Techniques

NEEDS TO BE DONE

4.4.2 Training

NEEDS TO BE DONE

4.4.3 Feature Selection

NEEDS TO BE DONE

4.4.4 Hyperparameter Optimization

NEEDS TO BE DONE

4.5 Systematic Uncertainties

NEEDS TO BE DONE

4.6 Results

CHAPTER 5
CONCLUSION

Appendices

BIBLIOGRAPHY

[1] The ATLAS Collaboration.

“The ATLAS Experiment at the CERN Large Hadron Collider”.

Journal of Instrumentation 3.08 (Aug. 2008), S08003–S08003.

DOI: 10.1088/1748-0221/3/08/s08003.

URL: <https://doi.org/10.1088/1748-0221/3/08/s08003>.