

MA_vax: vaccination-impact analysis report

Massachusetts, 2025–2026 influenza season. Age-structured SEIR model with a parallel vaccinated arm, calibrated to daily hospital admissions by age group via Bayesian MCMC. This report documents the model, the fit, and the resulting vaccination-impact analysis.

Every count in this report is a **hospital admission over the whole 250-day season** unless stated otherwise. Tables report the absolute number of hospitalizations averted alongside the percentage and the per-100,000 rates, so the size of the burden each figure describes is visible next to the rate.


1. Model structure

1.1 Compartments

Nine compartments, each stratified by 7 age groups (0, 1–4, 5–12, 13–17, 18–49, 50–64, 65+):

Compartment	Meaning
S	Susceptible, unvaccinated
E	Exposed (latent), unvaccinated
I	Infectious, unvaccinated
R	Recovered/removed (either arm)
SV	Susceptible, vaccinated
EV	Exposed (latent), vaccinated
IV	Infectious, vaccinated
H	Hospitalized (either arm)
D	Dead (either arm)

The vaccinated arm (SV → EV → IV → H/R) mirrors the unvaccinated arm (S → E → I → H/R), with its own susceptibility and severity parameters. H, R, D are shared, pooled compartments — once hospitalized (or recovered), an individual's vaccination history is no longer tracked.



1.2 Force of infection

Contacts come from fixed age×age contact matrices (Mistry et al. 2021 synthetic contact matrices), with school/work contacts removed on non-school/work days:

```

C(t) = total_C - (1 - is_school(t))·school_C - (1 - is_work(t))·work_C
beta_adj(t) = beta_baseline · m(t) · (1 + humidity_impact · exp(-180 · humidity(t)))
wtd_inf_prop(t) = (I·I_relative_infectiousness + IV·IV_relative_infectiousness) / population
foi(t) = beta_adj(t) · (C(t) @ wtd_inf_prop(t))
S_to_E = foi(t) · relative_suscept · S
SV_to_EV = foi(t) · vax_susceptibility · SV
  
```

$m(t)$ is a smoothly time-varying transmission multiplier (§2.3) that absorbs behavioral/seasonal variation the mechanistic terms above don't otherwise capture. $IV_relative_infectiousness = 1.0$ — breakthrough infections (in vaccinated individuals) are assumed exactly as transmissible as unvaccinated infections. $vax_susceptibility$ (age-specific, ≤ 1) is the residual susceptibility of a vaccinated individual; $1 - vax_susceptibility$ is the model's implied vaccine effectiveness (VE) against infection.

1.3 Progression, hospitalization, death

$$\begin{aligned} E_to_I &= E_to_I_rate \cdot E & EV_to_IV &= EV_to_IV_rate \cdot EV \\ I_to_H &= I_out_rate \cdot I_to_H_prop \cdot I & IV_to_H &= I_out_rate \cdot IV_to_H_prop \cdot IV \\ I_to_R &= I_out_rate \cdot (1-I_to_H_prop) \cdot I & IV_to_R &= I_out_rate \cdot (1-IV_to_H_prop) \cdot IV \\ H_to_D &= H_out_rate \cdot H_to_D_prop \cdot H \\ H_to_R &= H_out_rate \cdot (1-H_to_D_prop) \cdot H \end{aligned}$$

$IV_to_H_prop < I_to_H_prop$ for every age group — vaccination reduces hospitalization risk *given* infection (severity VE), on top of reducing infection risk.

1.4 Vaccination flow

Daily doses (an age-specific proportion of the population, with a 14-day delay between dose and effective immunity) are applied as an exact $S \rightarrow SV$ count each day, capped at whatever remains in S :

$$\begin{aligned} base(t) &= S + SV \\ S_to_SV(t) &= \min(\text{round}(vax_prop(t) \cdot base(t)), S) \end{aligned}$$

The base pool ($S + SV$) is not eroded by vaccination itself — only infection depletes it — so a roughly flat input proportion vaccinates a roughly constant head-count per day, until S starts running low late in the season.

1.5 Numerical scheme

Simulations use explicit Euler integration with 7 sub-steps per day. The ensemble behind every interval in this report is a **posterior parameter ensemble**: one deterministic simulation per accepted posterior draw (§2.3), not chain-binomial transition noise. Vaccination is a deterministic scheduled count in either case, since it comes from an external delivery schedule rather than a hazard rate.

1.6 Initial conditions

At the start of the simulation (2025-09-01): $S = population - E_0$, $E = E_0$ (age-specific seed counts, scaled by a single fitted multiplier, §2.3), and all other compartments start at zero.

2. Parameters

2.1 Fixed scalar parameters

Parameter	Value	Meaning
num_days	250	Simulation length (2025-09-01 through 2026-05-08)
$relative_suscept$	1.0	Susceptibility multiplier, unvaccinated arm
$I_relative_infectiousness$	1.0	Infectiousness weight, unvaccinated I
$IV_relative_infectiousness$	1.0	Infectiousness weight, vaccinated IV
$E_to_I_rate$	0.5 /day	~2-day latent period
$EV_to_IV_rate$	0.5 /day	Same, vaccinated arm
I_out_rate	0.333 /day	~3-day infectious period

Parameter	Value	Meaning
<code>H_out_rate</code>	0.17 /day	~6-day hospital stay
<code>vax_transfer_delay_days</code>	14	Days from dose to modeled immunity

2.2 Age-stratified fixed parameters

Age group	Population	Hospitalization risk, given infection ¹	Hospitalization risk, given breakthrough infection ¹	Death risk, given hospitalization	Residual susceptibility if vaccinated	Initial infections seeded ¹
0	70,067	0.697%	0.636%	1.74%	0.57	2
1-4	280,268	0.697%	0.636%	1.74%	0.57	8
5-12	606,291	0.274%	0.250%	1.17%	0.57	17
13-17	411,782	0.274%	0.250%	1.17%	0.57	12
18-49	2,978,204	0.561%	0.511%	2.63%	0.79	85
50-64	1,424,434	1.060%	0.966%	6.30%	0.79	41
65+	1,221,349	9.091%	6.273%	7.99%	1.00	35

¹ These are pre-fit baseline values — the calibration scales both hospitalization-risk columns by a fitted age-specific multiplier (§2.3), so the values actually used in the calibrated model are these figures × that multiplier and the vaccine effectiveness against hospitalization remains the same. A residual susceptibility of 1.00 for 65+ means the fitted baseline assumes **no infection-blocking effect** of vaccination in that age group (only the severity effect applies there). This holds for the **Low VE** sensitivity scenario too, but not for **High VE**, which scales it to 0.86 — see Table S.A.4.

2.3 Fitting method

Free parameters were estimated with an affine-invariant ensemble Markov Chain Monte Carlo sampler (40 walkers, 4000 iterations per walker), run in parallel over the full 251-day target window. The free parameters and their priors:

Parameter	Prior	Role
<code>beta_baseline</code>	Uniform(0.015, 0.06)	Baseline transmission rate
<code>humidity_impact</code>	Uniform(0, 1)	Strength of humidity forcing
Initial-seed multiplier	Log-uniform(0.1, 10)	Multiplies the age-specific initial-infection seed counts
Hospitalization-risk multiplier (× 7, one per age group)	Uniform(0.1, 2.0)	Multiplies both hospitalization-risk columns (§2.2) for that age group
<code>m(t)</code> log-increments (× 18, one per 14-day knot)	Normal(0, 0.25)	Month-to-month-ish random-walk steps in log-transmission (§1.2)
<code>phi</code>	—	Negative-Binomial dispersion (likelihood nuisance parameter, not a model input)

Likelihood: a Negative-Binomial (NB2) observation model, chosen over Poisson to allow overdispersion, jointly across 8 targets — daily hospital admissions **by age group** (7 time series) plus a single scalar **end-of-season cumulative hospitalizations by age** target. The targets are not weighted equally: the three age groups carrying most of the burden are up-weighted (18-49 and 50-64 at weight 2, 65+ at weight 5, the four younger groups at 1), and the end-of-season cumulative target at weight 10, so the fit is pulled towards reproducing the season's total burden and its concentration in older ages rather than treating every age curve alike.

Posterior sampling: the first 1700 iterations were discarded as burn-in, and the remaining chain thinned to every 200th sample, leaving 627 posterior draws. Two point estimates are used in this report:

- **Posterior mean:** the marginal mean of each parameter. Cheap and stable, but for a correlated posterior can land on a parameter *combination* the sampler never actually visited.
- **"Best" point:** the single posterior draw with the highest log-posterior — not the mean of anything, so it preserves whatever correlation structure exists between parameters.

These two can disagree materially (§2.4) — a reminder that the marginal mean of a correlated posterior needn't correspond to a jointly plausible parameter combination. $m(t)$'s random-walk prior is a smoothness assumption, not a mechanistic one — it should not be over-interpreted as an independently-measured behavioral signal, only as "how much transmission needs to have moved, beyond what humidity/contacts explain, to reconcile the model with the data."

2.4 Fitted parameters

Posterior mean $\pm$ 90% credible interval (5th–95th percentile) across the 627 posterior draws, vs. the "best" (highest-posterior-density) point:

Parameter	Posterior mean	5%	95%	"Best" point
β_{baseline}	0.0305	0.0261	0.0359	0.0281
humidity_impact	0.591	0.234	0.916	0.881
Initial-seed multiplier	1.70	0.47	3.39	1.47
Hospitalization-risk multiplier — 0	1.51	0.95	1.93	1.63
Hospitalization-risk multiplier — 1-4	1.51	0.95	1.94	1.56
Hospitalization-risk multiplier — 5-12	1.18	0.78	1.63	1.22
Hospitalization-risk multiplier — 13-17	1.03	0.62	1.48	0.99
Hospitalization-risk multiplier — 18-49	0.59	0.39	0.81	0.51
Hospitalization-risk multiplier — 50-64	0.78	0.51	1.06	0.79
Hospitalization-risk multiplier — 65+	1.06	0.73	1.40	1.00
ϕ (NB dispersion)	78	18	223	344

The 18 $m(t)$ log-increments aren't independently interpretable as a table — they're summarized visually through the fit check in §4 instead.

2.5 Cumulative vaccination coverage

Cumulative proportion of each age group actually vaccinated over the season (reported vaccination-schedule data, summed over the season):

Age group	Population	Cumulative coverage
0	70,067	24.1%
1-4	280,268	48.3%
5-12	606,291	36.2%
13-17	411,782	36.5%
18-49	2,978,204	28.1%
50-64	1,424,434	38.2%
65+	1,221,349	60.6%
All (population-weighted)	6,992,395	37.8%

Coverage is well below a 70% mark in **every** age group: 65+ comes closest at 60.6%, then 1-4 (48.3%), 50-64 (38.2%), 13-17 (36.5%) and 5-12 (36.2%), with 18-49 (28.1%) and age 0 (24.1%) lowest. This matters for how the "scale to 70% coverage" rows in the appendix (Tables S.A.3/S.A.6) should be read: unlike in earlier versions of this analysis, all seven groups are scaled up, so no column is structurally zero — though the 65+ column still comes close, because vaccinating 65+ blocks no infection in the fitted baseline (§2.2) and so only helps 65+ itself.

The figures above are what the vaccination **schedule** reports. The doses that actually land are fewer — the §1.4 cap declines to vaccinate someone who has already been infected — putting simulated coverage 0.2 to 1.6 percentage points lower per age group. Those undelivered doses are not saved: in the real world they are still bought and administered, into arms that no longer benefit, at least in our model where the recovered and deceased compartments are final. The appendix reconciles the two figures age group by age group, and every "per 100,000 doses" panel in this report divides by the **scheduled** count for that reason.


3. Vaccination-impact results

Every table in this section reports a **median and 95% interval across 627 simulations** — one simulation per posterior parameter draw (§2.3), re-using the same draw for every scenario being compared within a table so that the comparison is paired (variance from the parameter draw itself cancels out of the *difference* between scenarios). These intervals therefore reflect **calibration uncertainty** — how much the vaccination- impact conclusions would change under a different (but similarly plausible) fit to the same data — not day-to-day epidemic-process noise.

For scale: over the season the fitted baseline produces a median of **6,070 hospitalizations** across all ages, against **13,370** with no vaccination at all.

New daily hospitalizations: baseline vs. no vaccination

Total-population new hospitalizations per day, posterior median and 95% interval across the 627 parameter draws (§2.3, same posterior-uncertainty basis as the rest of this section), comparing the fitted baseline vaccination schedule against a counterfactual with no vaccination at all:



The fitted vaccination program cuts the peak in daily new hospitalizations to roughly **45%** of what it would otherwise be (median peak ≈ 165 /day baseline vs. ≈ 365 /day with no vaccination, both around 2025-12-28) and reduces the epidemic's height without changing its timing.

Table S.A.1 — Hospitalizations averted, infection vs. severity protection

Decomposes the total hospitalizations averted by vaccination into two channels: protection against getting infected at all, and — *given* a breakthrough infection still happens — protection against that infection becoming severe enough to need hospitalization.

Infection protection (no vaccination → infection-protection-only, i.e. VE against infection retained but VE against severity zeroed out)

Age group	Hospitalizations Averted	% Hospitalizations Averted	Averted per 100,000 Population	Averted per 100,000 Doses
0	40 [24 – 52]	52.7% [48.2% – 55.3%]	57.1 [34.2 – 74.3]	236.6 [141.8 – 307.6]
1-4	208 [140 – 261]	56.9% [52.5% – 59.5%]	74.3 [49.9 – 93.2]	153.9 [103.3 – 193.0]
5-12	188 [126 – 251]	54.7% [49.4% – 57.7%]	31.0 [20.9 – 41.5]	85.6 [57.5 – 114.4]
13-17	115 [74 – 158]	54.7% [49.5% – 57.8%]	27.8 [17.9 – 38.4]	76.3 [49.0 – 105.2]
18-49	743 [548 – 950]	49.8% [44.6% – 52.7%]	24.9 [18.4 – 31.9]	88.8 [65.5 – 113.5]
50-64	792 [584 – 981]	50.7% [45.9% – 53.5%]	55.6 [41.0 – 68.9]	145.6 [107.4 – 180.4]
65+	4,433 [3,698 – 5,162]	48.1% [43.5% – 50.9%]	362.9 [302.8 – 422.6]	598.9 [499.6 – 697.4]
All	6,583 [5,440 – 7,404]	49.2% [44.4% – 52.0%]	94.1 [77.8 – 105.9]	249.0 [205.8 – 280.1]

Severity protection (infection-protection-only → full baseline, i.e. adding back VE against severity)

Age group	Hospitalizations Averted	% Hospitalizations Averted	Averted per 100,000 Population	Averted per 100,000 Doses
0	0 [0 – 1]	0.5% [0.5% – 0.5%]	0.5 [0.3 – 0.7]	2.2 [1.4 – 3.0]
1-4	4 [2 – 5]	1.0% [0.9% – 1.1%]	1.3 [0.8 – 1.7]	2.6 [1.8 – 3.4]
5-12	3 [2 – 3]	0.7% [0.7% – 0.8%]	0.4 [0.3 – 0.5]	1.1 [0.8 – 1.5]
13-17	2 [1 – 2]	0.8% [0.7% – 0.9%]	0.4 [0.3 – 0.5]	1.1 [0.7 – 1.4]
18-49	11 [9 – 14]	0.8% [0.7% – 0.8%]	0.4 [0.3 – 0.5]	1.4 [1.1 – 1.6]
50-64	16 [13 – 20]	1.1% [1.0% – 1.2%]	1.2 [0.9 – 1.4]	3.0 [2.4 – 3.6]
65+	720 [647 – 801]	7.7% [7.3% – 8.4%]	59.0 [53.0 – 65.6]	97.3 [87.4 – 108.2]
All	756 [681 – 837]	5.6% [5.2% – 6.2%]	10.8 [9.7 – 12.0]	28.6 [25.8 – 31.7]

Total (no vaccination → full baseline)

Age group	Hospitalizations Averted	% Hospitalizations Averted	Averted per 100,000 Population	Averted per 100,000 Doses
0	40 [24 – 53]	53.2% [48.7% – 55.8%]	57.7 [34.6 – 75.0]	238.7 [143.2 – 310.6]
1-4	212 [142 – 266]	57.9% [53.5% – 60.4%]	75.6 [50.8 – 94.9]	156.5 [105.1 – 196.5]
5-12	191 [128 – 255]	55.4% [50.2% – 58.4%]	31.4 [21.1 – 42.0]	86.8 [58.3 – 115.8]
13-17	116 [75 – 160]	55.5% [50.3% – 58.5%]	28.2 [18.2 – 38.9]	77.4 [49.9 – 106.6]
18-49	754 [558 – 963]	50.5% [45.5% – 53.4%]	25.3 [18.8 – 32.3]	90.1 [66.7 – 115.1]
50-64	809 [599 – 1,000]	51.8% [47.0% – 54.5%]	56.8 [42.0 – 70.2]	148.7 [110.0 – 183.9]
65+	5,147 [4,398 – 5,959]	55.9% [51.8% – 58.2%]	421.4 [360.1 – 487.9]	695.4 [594.3 – 805.2]
All	7,327 [6,169 – 8,190]	54.8% [50.5% – 57.4%]	104.8 [88.2 – 117.1]	277.2 [233.4 – 309.9]

Almost all of the averted burden comes from **blocking infection**, not from reducing severity given a breakthrough: of the 7,327 hospitalizations averted in total, 6,583 come from the infection channel and 756 from the severity channel — and 720 of those 756 are in 65+, the one group where the fitted model gives vaccination no infection-blocking effect at all.

Table S.A.2 — Hospitalizations averted by age group vaccinated

Each column vaccinates a single age group only (all others left unvaccinated) and compares to no vaccination at all; "All" is the full baseline schedule. Rows are the age group in which hospitalizations are counted, so off-diagonal cells show the indirect (transmission-blocking) benefit to *other* age groups from vaccinating this one.

Hospitalizations averted (count)

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	7 [4 – 9]	4 [3 – 5]	12 [7 – 16]	9 [5 – 12]	12 [7 – 15]	5 [3 – 7]	0 [-0 – 0]	40 [24 – 53]
1-4	1 [1 – 1]	83 [56 – 104]	63 [41 – 79]	43 [27 – 53]	53 [35 – 67]	24 [16 – 31]	0 [-0 – 0]	212 [142 – 266]
5-12	1 [1 – 1]	17 [11 – 23]	102 [69 – 137]	41 [26 – 55]	46 [29 – 62]	21 [14 – 29]	0 [-0 – 0]	191 [128 – 255]
13-17	0 [0 – 1]	9 [5 – 13]	31 [19 – 46]	57 [37 – 80]	28 [17 – 40]	14 [8 – 20]	0 [-0 – 0]	116 [75 – 160]
18-49	4 [3 – 6]	72 [50 – 94]	226 [157 – 298]	178 [123 – 235]	295 [214 – 380]	109 [76 – 143]	0 [-0 – 0]	754 [558 – 963]
50-64	4 [3 – 5]	71 [49 – 91]	232 [159 – 297]	187 [129 – 241]	232 [162 – 294]	241 [179 – 299]	0 [-0 – 0]	809 [599 – 1,000]
65+	26 [20 – 31]	445 [353 – 535]	1,437 [1,134 – 1,728]	1,137 [896 – 1,373]	1,382 [1,099 – 1,645]	740 [592 – 881]	1,404 [1,245 – 1,591]	5,147 [4,398 – 5,959]
All	44 [35 – 50]	708 [556 – 807]	2,125 [1,648 – 2,465]	1,668 [1,288 – 1,950]	2,065 [1,623 – 2,396]	1,161 [920 – 1,338]	1,404 [1,245 – 1,591]	7,327 [6,169 – 8,190]

% reduction in hospitalizations

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	9.2% [9.0% – 9.4%]	5.6% [4.8% – 6.1%]	16.2% [13.8% – 17.8%]	12.0% [10.1% – 13.3%]	15.4% [13.2% – 16.9%]	7.0% [5.9% – 7.7%]	0.0% [-0.0% – 0.0%]	53.2% [48.7% – 55.8%]
1-4	0.3% [0.3% – 0.3%]	22.6% [21.5% – 23.4%]	17.3% [14.7% – 19.1%]	11.7% [9.7% – 13.1%]	14.6% [12.3% – 16.2%]	6.7% [5.6% – 7.5%]	0.0% [-0.0% – 0.0%]	57.9% [53.5% – 60.4%]
5-12	0.3% [0.2% – 0.3%]	4.9% [4.1% – 5.5%]	29.7% [26.8% – 31.7%]	11.7% [9.6% – 13.3%]	13.1% [10.7% – 14.9%]	6.2% [5.0% – 7.1%]	0.0% [-0.0% – 0.0%]	55.4% [50.2% – 58.4%]
13-17	0.2% [0.2% – 0.3%]	4.2% [3.4% – 4.8%]	15.0% [12.3% – 16.9%]	27.4% [25.0% – 29.1%]	13.2% [10.7% – 15.0%]	6.5% [5.2% – 7.4%]	0.0% [-0.0% – 0.0%]	55.5% [50.3% – 58.5%]

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
18-49	0.3% [0.2% – 0.3%]	4.8% [4.0% – 5.3%]	15.1% [12.7% – 16.9%]	11.9% [9.9% – 13.3%]	19.7% [17.4% – 21.3%]	7.3% [6.1% – 8.1%]	0.0% [-0.0% – 0.0%]	50.5% [45.5% – 53.4%]
50-64	0.3% [0.2% – 0.3%]	4.5% [3.8% – 5.1%]	14.9% [12.5% – 16.6%]	12.0% [10.1% – 13.4%]	14.9% [12.6% – 16.4%]	15.4% [14.1% – 16.3%]	0.0% [-0.0% – 0.0%]	51.8% [47.0% – 54.5%]
65+	0.3% [0.2% – 0.3%]	4.8% [4.1% – 5.3%]	15.5% [13.3% – 17.1%]	12.3% [10.5% – 13.6%]	15.0% [12.8% – 16.4%]	8.0% [6.9% – 8.8%]	15.2% [15.0% – 15.3%]	55.9% [51.8% – 58.2%]
All	0.3% [0.3% – 0.4%]	5.3% [4.5% – 5.7%]	15.8% [13.5% – 17.4%]	12.4% [10.5% – 13.7%]	15.4% [13.3% – 16.9%]	8.7% [7.5% – 9.5%]	10.5% [10.0% – 11.1%]	54.8% [50.5% – 57.4%]

Hospitalizations averted per 100,000 population

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	10.0 [6.1 – 13.2]	6.0 [3.6 – 7.8]	17.5 [10.3 – 22.5]	13.0 [7.7 – 16.6]	16.7 [9.9 – 21.5]	7.5 [4.5 – 9.7]	0.0 [-0.0 – 0.0]	57.7 [34.6 – 75.0]
1-4	0.4 [0.3 – 0.5]	29.5 [19.9 – 37.2]	22.6 [14.6 – 28.2]	15.2 [9.6 – 19.0]	19.0 [12.3 – 23.9]	8.7 [5.6 – 11.0]	0.0 [-0.0 – 0.0]	75.6 [50.8 – 94.9]
5-12	0.1 [0.1 – 0.2]	2.8 [1.8 – 3.8]	16.9 [11.3 – 22.6]	6.7 [4.3 – 9.1]	7.5 [4.8 – 10.2]	3.5 [2.3 – 4.8]	0.0 [-0.0 – 0.0]	31.4 [21.1 – 42.0]
13-17	0.1 [0.1 – 0.2]	2.1 [1.3 – 3.1]	7.6 [4.5 – 11.1]	14.0 [9.0 – 19.3]	6.7 [4.0 – 9.8]	3.3 [2.0 – 4.8]	0.0 [-0.0 – 0.0]	28.2 [18.2 – 38.9]
18-49	0.1 [0.1 – 0.2]	2.4 [1.7 – 3.2]	7.6 [5.3 – 10.0]	6.0 [4.1 – 7.9]	9.9 [7.2 – 12.8]	3.7 [2.5 – 4.8]	0.0 [-0.0 – 0.0]	25.3 [18.8 – 32.3]
50-64	0.3 [0.2 – 0.4]	5.0 [3.4 – 6.4]	16.3 [11.2 – 20.9]	13.2 [9.0 – 16.9]	16.3 [11.4 – 20.7]	16.9 [12.5 – 21.0]	0.0 [-0.0 – 0.0]	56.8 [42.0 – 70.2]
65+	2.1 [1.7 – 2.5]	36.4 [28.9 – 43.8]	117.7 [92.8 – 141.5]	93.1 [73.4 – 112.4]	113.1 [90.0 – 134.7]	60.6 [48.5 – 72.1]	114.9 [101.9 – 130.3]	421.4 [360.1 – 487.9]
All	0.6 [0.5 – 0.7]	10.1 [7.9 – 11.5]	30.4 [23.6 – 35.3]	23.9 [18.4 – 27.9]	29.5 [23.2 – 34.3]	16.6 [13.2 – 19.1]	20.1 [17.8 – 22.8]	104.8 [88.2 – 117.1]

Hospitalizations averted per 100,000 doses

Each cell divides hospitalizations averted in the **row's** age group by the doses scheduled for the **column's** age group, so off-diagonal cells are the indirect benefit per dose spent on the vaccinated group. The **All vaccinated** column has no single targeted group, so there each row keeps its own age group's dose count, matching Table S.A.1's per-dose columns.

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	41.4 [25.2 – 54.7]	3.1 [1.8 – 4.0]	5.6 [3.3 – 7.2]	6.0 [3.6 – 7.8]	1.4 [0.8 – 1.8]	1.0 [0.6 – 1.3]	0.0 [-0.0 – 0.0]	238.7 [143.2 – 310.6]
1-4	6.7 [4.4 – 8.5]	61.2 [41.2 – 77.0]	28.8 [18.6 – 36.0]	28.4 [17.9 – 35.5]	6.4 [4.1 – 8.0]	4.5 [2.9 – 5.6]	0.0 [-0.0 – 0.0]	156.5 [105.1 – 196.5]
5-12	5.2 [3.3 – 7.0]	12.7 [8.1 – 17.1]	46.6 [31.3 – 62.2]	27.2 [17.4 – 36.7]	5.5 [3.5 – 7.4]	4.0 [2.5 – 5.4]	0.0 [-0.0 – 0.0]	86.8 [58.3 – 115.8]
13-17	2.9 [1.7 – 4.3]	6.4 [3.8 – 9.5]	14.3 [8.5 – 20.7]	38.2 [24.7 – 53.0]	3.3 [2.0 – 4.8]	2.5 [1.5 – 3.7]	0.0 [-0.0 – 0.0]	77.4 [49.9 – 106.6]
18-49	24.9 [17.4 – 32.7]	52.8 [36.8 – 69.5]	103.1 [71.3 – 135.5]	118.5 [81.7 – 156.2]	35.3 [25.5 – 45.4]	20.0 [13.9 – 26.3]	0.0 [-0.0 – 0.0]	90.1 [66.7 – 115.1]
50-64	23.8 [16.5 – 30.6]	52.3 [35.9 – 66.9]	105.6 [72.4 – 135.3]	124.7 [85.5 – 160.2]	27.7 [19.3 – 35.2]	44.3 [32.9 – 54.9]	0.0 [-0.0 – 0.0]	148.7 [110.0 – 183.9]
65+	151.5 [120.2 – 181.7]	328.8 [260.9 – 394.9]	654.1 [515.9 – 786.4]	756.9 [596.4 – 913.8]	165.1 [131.3 – 196.6]	136.0 [108.8 – 162.0]	189.6 [168.2 – 215.0]	695.4 [594.3 – 805.2]
All	258.8 [204.7 – 295.1]	522.7 [410.5 – 596.4]	967.2 [749.8 – 1121.8]	1109.9 [857.3 – 1297.4]	246.8 [193.9 – 286.3]	213.4 [169.2 – 245.9]	189.6 [168.2 – 215.0]	277.2 [233.4 – 309.9]

Off-diagonal entries confirm real indirect effects — e.g. vaccinating 5-12 alone reduces hospitalizations in 0 by 16.2% and in 1-4 by 17.3%, both larger than several of those groups' own-age direct effects, consistent with school-age children acting as a major transmission hub in the contact structure. 65+ is the only group with zero indirect effect on every other group.

In absolute terms the indirect benefit dwarfs the direct one for the school-age columns: vaccinating 5-12 alone averts 2,125 hospitalizations across the population, only 102 of them in 5-12 itself and 1,437 of them in 65+.

Per dose, the same story: vaccinating 13-17 averts 756.9 [596.4 – 913.8] hospitalizations per 100,000 doses in the 65+ group alone, against 38.2 [24.7 – 53.0] in 13-17 itself. This is why the **All** row — hospitalizations averted across the whole population per dose — ranks 13-17 (1109.9) and 5-12 (967.2) far above the groups that carry the burden directly.

Table S.A.4 — Vaccine-effectiveness sensitivity scenarios

Implied vaccine effectiveness under three illustrative VE presets — a parameter table, not a simulation result. **Baseline VE (fitted)** is the model's fitted VE (included here as the reference point, not itself a sensitivity scenario); **Low VE/High VE** bracket it and correspond to the estimated lower and upper bound of VE. The ratio of low/high VE to baseline remains constant across parameter sets.

Scenario	Age group	VE against infection	VE against hospitalization (overall)	VE against hospitalization, given infection
Low VE	0	33%	33%	0%
Low VE	1-4	33%	33%	0%
Low VE	5-12	33%	33%	0%
Low VE	13-17	33%	33%	0%
Low VE	18-49	6%	10%	4%

Scenario	Age group	VE against infection	VE against hospitalization (overall)	VE against hospitalization, given infection
Low VE	50-64	6%	10%	4%
Low VE	65+	0%	21%	21%
Baseline VE (fitted)	0	43%	48%	9%
Baseline VE (fitted)	1-4	43%	48%	9%
Baseline VE (fitted)	5-12	43%	48%	9%
Baseline VE (fitted)	13-17	43%	48%	9%
Baseline VE (fitted)	18-49	21%	28%	9%
Baseline VE (fitted)	50-64	21%	28%	9%
Baseline VE (fitted)	65+	0%	31%	31%
High VE	0	51%	78%	55%
High VE	1-4	51%	78%	55%
High VE	5-12	51%	78%	55%
High VE	13-17	51%	78%	55%
High VE	18-49	34%	44%	16%
High VE	50-64	34%	44%	16%
High VE	65+	14%	39%	29%

Also note **65+ has 0% VE against infection in the fitted baseline and in Low VE** — residual susceptibility is 1.00 for that age group in both (§2.2), so only the severity-protection channel operates there. **High VE** is the exception: it scales residual susceptibility to 0.86, giving 65+ a 14% VE against infection, so that scenario is the only one in which vaccinating 65+ blocks any transmission at all. This is why 65+ shows an all-zero indirect effect on every other age group throughout Table S.A.2, which is built on the fitted baseline.

Table S.A.5 — Hospitalizations averted across VE scenarios

Compares each VE sensitivity scenario (§ above) against no vaccination at all.

Hospitalizations averted (count)

Age group	Low VE	Baseline VE (fitted)	High VE
0	27 [16 – 34]	40 [24 – 53]	51 [31 – 66]
1-4	145 [97 – 181]	212 [142 – 266]	264 [179 – 334]
5-12	130 [87 – 174]	191 [128 – 255]	239 [162 – 316]
13-17	79 [49 – 111]	116 [75 – 160]	146 [94 – 198]
18-49	464 [331 – 599]	754 [558 – 963]	952 [719 – 1,211]
50-64	490 [349 – 616]	809 [599 – 1,000]	1,024 [777 – 1,256]

Age group	Low VE	Baseline VE (fitted)	High VE
65+	3,435 [2,876 – 4,002]	5,147 [4,398 – 5,959]	6,325 [5,483 – 7,281]
All	4,810 [3,959 – 5,471]	7,327 [6,169 – 8,190]	9,050 [7,766 – 10,049]

% reduction in hospitalizations

Age group	Low VE	Baseline VE (fitted)	High VE
0	35.0% [31.2% – 37.4%]	53.2% [48.7% – 55.8%]	66.8% [63.0% – 69.1%]
1-4	39.7% [35.8% – 42.1%]	57.9% [53.5% – 60.4%]	72.4% [68.9% – 74.4%]
5-12	37.8% [33.3% – 40.7%]	55.4% [50.2% – 58.4%]	69.0% [64.7% – 71.6%]
13-17	37.7% [33.2% – 40.6%]	55.5% [50.3% – 58.5%]	69.4% [65.1% – 71.9%]
18-49	31.0% [26.8% – 33.7%]	50.5% [45.5% – 53.4%]	63.9% [59.3% – 66.5%]
50-64	31.4% [27.4% – 34.0%]	51.8% [47.0% – 54.5%]	65.5% [61.3% – 67.9%]
65+	37.3% [33.9% – 39.5%]	55.9% [51.8% – 58.2%]	68.6% [65.1% – 70.7%]
All	36.0% [32.3% – 38.3%]	54.8% [50.5% – 57.4%]	67.8% [64.1% – 70.0%]

Hospitalizations averted per 100,000 population

Age group	Low VE	Baseline VE (fitted)	High VE
0	38.1 [22.5 – 49.0]	57.7 [34.6 – 75.0]	72.4 [43.8 – 94.7]
1-4	51.8 [34.5 – 64.6]	75.6 [50.8 – 94.9]	94.3 [63.8 – 119.2]
5-12	21.5 [14.3 – 28.7]	31.4 [21.1 – 42.0]	39.4 [26.7 – 52.1]
13-17	19.1 [12.0 – 26.9]	28.2 [18.2 – 38.9]	35.5 [22.9 – 48.1]
18-49	15.6 [11.1 – 20.1]	25.3 [18.8 – 32.3]	32.0 [24.1 – 40.7]
50-64	34.4 [24.5 – 43.2]	56.8 [42.0 – 70.2]	71.9 [54.6 – 88.2]
65+	281.3 [235.5 – 327.7]	421.4 [360.1 – 487.9]	517.8 [448.9 – 596.1]
All	68.8 [56.6 – 78.2]	104.8 [88.2 – 117.1]	129.4 [111.1 – 143.7]

Even under the pessimistic **Low VE** assumption, the current vaccination schedule still averts ~36% of hospitalizations overall (4,810 admissions) — against ~55% (7,327) at fitted VE and ~68% (9,050) at high VE. The schedule's *coverage* (§2.5) matters roughly as much as the assumed per-dose effectiveness across this VE range.

4. Baseline fit check — posterior-uncertainty simulation vs. raw data

This section reads the baseline (fitted-vaccination) scenario from the posterior parameter ensemble — all 627 draws (§2.3) — and reports the median and 95% interval across draws, to check the calibrated model against the raw data it was fit to.

Cumulative hospitalizations, by age group


Sum of simulated (posterior median and 95% interval) vs. raw daily hospital admissions, over the range of dates common to both series (2025-09-01 – 2026-05-08):

Age group	Simulated (median)	Simulated 95% interval	Raw data	% difference (median)
0	35.7	22.5 – 48.5	41.1	-13.0%
1-4	153.5	103.8 – 200.6	164.6	-6.8%
5-12	156.4	110.6 – 203.8	160.5	-2.6%


Age group	Simulated (median)	Simulated 95% interval	Raw data	% difference (median)
13-17	95.1	63.0 – 122.7	90.3	+5.3%
18-49	746.3	581.6 – 894.8	726.0	+2.8%
50-64	753.2	594.6 – 911.0	749.6	+0.5%
65+	4,124.8	3,699.5 – 4,554.8	4,222.3	-2.3%
All	6,070.0	5,569.0 – 6,549.0	6,154.4	-1.4%

The fit tracks the data closely overall (-1.4% on the total, raw value within the 95% interval for every age group) — the largest relative miss is in the smallest group (age 0, -13.0%, but only ~5 admissions off in absolute terms).

Daily new hospitalizations by age group



Cumulative hospitalizations by age group



The posterior interval captures the single main epidemic wave (peaking around late December/early January) well across all age groups.

5. Data sources

Input	Source
Age-specific vaccination coverage	MIDAS Flu Scenario Modeling Hub resources, age-specific coverage dataset
Hospital admissions (calibration target)	MIDAS Flu Scenario Modeling Hub, target-data time series
Population by age group	US Census (tidycensus)
Contact matrices	Mistry et al. 2021 synthetic contact matrices

Input	Source
Absolute humidity	gridMET daily specific-humidity data, averaged over Massachusetts
School/work calendar	Constructed school/work-day calendar for the state, for each day of the season

6. Notes

- **The confidence intervals in §3 reflect calibration uncertainty, not epidemic-process noise** — they come from re-running each scenario under 627 different (but similarly plausible) posterior parameter draws, each with deterministic transitions.
- **$m(t)$ is a statistical smoothing device, not a mechanistic term** — the 14-day-knot random walk in this fit absorbs whatever transmission variation the mechanistic model (contacts, humidity) doesn't explain; it should not be read as an independently-measured behavioral signal.
- **The "scale to 70% coverage" scenarios (Tables S.A.3/S.A.6, appendix) now raise every age group** — at the current schedule no age group reaches 70% (§2.5), so unlike in earlier versions of this analysis none of them is left at its baseline schedule, and the appendix's per-additional-dose denominators are non-zero for all seven.
- **The 50%/55% analogs of Tables S.A.3/S.A.6 (appendix) leave 65+ untouched** — 65+'s baseline coverage (60.6%, §2.5) is already above both lower targets, so (unlike at 70%) its column is exactly zero in both, not just small.
- **"Per 100,000 doses" counts doses the schedule reports, not doses the model delivers** — 3.2% of scheduled doses go to people already infected (see the dose-accounting appendix). Those are wasted, not unspent, so they belong in a cost-effectiveness denominator; excluding them would flatter each scenario in proportion to how large its epidemic was.

Appendix: Tables S.A.3 and S.A.6

Table S.A.3 — Additional hospitalizations averted at 70% coverage

Each column scales a single age group's vaccination schedule up to 70% cumulative coverage; "All" scales every age group. Compared against the baseline vaccination scenario. Per §2.5, every one of the seven groups is below 70% at baseline, so every column raises real doses — the 65+ column is small not because it is already covered but because vaccinating 65+ blocks no infection in the fitted baseline, leaving only the severity channel and no benefit to any other age group.

Hospitalizations averted (count)

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	7 [4 – 9]	1 [1 – 1]	6 [4 – 8]	4 [3 – 6]	9 [6 – 12]	2 [2 – 3]	0 [-0 – 0]	21 [14 – 29]
1-4	1 [1 – 1]	19 [13 – 25]	27 [18 – 34]	19 [13 – 24]	37 [25 – 47]	10 [7 – 13]	0 [-0 – 0]	85 [57 – 110]
5-12	1 [1 – 1]	4 [3 – 6]	47 [33 – 62]	19 [13 – 25]	35 [24 – 45]	10 [7 – 13]	0 [-0 – 0]	89 [63 – 116]
13-17	1 [0 – 1]	2 [1 – 3]	15 [10 – 20]	27 [18 – 35]	21 [14 – 28]	6 [4 – 8]	0 [-0 – 0]	55 [36 – 71]
18-49	5 [4 – 6]	19 [15 – 24]	115 [89 – 142]	92 [71 – 114]	231 [183 – 280]	54 [42 – 67]	0 [-0 – 0]	409 [325 – 493]
50-64	4 [3 – 5]	18 [14 – 22]	114 [90 – 137]	94 [73 – 113]	183 [144 – 220]	109 [86 – 131]	0 [-0 – 0]	406 [321 – 493]
65+	25 [23 – 28]	104 [93 – 117]	633 [566 – 717]	510 [455 – 577]	981 [881 – 1,100]	317 [284 – 356]	102 [91 – 113]	2,113 [1,895 – 2,359]

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
All	44 [39 – 48]	169 [151 – 182]	961 [866 – 1,050]	768 [689 – 840]	1,501 [1,369 – 1,636]	511 [462 – 555]	102 [91 – 113]	3,184 [2,929 – 3,452]

% reduction in hospitalizations

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	19.1% [18.8% – 19.3%]	2.9% [2.7% – 3.1%]	16.3% [14.9% – 17.1%]	12.3% [11.2% – 13.0%]	25.0% [23.3% – 26.1%]	6.9% [6.3% – 7.3%]	0.0% [-0.0% – 0.0%]	59.9% [58.2% – 61.4%]
1-4	0.7% [0.6% – 0.7%]	12.4% [12.1% – 12.7%]	17.5% [16.1% – 18.3%]	12.1% [11.0% – 12.9%]	23.9% [22.2% – 25.1%]	6.7% [6.1% – 7.1%]	0.0% [-0.0% – 0.0%]	55.5% [53.5% – 57.2%]
5-12	0.6% [0.5% – 0.6%]	2.7% [2.5% – 2.9%]	30.2% [28.8% – 31.3%]	12.5% [11.3% – 13.3%]	22.2% [20.3% – 23.5%]	6.4% [5.7% – 6.8%]	0.0% [-0.0% – 0.0%]	57.2% [55.0% – 59.0%]
13-17	0.6% [0.5% – 0.6%]	2.3% [2.1% – 2.5%]	15.7% [14.2% – 16.7%]	28.4% [27.0% – 29.3%]	22.4% [20.4% – 23.7%]	6.7% [6.0% – 7.2%]	0.0% [-0.0% – 0.0%]	57.7% [55.6% – 59.6%]
18-49	0.6% [0.6% – 0.7%]	2.6% [2.3% – 2.8%]	15.6% [14.1% – 16.4%]	12.4% [11.2% – 13.2%]	31.0% [29.4% – 32.2%]	7.3% [6.7% – 7.8%]	0.0% [-0.0% – 0.0%]	54.8% [52.6% – 56.5%]
50-64	0.6% [0.5% – 0.6%]	2.5% [2.2% – 2.6%]	15.2% [13.8% – 16.0%]	12.5% [11.3% – 13.2%]	24.3% [22.6% – 25.5%]	14.5% [13.8% – 14.9%]	0.0% [-0.0% – 0.0%]	53.9% [51.8% – 55.5%]
65+	0.6% [0.6% – 0.7%]	2.5% [2.3% – 2.7%]	15.5% [14.2% – 16.3%]	12.5% [11.4% – 13.1%]	24.0% [22.4% – 25.1%]	7.8% [7.2% – 8.1%]	2.5% [2.4% – 2.5%]	51.5% [49.5% – 53.1%]
All	0.7% [0.7% – 0.8%]	2.8% [2.6% – 2.9%]	15.9% [14.6% – 16.7%]	12.7% [11.6% – 13.4%]	24.8% [23.2% – 26.0%]	8.5% [7.8% – 8.9%]	1.7% [1.6% – 1.8%]	52.6% [50.6% – 54.2%]

Hospitalizations averted per 100,000 population

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	9.7 [6.1 – 13.1]	1.5 [0.9 – 2.0]	8.3 [5.1 – 11.0]	6.3 [3.8 – 8.3]	12.7 [7.9 – 17.0]	3.5 [2.2 – 4.7]	0.0 [-0.0 – 0.0]	30.5 [19.3 – 41.1]
1-4	0.4 [0.3 – 0.5]	6.8 [4.6 – 8.8]	9.5 [6.5 – 12.2]	6.6 [4.5 – 8.5]	13.1 [8.8 – 16.7]	3.7 [2.5 – 4.7]	0.0 [-0.0 – 0.0]	30.3 [20.5 – 39.4]
5-12	0.2 [0.1 – 0.2]	0.7 [0.5 – 0.9]	7.8 [5.5 – 10.1]	3.2 [2.2 – 4.2]	5.7 [4.0 – 7.4]	1.6 [1.1 – 2.1]	0.0 [-0.0 – 0.0]	14.7 [10.5 – 19.2]
13-17	0.1 [0.1 – 0.2]	0.5 [0.3 – 0.7]	3.6 [2.4 – 4.8]	6.5 [4.3 – 8.5]	5.1 [3.4 – 6.8]	1.5 [1.0 – 2.0]	0.0 [-0.0 – 0.0]	13.3 [8.8 – 17.3]

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
18-49	0.2 [0.1 – 0.2]	0.6 [0.5 – 0.8]	3.9 [3.0 – 4.8]	3.1 [2.4 – 3.8]	7.8 [6.2 – 9.4]	1.8 [1.4 – 2.2]	0.0 [-0.0 – 0.0]	13.7 [10.9 – 16.5]
50-64	0.3 [0.2 – 0.4]	1.3 [1.0 – 1.6]	8.0 [6.3 – 9.6]	6.6 [5.2 – 7.9]	12.9 [10.1 – 15.4]	7.6 [6.1 – 9.2]	0.0 [-0.0 – 0.0]	28.5 [22.5 – 34.6]
65+	2.1 [1.8 – 2.3]	8.5 [7.6 – 9.6]	51.8 [46.4 – 58.7]	41.7 [37.2 – 47.3]	80.3 [72.2 – 90.1]	25.9 [23.2 – 29.2]	8.3 [7.5 – 9.3]	173.0 [155.1 – 193.1]
All	0.6 [0.6 – 0.7]	2.4 [2.2 – 2.6]	13.7 [12.4 – 15.0]	11.0 [9.9 – 12.0]	21.5 [19.6 – 23.4]	7.3 [6.6 – 7.9]	1.5 [1.3 – 1.6]	45.5 [41.9 – 49.4]

Hospitalizations averted per 100,000 additional doses

Denominators are the *additional* doses each scenario schedules, which reduces to $\max(0, 70\% - \text{baseline coverage}) \times \text{population}$: 32,126 doses for age 0, 60,818 for 1-4, 204,670 for 5-12, 137,978 for 13-17, 1,247,872 for 18-49, 453,063 for 50-64 and 114,828 for 65+. Being a property of the schedule, this is exact and identical across all 627 parameter draws. The **All ages** column raises all seven groups at once, so its denominator is the **total** 2,251,355 additional doses.

Because every column divides all of its rows by that one dose count, the age rows within a column decompose its **All** row by where the averted burden lands — and so sum to it, up to small differences from these cells being medians of per-replicate ratios rather than ratios of medians.

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	21.2 [13.4 – 28.7]	1.7 [1.1 – 2.3]	2.8 [1.7 – 3.8]	3.2 [2.0 – 4.2]	0.7 [0.4 – 1.0]	0.5 [0.3 – 0.7]	0.0 [-0.0 – 0.0]	0.9 [0.6 – 1.3]
1-4	3.4 [2.3 – 4.3]	31.2 [21.1 – 40.6]	13.0 [8.9 – 16.7]	13.5 [9.2 – 17.2]	2.9 [2.0 – 3.7]	2.3 [1.5 – 2.9]	0.0 [-0.0 – 0.0]	3.8 [2.6 – 4.9]
5-12	2.9 [2.0 – 3.8]	6.9 [4.8 – 9.1]	23.0 [16.3 – 30.0]	14.1 [9.7 – 18.3]	2.8 [1.9 – 3.6]	2.2 [1.5 – 2.9]	0.0 [-0.0 – 0.0]	4.0 [2.8 – 5.2]
13-17	1.6 [1.1 – 2.2]	3.6 [2.4 – 4.8]	7.3 [4.7 – 9.6]	19.5 [12.9 – 25.5]	1.7 [1.1 – 2.2]	1.4 [0.9 – 1.9]	0.0 [-0.0 – 0.0]	2.4 [1.6 – 3.2]
18-49	14.9 [11.5 – 18.4]	31.6 [24.3 – 39.1]	56.1 [43.4 – 69.3]	66.7 [51.3 – 82.4]	18.5 [14.7 – 22.4]	12.0 [9.3 – 14.8]	0.0 [-0.0 – 0.0]	18.2 [14.4 – 21.9]
50-64	14.0 [10.9 – 16.9]	30.4 [23.7 – 36.6]	55.8 [43.8 – 66.9]	67.9 [53.3 – 81.6]	14.7 [11.6 – 17.6]	24.0 [19.0 – 29.0]	0.0 [-0.0 – 0.0]	18.0 [14.3 – 21.9]
65+	78.6 [70.2 – 88.5]	170.5 [152.2 – 192.7]	309.2 [276.8 – 350.3]	369.3 [329.5 – 418.4]	78.6 [70.6 – 88.2]	69.9 [62.6 – 78.6]	88.7 [79.6 – 98.6]	93.9 [84.2 – 104.8]
All	136.9 [122.8 – 148.2]	277.4 [248.7 – 300.1]	469.5 [423.3 – 512.9]	556.5 [499.2 – 609.1]	120.3 [109.7 – 131.1]	112.9 [102.0 – 122.5]	88.7 [79.6 – 98.6]	141.4 [130.1 – 153.3]

Per additional dose, raising **13-17** to 70% is the best buy: 556.5 [499.2 – 609.1] hospitalizations averted per 100,000 doses, followed by 5-12 (469.5) — against 141.4 for the all-ages push, 136.9 for age 0, 120.3 for 18-49 and 88.7 for 65+. And 510 of the 768 admissions that the 13-17 push averts land in **65+**, not in 13-17 itself. 18-49 dominates the *absolute* totals (1,501 averted) only because it absorbs 1,247,872 of the 2,251,355 additional doses.

Table S.A.6 — Additional hospitalizations averted at 70% coverage, across VE scenarios

For each VE sensitivity scenario, compares that scenario's own baseline vaccination to the 70%-coverage target applied to every age group.

Hospitalizations averted (count)

Age group	Low VE	Baseline VE (fitted)	High VE
0	19 [12 – 26]	21 [14 – 29]	19 [12 – 25]
1-4	78 [53 – 99]	85 [57 – 110]	68 [46 – 90]
5-12	82 [56 – 107]	89 [63 – 116]	75 [53 – 98]
13-17	50 [32 – 67]	55 [36 – 71]	46 [30 – 59]
18-49	329 [249 – 411]	409 [325 – 493]	366 [285 – 438]
50-64	335 [257 – 406]	406 [321 – 493]	356 [283 – 430]
65+	1,794 [1,567 – 2,056]	2,113 [1,895 – 2,359]	1,845 [1,644 – 2,055]
All	2,699 [2,359 – 2,969]	3,184 [2,929 – 3,452]	2,772 [2,540 – 3,005]

% reduction in hospitalizations

Age group	Low VE	Baseline VE (fitted)	High VE
0	39.1% [36.5% – 40.7%]	59.9% [58.2% – 61.4%]	73.5% [71.9% – 74.9%]
1-4	35.3% [32.4% – 37.1%]	55.5% [53.5% – 57.2%]	67.6% [65.7% – 69.4%]
5-12	37.8% [34.6% – 39.7%]	57.2% [55.0% – 59.0%]	69.0% [67.0% – 70.9%]
13-17	38.0% [34.8% – 40.0%]	57.7% [55.6% – 59.6%]	69.9% [67.9% – 71.8%]
18-49	31.9% [28.8% – 33.8%]	54.8% [52.6% – 56.5%]	67.0% [65.2% – 68.7%]
50-64	31.3% [28.3% – 33.1%]	53.9% [51.8% – 55.5%]	66.1% [64.3% – 67.7%]
65+	31.0% [28.3% – 32.7%]	51.5% [49.5% – 53.1%]	62.8% [61.1% – 64.5%]
All	31.6% [28.7% – 33.3%]	52.6% [50.6% – 54.2%]	64.2% [62.5% – 65.9%]

Hospitalizations averted per 100,000 population

Age group	Low VE	Baseline VE (fitted)	High VE
0	27.5 [16.9 – 36.5]	30.5 [19.3 – 41.1]	26.6 [17.0 – 36.0]
1-4	27.7 [18.9 – 35.1]	30.3 [20.5 – 39.4]	24.3 [16.4 – 32.1]
5-12	13.5 [9.2 – 17.7]	14.7 [10.5 – 19.2]	12.4 [8.8 – 16.2]
13-17	12.2 [7.9 – 16.3]	13.3 [8.8 – 17.3]	11.1 [7.4 – 14.3]
18-49	11.0 [8.4 – 13.8]	13.7 [10.9 – 16.5]	12.3 [9.6 – 14.7]
50-64	23.5 [18.0 – 28.5]	28.5 [22.5 – 34.6]	25.0 [19.8 – 30.2]
65+	146.9 [128.3 – 168.3]	173.0 [155.1 – 193.1]	151.1 [134.6 – 168.2]
All	38.6 [33.7 – 42.5]	45.5 [41.9 – 49.4]	39.6 [36.3 – 43.0]

The absolute gain from the 70% push peaks at the *fitted* VE rather than at high VE (3,184 vs. 2,772 averted): under **High VE** the scenario's own baseline schedule already suppresses so much of the epidemic that there is less burden left for the extra doses to avert.

Table S.A.3 (50% coverage) — Additional hospitalizations averted at 50% coverage

Same construction as Table S.A.3 above, with every age group's schedule scaled to 50% cumulative coverage instead of 70%. Per §2.5, 65+ (60.6%) is the only age group already above a 50% target, so unlike the 70% table its column is now **exactly zero** — vaccinating 65+ further is the one scenario here that is structurally identical to baseline, not merely small. All six other groups are still raised (none of them reach 50% at baseline).

Hospitalizations averted (count)

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	4 [2 – 5]	0 [0 – 0]	3 [2 – 3]	2 [1 – 3]	5 [3 – 7]	1 [1 – 1]	0 [0 – 0]	12 [8 – 17]
1-4	1 [0 – 1]	2 [1 – 2]	12 [8 – 15]	8 [6 – 10]	21 [14 – 26]	4 [3 – 5]	0 [0 – 0]	43 [29 – 55]
5-12	1 [0 – 1]	0 [0 – 0]	22 [15 – 28]	8 [6 – 11]	19 [14 – 25]	4 [3 – 5]	0 [0 – 0]	49 [34 – 63]
13-17	0 [0 – 0]	0 [0 – 0]	7 [4 – 9]	12 [8 – 16]	12 [8 – 16]	2 [2 – 3]	0 [0 – 0]	30 [20 – 39]
18-49	3 [2 – 3]	2 [1 – 2]	52 [40 – 64]	40 [31 – 50]	133 [106 – 162]	21 [16 – 26]	0 [0 – 0]	227 [180 – 277]
50-64	3 [2 – 3]	1 [1 – 2]	51 [40 – 62]	41 [32 – 49]	104 [82 – 125]	43 [34 – 51]	0 [0 – 0]	218 [173 – 263]
65+	14 [13 – 16]	8 [7 – 9]	286 [254 – 324]	222 [197 – 253]	557 [499 – 627]	122 [109 – 138]	0 [0 – 0]	1,104 [992 – 1,242]
All	25 [22 – 27]	13 [12 – 15]	434 [388 – 475]	335 [299 – 368]	856 [775 – 932]	198 [178 – 215]	0 [0 – 0]	1,689 [1,539 – 1,840]

% reduction in hospitalizations

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	10.8% [10.6% – 10.9%]	0.2% [0.2% – 0.2%]	7.4% [6.7% – 7.8%]	5.4% [4.8% – 5.7%]	14.2% [13.1% – 14.9%]	2.7% [2.4% – 2.8%]	0.0% [0.0% – 0.0%]	35.0% [33.4% – 36.1%]
1-4	0.4% [0.4% – 0.4%]	1.0% [1.0% – 1.0%]	7.9% [7.2% – 8.4%]	5.3% [4.7% – 5.6%]	13.6% [12.5% – 14.3%]	2.6% [2.3% – 2.8%]	0.0% [0.0% – 0.0%]	27.9% [26.1% – 29.1%]
5-12	0.3% [0.3% – 0.4%]	0.2% [0.2% – 0.2%]	14.1% [13.2% – 14.6%]	5.5% [4.9% – 5.8%]	12.6% [11.3% – 13.4%]	2.5% [2.2% – 2.6%]	0.0% [0.0% – 0.0%]	31.3% [29.3% – 32.7%]
13-17	0.3% [0.3% – 0.3%]	0.2% [0.2% – 0.2%]	7.1% [6.3% – 7.6%]	12.7% [12.0% – 13.2%]	12.7% [11.4% – 13.5%]	2.6% [2.3% – 2.8%]	0.0% [0.0% – 0.0%]	31.6% [29.6% – 33.0%]
18-49	0.4% [0.3% – 0.4%]	0.2% [0.2% – 0.2%]	7.0% [6.3% – 7.4%]	5.4% [4.8% – 5.8%]	17.9% [16.8% – 18.7%]	2.8% [2.6% – 3.0%]	0.0% [0.0% – 0.0%]	30.5% [28.7% – 31.8%]
50-64	0.3% [0.3% – 0.4%]	0.2% [0.2% – 0.2%]	6.8% [6.1% – 7.2%]	5.4% [4.9% – 5.8%]	13.8% [12.7% – 14.6%]	5.7% [5.4% – 5.9%]	0.0% [0.0% – 0.0%]	29.0% [27.2% – 30.3%]

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
65+	0.3% [0.3% – 0.4%]	0.2% [0.2% – 0.2%]	7.0% [6.4% – 7.4%]	5.4% [4.9% – 5.8%]	13.6% [12.6% – 14.3%]	3.0% [2.8% – 3.2%]	0.0% [0.0% – 0.0%]	27.0% [25.3% – 28.2%]
All	0.4% [0.4% – 0.4%]	0.2% [0.2% – 0.2%]	7.2% [6.5% – 7.6%]	5.5% [5.0% – 5.9%]	14.1% [13.1% – 14.9%]	3.3% [3.0% – 3.5%]	0.0% [0.0% – 0.0%]	27.9% [26.1% – 29.2%]

Hospitalizations averted per 100,000 population

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	5.5 [3.5 – 7.4]	0.1 [0.1 – 0.2]	3.7 [2.3 – 4.9]	2.7 [1.7 – 3.6]	7.2 [4.4 – 9.6]	1.4 [0.8 – 1.8]	0.0 [0.0 – 0.0]	17.7 [11.1 – 23.9]
1-4	0.2 [0.1 – 0.3]	0.5 [0.4 – 0.7]	4.3 [3.0 – 5.5]	2.9 [2.0 – 3.7]	7.4 [5.0 – 9.4]	1.4 [1.0 – 1.8]	0.0 [0.0 – 0.0]	15.2 [10.3 – 19.5]
5-12	0.1 [0.1 – 0.1]	0.1 [0.0 – 0.1]	3.6 [2.5 – 4.7]	1.4 [1.0 – 1.8]	3.2 [2.2 – 4.2]	0.6 [0.4 – 0.8]	0.0 [0.0 – 0.0]	8.0 [5.7 – 10.4]
13-17	0.1 [0.0 – 0.1]	0.0 [0.0 – 0.1]	1.6 [1.1 – 2.1]	2.9 [1.9 – 3.8]	2.9 [1.9 – 3.8]	0.6 [0.4 – 0.8]	0.0 [0.0 – 0.0]	7.3 [4.8 – 9.5]
18-49	0.1 [0.1 – 0.1]	0.1 [0.0 – 0.1]	1.7 [1.3 – 2.2]	1.3 [1.0 – 1.7]	4.5 [3.5 – 5.5]	0.7 [0.5 – 0.9]	0.0 [0.0 – 0.0]	7.6 [6.0 – 9.3]
50-64	0.2 [0.1 – 0.2]	0.1 [0.1 – 0.1]	3.6 [2.8 – 4.3]	2.9 [2.2 – 3.5]	7.3 [5.7 – 8.8]	3.0 [2.4 – 3.6]	0.0 [0.0 – 0.0]	15.3 [12.1 – 18.5]
65+	1.2 [1.0 – 1.3]	0.7 [0.6 – 0.8]	23.4 [20.8 – 26.5]	18.2 [16.1 – 20.7]	45.6 [40.9 – 51.3]	10.0 [9.0 – 11.3]	0.0 [0.0 – 0.0]	90.4 [81.2 – 101.7]
All	0.4 [0.3 – 0.4]	0.2 [0.2 – 0.2]	6.2 [5.6 – 6.8]	4.8 [4.3 – 5.3]	12.2 [11.1 – 13.3]	2.8 [2.5 – 3.1]	0.0 [0.0 – 0.0]	24.2 [22.0 – 26.3]

Hospitalizations averted per 100,000 additional doses

Denominators reduce to $\max(0, 50\% - \text{baseline coverage}) \times \text{population}$: 18,112 doses for age 0, 4,765 for 1-4, 83,411 for 5-12, 55,621 for 13-17, 652,232 for 18-49, 168,176 for 50-64 and **zero** for 65+ (already above 50%). The **All ages** column's denominator is the total 982,317 additional doses.

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	21.3 [13.5 – 28.8]	1.7 [1.1 – 2.3]	3.1 [1.9 – 4.2]	3.4 [2.1 – 4.5]	0.8 [0.5 – 1.0]	0.6 [0.3 – 0.8]	—	1.3 [0.8 – 1.7]
1-4	3.4 [2.3 – 4.3]	32.0 [21.7 – 41.6]	14.5 [9.9 – 18.5]	14.5 [9.9 – 18.6]	3.2 [2.2 – 4.1]	2.3 [1.6 – 3.0]	—	4.3 [3.0 – 5.6]
5-12	2.9 [2.0 – 3.8]	7.0 [4.9 – 9.2]	26.2 [18.5 – 34.1]	15.3 [10.4 – 19.8]	3.0 [2.1 – 3.9]	2.3 [1.6 – 3.0]	—	5.0 [3.5 – 6.4]
13-17	1.6 [1.1 – 2.2]	3.7 [2.4 – 4.9]	8.0 [5.2 – 10.6]	21.7 [14.4 – 28.4]	1.8 [1.2 – 2.4]	1.4 [0.9 – 1.9]	—	3.1 [2.0 – 4.0]

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
18-49	14.9 [11.5 – 18.4]	32.0 [24.6 – 39.5]	62.1 [47.7 – 76.8]	72.1 [55.5 – 89.2]	20.4 [16.2 – 24.9]	12.4 [9.6 – 15.4]	—	23.1 [18.3 – 28.2]
50-64	14.0 [10.9 – 16.9]	30.8 [24.0 – 37.1]	61.7 [48.1 – 74.3]	73.6 [57.4 – 88.6]	16.0 [12.5 – 19.1]	25.3 [20.1 – 30.6]	—	22.2 [17.6 – 26.8]
65+	78.8 [70.4 – 88.8]	172.9 [154.1 – 195.9]	342.6 [304.0 – 388.6]	399.9 [354.5 – 454.3]	85.5 [76.5 – 96.2]	72.7 [65.1 – 81.9]	—	112.4 [101.0 – 126.4]
All	137.3 [123.0 – 148.6]	281.4 [252.0 – 305.3]	520.0 [465.6 – 568.9]	603.1 [536.9 – 660.8]	131.2 [118.9 – 142.9]	117.7 [105.9 – 127.8]	—	171.9 [156.7 – 187.3]

Same ranking as the 70% table: 13-17 is still the best per-dose buy (603.1 averted per 100,000 doses), followed by 5-12 (520.0) — against 171.9 for the all-ages push. The all-ages push now averts 1,689 [1,539 – 1,840] hospitalizations overall — a 27.9% reduction — against 3,184 at 70% coverage (§3, Table S.A.3 above).

Table S.A.3 (55% coverage) — Additional hospitalizations averted at 55% coverage

Same construction, scaled to 55% cumulative coverage. 65+ is again the only group already above target (60.6% > 55%), so its column is again exactly zero.

Hospitalizations averted (count)

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	5 [3 – 6]	0 [0 – 0]	3 [2 – 5]	3 [2 – 3]	6 [4 – 8]	1 [1 – 2]	0 [0 – 0]	15 [10 – 20]
1-4	1 [0 – 1]	6 [4 – 8]	16 [11 – 21]	11 [7 – 14]	25 [17 – 32]	6 [4 – 7]	0 [0 – 0]	56 [38 – 72]
5-12	1 [0 – 1]	1 [1 – 2]	29 [21 – 38]	11 [8 – 15]	24 [17 – 31]	5 [4 – 7]	0 [0 – 0]	61 [43 – 80]
13-17	0 [0 – 0]	1 [0 – 1]	9 [6 – 12]	16 [11 – 21]	14 [9 – 19]	3 [2 – 5]	0 [0 – 0]	38 [25 – 49]
18-49	3 [2 – 4]	6 [5 – 7]	69 [53 – 86]	54 [42 – 67]	160 [127 – 195]	30 [23 – 37]	0 [0 – 0]	282 [224 – 343]
50-64	3 [2 – 4]	6 [4 – 7]	69 [54 – 83]	55 [43 – 66]	126 [99 – 151]	60 [48 – 72]	0 [0 – 0]	276 [218 – 333]
65+	17 [15 – 19]	32 [29 – 37]	383 [341 – 434]	300 [267 – 341]	674 [602 – 757]	173 [155 – 195]	0 [0 – 0]	1,393 [1,251 – 1,563]
All	30 [27 – 32]	53 [47 – 57]	581 [521 – 636]	453 [404 – 496]	1,033 [938 – 1,125]	280 [252 – 304]	0 [0 – 0]	2,127 [1,942 – 2,312]

% reduction in hospitalizations

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	12.9% [12.7% – 13.0%]	0.9% [0.8% – 1.0%]	9.9% [9.0% – 10.4%]	7.2% [6.5% – 7.7%]	17.1% [15.9% – 18.0%]	3.8% [3.4% – 4.0%]	0.0% [0.0% – 0.0%]	42.7% [40.9% – 43.9%]
1-4	0.5% [0.4% – 0.5%]	3.9% [3.8% – 4.0%]	10.6% [9.7% – 11.2%]	7.1% [6.4% – 7.6%]	16.4% [15.1% – 17.3%]	3.7% [3.3% – 3.9%]	0.0% [0.0% – 0.0%]	36.4% [34.4% – 37.8%]
5-12	0.4% [0.4% – 0.4%]	0.9% [0.8% – 0.9%]	18.7% [17.6% – 19.4%]	7.4% [6.6% – 7.9%]	15.2% [13.8% – 16.2%]	3.5% [3.1% – 3.7%]	0.0% [0.0% – 0.0%]	39.4% [37.2% – 41.0%]
13-17	0.4% [0.3% – 0.4%]	0.7% [0.6% – 0.8%]	9.5% [8.5% – 10.1%]	17.0% [16.1% – 17.7%]	15.3% [13.9% – 16.3%]	3.6% [3.3% – 3.9%]	0.0% [0.0% – 0.0%]	39.7% [37.5% – 41.3%]
18-49	0.4% [0.4% – 0.5%]	0.8% [0.7% – 0.9%]	9.4% [8.4% – 10.0%]	7.3% [6.6% – 7.8%]	21.6% [20.2% – 22.5%]	4.0% [3.6% – 4.2%]	0.0% [0.0% – 0.0%]	38.0% [35.9% – 39.4%]
50-64	0.4% [0.4% – 0.4%]	0.8% [0.7% – 0.8%]	9.2% [8.3% – 9.7%]	7.3% [6.6% – 7.8%]	16.7% [15.4% – 17.6%]	8.0% [7.6% – 8.2%]	0.0% [0.0% – 0.0%]	36.6% [34.6% – 38.0%]
65+	0.4% [0.4% – 0.4%]	0.8% [0.7% – 0.8%]	9.4% [8.5% – 9.9%]	7.4% [6.7% – 7.8%]	16.5% [15.3% – 17.3%]	4.2% [3.9% – 4.5%]	0.0% [0.0% – 0.0%]	34.0% [32.1% – 35.4%]
All	0.5% [0.4% – 0.5%]	0.9% [0.8% – 0.9%]	9.6% [8.8% – 10.2%]	7.5% [6.8% – 7.9%]	17.1% [15.8% – 17.9%]	4.6% [4.3% – 4.9%]	0.0% [0.0% – 0.0%]	35.2% [33.1% – 36.6%]

Hospitalizations averted per 100,000 population

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	6.6 [4.1 – 8.9]	0.5 [0.3 – 0.6]	5.0 [3.1 – 6.6]	3.7 [2.3 – 4.9]	8.7 [5.4 – 11.6]	1.9 [1.2 – 2.5]	0.0 [0.0 – 0.0]	21.6 [13.6 – 29.1]
1-4	0.3 [0.2 – 0.3]	2.1 [1.4 – 2.8]	5.8 [4.0 – 7.4]	3.9 [2.7 – 5.0]	9.0 [6.1 – 11.4]	2.0 [1.4 – 2.5]	0.0 [0.0 – 0.0]	19.9 [13.5 – 25.6]
5-12	0.1 [0.1 – 0.1]	0.2 [0.2 – 0.3]	4.8 [3.4 – 6.2]	1.9 [1.3 – 2.4]	3.9 [2.7 – 5.1]	0.9 [0.6 – 1.2]	0.0 [0.0 – 0.0]	10.1 [7.1 – 13.2]
13-17	0.1 [0.1 – 0.1]	0.2 [0.1 – 0.2]	2.2 [1.4 – 2.9]	3.9 [2.6 – 5.1]	3.5 [2.3 – 4.6]	0.8 [0.5 – 1.1]	0.0 [0.0 – 0.0]	9.2 [6.1 – 12.0]
18-49	0.1 [0.1 – 0.1]	0.2 [0.2 – 0.2]	2.3 [1.8 – 2.9]	1.8 [1.4 – 2.2]	5.4 [4.3 – 6.6]	1.0 [0.8 – 1.2]	0.0 [0.0 – 0.0]	9.5 [7.5 – 11.5]
50-64	0.2 [0.2 – 0.3]	0.4 [0.3 – 0.5]	4.8 [3.8 – 5.8]	3.9 [3.0 – 4.7]	8.8 [6.9 – 10.6]	4.2 [3.3 – 5.1]	0.0 [0.0 – 0.0]	19.4 [15.3 – 23.4]
65+	1.4 [1.2 – 1.6]	2.6 [2.4 – 3.0]	31.4 [27.9 – 35.6]	24.6 [21.8 – 27.9]	55.1 [49.3 – 62.0]	14.2 [12.7 – 15.9]	0.0 [0.0 – 0.0]	114.0 [102.5 – 128.0]

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
All	0.4 [0.4 – 0.5]	0.8 [0.7 – 0.8]	8.3 [7.5 – 9.1]	6.5 [5.8 – 7.1]	14.8 [13.4 – 16.1]	4.0 [3.6 – 4.3]	0.0 [0.0 – 0.0]	30.4 [27.8 – 33.1]

Hospitalizations averted per 100,000 additional doses

Denominators: 21,616 doses for age 0, 18,778 for 1-4, 113,726 for 5-12, 76,210 for 13-17, 801,142 for 18-49, 239,398 for 50-64 and zero for 65+. The **All ages** column's denominator is the total 1,270,870 additional doses.

Age group (counted)	0 vaccinated	1-4 vaccinated	5-12 vaccinated	13-17 vaccinated	18-49 vaccinated	50-64 vaccinated	65+ vaccinated	All vaccinated
0	21.3 [13.4 – 28.7]	1.7 [1.1 – 2.3]	3.1 [1.9 – 4.1]	3.4 [2.1 – 4.5]	0.8 [0.5 – 1.0]	0.6 [0.3 – 0.7]	—	1.2 [0.7 – 1.6]
1-4	3.4 [2.3 – 4.3]	31.8 [21.5 – 41.4]	14.3 [9.8 – 18.2]	14.3 [9.8 – 18.3]	3.1 [2.1 – 4.0]	2.3 [1.6 – 3.0]	—	4.4 [3.0 – 5.6]
5-12	2.9 [2.0 – 3.8]	7.0 [4.9 – 9.2]	25.6 [18.0 – 33.3]	15.0 [10.3 – 19.5]	3.0 [2.1 – 3.8]	2.3 [1.6 – 3.0]	—	4.8 [3.4 – 6.3]
13-17	1.6 [1.1 – 2.2]	3.7 [2.4 – 4.9]	7.9 [5.1 – 10.4]	21.2 [14.1 – 27.8]	1.8 [1.2 – 2.4]	1.4 [0.9 – 1.9]	—	3.0 [2.0 – 3.9]
18-49	14.9 [11.5 – 18.4]	31.9 [24.6 – 39.5]	61.1 [46.9 – 75.5]	71.1 [54.7 – 87.9]	20.0 [15.9 – 24.4]	12.4 [9.6 – 15.3]	—	22.2 [17.6 – 27.0]
50-64	14.0 [10.9 – 16.9]	30.7 [24.0 – 37.0]	60.7 [47.3 – 72.9]	72.5 [56.6 – 87.3]	15.7 [12.3 – 18.8]	25.0 [19.9 – 30.3]	—	21.7 [17.2 – 26.2]
65+	78.7 [70.3 – 88.6]	172.3 [153.8 – 195.1]	336.7 [299.6 – 381.8]	394.2 [349.8 – 447.3]	84.1 [75.2 – 94.5]	72.2 [64.6 – 81.3]	—	109.6 [98.5 – 123.0]
All	137.1 [123.0 – 148.4]	280.7 [251.5 – 303.8]	511.0 [458.3 – 558.9]	594.1 [530.5 – 650.8]	129.0 [117.1 – 140.4]	116.8 [105.2 – 126.8]	—	167.4 [152.8 – 181.9]

Same ranking again: 13-17 (594.1) then 5-12 (511.0) remain the best per-dose buys, against 167.4 for the all-ages push. The 50%/55%/70% coverage series is monotonic as expected — 27.9% → 35.2% → 52.6% overall reduction. Of the 1,057 additional hospitalizations averted going from the 55% to the 70% push, 720 (68%) are counted in **65+** alone — the one age group the 70% target newly raises above its own baseline schedule (60.6%) that the 50%/55% targets leave untouched (\$2.5), so it is the single largest source of *additional* benefit from pushing past 55% even though its own direct dose count barely changes.

Table S.A.6 (50%/55% coverage) — Additional hospitalizations averted, across VE scenarios

50%/55% analogs of Table S.A.6: each VE preset's own baseline compared against that preset's own 50%- or 55%-coverage (all ages) target (same VE × coverage combinations as the 70% table, just at the two lower targets).

50% coverage

Hospitalizations averted (count)

Age group	Low VE	Baseline VE (fitted)	High VE
0	10 [6 – 13]	12 [8 – 17]	12 [8 – 16]
1-4	32 [22 – 41]	43 [29 – 55]	37 [25 – 48]
5-12	39 [26 – 51]	49 [34 – 63]	45 [32 – 59]

Age group	Low VE	Baseline VE (fitted)	High VE
13-17	24 [15 – 32]	30 [20 – 39]	27 [18 – 35]
18-49	156 [117 – 197]	227 [180 – 277]	224 [176 – 267]
50-64	155 [117 – 189]	218 [173 – 263]	210 [166 – 254]
65+	801 [687 – 926]	1,104 [992 – 1,242]	1,054 [945 – 1,179]
All	1,224 [1,039 – 1,353]	1,689 [1,539 – 1,840]	1,614 [1,482 – 1,743]

% reduction in hospitalizations

Age group	Low VE	Baseline VE (fitted)	High VE
0	20.1% [18.5% – 21.1%]	35.0% [33.4% – 36.1%]	47.6% [46.3% – 48.9%]
1-4	14.7% [13.0% – 15.8%]	27.9% [26.1% – 29.1%]	36.7% [35.1% – 38.1%]
5-12	17.9% [15.9% – 19.1%]	31.3% [29.3% – 32.7%]	41.5% [39.9% – 43.1%]
13-17	18.0% [15.9% – 19.2%]	31.6% [29.6% – 33.0%]	42.1% [40.4% – 43.7%]
18-49	15.1% [13.3% – 16.2%]	30.5% [28.7% – 31.8%]	41.0% [39.5% – 42.4%]
50-64	14.5% [12.7% – 15.5%]	29.0% [27.2% – 30.3%]	39.0% [37.5% – 40.3%]
65+	13.9% [12.2% – 14.8%]	27.0% [25.3% – 28.2%]	36.0% [34.6% – 37.3%]
All	14.3% [12.7% – 15.3%]	27.9% [26.1% – 29.2%]	37.3% [35.9% – 38.6%]

Hospitalizations averted per 100,000 population

Age group	Low VE	Baseline VE (fitted)	High VE
0	14.1 [8.6 – 18.7]	17.7 [11.1 – 23.9]	17.2 [11.0 – 23.3]
1-4	11.6 [7.9 – 14.7]	15.2 [10.3 – 19.5]	13.2 [8.8 – 17.3]
5-12	6.4 [4.3 – 8.4]	8.0 [5.7 – 10.4]	7.4 [5.3 – 9.7]
13-17	5.7 [3.7 – 7.7]	7.3 [4.8 – 9.5]	6.7 [4.4 – 8.6]
18-49	5.2 [3.9 – 6.6]	7.6 [6.0 – 9.3]	7.5 [5.9 – 9.0]
50-64	10.9 [8.2 – 13.3]	15.3 [12.1 – 18.5]	14.7 [11.7 – 17.9]
65+	65.6 [56.2 – 75.8]	90.4 [81.2 – 101.7]	86.3 [77.4 – 96.5]
All	17.5 [14.9 – 19.3]	24.2 [22.0 – 26.3]	23.1 [21.2 – 24.9]

55% coverage

Hospitalizations averted (count)

Age group	Low VE	Baseline VE (fitted)	High VE
0	13 [8 – 17]	15 [10 – 20]	14 [9 – 19]
1-4	45 [31 – 57]	56 [38 – 72]	47 [32 – 62]
5-12	51 [35 – 67]	61 [43 – 80]	55 [39 – 72]
13-17	31 [20 – 42]	38 [25 – 49]	34 [22 – 43]
18-49	204 [154 – 257]	282 [224 – 343]	270 [212 – 322]
50-64	206 [156 – 250]	276 [218 – 333]	257 [204 – 312]
65+	1,060 [915 – 1,225]	1,393 [1,251 – 1,563]	1,293 [1,158 – 1,445]

Age group	Low VE	Baseline VE (fitted)	High VE
All	1,620 [1,385 – 1,788]	2,127 [1,942 – 2,312]	1,971 [1,809 – 2,134]

% reduction in hospitalizations

Age group	Low VE	Baseline VE (fitted)	High VE
0	25.5% [23.5% – 26.7%]	42.7% [40.9% – 43.9%]	56.1% [54.7% – 57.5%]
1-4	20.6% [18.4% – 21.9%]	36.4% [34.4% – 37.8%]	46.8% [45.2% – 48.5%]
5-12	23.6% [21.2% – 25.1%]	39.4% [37.2% – 41.0%]	50.7% [48.9% – 52.4%]
13-17	23.7% [21.2% – 25.2%]	39.7% [37.5% – 41.3%]	51.3% [49.6% – 53.1%]
18-49	19.9% [17.6% – 21.2%]	38.0% [35.9% – 39.4%]	49.5% [47.8% – 51.0%]
50-64	19.2% [17.0% – 20.5%]	36.6% [34.6% – 38.0%]	47.8% [46.2% – 49.3%]
65+	18.4% [16.3% – 19.6%]	34.0% [32.1% – 35.4%]	44.1% [42.6% – 45.6%]
All	19.0% [16.9% – 20.2%]	35.2% [33.1% – 36.6%]	45.6% [44.1% – 47.1%]

Hospitalizations averted per 100,000 population

Age group	Low VE	Baseline VE (fitted)	High VE
0	17.9 [10.9 – 23.7]	21.6 [13.6 – 29.1]	20.3 [13.0 – 27.5]
1-4	16.1 [11.0 – 20.5]	19.9 [13.5 – 25.6]	16.8 [11.3 – 22.2]
5-12	8.5 [5.7 – 11.0]	10.1 [7.1 – 13.2]	9.1 [6.4 – 11.8]
13-17	7.6 [4.9 – 10.1]	9.2 [6.1 – 12.0]	8.1 [5.4 – 10.5]
18-49	6.9 [5.2 – 8.6]	9.5 [7.5 – 11.5]	9.1 [7.1 – 10.8]
50-64	14.5 [11.0 – 17.6]	19.4 [15.3 – 23.4]	18.0 [14.3 – 21.9]
65+	86.8 [74.9 – 100.3]	114.0 [102.5 – 128.0]	105.9 [94.8 – 118.3]
All	23.2 [19.8 – 25.6]	30.4 [27.8 – 33.1]	28.2 [25.9 – 30.5]

Same pattern as the 70% table at both lower targets: the **absolute** gain peaks at the *fitted* VE rather than at **High VE** (1,689 vs. 1,614 at 50%; 2,127 vs. 1,971 at 55%) — **High VE**'s own baseline already suppresses enough of the epidemic that less burden is left for the extra doses to avert, the same mechanism as at 70%. **% reduction** and **per-100,000-population** rates, by contrast, remain monotonic in VE at every target (Low < Baseline < High) because those are normalized to each scenario's own denominator rather than reflecting the absolute size of the epidemic being cut into.

Appendix: dose accounting — scheduled vs. delivered doses

Two different dose counts appear in this model, and the per-100,000-doses panels depend on which one is used.

Scheduled doses are what the vaccination schedule reports: the daily vaccination proportions of §2.5, summed over the season and multiplied by population. **Delivered** doses are what the model records as an actual **S** → **SV** transition. They differ because of the cap in §1.4 — a dose is only delivered if the intended recipient is still in **S**. Someone already infected, recovered, or hospitalized is skipped.

That gap is not a saving. In the real world the dose is still bought, shipped and injected; it simply arrives after the recipient has already been infected and buys no protection. Counting only delivered doses would therefore understate the true cost of every scenario — and understate it *most* for the scenarios with the largest epidemics, which is exactly backwards for a cost-effectiveness denominator. Every "per 100,000 doses" figure in this report divides by the scheduled count.

Baseline schedule, median across the 627 posterior draws:


Age group	Population	Scheduled doses	Delivered doses	Wasted doses	% wasted	Scheduled coverage	Delivered coverage
0	70,067	16,921	16,726	195	1.2%	24.1%	23.9%
1-4	280,268	135,369	133,722	1,647	1.2%	48.3%	47.7%
5-12	606,291	219,734	213,727	6,007	2.7%	36.2%	35.3%
13-17	411,782	150,270	147,131	3,139	2.1%	36.5%	35.7%
18-49	2,978,204	836,870	802,991	33,879	4.0%	28.1%	27.0%
50-64	1,424,434	544,041	525,012	19,029	3.5%	38.2%	36.9%
65+	1,221,349	740,116	720,282	19,834	2.7%	60.6%	59.0%
All	6,992,395	2,643,322	2,559,591	83,731	3.2%	37.8%	36.6%


Waste tracks infection attack rate, as expected: it is lowest in the youngest groups (1.2% in 0 and 1-4) and highest in 18-49 (4.0%) and 50-64 (3.5%), which between them account for 52,908 of the 83,731 wasted doses. Overall 3.2% of the season's scheduled doses land in arms that no longer benefit, pulling realized coverage from 37.8% down to 36.6%.

A second consequence, relevant to why the per-dose tables are built the way they are: delivered doses are an *output* of the simulation, so they vary with the parameter draw even when the schedule is identical, and they leak between age groups, since a milder epidemic in one group leaves more susceptibles for the cap to reach in another. Scheduled doses have neither property.

Vaccination coverage over time, by age group

The proportion of each age group's population the **schedule** vaccinates per day, and the cumulative version of the same series — both taken directly from the vaccination-schedule input (`daily_vaccines_df`), with no adjustment for the §1.4 delivery cap above. The cumulative series' final value reproduces the §2.5 coverage table exactly, by construction (it is that table, read out day by day rather than just at the end of the season).





The schedule is front-loaded, with a pronounced spike in early November (all age groups peak together, 65+ reaching nearly 1.3% of its population in a single day) and a long tail of small, infrequent top-ups from December onward. 65+ and 1-4 pull ahead of the other five age groups almost immediately and stay ahead throughout the season, consistent with their higher final coverage (60.6% and 48.3% respectively, §2.5); 18-49 lags visibly in the first six weeks before converging toward the pack's middle by January.