The background of the image is a world map centered on the Northern Hemisphere. It features a grid of latitude and longitude lines. The grid consists of approximately 15 longitude lines and 8 latitude lines, starting from the North Pole at the top and extending down to the equator. The continents are shown as irregular outlines. The title text is overlaid on this grid.

An Album of Map Projections

**U.S. Geological Survey
Professional Paper 1453**

An Album of Map Projections

**By John P. Snyder,
U.S. Geological Survey,
and Philip M. Voxland,
University of Minnesota**

**Introduction by Joel L. Morrison,
U.S. Geological Survey**

**U.S. Geological Survey
Professional Paper 1453**

Department of the Interior

MANUEL LUJAN, JR., Secretary

U.S. Geological Survey

Dallas L. Peck, Director

Any use of trade names in this publication is for descriptive purposes
only and does not imply endorsement by the U.S. Geological Survey

Library of Congress Cataloging in Publication Data

Snyder, John Parr, 1926—
An album of map projections.

(U.S. Geological Survey professional paper ; 1453)

Bibliography: p.

Includes index.

Supt. of Docs. no.: I 19.16:1453

1. Map-projection. I. Voxland, Philip M. II. Title. III. Series.


GA110.S575 1989 526.8 86-600253

For sale by the Books and Open-File Reports Section
U.S. Geological Survey, Federal Center,
Box 25425, Denver CO 80225

United States Government Printing Office : 1989

CONTENTS

Preface	vii
Introduction	1
Glossary	2
Guide to selecting map projections	5
Distortion diagrams	8
Cylindrical projections	
Mercator	10
Transverse Mercator	12
Oblique Mercator	14
Lambert Cylindrical Equal-Area	16
Behrmann Cylindrical Equal-Area	19
Plate Carrée	22
Equirectangular	24
Cassini	26
Oblique Plate Carrée	28
Central Cylindrical	30
Gall	33
Miller Cylindrical	35
Pseudocylindrical projections	
Sinusoidal	37
McBryde-Thomas Flat-Polar Sinusoidal	44
Eckert V	46
Winkel I	48
Eckert VI	50
McBryde S3	52
Mollweide	54
Eckert III	58
Eckert IV	60
Putniņš P _{2'}	62
Hatano Asymmetrical Equal-Area	64
Goode Homolosine	66
Boggs Eumorphic	68
Craster Parabolic	70
McBryde-Thomas Flat-Polar Parabolic	72
Quartic Authalic	74
McBryde-Thomas Flat-Polar Quartic	76
Putniņš P ₅	78
DeNoyer Semi-Elliptical	80
Robinson	82
Collignon	84
Eckert I	86
Eckert II	88
Loximuthal	90
Conic projections	
Equidistant Conic	92
Lambert Conformal Conic	95
Bipolar Oblique Conic Conformal	99
Albers Equal-Area Conic	100
Lambert Equal-Area Conic	102
Perspective Conic	104
Polyconic	106
Rectangular Polyconic	110
Modified Polyconic, International Map of the World series	111
Bonne	112
Werner	114
Azimuthal projections	
Perspective	
Gnomonic	116
Stereographic	120
Orthographic	124
General Vertical Perspective	128
Nonperspective	
Azimuthal Equidistant	132
Lambert Azimuthal Equal-Area	136
Airy	140



CONTENTS

Modified	
Two-Point Azimuthal	144
Two-Point Equidistant	146
Miller Oblated Stereographic	148
Wiechel	149
Craig Retroazimuthal	150
Hammer Retroazimuthal	152
Littrow	154
Berghaus Star	156
Aitoff	158
Hammer	160
Briesemeister	162
Winkel Tripel	164
Eckert-Greifendorff	166
Wagner VII	168
Chamberlin Trimetric	170
Tilted Perspective	172
Miscellaneous	
Bacon Globular	174
Fournier Globular	175
Nicolosi Globular	176
Apian Globular I	177
Ortelius Oval	178
Lagrange	180
Eisenlohr	184
August Epicycloidal	186
Guyou	188
Peirce Quincuncial	190
Adams Hemisphere in a Square	192
Adams World in a Square I	194
Adams World in a Square II	195
Lee Conformal World in a Triangle	196
Conformal World in an Ellipse	198
Van der Grinten I	200
Van der Grinten II	202
Van der Grinten III	203
Van der Grinten IV	204
Armadillo	206
GS50	208
Space Oblique Mercator	211
Satellite Tracking	213
Appendix: Formulas for projections	216
Cylindrical projections	218
Pseudocylindrical projections	220
Conic projections	224
Pseudoconic projections	225
Azimuthal projections	
Perspective	226
Nonperspective	228
Modified	230
Miscellaneous projections	234
Selected references	240
Index	242

Preface

Every map must begin, either consciously or unconsciously, with the choice of a map projection, the method of transforming the spherelike surface being mapped onto a flat plane. Because this transformation can become complicated, there is a tendency to choose the projection that has been used for a suitable available base map, even if it does not emphasize the desired properties.

This album has been prepared to acquaint those in the cartographic profession and other cartophiles with the wide range of map projections that have been developed during the past few centuries. Because including all of the hundreds of known projection systems would be both difficult and expensive, this volume presents some 90 basic projections in over 130 different modifications and aspects. Dozens of other projections are mentioned briefly. All the popular projections have been included, along with many others that have interesting appearances or features, even if their usefulness for any serious mapping is limited.

The most important question to raise about a map projection concerns distortion. How accurately does the flat projection show the desired portion of the spherelike Earth (or heavens or planetary body)? Since accuracy is desired in area, shape, and distance but cannot be achieved in all three simultaneously and since the region mapped may vary in overall extent and shape, the answer is not clear cut. Because this album emphasizes visual comparisons rather than tables of scale variation, the Tissot indicatrix has been adopted as the most common means of comparing distortion on different parts of the same projection and on different projections.

Computerized plotting has facilitated the preparation of many outline map projections

on a standardized basis, so that comparisons are more realistic. These maps have been plotted on film as lines 0.005 in. wide by means of a photo-head computer-driven plotter and World Data Bank I shoreline files (national boundaries included where appropriate). Central meridians and other parameters have been given the same values in so far as it was practical.

Some smaller maps have been prepared with the Dahlgren (Hershey) shoreline file. The map projection software for any given map consisted of one of three programs. WORLD was developed for a mainframe computer by Philip Voxland at the University of Minnesota. CYLIND and AZIMUTH were developed for a personal microcomputer and converted to mainframe computer programs at the U.S. Geological Survey (USGS) by John Snyder, with the guidance of Michael K. Linck, Jr., and John F. Waananen. The authors thank Joel L. Morrison, past president of the International Cartographic Association and Assistant Division Chief for Research, National Mapping Division, USGS, for writing the introduction to this book, as well as Tau Rho Alpha (USGS) and Arthur H. Robinson (University of Wisconsin/Madison) for many constructive comments on the manuscript. The authors also thank Christopher J. Cialek, Kathie R. Fraser, Carolyn S. Hulett, and Ann Yaeger of the USGS for their assistance.

**John P. Snyder
Philip M. Voxland**

=====

Introduction

The puzzling world of map projections has fascinated humans for at least 2,500 years. Perhaps no single aspect of the cartographic discipline has held the cartographer's interest so consistently throughout history. How does one portray on a nonspherical surface the essentially spherical shape of the planet on which we live? It cannot be done without distortion, but the number of ways in which that distortion can be handled is infinite.

Although the subject is inherently mathematical and very complex in some instances, it also is highly visual and readily calls forth human reactions ranging from near disbelief to unquestioned acceptance as truth, from confusion to a clear understanding of information. Map projections have been devised for propaganda purposes, as evidence in court, for use in detailed planning, and to attract attention.

Most people are affected more than they realize by map projections. Cognition of the relative areas and shapes on the Earth's surface and knowledge (correct or incorrect) of angles, sizes, distances, and other relationships among places on the globe are determined by the cartographer's choice of map projections. "An Album of Map Projections" should enable readers to sharpen and in some cases to correct their knowledge of the distances, directions, and relative areas of the world.

Although there has been no dearth of valuable books describing a wide range of map projections, "An Album of Map Projections" is the first attempt to set forth in a comprehensive manner most of the useful and commonly known map projections. Consistent and concise non-mathematical textual descriptions are accompanied by standardized visual portrayals. Descriptions of graticules, scale, distortion characteristics, and aspects are carefully worded. Supplemental information on usage, origin, and similar projections is also presented.

No other collection of world portrayals is comparable, because of the standard-format illustrations and the accompanying distortion diagrams presented in this volume. These illustrations will be useful for classroom teaching, for selecting a projection to fit an output format, and for making visual comparisons between projections.

An appendix contains forward formulas for all but a few highly complex projections to transform from latitude and longitude positions on a sphere to rectangular coordinates. Inverse formulas and formulas treating the Earth as an ellipsoid have not been included. Standardized notation is introduced and used throughout.

Finally, it must be noted that this volume could not have been realistically produced without the aid of a computer. Although a compendium of this nature has been needed for decades (to avoid some of the confusion resulting from identical projections' being called by different names, if nothing else), the enormity of manually compiling the necessary information prevented its production. The computer's ability to calculate and plot the portrayals made this album possible.

Joel L. Morrison

Glossary

Aspect Conceptual placement of a projection system in relation to the Earth's axis (direct, normal, polar, equatorial, oblique, and so on).

Authalic projection See Equal-area projection.

Azimuthal projection Projection on which the azimuth or direction from a given central point to any other point is shown correctly. Also called a zenithal projection. When a pole is the central point, all meridians are spaced at their true angles and are straight radii of concentric circles that represent the parallels.

Central meridian Meridian passing through the center of a projection, often a straight line about which the projection is symmetrical.

Central projection Projection in which the Earth is projected geometrically from the center of the Earth onto a plane or other surface. The Gnomonic and Central Cylindrical projections are examples.

Complex algebra Branch of algebra that deals with complex numbers (combinations of real and imaginary numbers using the square root of -1).

Complex curves Curves that are not elementary forms such as circles, ellipses, hyperbolas, parabolas, and sine curves.

Composite projection Projection formed by connecting two or more projections along common lines such as parallels of latitude, necessary adjustments being made to achieve fit. The Goode Homolosine projection is an example.

Conformal projection Projection on which all angles at each point are preserved. Also called an orthomorphic projection.

Conceptually projected Convenient way to visualize a projection system, although it may not correspond to the actual mathematical projection method.

Conic projection Projection resulting from the conceptual projection of the Earth onto a

tangent or secant cone, which is then cut lengthwise and laid flat. When the axis of the cone coincides with the polar axis of the Earth, all meridians are straight equidistant radii of concentric circular arcs representing the parallels, but the meridians are spaced at less than their true angles. Mathematically, the projection is often only partially geometric.

Constant scale Linear scale that remains the same along a particular line on a map, although that scale may not be the same as the stated or nominal scale of the map.

Conventional aspect See Direct aspect.

Correct scale Linear scale having exactly the same value as the stated or nominal scale of the map, or a scale factor of 1.0. Also called true scale.

Cylindrical projection Projection resulting from the conceptual projection of the Earth onto a tangent or secant cylinder, which is then cut lengthwise and laid flat. When the axis of the cylinder coincides with the axis of the Earth, the meridians are straight, parallel, and equidistant, while the parallels of latitude are straight, parallel, and perpendicular to the meridians. Mathematically, the projection is often only partially geometric.

Direct aspect Form of a projection that provides the simplest graticule and calculations. It is the polar aspect for azimuthal projections, the aspect having a straight Equator for cylindrical and pseudocylindrical projections, and the aspect showing straight meridians for conic projections. Also called conventional or normal aspect.

Distortion Variation of the area or linear scale on a map from that indicated by the stated map scale, or the variation of a shape or angle on a map from the corresponding shape or angle on the Earth.

Ellipsoid When used to represent the Earth, a

solid geometric figure formed by rotating an ellipse about its minor (shorter) axis. Also called spheroid.

Equal-area projection Projection on which the areas of all regions are shown in the same proportion to their true areas. Also called an equivalent or authalic projection. Shapes may be greatly distorted.

Equatorial aspect Aspect of an azimuthal projection on which the center of projection or origin is some point along the Equator. For cylindrical and pseudocylindrical projections, this aspect is usually called conventional, direct, normal, or regular rather than equatorial.

Equidistant projection Projection that maintains constant scale along all great circles from one or two points. When the projection is centered on a pole, the parallels are spaced in proportion to their true distances along each meridian.

Equivalent projection See Equal-area projection.

Flat-polar projection Projection on which, in direct aspect, the pole is shown as a line rather than as a point.

Free of distortion Having no distortion of shape, area, or linear scale. On a flat map, this condition can exist only at certain points or along certain lines.

Geometric projection See Perspective projection.

Globular projection Generally, a nonazimuthal projection developed before 1700 on which a hemisphere is enclosed in a circle and meridians and parallels are simple curves or straight lines.

Graticule Network of lines representing a selection of the Earth's parallels and meridians.

Great circle Any circle on the surface of a sphere, especially when the sphere represents the Earth, formed by the intersection of the surface with a plane passing through

the center of the sphere. It is the shortest path between any two points along the circle and therefore important for navigation. All meridians and the Equator are great circles on the Earth taken as a sphere.

Indicatrix Circle or ellipse having the same shape as that of an infinitesimally small circle (having differential dimensions) on the Earth when it is plotted with finite dimensions on a map projection. Its axes lie in the directions of and are proportional to the maximum and minimum scales at that point. Often called a Tissot indicatrix after the originator of the concept.

Interrupted projection Projection designed to reduce peripheral distortion by making use of separate sections joined at certain points or along certain lines, usually the Equator in the normal aspect, and split along lines that are usually meridians. There is normally a central meridian for each section.

Large-scale mapping Mapping at a scale larger than about 1:75,000, although this limit is somewhat flexible.

Latitude (geographic) Angle made by a perpendicular to a given point on the surface of a sphere or ellipsoid representing the Earth and the plane of the Equator (+ if the point is north of the Equator, - if it is south). One of the two common geographic coordinates of a point on the Earth.

Latitude of opposite sign See Parallel of opposite sign.

Limiting forms Form taken by a system of projection when the parameters of the formulas defining that projection are allowed to reach limits that cause it to be identical with another separately defined projection.

Longitude Angle made by the plane of a meridian passing through a given point on the Earth's surface and the plane of the (prime) meridian passing through Greenwich, England, east or west to 180° (+ if the point is

east, - if it is west). One of the two common geographic coordinates of a point on the Earth.

Loxodrome Complex curve (a spherical helix) on the Earth's surface that crosses every meridian at the same oblique angle; in navigation, called a rhumb line. A navigator can proceed between any two points along a rhumb line by maintaining a constant bearing. A loxodrome is a straight line on the Mercator projection.

Meridian Reference line on the Earth's surface formed by the intersection of the surface with a plane passing through both poles and some third point on the surface. This line is identified by its longitude. On the Earth as a sphere, this line is half a great circle; on the Earth as an ellipsoid, it is half an ellipse.

Minimum-error projection Projection having the least possible total error of any projection in the designated classification, according to a given mathematical criterion. Usually, this criterion calls for the minimum sum of squares of deviations of linear scale from true scale throughout the map ("least squares").

Nominal scale Stated scale at which a map projection is constructed.

Normal aspect See Direct aspect.

Oblique aspect Aspect of a projection on which the center of projection or origin is located at a point which is neither at a pole nor along the Equator.

Orthoapsidal projection Projection on which the surface of the Earth taken as a sphere is transformed onto a solid other than the sphere and then projected orthographically and obliquely onto a plane for the map.

Orthographic projection Specific azimuthal projection or a type of projection in which the Earth is projected geometrically onto a

surface by means of parallel projection lines.

Orthomorphic projection See Conformal projection.

Parallel Small circle on the surface of the Earth formed by the intersection of the surface of the reference sphere or ellipsoid with a plane parallel to the plane of the Equator. This line is identified by its latitude. The Equator (a great circle) is usually also treated as a parallel.

Parallel of opposite sign Parallel that is equally distant from but on the opposite side of the Equator. For example, for lat 30° N. (or +30°), the parallel of opposite sign is lat 30° S. (or -30°). Also called latitude of opposite sign.

Parameters Values of constants as applied to a map projection for a specific map; examples are the values of the scale, the latitudes of the standard parallels, and the central meridian. The required parameters vary with the projection.

Perspective projection Projection produced by projecting straight lines radiating from a selected point (or from infinity) through points on the surface of a sphere or ellipsoid and then onto a tangent or secant plane. Other perspective maps are projected onto a tangent or secant cylinder or cone by using straight lines passing through a single axis of the sphere or ellipsoid. Also called geometric projection.

Planar projection Projection resulting from the conceptual projection of the Earth onto a tangent or secant plane. Usually, a planar projection is the same as an azimuthal projection. Mathematically, the projection is often only partially geometric.

Planimetric map Map representing only the horizontal positions of features (without their elevations).

Glossary

Polar aspect Aspect of a projection, especially an azimuthal one, on which the Earth is viewed from the polar axis. For cylindrical or pseudocylindrical projections, this aspect is called transverse.

Polyconic projection Specific projection or member of a class of projections in which, in the normal aspect, all the parallels of latitude are nonconcentric circular arcs, except for a straight Equator, and the centers of these circles lie along a central axis.

Pseudoconic projection Projection that, in the normal aspect, has concentric circular arcs for parallels and on which the meridians are equally spaced along the parallels, like those on a conic projection, but on which meridians are curved.

Pseudocylindrical projection Projection that, in the normal aspect, has straight parallel lines for parallels and on which the meridians are (usually) equally spaced along parallels, as they are on a cylindrical projection, but on which the meridians are curved.

Regional map Small-scale map of an area covering at least 5 or 10 degrees of latitude and longitude but less than a hemisphere.

Regular aspect See Direct aspect.

Retroazimuthal projection Projection on which the direction or azimuth from every point on the map to a given central point is shown correctly with respect to a vertical line parallel to the central meridian. The reverse of an azimuthal projection.

Rhumb line See Loxodrome.

Scale Ratio of the distance on a map or globe to the corresponding distance on the Earth; usually stated in the form 1:5,000,000, for example.

Scale factor Ratio of the scale at a particular location and direction on a map to the stated scale of the map. At a standard parallel,

or other standard line, the scale factor is 1.0.

Secant cone, cylinder, or plane A secant cone or cylinder intersects the sphere or ellipsoid along two separate lines; these lines are parallels of latitude if the axes of the geometric figures coincide. A secant plane intersects the sphere or ellipsoid along a line that is a parallel of latitude if the plane is at right angles to the axis.

Similar (projection) Subjective and qualitative term indicating a moderate or strong resemblance.

Singular points Certain points on most but not all conformal projections at which conformality fails, such as the poles on the normal aspect of the Mercator projection.

Small circle Circle on the surface of a sphere formed by intersection with a plane that does not pass through the center of the sphere. Parallels of latitude other than the Equator are small circles on the Earth taken as a sphere.

Small-scale mapping Mapping at a scale smaller than about 1:1,000,000, although the limiting scale sometimes has been made as large as 1:250,000.

Spheroid See Ellipsoid.

Standard parallel In the normal aspect of a projection, a parallel of latitude along which the scale is as stated for that map. There are one or two standard parallels on most cylindrical and conic map projections and one on many polar stereographic projections.

Stereographic projection Specific azimuthal projection or type of projection in which the Earth is projected geometrically onto a surface from a fixed (or moving) point on the opposite face of the Earth.

Tangent cone or cylinder Cone or cylinder that just touches the sphere or ellipsoid along a single line. This line is a parallel of latitude

if the axes of the geometric figures coincide.

Thematic map Map designed to portray primarily a particular subject, such as population, railroads, or croplands.

Tissot indicatrix See Indicatrix.

Topographic map Map that usually represents the vertical positions of features as well as their horizontal positions.

Transformed latitudes, longitudes, or poles

Graticule of meridians and parallels on a projection after the Earth has been turned with respect to the projection so that the Earth's axis no longer coincides with the conceptual axis of the projection. Used for oblique and transverse aspects of many projections.

Transverse aspect Aspect of a map projection on which the axis of the Earth is rotated so that it is at right angles to the conceptual axis of the map projection. For azimuthal projections, this aspect is usually called equatorial rather than transverse.

True scale See Correct scale.

Zenithal projection See Azimuthal projection.

Guide to Selecting Map Projections

This publication displays the great variety of map projections from which the choice for a particular map can be made. Although most of the projections illustrated in this album have been in existence for many years, cartographers have been content with using only a few of them.

This past lack of innovation is easily understood. The difficulty and expense of creating a new base map using a different projection often outweighed the perceived benefits. The advent of computer-assisted cartography has now made it much easier to prepare base maps. A map can now be centered anywhere on the globe, can be drawn according to any one of many projection formulas, and can use cartographic data files (coastline boundaries, for example) having a level of generalization appropriate to the scale of the map.

However, now that more choices can be made, the actual decision may be more difficult. Some of the traditional rules for selection are now merely guidelines. This chapter discusses classifications of maps and presents well-established principles for their evaluation. Technical and mathematical methods also exist for determining exactly the characteristics of a specific projection, but those methods are not used here.

First, we should define what is meant by a map projection. A projection is a systematic transformation of the latitudes and longitudes of locations on the surface of a sphere or an ellipsoid into locations on a plane. That is, locations in three-dimensional space are made to correspond to a two-dimensional representation.

If we assume for simplicity that a globe, which is a sphere, can perfectly represent the surface of the Earth, then all of the following characteristics must be true of that globe:

1. Areas are everywhere correctly represented.

2. All distances are correctly represented.
3. All angles are correctly represented.
4. The shape of any area is faithfully represented.

When the sphere is projected onto a plane, the map will no longer have all of these characteristics simultaneously. Indeed, the map may have none of them. One way of classifying maps and their projections is through terms describing the extent to which the map preserves any of those properties.

Properties of Map Projections

An **equal-area** map projection correctly represents areas of the sphere on the map. If a coin is placed on any area of such a map, it will cover as much of the area of the surface of the sphere as it would if it were placed elsewhere on the map. When this type of projection is used for small-scale maps showing larger regions, the distortion of angles and shapes increases as the distance of an area from the projection origin increases.

An **equidistant** map projection is possible only in a limited sense. That is, distances can be shown at the nominal map scale along a line from only one or two points to any other point on the map. The focal points usually are at the map center or some central location. The term is also often used to describe maps on which the scale is shown correctly along all meridians.

An **azimuthal** map likewise is limited in the sense that it can correctly show directions or angles to all other points on the map only with respect to one (or rarely two) central point(s).

A **conformal** map is technically defined as a map on which all angles at infinitely small locations are correctly depicted. A conformal projection increasingly distorts areas away from the map's point or lines of true scale and

increasingly distorts shapes as the region becomes larger but distorts the shapes of moderately small areas only slightly.

Consistent with these definitions, maps simultaneously exhibiting several of these properties can be devised:

	Conformal	Equal area	Equidistant	Azimuthal
Conformal	—	No	No	Yes
Equal area	No	—	No	Yes
Equidistant	No	No	—	Yes
Azimuthal	Yes	Yes	Yes	—

In summary, any map will distort areas, angles, directions, or distances to some extent. Since all of these distortions can be measured or estimated, one rigorous selection rule would be "to select a projection in which the extreme distortions are smaller than would occur in any other projection used to map the same area" (Maling, 1973, p. 159).

A map projection may have none of these general properties and still be satisfactory. A map projection possessing one of these properties may nevertheless be a poor choice. As an example, the Mercator projection continues to be used inappropriately for worldwide thematic data. The Mercator map projection is conformal and has a valid use in navigation but very seriously distorts areas near the poles, which it cannot even show. It should not be used (although it frequently is) for depicting general information or any area-related subjects.

Classification Based on Construction

From the perspective of design as well as distortion reduction, a projection may be selected because of the characteristic curves formed by the meridians and parallels. L.P. Lee preferred terms based on the pattern formed by the meridians and parallels in the normal aspect or orientation. The following definitions

Guide to Selecting Map Projections

are quoted from his 1944 paper, "The nomenclature and classification of map projections" (Lee, 1944, p. 193):

Cylindric: Projections in which the meridians are represented by a system of equidistant parallel straight lines, and the parallels by a system of parallel straight lines at right angles to the meridians.

Pseudocylindric: Projections in which the parallels are represented by a system of parallel straight lines, and the meridians by concurrent curves.

Conic: Projections in which the meridians are represented by a system of equally inclined concurrent straight lines, and the parallels by concentric circular arcs, the angle between any two meridians being less than their true difference of longitude.

Pseudoconic: Projections in which the parallels are represented by concentric circular arcs, and the meridians by concurrent curves.

Polyconic: Projections in which the parallels are represented by a system of nonconcentric circular arcs with their centres lying on the straight line representing the central meridian.

Azimuthal: Projections in which the meridians are represented by a system of concurrent straight lines inclined to each other at their true difference of longitude, and the parallels by a system of concentric circles with their common centre at the point of concurrency of the meridians.

D.H. Maling used an additional term consistent with Lee's definitions:

Pseudoazimuthal: Projections comprised of "concentric circular parallels and curved meridians which converge at the pole at their true angular values" (Maling, 1960, p. 209).

Finally, a small class of projections has also been identified by the term **retroazimuthal**, denoting projections in which the direction to a

central point from every point on the map is shown correctly.

Philosophy of Map-Projection Selection

On the basis of these concepts, three traditional rules for choosing a map projection were at one time recommended:

1. For low-latitude areas: cylindrical.
2. For middle-latitude areas: conical.
3. For polar regions: azimuthal.

Inherent in these guidelines was the idea that it would be difficult to recenter a map so that the area of main interest was near the area of the map that has the least areal or angular distortion. On the other hand, it is now much less difficult than it formerly was to rotate latitude and longitude coordinates so that any point is moved to any other new location. As a result, a projection does not need to be rejected merely because the only available copy is centered inappropriately for a new application. In fact, many projections permit the user to alter the form of the map to reduce the distortions within a certain area. Most commonly, such alteration is accomplished by establishing standard lines along which distortion is absent; often, these lines are parallels of latitude. However, most properties of the map projection are affected when a standard line is changed. Another way to alter the relationships on a map is by using different aspects, which involves moving the center of a projection from the normal position at a pole or along the Equator to some other position.

"Map makers must first define the purpose of a map which in turn can provide answers to many questions essential to decisions concerning the projections" (Hsu, 1981, p. 170). When a map requires a general property, the choice of a projection becomes limited. For example, because conformal projections correctly show angles at every location, they are advisable for

maps displaying the flow of oceanic or atmospheric currents. The risk of using a conformal projection for a worldwide map is that the distortion of areas greatly enlarges the outer boundaries, and a phenomenon may seem to take on an importance that the mapmaker did not intend. Equal-area maps should be considered for displaying area-related subjects or themes, such as crop-growing regions.

"Once the purpose of a map has been decided, the geographical area to be included on the map must be determined. Let us call it the *map area* for brevity. The map area can be a region or the entire world. The larger the area covered, the greater is the earth's curvature involved on the map. . . . If the map area is a region, then its shape, size, and location are important determinants in making decisions concerning projections" (Hsu, 1981, p. 172).

Bearing in mind these determinants, mapmakers can apply traditional rules of choice, such as those mentioned above, or they can study the patterns of distortion associated with particular projections. For example, azimuthal projections have a circular pattern for lines of constant distortion characteristics, centered on the map origin or projection center. Thus, if an area is approximately circular and if its center is made the origin of the projection, it is possible to create a map that minimizes distortion for that map area. Ideally, the general shape of a geographic region should be matched with the distortion pattern of a specific projection.

An appropriate map projection can be selected on the basis of these principles and classifications. Although there may be no absolutely correct choice, it is clearly possible to make a bad judgment.

Edge Matching

One problem encountered by many map users is the edge matching of adjacent regions. For two or more maps to fit exactly along their

edges, whether these be common meridians, parallels, or rectangular coordinate gridlines, not only must they be cast on the same projection and at the same scale, but the projection for each map must also have the same critical parameters. The critical parameters, which vary somewhat with each projection, are those specifications that affect the shape and size but not the position or orientation of a projection.

The Mercator projection, which is commonly used in edge-matching operations, must be based on the same ellipsoid or sphere and must use the same scale at the same latitude for each map. Conic projections such as the Albers, Lambert Conformal, or Equidistant will match only if the standard parallels and the ellipsoid are the same for each map. For the Mercator and conic projections, the central meridian and the latitude of origin do not have to be identical for edge matching. For the Transverse Mercator projection, however, the ellipsoid, the central meridian, and the scale along the central meridian must be the same, although the latitude of origin may vary.

In practice, edge matching is hampered by the dimensional instability of the paper or other material that the maps are printed on and also by cartographic or surveying errors in extending roads and streams to the edges of the maps. The differences between the projections used for large-scale maps may be small enough to permit a satisfactory fit, depending on the required accuracy and purpose of the maps.

Distortion Diagrams

The most important characteristics of a map projection are the magnitude of distortion and the effect of that distortion on the intended use of a map. To assist in evaluating these features, many of the graticules for the projections described herein are presented on two illustrations.

The larger of the two plots uses the World Data Bank I shoreline file (occasionally including national boundaries). It also uses the same central meridian (90° W.), unless doing so would defeat the purpose of the projection, as it would with the Briesemeister. Some evaluation of distortion can be achieved by comparing familiar shapes of islands and shorelines. A more uniform comparison can be made by using the smaller plots, which include no shorelines but which show Tissot indicatrices at every 30° of latitude and longitude, except at the poles (arbitrarily omitted because of plotting complications).


The Tissot indicatrix, devised by French cartographer Nicolas Auguste Tissot in the 19th century, shows the shape of infinitesimally small circles on the Earth as they appear when they are plotted by using a fixed finite scale at the same locations on a map. Every circle is plotted as a circle or an ellipse or, in extreme cases, as a straight line. On an equal-area projection, all these ellipses and circles are shown as having the same area. The flattening of the ellipse shows the extent of local shape distortion and how much the scale is changed and in what direction. On conformal map projections, all indicatrices remain circles, but the areas change. On other projections, both the areas and the shapes of the indicatrices change.

For example, for the conformal Mercator


projection, all indicatrices are circles (fig. 1A, p. 10). Scale is constant along every parallel; thus, every circle along a given parallel is the same size. On the other hand, the farther the parallel from the Equator, the greater the size of the circle. Quantitatively, the diameter of the circle is proportional to the linear scale at that point, and the area of the circle is proportional to the area scale. The shape of the circle also indicates that the (linear) scale is the same in every direction at a given point.

Figure 13A (p. 37) represents indicatrices for the equal-area Sinusoidal projection. The indicatrices are circles along the Equator and along the central meridian, because there is no local shape distortion there. As it did in figure 1A, the shape of the circles indicates that the scale is the same in all directions at each location. The indicatrices elsewhere are ellipses, characterized by varying degrees of flattening and by axes pointed in various directions. Because the areas of all these ellipses are equal, the projection is equal area. The orientation and flattening of a given ellipse provide additional information. The major axis represents the magnitude and direction of the greatest scale at that point. The magnitude can be determined by multiplying the map scale by the ratio of the length of the major axis of the ellipse to the diameter of the circle at a point of no distortion, such as a point along the Equator of this projection. The minor axis represents the magnitude and direction of the smallest scale at that point, determined in the same manner. The angular distortion at each point is similarly indicated by the indicatrix. At every point,

there is no angular distortion in plotting the intersection of two particular lines on the Earth, one in the direction of the major axis of the ellipse and the other in the direction of the minor axis. These two lines intersect at right angles on the Earth (lines AB and CD) and also at right angles on the map (lines A'B' and C'D'). A pair of perpendicular lines inclined at 45° to these lines on the Earth (lines EF and GH) will not intersect at right angles on the map if the indicatrix is an ellipse (lines E'F' and G'H'). This statement is true for every pair of lines other than the two axes mentioned. The scale in these various directions is proportional to the length of the diameter across the ellipse in the same direction. In two particular



An infinitely small circle on the Earth, circumscribed by a square.



The circle shown on the preceding page as plotted on a non-conformal map, circumscribed by a rectangle.

directions, between the directions of the major and minor axes on an elliptical indicatrix of an equal-area map projection, the scale is the same as that of the map.

Figure 32A (p. 82) illustrates distortions on the Robinson, a projection that is neither equal area nor conformal. Both the size and the shape of the indicatrices change. No point is completely free of distortion; therefore, all indicatrices are at least slightly elliptical. Their areas are proportional to the area scale, and linear scale and angular distortion can be interpreted just as they were for the ellipses of the Sinusoidal plot, except that there is no circular indicatrix. The indicatrices as shown in this volume are obviously too small to permit effective quantitative scale and distortion measurements. Nevertheless, they provide more certainty in this evaluation than do the larger plots of shorelines.

MERCATOR Projection

Classifications

Cylindrical
Conformal

Graticule

Meridians: Equally spaced straight parallel lines
Parallels: Unequally spaced straight parallel lines, closest near the Equator, perpendicular to meridians
Poles: Cannot be shown
Symmetry: About any meridian or the Equator

Scale

True along the Equator or along two parallels equidistant from the Equator
Increases with distance from the Equator to infinity at the poles
Constant along any given parallel; same scale at parallel of opposite sign (north +, south -)
Same in all directions near any given point

Distortion

Infinitesimally small circles (indicatrices) of equal size on the globe appear as circles on the map (indicating conformality) but increase in size away from the Equator (indicating area distortion) (fig. 1A). Great distortion of area in polar regions. Conformality (and therefore local angle preservation) fails at the poles.

Other features

All loxodromes or rhumb lines (lines that make equal angles with all meridians and are therefore lines of constant true bearing) are straight lines.

Meridians can be geometrically projected onto a cylinder, the axis of which is the same as that of the globe.

Parallels cannot be geometrically (or perspectively) projected.

Meridians cannot be compressed relative to parallels, as they can on Cylindrical Equal-Area and Equirectangular projections, since conformality would be lost.

Usage

Designed and recommended for navigational usage because of straight rhumb lines; standard for marine charts

Recommended and used for conformal mapping of regions predominantly bordering the Equator

Often and inappropriately used as a world map in atlases and for wall charts. It presents a misleading view of the world because of the excessive distortion of area.

Origin

Presented by Gerardus Mercator (1512–94) of Flanders in 1569 on a large world map "for use in navigation"

Aspects


Normal is described here.

Transverse and Oblique aspects are listed separately (p. 12–15) because of importance and common treatment as separate projections.

Other names

Wright (rare) (after Edward Wright of England, who developed the mathematics in 1599)

Figure 1A.—Mercator projection with Tissot indicatrices, 30° graticule. All indicatrices are circular (indicating conformality), but areas vary.




Similar projections

Central Cylindrical projection (p. 30) also cannot show poles, but it is not conformal, and the spacing of parallels changes much more rapidly.

Miller Cylindrical projection (p. 35) shows the poles, is not conformal, and has more gradual spacing of parallels.

Gall projection (p. 33) shows the poles, is not conformal, and has more gradual spacing of parallels.

Figure 1B.—Mercator projection, with shorelines, 15° graticule. Central meridian 90° W.



TRANSVERSE MERCATOR Projection

Classifications

Transverse aspect of Mercator projection

Cylindrical

Conformal

Graticule

Meridians and parallels: Central meridian, each meridian 90° from central meridian, and the Equator are straight lines. Other meridians and parallels are complex curves, concave toward the central meridian and the nearest pole, respectively.

Poles: Points along the central meridian

Symmetry: About any straight meridian or the Equator

Scale

True along the central meridian or along two straight lines on the map equidistant from and parallel to the central meridian

Constant along any straight line on the map parallel to the central meridian. (These lines are only approximately straight for the projection of the ellipsoid.)

Increases with distance from the central meridian

Becomes infinite 90° from the central meridian

Distortion

At a given distance from the central meridian in figure 2A, the distortion in area is identical with that at the same distance from the Equator in figure 1A.

Other features

Conceptually projected onto a cylinder wrapped around the globe tangent to the central meridian or secant along two small circles equidistant from the central meridian

Cannot be geometrically (or perspectively) projected

Rhumb lines generally are not straight lines.

Usage

Many of the topographic and planimetric map quadrangles throughout the world at scales of 1:24,000 to 1:250,000

Basis for Universal Transverse Mercator (UTM) grid and projection

Basis for State Plane Coordinate System in U.S. States having predominantly north-south extent

Recommended for conformal mapping of regions having predominantly north-south extent

Origin

Presented by Johann Heinrich Lambert (1728–77) of Alsace in 1772. Formulas for ellipsoidal use developed by Carl Friedrich Gauss of Germany in 1822 and by L. Krüger of Germany, L.P. Lee of New Zealand, and others in the 20th century.

Other names

Gauss Conformal (ellipsoidal form only)

Gauss-Krüger (ellipsoidal form only)

Transverse Cylindrical Orthomorphic

Figure 2A.—Transverse Mercator projection, with Tissot indicatrices, 30° graticule.





Figure 2B.—Transverse Mercator projection, with shorelines, 15° graticule. Central meridian 90° E. and W. North Pole at -90° longitude on base projection.



OBLIQUE MERCATOR Projection

Classifications

Oblique aspect of Mercator projection
Cylindrical
Conformal

Graticule

Meridians and parallels: Two meridians 180° apart are straight lines. Other meridians and parallels are complex curves.

Poles: Points not on the central line

Symmetry: About either straight meridian

Scale

True along a chosen central line (a great circle at an oblique angle) or along two straight lines on the map parallel to the central line

Constant along any straight line parallel to the central line

(The scale for the projection of the ellipsoid varies slightly from these patterns.)

Increases with distance from the central line

Becomes infinite 90° from the central line

Distortion

At a given distance from the central line, distortion on figure 3A is the same as that on the regular Mercator projection (fig. 1A).

Other features

Conceptually projected onto a cylinder wrapped around the globe tangent to an oblique great circle or secant along two small circles equidistant from and on each side of the central great circle

Cannot be geometrically (or perspectively) projected

There are various means of adapting to the ellipsoid, but none can simultaneously maintain both perfect conformality and constant scale along the central line.

Usage

Large-scale mapping in Switzerland, Madagascar, and Borneo

Atlas maps of regions having greater extent in an oblique direction, such as Hawaii

Recommended for conformal mapping of regions having predominant extent in oblique direction, neither east-west nor north-south

Origin

Developed for various applications, chiefly large-scale mapping of the ellipsoid, by M. Rosenmund of Switzerland in 1903, J. Laborde of France in 1928, Martin Hotine of England in 1947, and others during the 20th century.

Other names

Rectified Skew Orthomorphic (when using Hotine's formulas)

Laborde (when using Laborde's formulas)

Hotine Oblique Mercator (when using Hotine's formulas)

Oblique Cylindrical Orthomorphic

Limiting forms

Mercator (p. 10), if the Equator is the central line

Transverse Mercator (p. 12), if a meridian is the central line

Figure 3A.—Oblique Mercator projection, with Tissot indicatrices, 30° graticule. North Pole at +30° latitude, -90° longitude on base projection.





Figure 3B.—Oblique Mercator projection, with shorelines, 15° graticule. Central meridian 90° W. North Pole at +30° latitude, -90° longitude on base projection.



LAMBERT CYLINDRICAL EQUAL-AREA Projection

Classifications

Cylindrical
Equal area
Perspective

Graticule

Meridians: Equally spaced straight parallel lines 0.32 as long as the Equator.
Parallels: Unequally spaced straight parallel lines, farthest apart near the Equator, perpendicular to meridians
Poles: Straight lines equal in length to the Equator
Symmetry: About any meridian or the Equator

Scale

True along the Equator
Increases with distance from the Equator in the direction of parallels and decreases in the direction of meridians to maintain equal area
Same scale at the parallel of opposite sign

Distortion

Infinitesimally small circles (indicatrices) of equal size on the globe are ellipses except at the Equator, where they are circles (fig. 4A). The areas of all the indicatrices are the same. Thus, there is shape distortion but no area distortion. Shape distortion in polar regions is extreme.

Other features

Simple graticule, perspectively projected in lines perpendicular to the axis onto a cylinder wrapped around the globe tangent to the Equator

Usage

Minimal except to describe basic principles in map projection texts




Figure 4A.—Lambert Cylindrical Equal-Area projection with Tissot indicatrices, 30° graticule. Standard parallel 0°. All ellipses have the same area, but shapes vary.

Prototype for Behrmann and other modified cylindrical equal-area projections (fig. 5B)

Recommended for equal-area mapping of regions predominantly bordering the Equator

Origin

Presented by Johann Heinrich Lambert (1728–77) of Alsace in 1772

Aspects

Normal is described here.

Transverse and oblique aspects are rarely used (figs. 4C, 4D) but are recommended for equal-area mapping of predominantly north-south regions or regions extending obliquely.

Other names

Cylindrical Equal-Area

Similar projections


If meridians are compressed relative to parallels and if the spacing of parallels is increased in inverse proportion, other cylindrical equal-area projections result (fig.

5B), and the standard parallel changes. The extreme case, in which the poles are standard parallels, consists of a single vertical line, infinitely long.

Named examples are as follows:

Behrmann
Gall Orthographic
Trystan Edwards
Peters
(See Behrmann Cylindrical Equal-Area projection, p. 19, for the differences.)

Figure 4B.—Lambert Cylindrical Equal-Area projection with shorelines, 15° graticule. Standard parallel 0°. Central meridian 90° W.



LAMBERT CYLINDRICAL EQUAL-AREA Projection

Figure 4C.—Transverse Lambert Cylindrical Equal-Area projection with shorelines, 15° graticule. Central meridians 90° E. and W., at true scale. North Pole at +90° longitude on base projection.





Figure 4D.—Oblique Lambert Cylindrical Equal-Area projection with shorelines, 15° graticule. Central meridian 90° W. Central great circle at true scale through latitude 60° N., longitude 180° and latitude 60° S., longitude 0°. North Pole at +30° latitude, -90° longitude on base projection.



BEHRMANN CYLINDRICAL EQUAL-AREA Projection

Classifications

Cylindrical
Equal area
Perspective

Graticule

Meridians: Equally spaced straight parallel lines 0.42 as long as the Equator.
Parallels: Unequally spaced straight lines, farthest apart near the Equator, perpendicular to meridians
Poles: Straight lines equal in length to the Equator
Symmetry: About any meridian or the Equator

Scale

True along latitudes 30° N. and S.
 Too small along the Equator but too large at the Equator along meridians
 Increases with distance from the Equator in the direction of parallels and decreases in the direction of meridians to maintain equal area
 Same scale at the parallel of opposite sign

Distortion

In contrast to figure 4A, figure 5A shows indicatrices as circles at latitudes 30° N. and S., where there is no distortion, instead of at the Equator. All others appear as ellipses, but their areas remain the same. They are compressed east to west and lengthened north to south between latitudes 30° N. and S. The opposite is true poleward of these latitudes.

Other features

Same as the Lambert Cylindrical Equal-Area projection except for horizontal compression and vertical expansion to achieve no distortion




Figure 5A.—Behrmann Cylindrical Equal-Area projection with Tissot indicatrices, 30° graticule. Standard parallels 30° N. and S.

at latitudes 30° N. and S. instead of at the Equator

Equivalent to a projection of the globe using parallel lines of projection onto a cylinder secant at 30° N. and S.

Origin

Presented by Walter Behrmann (1882–1955) of Berlin in 1910

Similar projections

Lambert Cylindrical Equal-Area (p. 16) by Johann Heinrich Lambert in 1772 (Standard parallel: Equator)

Gall Orthographic (fig. 5C) by James Gall in 1855 (Standard parallels: 45° N. and S.)

Trystan Edwards in 1953 (Standard parallels 37°24' N. and S.)

Peters (fig. 5C) by Arno Peters in 1967 (Standard parallels: approximately 45° N. and S., thus essentially identical with the Gall Orthographic)

BEHRMANN CYLINDRICAL EQUAL-AREA Projection

Figure 5B.—Behrmann Cylindrical Equal-Area projection with shorelines, 15° graticule. Central meridian 90° W. Standard parallels 30° N. and S.




Figure 5C.—Gall Orthographic or Peters projection with shorelines, 15° graticule. A cylindrical equal-area projection with standard parallels 45° N. and S. Central meridian 90° W.




PLATE CARREE Projection

Classifications

Cylindrical
Equidistant

Graticule

Meridians: Equally spaced straight parallel lines half as long as the Equator.

Parallels: Equally spaced straight parallel lines, perpendicular to and having same spacing as meridians

Poles: Straight lines equal in length to the Equator

Symmetry: About any meridian or the Equator

Scale

True along the Equator and along all meridians

Increases with the distance from the Equator along parallels

Constant along any given parallel; same scale at the parallel of opposite sign

Distortion

Infinitesimally small circles of equal size on the globe (indicatrices) are ellipses except along the Equator, where they remain circles (fig. 6A). Areas of the ellipses also vary. Thus, there is distortion of both shape and area.

Other features

Most simply constructed graticule of any projection


Conceptually projected onto a cylinder wrapped around the globe tangent to the Equator

Not perspective

Usage

Many maps during the 15th and 16th centuries

Figure 6A.—Plate Carrée projection with Tissot indicatrices, 30° graticule.



Simple outline maps of regions or of the world or index maps

Used only for the Earth taken as a sphere

Origin

May have been originated by Eratosthenes (275?–195? B.C.)

Marinus of Tyre also credited with its invention about A.D. 100

Aspects

Normal is described here.

Transverse aspect is the Cassini projection (figs. 8A, 8B), which is also applied to the ellipsoid.

Oblique aspect is rarely used (see fig. 9B).

Other names


Simple Cylindrical

Equidistant Cylindrical (particular form)

Similar projections

If meridians are compressed relative to parallels, the Equirectangular projection (p. 24) results.

Figure 6B.—Plate Carrée projection with shorelines, 15° graticule. Central meridian 90° W.



EQUIRECTANGULAR Projection

Classifications

Cylindrical
Equidistant

Graticule

Meridians: Equally spaced straight parallel lines more than half as long as the Equator.

Parallels: Equally spaced straight parallel lines, perpendicular to and having wider spacing than meridians

Poles: Straight lines equal in length to the Equator

Symmetry: About any meridian or the Equator

Scale

True along two standard parallels equidistant from the Equator and along all meridians

Too small along the Equator but increases with distance from the Equator along the parallels

Constant along any given parallel; same scale at the parallel of opposite sign

Distortion

Infinitesimally small circles on the globe (indicatrices) are circles on the map at latitudes 30° N. and S. for this choice of standard parallels (fig. 7A). Elsewhere, area and local shape are distorted.

Other features

Simple modification of Plate Carrée (p. 22) having east-west compression

Conceptually projected onto a cylinder secant to the globe along the chosen standard parallels

Not perspective




Figure 7A.—Equirectangular projection with Tissot indicatrices, 30° graticule. Standard parallels 30° N. and S.

Usage

Simple outline maps of regions or of the world or for index maps

Used only in the spherical form

Die Rechteckige Plattkarte

Projection of Marinus

Gall Isographic (if standard parallels are latitudes 45° N. and S.)

Origin

Marinus of Tyre about A.D. 100


Other names

Equidistant Cylindrical

Rectangular

La Carte Parallélogrammatique

Figure 7B.—Equirectangular projection with shorelines, 15° graticule. Central meridian 90° W. Standard parallels 30° N. and S.



CASSINI Projection

Classifications

Transverse aspect of Plate Carrée (p. 22)

Cylindrical

Equidistant

Graticule

Central meridian, each meridian 90° from central meridian, and the Equator are straight lines.

Other meridians and parallels are complex curves, concave toward the central meridian and the nearest pole, respectively.

Poles: Points along the central meridian

Symmetry: About any straight meridian or the Equator

Scale

True along the central meridian and along any straight line perpendicular to the central meridian

Increases with distance from the central meridian, along a direction parallel to the central meridian

Distortion

Function of the distance from the central meridian. No distortion occurs along the central meridian, but there is both area and local shape distortion elsewhere. Long horizontal straight lines near the upper and lower limits of figure 8A represent infinitesimal circles on the globe 90° from the central meridian.

Other features

Conceptually projected onto a cylinder tangent to the globe at the central meridian




Figure 8A.—Cassini projection with Tissot indicatrices, 30° graticule.

Can be compressed north-south to provide a Transverse Equirectangular projection, but rarely done

Usage


Topographic mapping (ellipsoidal form) of British Isles before the 1920's; replaced by the Transverse Mercator

Topographic mapping of a few countries currently

Origin

Developed by César François Cassini de Thury (1714–84) for topographic mapping of France in the middle 18th century

Figure 8B.—Cassini projection with shorelines, 15° graticule. Central meridian 90° W.



OBLIQUE PLATE CARREE Projection

Classifications

Oblique aspect of Plate Carrée

Cylindrical

Equidistant

Graticule

Two meridians 180° apart are straight lines.
Other meridians and parallels are complex curves.

Poles: Points away from central line

Symmetry: About either of the straight meridians

Scale

True along the chosen central line (a great circle at an oblique angle) and along any straight line perpendicular to the central line. Increases with distance from the central line along a direction parallel to the central line.

Distortion

At a given distance from the central line, distortion on figure 9A is the same as that on the Plate Carrée (fig. 6A).

Limiting forms

Plate Carrée (p. 22), if the Equator is the central line

Cassini (p. 26), if a meridian is the central line





Figure 9A.—Oblique Plate Carrée projection with Tissot indicatrices, 30° graticule. North Pole at $+30^\circ$ latitude, 0° longitude on base projection.

Figure 9B.—Oblique Plate Carrée projection with shorelines, 15° graticule. Central meridian 90° W. North Pole at +30° latitude, 0° longitude on base projection.



CENTRAL CYLINDRICAL Projection

Classifications

Cylindrical

Perspective

Neither conformal nor equal area

Graticule

Meridians: Equally spaced straight parallel lines

Parallels: Unequally spaced straight parallel lines, closest near the Equator, but spacing increases poleward at a much greater rate than it does on the Mercator. Perpendicular to meridians.

Poles: Cannot be shown

Symmetry: About any meridian or the Equator

Scale

True along the Equator

Increases with distance from the Equator to infinity at the poles

Changes with direction at any given point, except at the Equator, but scale in a given direction is constant at any given latitude or at the latitude of opposite sign

Distortion

Shape, area, and scale distortion increase rapidly away from the Equator, where there is no distortion (fig. 10A).

Other features

Projection is produced geometrically by projecting the Earth's surface perspective

from its center onto a cylinder tangent at the Equator. Should not be confused with the Mercator (p. 10), which is not perspective.

Usage

Distortion is too great for any usage except showing the appearance of the Earth when so projected and contrasting with the Mercator.

Origin

Uncertain

Aspects

Normal is described above.

Transverse aspect (fig. 10C) is called the Wetch projection, since it was discussed by J. Wetch in the early 19th century.

Other names

Simple Perspective Cylindrical

Similar projections

Mercator projection (p. 10), which is not perspective, also cannot show the poles, but the poleward increase in the spacing of the parallels does not occur as rapidly as it does on the Central Cylindrical projection.

Gall (p. 33) and other perspective cylindrical projections can be produced by moving the point of perspective away from the center of the Earth. The poles can then be shown.

Gnomonic projection (p. 116) is projected perspective from the center of the Earth onto a tangent plane rather than a cylinder and is very different in appearance.

Figure 10A.—Central Cylindrical projection with Tissot indicatrices, 30° graticule.





Figure 10B.—Central Cylindrical projection with shorelines, 15° graticule. Central meridian 90° W.



CENTRAL CYLINDRICAL Projection




Figure 10C.—Transverse Central Cylindrical projection with shorelines, 15° graticule. Central meridian 90° E. and W. at true scale.

GALL Projection

Classifications

Cylindrical

Perspective

Neither conformal nor equal area

Graticule

Meridians: Equally spaced straight parallel lines 0.77 as long as the Equator

Parallels: Unequally spaced straight parallel lines, closest near the Equator, but spacing does not increase poleward as fast as it does on the Mercator. Perpendicular to meridians.

Poles: Straight lines equal in length to the Equator

Symmetry: About any meridian or the Equator

Scale

True along latitudes 45° N. and S. in all directions.

Constant in any given direction along any other given latitude or the latitude of opposite sign

Changes with latitude and direction but is always too small between latitudes 45° N. and S. and too large beyond them

Distortion

None at latitudes 45° N. and S. Shape, area, and scale distortion increases moderately away from these latitudes but becomes severe at poles (fig. 11A).

Other features

Projection is produced geometrically by projecting the Earth perspective from the point on the Equator opposite a given meridian onto a secant cylinder cutting the globe at latitudes 45° N. and S.

Usage

World maps in British atlases and some other

atlases, as a projection somewhat resembling the Mercator but having less distortion of area and scale near the poles

Origin

Presented by James Gall of Edinburgh in 1855 as his Stereographic, which he preferred to his Orthographic (equal-area) and Isographic (equidistant) cylindrical projections presented at the same time and also based on cylinders secant at latitudes 45° N. and S.

Aspects

Only the normal aspect is used.

Other names

Gall Stereographic

Similar projections

Miller Cylindrical projection (p. 35) has different spacing of the parallels, and the line of no distortion is the Equator rather than latitudes 45° N. and S.


B.S.A.M. (Great Soviet World Atlas) projection of 1937 is the same, except that the cylinder is secant at latitudes 30° N. and S.

V.A. Kamenetskiy used an identical projection for Russian population density in 1929, except that the cylinder was made secant at latitudes 55° N. and S.

Guy Bomford of Oxford University in England about 1950 devised a Modified Gall projection, which is like the regular Gall in spacing except that meridians are slightly curved at higher latitudes to decrease scale exaggeration at the poles.

Moir devised "The Times" projection in which the straight parallels are spaced as they are on the Gall projection but the meridians are distinctly curved.

Figure 11A.—Gall projection with Tissot indicatrices, 30° graticule.



GALL Projection




Figure 11B.—Gall projection with shorelines, 15° graticule. Central meridian 90° W.

MILLER CYLINDRICAL Projection

Classifications

Cylindrical

Neither conformal nor equal area

Graticule

Meridians: Equally spaced straight parallel lines 0.73 as long as the Equator

Parallels: Unequally spaced straight parallel lines, closest near the Equator, but spacing does not increase poleward as fast as it does on the Mercator. Perpendicular to meridians.

Poles: Straight lines equal in length to the Equator

Symmetry: About any meridian or the Equator

Scale

True along the Equator in all directions

Constant in any given direction along any other given latitude; same scale at the latitude of opposite sign

Changes with latitude and direction

Distortion

None at the Equator. Shape, area, and scale distortion increases moderately away from the Equator but becomes severe at the poles (fig. 12A).

Other features

Parallels are spaced from the Equator by calculating the distance for 0.8 of the same latitude on the Mercator and dividing the result by 0.8. Therefore, the two projections are almost identical near the Equator.

Usage

World maps in numerous American atlases and some other atlases, as a projection resembling the Mercator but having less distortion of area and scale, especially near the poles

Origin

Presented by Osborn Maitland Miller (1897–1979) of the American Geographical Society in 1942

Aspects

Normal aspect is commonly used.


Oblique aspect has been used by the National Geographic Society.

Similar Projections

Gall projection (p. 33) has different spacing of parallels. The lines of no distortion are at latitudes 45° N. and S. rather than at the Equator.

Miller proposed other alternates in 1942, including one identical with his preferred cylindrical but using two-thirds instead of 0.8.

Figure 12A.—Miller Cylindrical projection with Tissot indicatrices, 30° graticule.



MILLER CYLINDRICAL Projection




Figure 12B.—Miller Cylindrical projection with shorelines, 15° graticule. Central meridian 90° W.

SINUSOIDAL Projection

Classifications

Pseudocylindrical

Equal area

Equally spaced parallels

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced sinusoidal curves intersecting at the poles and concave toward the central meridian.

Parallels: Equally spaced straight parallel lines perpendicular to the central meridian

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along every parallel and along the central meridian


Distortion

Severe near outer meridians at high latitudes (fig. 13A) but can be substantially reduced by interruption with several central meridians (fig. 13G). Free of distortion along the Equator and along the central meridian.

Usage

Atlas maps of South America and Africa. Occasionally used for world maps. Formerly used for other continental maps and star maps. Combined with Mollweide projection to develop other projections such as the Homolosine and the Boggs.

Figure 13A.—Sinusoidal projection with Tissot indicatrices, 30° graticule.



Origin

Developed in the 16th century. Used by J. Cossin in 1570 and by J. Hondius in Mercator atlases of the early 17th century. Often called Sanson-Flamsteed projection after later users. Oldest current pseudocylindrical projection.

Aspects

For educational purposes, it has been shown in various aspects as examples of normal, transverse, and oblique aspects of almost any pseudocylindrical projection (figs. 13C–13F).

Other names

Sanson-Flamsteed

Mercator Equal-Area

Similar projections

Several other pseudocylindrical projections, such as Craster Parabolic (p. 70) and Boggs Eumorphic (p. 68), are very similar, but parallels are not equally spaced, and meridians are curved differently. Eckert V (p. 46) and VI (p. 50) have sinusoidal meridians but have lines for poles.

SINUSOIDAL Projection

Figure 13B.—Sinusoidal projection with shorelines, 15° graticule. Central meridian 90° W.





Figure 13C.—Transverse Sinusoidal projection with shorelines, 15° graticule. Central meridian 90° W. North Pole centered on base projection (at 0° longitude).



SINUSOIDAL Projection

Figure 13D.—Transverse Sinusoidal projection with shorelines, 15° graticule. Central meridian 90° W. North Pole at -90° longitude on base projection.





Figure 13E.—Oblique Sinusoidal projection with shorelines, 15° graticule. Central meridian 90° W. North Pole at +45° latitude, 0° longitude on base projection.



SINUSOIDAL Projection

Figure 13F.—Oblique Sinusoidal projection with shorelines, 15° graticule. Central meridian 90° W. North Pole at +45° latitude, -90° longitude on base projection.





Figure 13G.—Interrupted Sinusoidal projection, with shorelines, 10° graticule. Interruptions symmetrical about Equator.



McBRYDE-THOMAS FLAT-POLAR SINUSOIDAL Projection

Classifications

Pseudocylindrical
Equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced sinusoids, concave toward the central meridian.

Parallels: Unequally spaced straight parallel lines, widest separation near the Equator. Perpendicular to the central meridian.

Poles: Lines one-third as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 55°51' N. and S.

Constant along any given latitude; same for the latitude of opposite sign

Distortion

Free of distortion only at latitudes 55°51' N. and S. on the central meridian (fig. 14A).

Usage

Basis of merged projections by McBryde (see McBryde S3, p. 52)

Origin

Presented by F. Webster McBryde and Paul D. Thomas through the U.S. Coast and Geodetic Survey in 1949

Similar projections

Sinusoidal projection (p. 37) uses sinusoids for meridians, but the poles are points.

Eckert VI projection (p. 50) uses sinusoids for meridians and is equal area, but the poles are lines half as long as the Equator.





Figure 14A.—McBryde-Thomas Flat-Polar Sinusoidal projection, with Tissot indicatrices, 30° graticule.

Figure 14B.—McBryde-Thomas Flat-Polar Sinusoidal projection, with shorelines, 15° graticule. Central meridian 90° W.



ECKERT V Projection

Classifications

Pseudocylindrical

Equally spaced parallels

Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced sinusoids, concave toward the central meridian.

Parallels: Equally spaced straight parallel lines. Perpendicular to the central meridian.

Poles: Lines half as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 37°55' N. and S., if the world map retains correct total area

Constant along any given latitude; same for the latitude of opposite sign


Distortion

No point is free of all distortion, but the Equator is free of angular distortion (fig. 15A).

Origin

Presented by Max Eckert (1868–1938) of Germany in 1906. The projection is an arithmetical average of the x and y coordinates

Figure 15A.—Eckert V projection, with Tissot indicatrices, 30° graticule.



of the Sinusoidal (p. 37) and Plate Carrée (p. 22) projections.


Similar projections

Eckert VI projection (p. 50) has meridians positioned identically, but parallels are spaced for equal area.

Winkel I (p. 48) is an average of coordinates of the Sinusoidal and Equirectangular projections.

Wagner III projection uses part rather than all of the sinusoidal curve for meridians.

Figure 15B.—Eckert V projection, with shorelines, 15° graticule. Central meridian 90° W.



WINKEL I Projection

Classifications

Pseudocylindrical

Equally spaced parallels

Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line 0.61 (or other value) as long as the Equator. Other meridians are equally spaced sinusoidal curves, concave toward the central meridian.

Parallels: Equally spaced straight parallel lines. Perpendicular to the central meridian.

Poles: Lines 0.61 (or other value) as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 50°28' N. and S. (or other chosen value)

Constant along any given latitude; same for the latitude of opposite sign

Distortion

Not free of distortion at any point


Origin

Developed by Oswald Winkel (1873–1953) of Germany in 1914 as the average of the Sinusoidal (p. 37) and Equirectangular (p. 24) projection in both x and y coordinates. When the standard parallels of the Equirectangular projection are varied, the standard parallels and appearance of Winkel I vary. Use of latitudes 50°28' N. and S. results in a map at the correct total-area scale, but the local-area scale varies.

Limiting form

Eckert V (p. 46), if the Equator is the standard parallel

Figure 16.—Winkel I projection, with shorelines, 15° graticule. Central meridian 90° W. Standard parallels 50°28' N. and S.



ECKERT VI Projection

Classifications

Pseudocylindrical
Equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced sinusoids, concave toward the central meridian.

Parallels: Equally spaced straight parallel lines, widest separation near the Equator. Perpendicular to the central meridian.

Poles: Lines half as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 49°16' N. and S.

Constant along any given latitude; same for the latitude of opposite sign

Distortion

Free of distortion only at latitudes 49°16' N. and S. at the central meridian (fig. 17A)

Usage


Thematic world maps in Soviet World Atlas of 1937

Some recent use for climatic maps by U.S. publishers

Origin

Presented by Max Eckert (1868–1938) of Germany in 1906.

Figure 17A.—Eckert VI projection, with Tissot indicatrices, 30° graticule.




Similar projections

Eckert V projection (p. 46) has meridians positioned identically, but parallels are equally spaced.

Wagner I projection (1932) is almost identical to Eckert VI, but Wagner I uses only part of the sinusoidal curve. Kavrayskiy VI projection (1936) is identical to Wagner I. Werenskiold II projection (1944) is the same as Wagner I, except for scale.

McBryde-Thomas Flat-Polar Sinusoidal (p. 44) uses the full sinusoid and is equal area, but the poles are one-third the length of the Equator.

Figure 17B.—Eckert VI projection, with shorelines, 15° graticule. Central meridian 90° W.



McBRYDE S3 Projection

Classifications

Pseudocylindrical composite

Equal area

Graticule

Meridians: Where the central meridian extends across the Equator, it is a straight line 0.44 as long as the Equator. Other central meridians in the usual interrupted form are straight and half as long. Other meridians are equally spaced sinusoidal curves, bending slightly at latitudes 55°51' N. and S., and all are concave toward the local central meridian.

Parallels: Straight parallel lines, perpendicular to the central meridian(s). Equally spaced between latitudes 55°51' N. and S.

Gradually closer together beyond these latitudes

Poles: Interrupted straight lines totaling 0.31 the length of the Equator

Symmetry: About the central meridian or the Equator (in uninterrupted form)

Scale

True along every latitude between 55°51' N. and S. and along the central meridian within the same latitude range. Constant along any given latitude; same for the latitude of opposite sign.

Distortion

Same as the Sinusoidal projection between latitude 55°51' N. and S. (see fig. 13A). Same as the McBryde-Thomas Flat-Polar Sinusoidal projection beyond this range (see fig. 14A). McBryde S3 is almost always used in the interrupted form and has several central meridians.

Usage

World map interrupted to show oceans or land masses, by McBryde

Origin


Developed by F. Webster McBryde (1908–) of Potomac, Md., in 1977 as a merging of the Sinusoidal with the McBryde-Thomas Flat-Polar Sinusoidal projection at the parallels of identical scale on the two projections, latitudes 55°51' N. and S. U.S. Patent by McBryde.

Similar projections

Identical with the Sinusoidal (p. 37) between latitudes 55°51' N. and S.; identical with the McBryde-Thomas Flat-Polar Sinusoidal (p. 44) poleward of those latitudes, except that those portions are closer to the Equator than they are on the McBryde-Thomas Flat-Polar Sinusoidal projection itself.

McBryde merged projections in several other similar combinations, also in 1977.

Figure 18.—Interrupted McBryde S3 projection, with shorelines, 10° graticule, to show oceans. Projection: U.S. Patent No. 4,315,747, held by F. Webster McBryde and used by permission.



MOLLWEIDE Projection

Classifications

Pseudocylindrical
Equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Meridians 90° E. and W. of the central meridian form a circle. Others are equally spaced semiellipses intersecting at the poles and concave toward the central meridian.

Parallels: Unequally spaced straight parallel lines, farthest apart near the Equator; spacing changes gradually. Perpendicular to the central meridian.

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 40°44' N. and S.

Constant along any given latitude; same for the latitude of opposite sign

Distortion

Severe near outer meridians at high latitudes (fig. 19A) but can be substantially reduced if interrupted with several central meridians (fig. 19D). Free of distortion only at latitudes 40°44' N. and S. on the central meridian.


Usage

Occasional world maps, especially thematic maps. Combined with Sinusoidal projection to develop other projections such as the Goode Homolosine and the Boggs.

Origin

Presented by Carl B. Mollweide (1774–1825) of Germany in 1805

Figure 19A.—Mollweide projection, with Tissot indicatrices, 30° graticule.



Aspects

For educational purposes, it has been shown in various aspects as examples of normal, transverse, and oblique aspects of almost any pseudocylindrical projection. The transverse aspect has also been used by John Bartholomew in *The Times Atlas* in England in 1958 (fig. 19C).

Other names

Homographic projection
Homalographic projection
Babinet projection
Elliptical projection


Similar projections

Goode Homolosine (p. 66) (Mollweide merged with Sinusoidal)
Boggs Eumorphic (p. 68) (Mollweide averaged with Sinusoidal)

Bromley, by Robert H. Bromley in 1965 (Mollweide compressed from north to south with east-west expansion to achieve no distortion along the Equator)

Hyperelliptical, by Waldo R. Tobler in 1973 (equal-area pseudocylindrical having "hyperelliptical" meridians that lie between the Mollweide ellipses and a circumscribed rectangle)

Figure 19B.—Mollweide projection, with shorelines, 15° graticule. Central meridian 90° W.



MOLLWEIDE Projection

Figure 19C.—“Atlantis” projection, a Transverse Mollweide used by John Bartholomew to highlight the Atlantic Ocean, with shorelines, 15° graticule. Central meridian 30° W. North Pole at –45° longitude on base projection.





Figure 19D.—Interrupted Mollweide projection, with shorelines, 10° graticule, to show oceans.



ECKERT III Projection

Classifications

Pseudocylindrical

Equally spaced parallels

Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced semiellipses, concave toward the central meridian. The outer meridians (180° east and west of the central meridian) are semicircles.

Parallels: Equally spaced straight parallel lines. Perpendicular to the central meridian.

Poles: Lines half as long as the Equator

Symmetry: About the central meridian or the Equator


Scale

True along latitudes 35°58' N. and S., if the world map retains correct total area (area varies locally). Constant along any given latitude; same for the latitude of opposite sign.

Distortion

No point is free of all scale distortion, but the Equator is free of angular distortion (fig. 20A).

Figure 20A.—Eckert III projection, with Tissot indicatrices, 30° graticule.



Origin


Presented by Max Eckert (1868–1938) of Germany in 1906

Similar projections

Eckert IV (p. 60) has meridians positioned identically, but parallels are spaced for equal area.

Putniņš P_{1'} has meridians that are only portions of semiellipses, but parallels are equidistant.

Figure 20B.—Eckert III projection, with shorelines, 15° graticule. Central meridian 90° W.



ECKERT IV Projection

Classifications

Pseudocylindrical

Equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced semiellipses, concave toward the central meridian. The outer meridians (180° east and west of the central meridian) are semicircles.

Parallels: Unequally spaced straight parallel lines, widest separation near the Equator. Perpendicular to the central meridian.

Poles: Lines half as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 40°30' N. and S.

Constant along any given latitude; same for the latitude of opposite sign

Distortion

Free of distortion only at latitudes 40°30' N. and S. at the central meridian (fig. 21A).


Usage

Thematic and other world maps in numerous atlases and textbooks and for sheet maps

Origin

Presented by Max Eckert (1868–1938) of Germany in 1906

Figure 21A.—Eckert IV projection, with Tissot indicatrices, 30° graticule.



Aspects


Like most projections, transverse and oblique aspects can be prepared, although they are seldom used in this case.

Similar projections

Eckert III (p. 58) has meridians positioned identically, but parallels are equally spaced.

Putniš P_{2'} (p. 62) has meridians that are only portions of semiellipses, but parallels are also spaced for equal area.

Figure 21B.—Eckert IV projection, with shorelines, 15° graticule. Central meridian 90° W.



PUTNINS P_{2'} Projection

Classifications

Pseudocylindrical
Equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced portions of ellipses (less than semiellipses) that would intersect beyond the poles and are concave toward the central meridian. The meridians at 103°55' east and west of the central meridian are circular arcs.

Parallels: Unequally spaced straight parallel lines, widest separation near the Equator.
Perpendicular to the central meridian.

Poles: Lines half as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 42°59' N. and S.

Constant along any given latitude; same for the latitude of opposite sign


Distortion

Distortion is not as extreme near outer meridians at high latitudes as it is on pointed-polar pseudocylindrical projections, but there is considerable distortion throughout polar regions (fig. 22A). Free of distortion only at latitudes 42°59' N. and S. at the central meridian.

Origin

Presented by Reinholds V. Putniņš of Latvia in 1934

Figure 22A.—Putniņš P_{2'} projection, with Tissot indicatrices, 30° graticule.



Similar projections

Wagner IV projection (1949) is identical.


Werenskiold III projection (1944) is identical, except that true scale is maintained along the Equator by enlarging the map.

Putniņš P₂ equal-area projection (1934) uses the same portions of ellipses, but the poles are points.

Putniņš P₁ and P_{1'} projections (1934) have meridians identical in shape and position to those of Putniņš P₂ and P_{2'}, respectively, but parallels are equally spaced, and the projections are not equal area.

Robinson projection (p. 82) is not equal area, meridians are not elliptical, and parallels are spaced differently.

Figure 22B.—Putniš P_2' projection, with shorelines, 15° graticule. Central meridian 90° W.



HATANO ASYMMETRICAL EQUAL-AREA Projection

Classifications

Pseudocylindrical
Equal area

Graticule

Meridians: The central meridian is a straight line 0.48 as long as the Equator. Other meridians are equally spaced elliptical arcs, concave toward the central meridian, but the eccentricity of each ellipse changes at the Equator.

Parallels: Unequally spaced straight parallel lines. Perpendicular to the central meridian. Spacing not symmetrical about the Equator.

Poles: Lines. The North Pole is two-thirds as long and the South Pole is three-fourths as long as the Equator.

Symmetry: About the central meridian but not about the Equator

Scale

True along latitudes 40°42' N. and 38°27' S.

Constant along any given latitude; normally different for the latitude of opposite sign


Distortion

Free of distortion only at latitudes 40°42' N. and 38°27' S. at the central meridian (fig. 23A).

Origin

Presented by Masataka Hatano of Japan in 1972 as a modification of the Mollweide and Putniš P₂' projections

Figure 23A.—Hatano Asymmetrical Equal-Area projection, with Tissot indicatrices, 30° graticule.




Similar projections

Mollweide projection (p. 54) uses full symmetrical semiellipses for meridians, and poles are points.

Putniš P₂' projection (p. 62) uses arcs less than semiellipses for meridians, but they are symmetrical about the Equator, and poles are lines half the length of the Equator.

Hatano Symmetrical Equal-Area projection (1972) uses the same graticule for the Northern Hemisphere. The Southern Hemisphere is symmetrical (the South Pole is also two-thirds the length of the Equator).

Figure 23B.—Hatano Asymmetrical Equal-Area projection, with shorelines, 15° graticule. Central meridian 90° W.



GOODE HOMOLOSINE Projection

Classifications

Pseudocylindrical composite
Equal area

Graticule

Meridians: In the interrupted form, there are six central meridians, each a straight line 0.22 as long as the Equator but not crossing the Equator. Other meridians are equally spaced sinusoidal curves between latitudes 40°44' N. and S. and elliptical arcs elsewhere, all concave toward the central meridian. There is a slight bend in meridians at the 40°44' latitudes.

Parallels: Straight parallel lines, perpendicular to the central meridians. Equally spaced between latitudes 40°44' N. and S. Gradually closer together beyond these latitudes as a pole is approached.

Poles: Points

Symmetry: None in the usual interrupted form. If uninterrupted, there is symmetry about the central meridian or the Equator.

Scale

True along every latitude between 40°44' N. and S. and along the central meridian within the same latitude range. At higher latitudes, scale varies but is the same for the latitude of opposite sign.

Distortion

Same as Sinusoidal projection between latitudes 40°44' N. and S. (see fig. 13A). Same as Mollweide projection beyond this range (see fig. 19A). Goode Homolosine projection is almost always presented interrupted, as the inventor originally intended.

Usage

Numerous world maps, especially in *Goode's Atlas* (Rand McNally)

Origin

Developed in 1923 by J. Paul Goode (1862–1932) of the University of Chicago as a merging of the Mollweide (or Homographic) with the Sinusoidal at the parallels of identical scale, latitudes 40°44' N. and S.; hence, the name Homolosine. Frequently plotted with some northern regions repeated on different interrupted sections to reduce discontinuities of land.

Other names

Homolosine


Similar projections

Identical with Sinusoidal (p. 37) between latitudes 40°44' N. and S. and identical with Mollweide (p. 54) beyond these latitudes, except that the Mollweide portions are closer to the Equator than they are on the original projection

Sinu-Mollweide, devised by Allen K. Philbrick in 1953, is a merging of the Mollweide, used from the North Pole to latitude 40°44' S., with the Sinusoidal, used only south of latitude 40°44' S. It is equal area.

In 1968, György Érdi-Kraus used a special Flat-Polar Sinusoidal projection between latitudes 60° N. and S. and the Mollweide projection poleward. It is equal area only within each of the two projections, which are at two different area scales.

Figure 24.—Interrupted Goode Homolosine projection, with shorelines, 10° graticule, to show land masses.



BOGGS EUMORPHIC Projection

Classifications

Pseudocylindrical

Equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced complex curves intersecting at the poles and concave toward the central meridian.

Parallels: Unequally spaced straight parallel lines perpendicular to the central meridian

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 40°15' N. and S.

Constant along any given latitude; same for the latitude of opposite sign


Distortion

Distortion is severe near outer meridians at high latitudes (fig. 25A) but can be substantially reduced by interruption. Free of distortion only at latitudes 40°15' N. and S. at the central meridian.

Other features

Designed as the arithmetic mean of the Sinusoidal and Mollweide projections for north-south (y) coordinates only, when drawn with the same area scale

Figure 25A.—Boggs Eumorphic projection, with Tissot indicatrices, 30° graticule.



Usage

Thematic world maps in atlases and textbooks, usually in interrupted form


Origin

Presented by S. Whittemore Boggs (1884–1954) of the U.S. Department of State in 1929

Similar projections

Several pseudocylindricals, such as the Sinusoidal (p. 37) and Craster Parabolic (p. 70).

Figure 25B.—Boggs Eumorphic projection, with shorelines, 15° graticule. Central meridian 90° W.



CRASTER PARABOLIC Projection

Classifications

Pseudocylindrical
Equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced parabolas intersecting at the poles and concave toward the central meridian.

Parallels: Unequally spaced straight parallel lines, farthest apart near the Equator; spacing changes very gradually. Perpendicular to the central meridian.

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 36°46' N. and S.
Constant along any given latitude; same for the latitude of opposite sign

Distortion

Distortion is severe near outer meridians at high latitudes (fig. 26A) but somewhat less than that of the Sinusoidal projection. Can be substantially reduced by interruption (see similar example for Sinusoidal, fig. 13G). Free of distortion only at latitudes 36°46' N. and S. at the central meridian.


Usage

Thematic world maps in textbooks

Origin

Presented by John Evelyn Edmund Craster

Figure 26A.—Craster Parabolic projection, with Tissot indicatrices, 30° graticule.



(1873–?) of England in 1929. Developed further by Charles H. Deetz and O.S. Adams in 1934

Aspects

Oblique aspect used for map of Asia by National Geographic Society

Other names


Putniņš P₄, independently presented in Latvia in 1934.

Similar projections

Several pseudocylindricals, such as the Sinusoidal (p. 37) and the Boggs Eumorphic (p. 68)

Putniņš P₃ projection (1934) has meridians, poles, and Equator identical to those of the Craster, but parallels are equally spaced.

Figure 26B.—Craster Parabolic projection, with shorelines, 15° graticule. Central meridian 90° W.



McBRYDE-THOMAS FLAT-POLAR PARABOLIC Projection

Classifications

Pseudocylindrical

Equal area

Graticule

Meridians: Central meridian is a straight line 0.48 as long as the Equator. Other meridians are equally spaced parabolic curves, concave toward the central meridian.

Parallels: Unequally spaced straight parallel lines, farthest apart near the Equator.

Perpendicular to the central meridian.

Poles: Lines one-third as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 45°30' N. and S.

Constant along any given latitude; same for the latitude of opposite sign

Distortion

Distortion is severe near outer meridians at high latitudes (fig. 27A) but less than the corresponding distortion on pointed-polar projections. Free of distortion only at latitudes 45°30' N. and S. at the central meridian.

Usage

Basis of merged projections by McBryde

Origin

Presented by F. Webster McBryde and Paul D. Thomas through the U.S. Coast and Geodetic Survey in 1949




Figure 27A.—McBryde-Thomas Flat-Polar Parabolic projection, with Tissot indicatrices, 30° graticule.

Similar projections

Craster Parabolic projection (p. 70), equal area with parabolic meridians but with pointed poles and a central meridian that is half the length of the Equator

Putniņš P_{4'} projection (1934), equal area with parabolic meridians but with poles and a central meridian that are all half the length of the Equator. Werenskiold I (1944) is identical to Putniņš P_{4'} except for the scale.

Putniņš P_{3'} projection (1934) has meridians, poles, and Equator identical to those of Putniņš P_{4'}, but parallels are equally spaced.

Figure 27B.—McBryde-Thomas Flat-Polar Parabolic projection, with shorelines, 15° graticule. Central meridian 90° W.



QUARTIC AUTHALIC Projection

Classifications

Pseudocylindrical

Equal area

Graticule

Meridians: Central meridian is a straight line 0.45 as long as the Equator. Other meridians are equally spaced curves fitting a fourth-order (quartic) equation and concave toward the central meridian.

Parallels: Unequally spaced straight parallel lines, farthest apart near the Equator. Spacing changes gradually. Perpendicular to the central meridian.

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along the equator

Constant along any given latitude; same for the latitude of opposite sign


Distortion

Distortion is severe near outer meridians at high latitudes (fig. 28A) but somewhat less than that of the Sinusoidal projection. Distortion can be substantially reduced by interruption (see similar example for Sinusoidal, fig. 13G). Free of distortion along the Equator.

Usage

Basis for McBryde-Thomas Flat-Polar Quartic projection (p. 76)

Figure 28A.—Quartic Authalic projection, with Tissot indicatrices, 30° graticule.



Origin


Presented by Karl Siemon (?–1937) of Germany in 1937. Independently presented by Oscar Sherman Adams (1874–1962) of the U.S. Coast and Geodetic Survey in 1945.

Similar projections

Several pseudocylindricals, such as the Sinusoidal (p. 37)

Eckert-Greifendorff (p. 166) has slightly curved parallels.

Figure 28B.—Quartic Authalic projection, with shorelines, 15° graticule. Central meridian 90° W.



McBRYDE-THOMAS FLAT-POLAR QUARTIC Projection

Classifications

Pseudocylindrical
Equal area

Graticule

Meridians: Central meridian is a straight line 0.45 as long as the Equator. Other meridians are equally spaced curves fitting a fourth-order (quartic) equation and concave toward the central meridian.

Parallels: Unequally spaced straight parallel lines, farthest apart near the Equator.

Perpendicular to the central meridian.

Poles: Lines one-third as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 33°45' N. and S.

Constant along any given latitude; same for the latitude of opposite sign

Distortion

Distortion is severe near outer meridians at high latitudes (fig. 29A) but less than the corresponding distortion on pointed-polar projections. Free of distortion only at latitudes 33°45' N. and S. at the central meridian.

Usage

Examples in various geography textbooks.

Basis of merged projections by McBryde.

Origin

Presented by F. Webster McBryde and Paul




Figure 29A.—McBryde-Thomas Flat-Polar Quartic projection, with Tissot indicatrices, 30° graticule.

D. Thomas through the U.S. Coast and Geodetic Survey in 1949

Other names


Flat-Polar Quartic projection

Similar projections

Quartic Authalic projection (p. 74), the basis of this projection, has pointed poles.

Other McBryde-Thomas projections also have poles one-third the length of the Equator and otherwise resemble the projection described here.

Figure 29B.—McBryde-Thomas Flat-Polar Quartic projection, with shorelines, 15° graticule. Central meridian 90° W.



PUTNIŅŠ P₅ Projection

Classifications

Pseudocylindrical

Equally spaced parallels

Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are equally spaced portions of hyperbolae, concave toward the central meridian.

Parallels: Equally spaced straight parallel lines, perpendicular to the central meridian.

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 21°14' N. and S.

Constant along any given latitude; same for the latitude of opposite sign. Constant along the central meridian.

Distortion

Similar to other pointed-polar cylindricals but not free of distortion at any point (fig. 30A).


Origin

Presented by Reinholds V. Putniņš of Latvia in 1934

Similar projections

Putniņš P₆ equal-area projection (1934) has

Figure 30A.—Putniņš P₅ projection, with Tissot indicatrices, 30° graticule.




hyperbolic meridians identical in shape to those of the pointed-polar P₅; parallels are straight but are spaced for equal area.

Putniņš P_{5'} projection (1934) has hyperbolic meridians, but the poles are straight lines half the length of the Equator. Parallels are equally spaced. The projection is not equal area.

Putniņš P_{6'} equal-area projection (1934) has hyperbolic meridians identical in shape to those of the flat-polar P_{5'}, but parallels are straight lines spaced for equal area.

Figure 30B.—Putniņš P_5 projection, with shorelines, 15° graticule. Central meridian 90° W.



DENOYER SEMI-ELLIPTICAL Projection

Classifications

Modified pseudocylindrical

Equally spaced parallels

Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line half as long as the Equator. Other meridians are curves that are nearly elliptical areas and nearly equally spaced, with some spreading near the outer limits; all are concave toward the central meridian.

Parallels: Equally spaced straight parallel lines. Perpendicular to the central meridian.

Poles: Lines about 0.31 times the length of the Equator

Symmetry: About the central meridian or the Equator

Scale

True along the central meridian and the Equator

Constant along any given latitude; same for the latitude of opposite sign

Distortion

Free of distortion along the Equator (fig. 31A)

Usage

By Denoyer-Geppert in thematic maps and atlases of the early and middle 20th century

Origin

Developed by L. Philip Denoyer of Chicago about 1920. Exact design basis not known.

Empirical approximations are used for computer plots.





Figure 31A.—Denoyer Semi-Elliptical projection approximation, with Tissot indicatrices, 30° graticule.

Figure 31B.—Denoyer Semi-Elliptical projection approximation, with shorelines, 15° graticule. Central meridian 90° W.



ROBINSON Projection

Classifications

Pseudocylindrical

Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line 0.51 as long as the Equator. Other meridians resemble elliptical arcs, are equally spaced, and are concave toward the central meridian.

Parallels: Straight parallel lines, equally spaced between latitudes 38° N. and S.; space decreases beyond these limits.

Poles: Lines 0.53 as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 38° N. and S.

Constant along any given latitude; same for the latitude of opposite sign

Distortion

No point is completely free of distortion, but distortion is very low within about 45° of the center and along the Equator (fig. 32B).

Considerable distortion near the poles.


Usage

By Rand McNally in *Goode's Atlas* for thematic world maps and in numerous other publications

Origin

Presented by Arthur H. Robinson (1915–) of the University of Wisconsin in 1963, at the request of Rand McNally and Company. Uses

Figure 32A.—Robinson projection, with Tissot indicatrices, 30° graticule.



tabular coordinates rather than mathematical formulas to make the world map "look" right.

Other names

Orthophanic ("right appearing")

Similar projections

Putniņš P_{2'} (p. 62), an equal-area projection on which meridians are elliptical arcs

In 1968, János Baranyi presented seven pseudocylindrical projections characterized by various graduated tabular spacings of the parallels. The poles are usually points but occasionally are lines, and the outer meridians 180° from the central meridian are circular arcs combined in some cases with straight lines. His projections are neither equal area nor conformal.

Figure 32B.—Robinson projection, with shorelines, 15° graticule. Central meridian 90° W.




Figure 32C.—Robinson projection, with shorelines, 30° graticule. Central meridian 0°.

COLLIGNON Projection

Classifications

Pseudocylindrical

Equal area

Graticule

Meridians: Equally spaced straight lines converging at the North Pole

Parallels: Unequally spaced straight parallel lines, farthest apart near the North Pole, closest near the South Pole. Perpendicular to the central meridian.

Poles: North Pole is a point; South Pole is a line 1.41 times as long as the Equator

Symmetry: About the central meridian

along each parallel (top and bottom parallels and perpendicular central meridian true to scale). It was used for numerous maps from the 15th to the 18th centuries but is neither equal area nor conformal.

Polyhedral projection, formerly used for large-scale mapping in various parts of central Europe. There were various forms, all resembling the Trapezoidal projection but applied to the ellipsoid.

Scale

True along latitudes 15°51' N.

Constant along any given latitude; different for each latitude

Distortion

Severe in many regions. Free of distortion only at latitude 15°51' N. at the central meridian.

Usage

Novelty showing straight-line equal-area graticule

Origin


Presented by Édouard Collignon of France in 1865

Similar projections

Eckert I (p. 86) and II (p. 88) also use straight lines for both meridians and parallels, but meridians are broken at the Equator.

The Trapezoidal projection has equidistant straight parallel lines for parallels and converging straight meridians equally spaced

Figure 33.—Collignon projection, with shorelines, 15° graticule. Central meridian 90° W.



ECKERT I Projection

Classifications

Pseudocylindrical
Equally spaced parallels
Neither conformal nor equal area

Graticule

Meridians: Equally spaced converging straight lines broken at the Equator. Central meridian is half as long as the Equator.

Parallels: Equally spaced straight parallel lines, perpendicular to the central meridian.

Poles: Lines half as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 47°10' N. and S.

Constant along any given latitude; same for the latitude of opposite sign. Constant along any given meridian.

Distortion

No point free of distortion (fig. 34A). Break at the Equator introduces excessive distortion there. Discontinuous ellipses are shown there owing to the plotting algorithm; actually, the shape is indeterminate on the Equator.

Usage

Novelty showing straight-line graticule

Origin

Presented by Max Eckert (1868–1938) of Germany in 1906

Similar projections

Eckert II (p. 88) has meridians positioned identically, but parallels are spaced for equal area.

Figure 34A.—Eckert I projection, with Tissot indicatrices, 30° graticule.





Figure 34B.—Eckert I projection, with shorelines, 15° graticule. Central meridian 90° W.



ECKERT II Projection

Classifications

Pseudocylindrical
Equal area

Graticule

Meridians: Equally spaced straight lines broken at the Equator. Central meridian is half as long as the Equator.

Parallels: Unequally spaced straight parallel lines, widest separation near the Equator.

Perpendicular to the central meridian.

Poles: Lines half as long as the Equator

Symmetry: About the central meridian or the Equator

Scale

True along latitudes 55°10' N. and S.

Constant along any given latitude; same for the latitude of opposite sign


Distortion

Free of distortion only at latitudes 55°10' N. and S. at the central meridian (fig. 35A). Break at the Equator introduces excessive distortion there. Discontinuous ellipses are shown there owing to the plotting algorithm; actually, the shape is indeterminate on the Equator.

Usage

Novelty showing straight-line equal-area graticule

Figure 35A.—Eckert II projection, with Tissot indicatrices, 30° graticule.



Origin


Presented by Max Eckert (1868–1938) of Germany in 1906

Similar projections

Eckert I (p. 86) has meridians positioned identically, but parallels are equidistant.

Collignon projection (p. 84) has straight meridians and is equal area, but meridians are not broken at the Equator, and the poles are different.

Figure 35B.—Eckert II projection, with shorelines, 15° graticule. Central meridian 90° W.



LOXIMUTHAL Projection

Classifications

Pseudocylindrical
Equally spaced parallels
Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line generally over half as long as the Equator, depending on the central latitude. If the central latitude is the Equator, the ratio is 0.5; if it is 40° N. or S., the ratio is 0.65. Other meridians are equally spaced complex curves intersecting at the poles and concave toward the central meridian.

Parallels: Equally spaced straight parallel lines. Perpendicular to the central meridian.

Poles: Points

Symmetry: About the central meridian.
Symmetry occurs about the Equator if the specified central latitude is the Equator.

Scale

True along the central meridian
Constant along any given latitude; normally different for the latitude of opposite sign

Distortion

Distortion is moderate to extreme, but the projection is intended for the special property noted below, not as a normal world map. Free of distortion only at the central latitude on the central meridian.

Special feature

Loxodromes (rhumb lines) from the central point (at the intersection of the central meridian and the central parallel) are shown straight, true to scale, and correct in azimuth from the center. Because of angular distortion on the map projection, the azimuths with respect to other points along a rhumb line are not shown correctly.

Origin


Presented by Karl Siemon (?–1937) of Germany in 1935. Independently presented as "Loximuthal" by Waldo R. Tobler (1930–) of the University of Michigan (1966).

Similar projections

Loxodromes are shown straight and at the correct azimuth between any two points on the regular Mercator projection, but they are not at true scale.

The Bordone Oval projection of 1520 is similar to the Loximuthal projection having the Equator as the central latitude.

Figure 36.—Loximuthal projection, with shorelines, 15° graticule. Central meridian 90° W. Central latitude 40° N.



EQUIDISTANT CONIC Projection

Classifications

Conic

Equally spaced parallels

Neither conformal nor equal area

Graticule

Meridians: Equally spaced straight lines converging at a common point, which is normally beyond the pole. The angles between them are less than the true angles.

Parallels: Equally spaced concentric circular arcs centered on the point of convergence of the meridians, which are therefore radii of the circular arcs

Poles: Normally circular arcs enclosing the same angle as that enclosed by the other parallels of latitude for a given range of longitude

Symmetry: About any meridian

Scale

True along each meridian and along one or two chosen standard parallels, usually but not necessarily on the same side of the Equator. As a rule of thumb, these parallels can be placed at one-sixth and five-sixths of the range of latitudes, but there are more refined means of selection. Scale is constant along any given parallel.

Distortion


Free of distortion only along the two standard parallels (fig. 37A). Distortion is constant along any given parallel. Compromise in distortion between equal-area and conformal conic projections.

Usage

The most common projection in atlases for small countries

Also used by the Soviet Union for mapping that nation

Figure 37A.—Equidistant Conic projection, with Tissot indicatrices, 30° graticule. Standard parallels: 20° and 60° N.



Origin

Rudimentary forms developed by Claudius Ptolemy (about A. D. 100). Improvements by Johannes Ruysch in 1508, Gerardus Mercator in the late 16th century, and Nicolas de l'Isle in 1745.

Other names

Simple Conic

Conic

Limiting forms

Polar Azimuthal Equidistant projection (fig. 37C and p. 133), if a pole is made the single standard parallel. The cone of projection thereby becomes a plane.

Plate Carrée (fig. 37D, also p. 22), if the single standard parallel is the Equator. The cone of projection thereby becomes a cylinder.

Equirectangular (Cylindrical) projection (p. 24), if two standard parallels are symmetrically placed north and south of the Equator.


Standard conic formulas must be rewritten for the second and third limiting forms.

Similar projections

Various methods of determining optimum standard parallels have been proposed by Patrick Murdoch (projections I, III) in 1758, Leonhard Euler in 1777, British Ordnance in the late 19th century, Dmitri I. Mendeleev in 1907, Wilhelm Schjerning (projection I) in 1882, V.V. Vitkovskiy (projection I) in 1907, and V.V. Kavrayskiy (projections II, IV) in 1934. Once the standard parallels are selected, all these projections are constructed by using formulas used for the Equidistant Conic with two standard parallels.

John Bartholomew combined the Equidistant Conic projection with the Bonne projection (see p. 112).

Figure 37B.—Equidistant Conic projection, with shorelines, 15° graticule. Central meridian 90° W. Standard parallels: 20° and 60° N.



EQUIDISTANT CONIC Projection




Figure 37C.—North Polar Azimuthal Equidistant projection, with shorelines, 30° graticule. Central meridian 90° W. Polar and planar limit of Equidistant Conic projection.




Figure 37D.—Plate Carrée projection, with shorelines, 30° graticule. Central meridian 90° W. Equatorial and cylindrical limit of Equidistant Conic projection.

LAMBERT CONFORMAL CONIC Projection

Classifications

Conic
Conformal

Graticule

Meridians: Equally spaced straight lines converging at a common point, which is one of the poles. The angles between them are less than the true angles.

Parallels: Unequally spaced concentric circular arcs centered on the pole of convergence of the meridians, which are therefore radii of the circular arcs. Spacing of parallels increases away from the central latitudes.

Poles: The pole nearest a standard parallel is a point; the other pole cannot be shown.

Symmetry: About any meridian

Scale

True along one or two chosen standard parallels, usually but not necessarily on the same side of the Equator. As a rule of thumb, these parallels can be placed at one-sixth and five-sixths of the range of latitudes, but there are more refined means of selection.

Scale is constant along any given parallel and is the same in all directions at a given point.

Distortion

Free of distortion only along the one or two standard parallels (fig. 38A). Distortion is constant along any given parallel. Conformal everywhere except at the poles.

Usage

Extensively used in ellipsoidal form for large-scale mapping of regions of predominantly east-west extent, including topographic quadrangles (1:24,000 and 1:62,500 scale) for many of the U.S. State Plane Coordinate




Figure 38A.—Lambert Conformal Conic projection, with Tissot indicatrices, 30° graticule. Standard parallels: 20° and 60° N.

System zones, many maps in the International Map of the World (1:1,000,000 scale) series, the U.S. State Base Maps (1:500,000 scale), and topographic mapping in many other nations

Also used for atlas maps of some countries
Recommended for conformal mapping of regions of predominantly east-west extent

LAMBERT CONFORMAL CONIC Projection



Figure 38B.—Lambert Conformal Conic projection, with shorelines, 15° graticule. Central meridian 90° W. Standard parallels: 20° and 60° N.

Origin

Presented by Johann Heinrich Lambert (1728–77) of Alsace in 1772

Aspects

Bipolar Oblique Conic Conformal projection (p. 99) is the only common oblique aspect. See figure 38E for another example applied to the world.

Other names

Conical Orthomorphic

Limiting forms

Polar Stereographic projection (fig. 38C and p. 121), if a pole is the single standard parallel. The cone of projection thereby becomes a plane.

Mercator projection (fig. 38D, also p. 10), if the single standard parallel is the Equator or if two standard parallels are symmetrically placed north and south of the Equator. The cone of projection thereby becomes a cylinder. Standard conic formulas must be rewritten.

Similar projections

Various methods of determining optimum standard parallels have been proposed by John Herschel in 1860, V.V. Vitkovskiy (projection III) in 1907, N.Ya. Tsinger in 1916, and V.V. Kavrayskiy (projection III) in 1934. Once the standard parallels are selected, all these projections are constructed by using the same formulas used for the Lambert Conformal Conic with two standard parallels.

Bipolar Oblique Conic Conformal projection (p. 99) consists of oblique aspects of two Lambert Conformal Conic projections laid side by side.




Figure 38C.—North Polar Stereographic projection, with shorelines, 30° graticule. Central meridian 90° W. Polar and planar limit of Lambert Conformal Conic projection.




Figure 38D.—Mercator projection, with shorelines, 30° graticule. Central meridian 90° W. Equatorial and cylindrical limit of Lambert Conformal Conic projection.

LAMBERT CONFORMAL CONIC Projection




Figure 38E.—Oblique Lambert Conformal Conic projection, with shorelines, 15° graticule. Central meridian 90° W. Standard parallels +20° and +60° on base projection. North Pole at +30° latitude, -90° longitude on base projection.

BIPOLAR OBLIQUE CONIC CONFORMAL Projection

Classifications

Conic
Conformal

Graticule

Meridians: Unequally spaced curved lines, except for two straight meridians from the transformed poles to the nearest geographic pole

Parallels: Unequally spaced curved lines

Poles: Points

Symmetry: None

Scale

True along four circular arcs joined to form two S curves. These arcs do not follow any parallel or meridian.

Scale is constant along circular arcs centered on each transformed pole for part of the map. The pole for most of North America is in the North Atlantic, and that for South America is in the South Pacific.

Distortion

Low throughout North and South America. None along the lines of true scale and constant along circular arcs centered on the transformed pole for the given part of the map. Slight nonconformality near the line joining the two transformed poles.

Usage

Designed specifically for a low-error map of North and South America constructed by the American Geographical Society. Other organizations, including the U.S. Geological Survey, have used the American Geographical Society map as a base for maps of North or South America alone, but, since the advent of computer plotting, other projections are preferable and practical for regional maps.

Origin

Presented by Osborn Maitland Miller (1897–1979) and William A. Briesemeister of the American Geographical Society in 1941



Figure 39.—Bipolar Oblique Conic Conformal projection, with shorelines, 10° graticule. Specially designed for North and South America.

ALBERS EQUAL-AREA CONIC Projection

Classifications

Conic

Equal area

Graticule

Meridians: Equally spaced straight lines converging at a common point, which is normally beyond the pole. The angles between them are less than the true angles.

Parallels: Unequally spaced concentric circular arcs centered on the point of convergence of the meridians. The meridians are therefore radii of the circular arcs. Spacing of parallels decreases away from the central latitudes.

Poles: Normally circular arcs enclosing the same angle as that enclosed by the other parallels of latitude for a given range of longitude

Symmetry: About any meridian

Scale

True along one or two chosen standard parallels, usually but not necessarily on the same side of the Equator. As a rule of thumb, these parallels can be placed at one-sixth and five-sixths of the range of latitudes, but there are more refined means of selection.

Scale is constant along any given parallel. The scale factor at any given point along the meridian is the reciprocal of that along the parallel, to preserve area.

Distortion

Free of angular and scale distortion only along the one or two standard parallels (fig. 40A)

Distortion is constant along any given parallel.

Usage

Frequently used in the ellipsoidal form for maps of the United States in the *National*

Atlas of the United States, for thematic maps, and for world atlases

Also used and recommended for equal-area maps of regions that are predominantly east-west in extent

Origin

Presented by Heinrich Christian Albers (1773–1833) of Germany in 1805

Aspects

Rarely used in oblique aspect but occasionally proposed in journal articles

Limiting forms

Polar Lambert Azimuthal Equal-Area projection (fig. 40C and p. 137), if a pole is made the single standard parallel. The cone of projection thereby becomes a plane.

Lambert Equal-Area Conic projection (p. 102), if the pole and another parallel are made the two standard parallels

Lambert Cylindrical Equal-Area projection (fig. 40D and p. 17), if the Equator is the single standard parallel. The cone of projection thereby becomes a cylinder.

Behrmann or other cylindrical equal-area projections (p. 19), if the two standard parallels are symmetrically placed north and south of the Equator.

Standard conic formulas must be rewritten for the third and fourth limiting forms.

Similar projections

Various methods of determining optimum standard parallels have been proposed by N.A. Tissot in 1881, V.V. Vitkovskiy (projection II) in 1907, N.Ya. Tsinger in 1916, and F.N. Krasovskiy (projection I) in 1922. Once the standard parallels are selected, all these projections are constructed by using the same formulas used for the Albers Equal-Area Conic with two standard parallels.

Figure 40A.—Albers Equal-Area Conic projection, with Tissot indicatrices, 30° graticule. Standard parallels: 20° and 60° N.




Figure 40B.—Albers Equal-Area Conic Projection, with shorelines, 15° graticule. Central meridian 90° W. Standard parallels 20° and 60° N.




Figure 40C.—North Polar Azimuthal Equal-Area projection, with shorelines, 30° graticule. Central meridian 90° W. Polar and planar limit of Albers Equal-Area projection.





Figure 40D.—Lambert Cylindrical Equal-Area projection, with shorelines, 30° graticule. Standard parallel 0°. Central meridian 90° W. Equatorial and cylindrical limit of Albers Equal-Area projection.



LAMBERT EQUAL-AREA CONIC Projection

Classifications

Conic

Equal area

Graticule

Meridians: Equally spaced straight lines converging at a common point, which is one of the poles. The angles between them are less than the true angles.

Parallels: Unequally spaced concentric circular arcs centered on the pole of convergence of the meridians. The meridians are therefore radii of the circular arcs. Spacing of parallels decreases away from the pole of convergence.

Poles: One pole is a point; the other pole is a circular arc enclosing the same angle as that enclosed by the other parallels of latitude for a given range of longitudes.

Symmetry: About any meridian

Scale

True along the chosen standard parallel

Scale is constant along any given parallel.

The scale factor at any given point along the meridian is the reciprocal of that along the parallel, to preserve area.

Distortion

Free of scale and angular distortion only along the standard parallel. Severe stretching near each pole.

Scale distortion and angular distortion are constant along any given parallel.

Origin

Presented by Johann Heinrich Lambert (1728–77) of Alsace in 1772

Limiting form


Polar Lambert Azimuthal Equal-Area

projection (p. 136), if the central pole is the standard parallel. The cone of projection thereby becomes a plane.

Similar projections

Albers Equal-Area Conic projection (p. 100) also has concentric circular arcs for parallels and straight meridians, but it has two standard parallels close to the latitudes of interest. If just one of its standard parallels is made a pole, it becomes the Lambert Equal-Area Conic projection, but that pole is not free of distortion, as the usual standard parallel is.

Figure 41.—Lambert Equal-Area Conic projection, with shorelines, 15° graticule. Central meridian 90° W. Standard parallel 0°.



PERSPECTIVE CONIC Projection

Classifications

Conic
Perspective
Neither conformal nor equal area

Graticule

Meridians: Equally spaced straight lines converging at a common point, which is one of the poles. The angles between them are less than the true angles.

Parallels: Unequally spaced concentric circular arcs centered on the pole of convergence of the meridians. The meridians are therefore radii of the circular arcs. Spacing of parallels increases away from the central latitudes.

Poles: One pole is a point; the other pole cannot generally be shown, but, under some conditions, it is a circular arc.

Symmetry: About any meridian





Figure 42A.—Perspective Conic projection, with shorelines, 15° graticule. Central meridian 90° W. Standard parallels 20° and 60° N.

Figure 42B.—North Polar Gnomonic projection, with shorelines, 30° graticule. Central meridian 90° W. Range 60° from center. Polar and planar limit of Perspective Conic projection.



Scale

True along one or two chosen standard parallels, which may be on the same side of or both sides of the Equator
Scale is constant along any given parallel.

Distortion

Free of distortion only along the one or two standard parallels. Distortion is constant along any given parallel but changes more rapidly in a north-south direction than it does on conformal or equal-area conics, and the Perspective Conic has no compensating advantage. Other projections should be used instead.

Other features

Projection is produced geometrically by projecting the Earth perspective from the center (or from some other point) onto a cone tangent or secant along the standard parallels.

Limiting forms

Polar Gnomonic projection (fig. 42B and p. 117), if the pole is the single standard parallel.
The cone of projection thereby becomes a plane.

Central Cylindrical projection (fig. 42C and p. 30), if the Equator is the single standard parallel. The cone of projection thereby becomes a cylinder. Standard conic formulas must be rewritten.

Similar projection

A stereographic perspective conic was presented by Carl Braun in 1867. The map is projected from the South Pole onto a cone tangent at latitude 30° N.




Figure 42C.—Central Cylindrical projection, with shorelines, 30° graticule. Central meridian 90° W. Equatorial and cylindrical limit of Perspective Conic projection.

POLYCONIC Projection

Classifications

Polyconic

Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line. All other meridians are complex curves, spaced equally along the Equator and along each parallel and concave toward the central meridian.

Parallels: Equator is a straight line, poles are points, and all other parallels are nonconcentric circular arcs spaced at true distances along the central meridian. Each parallel has a curvature identical to its curvature on a cone tangent at that latitude. Many cones are involved; hence, the name "Polyconic projection."

Symmetry: About the central meridian and the Equator

Scale

True along the central meridian and along each parallel

Distortion

Free of distortion only along the central meridian (fig. 43A). Extensive distortion if the range extends very far east or west.

Usage

The sole projection used for large-scale mapping in the ellipsoidal form (topographic quadrangles) of the United States by the U.S. Geological Survey until the 1950's

Basis for Progressive Military Grid used by the U.S. Army until the 1940's

Projection for many early coastal charts by the U.S. Coast and Geodetic Survey

Not recommended for regional maps, because other projections are better

Origin

Apparently originated about 1820 by Ferdinand Rudolph Hassler (1770–1843), first director of the Survey of the Coast (later the U.S. Coast and Geodetic Survey)

Aspects

Transverse aspect (fig. 43D) was proposed by Charles H. Deetz of the U.S. Coast and Geodetic Survey in the early 20th century. The National Geographic Society has used the transverse aspect for maps of Eurasia and the Soviet Union.

Other names

American Polyconic

Ordinary Polyconic

Similar projections

Rectangular Polyconic projection (p. 110) has identical parallels of latitude, but meridians are plotted to intersect parallels at right angles.

The Modified Polyconic projection for the International Map of the World (p. 111) series has two meridians that are true to scale.


An "Equidistant Polyconic" projection was used by the U.S. Coast and Geodetic Survey for coastal charts in the 19th century. Its construction is unclear, but it was apparently different from other polyconics.

Equal-area polyconic projections were presented by Hans Maurer in 1935 in Germany and by Albert H. Bumstead of the National Geographic Society in 1937.

Other modified Polyconics have been presented by G.T. McCaw in England in 1921, by G.A. Ginzburg and T.D. Salmanova in the Soviet Union in 1949 and 1951, and by G.E. Bousfield in Canada in 1950.


"Polyconic" is sometimes applied to all

Figure 43A.—Polyconic projection, with Tissot indicatrices, 30° graticule.



projections having circular arcs for parallels of latitude, including regular conics and the Stereographic, but this usage is not common.

Figure 43B.—Polyconic projection, with shorelines, 10° graticule. Central meridian 90° W.



POLYCONIC Projection

Figure 43C.—Polyconic projection, with shorelines (world map), 30° graticule. Central meridian 90° W.





Figure 43D.—Transverse Polyconic projection, with shorelines, 10° graticule. Central meridian 160° W. North Pole at -40° longitude on base projection, showing North Pacific Ocean.



RECTANGULAR POLYCONIC Projection

Classifications

Polyconic

Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line. All other meridians are complex curves, equally spaced along the Equator or along two chosen parallels symmetrical about the Equator. Meridians intersect all parallels at right angles.

Parallels: Equator is a straight line, poles are points, and all other parallels are nonconcentric circular arcs spaced at true distances along the central meridian. Each parallel has a curvature identical to its curvature on a cone tangent at that latitude. Many cones are involved; hence, the name "Polyconic projection."

Symmetry: About the central meridian or the Equator

Scale

True along the central meridian and along the Equator or two chosen parallels of true scale

Distortion

Free of distortion only along the central meridian. Although meridians intersect parallels at right angles, there is some angular distortion at all points other than along the central meridian.

Usage

Large-scale military mapping in Great Britain

Other names

War Office (British)



Figure 44.—Rectangular Polyconic projection, with shorelines, 10° graticule. Central meridian 90° W. Equator at true scale.

Similar projections

Listed under Polyconic projection (p. 106)

MODIFIED POLYCONIC Projection for the International Map of the World series

Classifications

Polyconic

Neither conformal nor equal area

Graticule

Meridians: Straight lines converging but not to a common point

Parallels: Equator is a straight line, poles are points, and all other parallels are nonconcentric circular arcs spaced at true distances along the two meridians 2° on either side of the central meridian (or 4° or 8° at latitudes beyond 60° N. or S.). Each parallel has a curvature identical to its curvature on a cone tangent at that latitude.

Poles: Points

Symmetry: About the central meridian.

Theoretically about the Equator, but the use is confined to zones that do not cross the Equator.

Scale

True along two meridians 2° on either side of the central meridian (or 4° or 8° at latitudes beyond 60° N. or S.) and along two parallels at the northern and southern edges of the quadrangles, which extend 4° in latitude, beginning at the Equator

Distortion

Not completely free of distortion anywhere, but distortion for the small quadrangles involved is very low

Other features

Quadrangles in the International Map of the World series using this projection can be joined at the edges from north to south or from east to west but not in all directions at once without creating gaps.

Usage

The official projection for all new quadrangle maps in the International Map of the World series from 1909 to 1962. It is still in use for existing quadrangles and for some new quadrangles. These quadrangles extend 4° in latitude and 6° in longitude, except that, beyond latitudes 60° N. or S., the width is 12° or 24° of longitude.

Origin

Presented by Charles Lallemand of France in 1909 specifically for the International Map of the World series

Similar projections

Listed under Polyconic projection (p. 106)




Figure 45.—Modified Polyconic projection for the International Map of the World series. Typical quadrant with 1° graticule. Lines of true scale are heavy lines.

BONNE Projection

Classifications

Pseudoconic

Equal area

Graticule

Meridians: Central meridian is a straight line. Other meridians are complex curves connecting points equally spaced along each parallel of latitude and concave toward the central meridian.

Parallels: Concentric circular arcs spaced at true distances along the central meridian.



The curvature of the central or standard parallel is identical to its curvature on a cone tangent at that latitude.

Poles: Points

Symmetry: About the central meridian

Scale

True along the central meridian and along each parallel



Distortion

Free of all distortion along the central meridian and the central parallel (fig. 46A)

Figure 46A.—Bonne projection, with Tissot indicatrices, 30° graticule.
Central parallel 40° N.

Figure 46C.—Werner projection, with shorelines, 30° graticule. Central meridian 90° W. Polar limit of Bonne projection.



Usage

Frequently used until recently for atlas maps of continents. Used in the ellipsoidal form for topographic mapping of France in the early 19th century.

Origin

Developed in rudimentary form by Claudius Ptolemy (about A. D. 100). Further developed by Bernardus Sylvanus in 1511. Used considerably by Rigobert Bonne (1727–95) of France, especially in 1752; the name is taken from him.

Limiting projections

Werner projection (fig. 46C and p. 114) if a pole is made the standard parallel


Sinusoidal projection (fig. 46D and p. 37) if the Equator is made the standard parallel.

The pseudoconic projection thereby becomes a pseudocylindrical projection, but the Bonne formulas must be rewritten.

Similar projection

John Bartholomew combined the Bonne projection with the Equidistant Conic projection with two standard parallels (p. 92). In his 1942 "Kite" and 1958 "Regional" projections, the Equidistant Conic is used in the north temperate zone, and the Bonne is used south of the Tropic of Cancer in interrupted form. The more northern regions are based on the Bonne ("Kite") and Conic ("Regional") projections. These two projections emphasize land masses, and Bartholomew's 1958 "Lotus" projection emphasizes the oceans by using a southern version of the "Regional" projection.

Figure 46D.—Sinusoidal projection, with shorelines, 30° graticule. Central meridian 90° W. Equatorial and pseudocylindrical limit of Bonne projection.



WERNER Projection

Classifications

Pseudoconic
Equal area
Cordiform (heart shaped)

Graticule

Meridians: Central meridian is a straight line. Other meridians are complex curves connecting points equally spaced along each parallel of latitude and concave toward the central meridian.

Parallels: Concentric circular arcs spaced at true distances along the central meridian.

One pole (usually the North Pole) is the center for these circular arcs.

Poles: Points

Symmetry: About the central meridian

Scale

True along the central meridian and along each parallel

Distortion

Free of all distortion only along the central meridian (fig. 47A)

Usage

Frequently used for world and some continental maps of the 16th and 17th centuries. Replaced by the Bonne projection for continental maps. No longer used except as a novelty.

Origin

Developed by Johannes Stabius (Stab) (?–1522) of Vienna about 1500 and promoted by Johannes Werner (1466–1528) of Nuremberg in 1514

Other names

Stab-Werner

Similar projections

Bonne projection is a general form on which the parallels are also circular arcs but are not generally centered on a pole. The Werner is the polar limiting form.

In 1904, Wilhelm Schjerning interrupted the Werner projection, centered on the South Pole, with three central meridians and three “petals” for a world map emphasizing the oceans.

Goode Polar Equal-Area projection, presented by John Paul Goode in 1928, is an interrupted Werner projection, centered on the North Pole and using several central meridians to display continents in a flowerlike pattern.

“Tetrahedral” projection, presented by John Bartholomew by 1944, is a combination of the Polar Azimuthal Equidistant projection (p. 133) from the pole to the tropic line of the same hemisphere and an interrupted Werner from there to the opposite pole, some adjustment being made to obtain a fit. The North Pole is the center for a map of land masses, and the South Pole is the center for an ocean map.

In 1968, W. William-Olsson of Sweden combined the northern Polar Lambert Azimuthal Equal-Area projection, extending to latitude 20° N., with four identical lobes extending to the South Pole. The lobes are based on the Werner projection but are compressed north-south and expanded east-west to fit the Lambert and to retain equality of area.

Figure 47A.—Werner projection, with Tissot indicatrices, 30° graticule.





Figure 47B.—Werner projection, with shorelines, 15° graticule. Central meridian 90° W.



GNOMONIC Projection

Classifications

Azimuthal

Perspective

Neither conformal nor equal area

Graticule

Polar aspect (fig. 48B):

Meridians: Equally spaced straight lines intersecting at the central pole. Angles between them are the true angles.

Parallels: Unequally spaced circles centered at the pole, which is a point. The Equator and the opposite hemisphere cannot be shown. Spacing increases rapidly away from the pole.

Symmetry: About any meridian

Equatorial aspect (fig. 48C):

Meridians: Unequally spaced straight parallel lines. Only meridians within 90° of the central meridian can be shown.

Spacing increases away from the central meridian.

Parallels: Equator is a straight line perpendicular to the meridians. Other parallels are hyperbolic arcs concave toward the nearest pole. Spacing increases rapidly away from the Equator. Poles cannot be shown.

Symmetry: About the central meridian or the Equator

Oblique aspect (fig. 48D):

Meridians: Unequally spaced straight lines intersecting at the pole nearest the center of the projection

Parallels: Equator is a straight line perpendicular to the central meridian. If the central parallel is a north latitude, its colatitude (90° – central latitude) is shown as a parabolic arc, more northern parallels are ellipses, and more southern parallels

are hyperbolas, all concave toward the nearest pole. If the central parallel is a south latitude, opposite signs apply.

Symmetry: About the central meridian

Range

Less than one hemisphere centered on a given pole of projection or map origin

Scale

True only where the central parallel crosses the central meridian. Increases rapidly with the distance from the center.

Distortion

Figure 48A shows distortion for the polar aspect. Other aspects have identical distortion at the same distance from the projection center. Only the center is free from distortion. Distortion increases rapidly away from the center.

Special features

All great circles (including all meridians and the Equator) are shown as straight lines. Therefore, the path of the shortest distance between any two points on the map is a straight line.

Perspective projection from the center of the globe onto a plane tangent at the central point of projection


Less than a hemisphere can be shown on one map.

Usage

To show great-circle paths as straight lines and thus to assist navigators and aviators in determining appropriate courses. Too much distortion of many kinds for regular geographic maps.

Used in the past for some star maps and for world globes in the shapes of polyhedra

Figure 48A.—Polar Gnomonic projection, with Tissot indicatrices, 30° graticule. Range 70°.



Origin

Possibly developed by Thales (c. 580 B.C.) of Greece. Name derived from the gnomon of a sundial, since meridians radiate like the hour markings on a sundial.

Other names


Gnomic

Central

Similar projection

Two-Point Azimuthal projection (p. 144) is a Gnomonic projection that has been compressed in one direction.

Figure 48B.—Polar Gnomonic projection, with shorelines, 10° graticule. Central meridian 90° W. Range 60°.



GNOMONIC Projection




Figure 48C.—Equatorial Gnomonic projection, with shorelines, 10° graticule. Central meridian 90° W. Range 60°.




Figure 48D.—Oblique Gnomonic projection, with shorelines, 10° graticule, central latitude 40° N., central meridian 90° W. Range 60°.

STEREOGRAPHIC Projection

Classifications

Azimuthal
Conformal
Perspective
Polyconic (general)

Graticule

Polar aspect (fig. 49B):

Meridians: Equally spaced straight lines intersecting at central pole. Angles between them are the true angles.

Parallels: Unequally spaced circles centered at the pole, which is a point. Opposite pole cannot be shown. Spacing gradually increases away from the pole.

Symmetry: About any meridian

Equatorial aspect (fig. 49C):

Meridians: Central meridian is a straight line. Other meridians are unequally spaced circular arcs intersecting at each pole. Spacing gradually increases away from the central meridian.

Parallels: Equator is a straight line. Other parallels are unequally spaced circular arcs concave toward the nearest pole. Spacing gradually increases away from the Equator along the central meridian, but spacing is equal along the meridians 90° from the central meridian.

Symmetry: About the central meridian or the Equator

Oblique aspect (fig. 49D):

Meridians: Central meridian is a straight line. Other meridians are circular arcs intersecting at each pole and having their centers on the straight parallel opposite in sign to the central latitude. Spacing gradually increases away from the central meridian.

Parallels: The parallel opposite in sign to the central latitude is a straight line perpendicular to the central meridian. All other parallels are circular arcs concave toward the pole on the same side of the straight parallel.

Symmetry: About the central meridian

Range

One hemisphere conveniently; most but not all of the other hemisphere at accelerating scale

Scale

True only where the central latitude crosses the central meridian or, alternatively, along a circle concentric about the projection center (or a parallel on the polar aspect). Scale is constant along any circle having its center at the projection center, but scale increases moderately with distance from the center within a hemisphere.

Distortion


Figure 49A shows distortion for the polar aspect. Other aspects have identical distortion at the same distance from the projection center. Only the center or the circle of true scale (if not the center) is free from all distortion.

Special features

All great or small circles (including all meridians and parallels) on the globe are shown as circular arcs or straight lines.

Perspective projection of the globe onto a tangent (or secant) plane from a point on the surface just opposite the point of tangency or the projection center. The straight parallel of latitude passes through this point.

Figure 49A.—Polar Stereographic projection, with Tissot indicatrices, 30° graticule. Range 100°.



Usage

Commonly used in the polar aspect for topographic maps of polar regions. The equatorial aspect was used regularly for maps of the Eastern and Western Hemispheres in the 17th and 18th centuries. Oblique aspects are used to show paths of solar eclipses.


Recommended for conformal mapping of regions approximately circular in shape

Basis of modified Stereographic conformal projections using complex algebra, such as the Miller Oblated Stereographic (p. 148)

Origin

Apparently developed in polar aspect by Egyptians and Greeks by the 2d century B.C.

Figure 49B.—Polar Stereographic projection, with shorelines, 10° graticule. Central meridian 90° W. Range 90°.



STEREOGRAPHIC Projection




Figure 49C.—Equatorial Stereographic projection, with shorelines, 10° graticule. Central meridian 90° W. Range 90°.




Figure 49D.—Oblique Stereographic projection, with shorelines, 10° graticule, central latitude 40° N. Central meridian 90° W. Range 90°.

ORTHOGRAPHIC Projection

Classifications

Azimuthal

Perspective

Neither conformal nor equal area

Graticule

Polar aspect (fig. 50B):

Meridians: Equally spaced straight lines intersecting at the central pole. Angles between them are the true angles.

Parallels: Unequally spaced circles centered at the pole, which is a point. Spacing decreases away from the pole.

Symmetry: About any meridian

Equatorial aspect (fig. 50C):

Meridians: Central meridian is a straight line. Outer meridians 90° away are circles. Other meridians are unequally spaced elliptical arcs intersecting at each pole and concave toward the central meridian. Spacing decreases away from the central meridian.

Parallels: Unequally spaced straight parallel lines perpendicular to the central meridian. Spacing decreases away from the Equator. Parallels intersect the outer meridian, however, at equal intervals.

Symmetry: About the central meridian or the Equator

Oblique aspect (fig. 50D):

Meridians: Central meridian is a straight line. Other meridians are semiellipses intersecting at each pole. Spacing

decreases away from the central meridian.

Parallels: Complete or partial ellipses, all of the same shape (or eccentricity); their minor axes lie along the central meridian. Spacing decreases away from the center of projection.

Symmetry: About the central meridian

Range

No more than one hemisphere at a time

Scale

True at the center and along any circle having its center at the projection center but only in the direction of the circumference of the circle. Scale decreases radially with distance from the center.

Distortion

Figure 50A shows distortion for the polar aspect. Other aspects have identical distortion at the same distance from the projection center. Only the center is free of distortion, which increases rapidly away from the center. Extreme distortion near the edge of the hemisphere.

Special features

Perspective projection of the globe onto a tangent plane from an infinite distance (that is, orthogonally); thus, the map has the look of a globe.

All great or small circles (including all meridians and parallels) are shown as elliptical arcs or straight lines.




Figure 50A.—Polar Orthographic projection, with Tissot indicatrices, 30° graticule. Range 80°.

Usage

Pictorial views of the Earth, resembling those seen from space


Origin

Apparently developed by Egyptians and Greeks by the 2d century B.C.

Similar projections

General Vertical Perspective (p. 128), when projected from more than a few thousand kilometers above the Earth

Figure 50B.—Polar Orthographic projection, with shorelines, 10° graticule. Central meridian 90° W.



ORTHOGRAPHIC Projection




Figure 50C.—Equatorial Orthographic projection, with shorelines, 10° graticule. Central meridian 90° W.



Figure 50D.—Oblique Orthographic projection, with shorelines, 10° graticule, central latitude 40° N. Central meridian 90° W.

GENERAL VERTICAL PERSPECTIVE Projection

Classifications

Azimuthal

Perspective

Neither conformal nor equal area (except in Stereographic form)

Graticule

Polar aspect (fig. 51A):

Meridians: Equally spaced straight lines intersecting at the central pole. Angles between them are the true angles.

Parallels: Unequally spaced circles centered at the pole, which is a point. If the usual perspective from above the surface is projected, the spacing of parallels decreases away from the pole, and the Equator cannot be shown. If point of perspective is beneath the surface, the spacing may increase or decrease away from the pole, and part of the opposite hemisphere can be shown.

Symmetry: About any meridian

Equatorial aspect (fig. 51B):

Meridians: Central meridian is a straight line. Other meridians are generally unequally spaced elliptical arcs intersecting at the poles (which may not be visible).

Parallels: Equator is a straight line. Other parallels are usually elliptical arcs. If point of perspective is beneath the surface, the parallels can be elliptical, parabolic, or hyperbolic arcs.

Symmetry: About the central meridian or the Equator

Oblique aspect (figs. 51C, 51D):

Meridians: Central meridian is a straight line. Other meridians are generally unequally spaced elliptical arcs intersecting at each pole.




Figure 51A.—Polar Vertical Perspective projection (La Hire form), with shorelines, 10° graticule. Central meridian 90°W.. Depth below surface 17,250 km.

Parallels: One parallel, normally nearer the pole than is the center of projection, depending on the position of the point of perspective, is a straight line. Other parallels are normally elliptical arcs.

Symmetry: About the central meridian

Range

If point of perspective is above the surface, less than one hemisphere

If point of perspective is on the opposite side of the Earth's center, more than one hemisphere

Scale

True at the center, if the projection is onto a tangent plane. Scale varies away from the center, depending on the point of perspective.

Distortion

Only the center is free of distortion, if projection is onto a tangent plane. Distortion is severe near the outer limit of the projection, although special locations for the points of projection on the opposite side of the Earth's center result in moderate distortion within a wide range of the center.

Special features

The General Vertical Perspective projection shows the Earth geometrically as it appears from any point above or below the surface when it is projected onto a tangent or secant plane perpendicular to the straight line connecting the point of perspective with the center of the Earth.

Usage

Pictorial views of the Earth, resembling those seen from space




Figure 51B.—Equatorial Vertical Perspective projection, with shorelines, 10° graticule. Central meridian 90°W. Height above surface 35,900 km (geosynchronous satellite position).

GENERAL VERTICAL PERSPECTIVE Projection

Origin

Aside from the earlier limiting forms listed below, the perspective projection from a point on the opposite side of the Earth's center was studied in the 18th and 19th centuries as a source of low-distortion projections having geometric construction.

Limiting forms

Gnomonic projection (p. 116), if the point of perspective is at the center of the Earth

Stereographic projection (p. 120), if the point of perspective is on the surface of the Earth opposite the center of projection

Orthographic projection (p. 124), if the point of perspective is at infinity

In turn, the Vertical Perspective is a limiting form of the Tilted Perspective projection (p. 172).

Other specific forms

In 1701, Philippe de la Hire of France suggested a point of projection at 1.7071 times the radius (fig. 51A). This distance is measured from the



Figure 51C.—Oblique Vertical Perspective projection, with shorelines, 5° graticule, central latitude 50°N. Central meridian 90°W. Height above surface 2,000 km.

center in a direction away from the surface of projection and gives the Equator of the polar aspect twice the radius of 45° latitude. Antoine Parent in 1702, Ernst Hammer about 1890, H.F. Gretschen in 1873, and Hans Maurer in 1935 were among others proposing points of projection for low-error perspectives.

A.R. Clarke and Henry James, in 1862 and later, calculated points of projection (on the far side, below the surface of projection) for minimum-error perspective azimuthal projections of regions, such as continents, bounded by a given great or small circle on the Earth. Figure 51D is Clarke's Twilight projection of 1879, bounded by a circle 108° in radius.

The Army Map Service AMS Lunar projection is a far-side perspective selected by Albert L. Nowicki in 1963 to show slightly more than a hemisphere of the Moon, as seen over a period of time from the Earth. The point of projection is 1.53748 times the radius from the center.

See also the Airy nonperspective azimuthal projection (p. 140).



Figure 51D.—Clarke's Twilight projection, with shorelines, 15° graticule, central latitude 23.5° N. Central meridian 0° . Depth below surface 15,290 km. Range 108° .

AZIMUTHAL EQUIDISTANT Projection

Classifications

Azimuthal
Equidistant
Nonperspective

Graticule

Polar aspect (fig. 52B):

Meridians: Equally spaced straight lines intersecting at the central pole. Angles between them are the true angles.

Parallels: Equally spaced circles, centered at the pole, which is a point. The entire Earth can be shown, but the opposite pole is a bounding circle having a radius twice that of the Equator.

Symmetry: About any meridian

Equatorial aspect (fig. 52C):

Meridians: Central meridian is a straight line. Meridian 90° away is a circle. Other meridians are complex curves, equally spaced along the Equator and intersecting at each pole.

Parallels: Equator is a straight line. Other parallels are complex curves concave toward the nearest pole and equally spaced along the central meridian and the meridian 90° from the central meridian.

Symmetry: About the central meridian or the Equator

Oblique aspect (fig. 52D):

Meridians: Central meridian is a straight line. Other meridians are complex curves intersecting at each pole.

Parallels: Complex curves equally spaced along the central meridian

Symmetry: About the central meridian

Range

Entire Earth

Scale

True along any straight line radiating from the center of projection. Increases in a direction perpendicular to the radius as the distance from the center increases.

Distortion

Figure 52A shows distortion for the polar aspect. Other aspects have identical distortion at the same distance from the projection center. Only the center is free from distortion. Distortion is moderate for one hemisphere but becomes extreme for a map of the entire Earth.

Special features

The distance between any two points on a straight line passing through the center of projection is shown at true scale; this feature is especially useful if one point is the center.

Compromise in distortion between Stereographic (conformal) and Lambert Azimuthal Equal-Area projections

Usage

Commonly used in the polar aspect for maps of polar regions, the Northern and Southern Hemispheres, and the "aviation-age" Earth. The oblique aspect is frequently used for world maps centered on important cities and occasionally for maps of continents. The ellipsoidal form is used for topographic mapping of Micronesia and Guam.

Origin

Possibly developed in the polar aspect by Egyptians for star charts. Several users during the 16th century.

Other names

Postel (in France and Russia, for Guillaume

Postel, who was considered an originator, although he first used it in 1581)

Zenithal Equidistant

Similar projections

Two-Point Azimuthal (p. 144) shows correct azimuths (but not distances) from either of two points to any other point.

Two-Point Equidistant (p. 146) shows correct distances (but not azimuths) from either of two points to any other point.

Chamberlin Trimetric (p. 170) shows approximately true distances from three chosen points to any other points (cannot be exact). The three points are placed near the edges of the region being mapped to reduce overall distortion.

Airy (p. 140) and Breusing (p. 143) azimuthal projections have spacings very similar to those of the Azimuthal Equidistant if the extent is less than one hemisphere.

Berghaus Star projection (p. 156) uses the Polar Azimuthal Equidistant projection for the Northern Hemisphere.

"Tetrahedral" projection (p. 114) combines the Polar Azimuthal Equidistant projection with an interrupted Werner projection.

Figure 52A.—Polar Azimuthal Equidistant projection, with Tissot indicatrices, 30° graticule. Range 170°.





Figure 52B.—Polar Azimuthal Equidistant projection, with shorelines, 15° graticule. Central meridian 90° W. Range 180° (world map).

AZIMUTHAL EQUIDISTANT Projection




Figure 52C.—Equatorial Azimuthal Equidistant projection, with shorelines, 15° graticule. Central meridian 90° W. Range 180° (world map).




Figure 52D.—Oblique Azimuthal Equidistant projection, with shorelines, 15° graticule, central latitude 40° N. Central meridian 90° W. Range 180° (world map).

LAMBERT AZIMUTHAL EQUAL-AREA Projection

Classifications

Azimuthal
Equal area
Nonperspective

Graticule

Polar aspect (fig. 53B):

Meridians: Equally spaced straight lines intersecting at the central pole. Angles between them are the true angles.

Parallels: Unequally spaced circles, centered at the pole, which is a point. Spacing of the circles gradually decreases away from the pole. The entire Earth can be shown, but the opposite pole is a bounding circle having a radius 1.41 times that of the Equator.

Symmetry: About any meridian

Equatorial aspect (fig. 53C):

Meridians: Central meridian is a straight line. Meridian 90° away is a circle. Other meridians are complex curves, unequally spaced along the Equator and intersecting at each pole. Spacing decreases away from the central meridian.

Parallels: Equator is a straight line. Other parallels are complex curves concave toward the nearest pole. They are unequally spaced along the central meridian, and spacing decreases away from the Equator. Along the meridian 90° from the central meridian, parallels are equally spaced.

Symmetry: About the central meridian or the Equator

Oblique aspect (fig. 53D):

Meridians: Central meridian is a straight line. Other meridians are complex curves intersecting at each pole shown.

Parallels: Complex curves unequally spaced along the central meridian; spacing decreases away from the center of projection

Symmetry: About the central meridian

Range

Entire Earth

Scale

True only at the center in all directions. Decreases with distance from the center along radii. Increases with distance from the center in a direction perpendicular to radii.

Distortion

Figure 53A shows distortion for the polar aspect. Other aspects have identical distortion at the same distance from the projection center. Only the center is free from distortion. Distortion is moderate for one hemisphere but becomes extreme for a map of the entire Earth.

Usage

Frequently used in the polar aspect in atlases for maps of polar regions and of Northern and Southern Hemispheres. The equatorial aspect is commonly used for atlas maps of the Eastern and Western Hemispheres. The oblique aspect is used for atlas maps of continents and oceans. The equatorial and oblique aspects are used by the U.S. Geological Survey in cooperation with others for maps of the Circum-Pacific Map Project.

Recommended for equal-area maps of regions approximately circular in extent

Origin

Presented by Johann Heinrich Lambert (1728-77) of Alsace in 1772




Figure 53A.—Polar Azimuthal Equal-Area projection, with Tissot indicatrices, 30° graticule. Range 100°.


Other names

Lorgna (for polar aspect, due to independent derivation, 1789)
Zenithal Equal-Area
Zenithal Equivalent

Similar projections

W. William-Olsson's projection (p. 114) combines the Lambert Azimuthal Equal-Area with a modified Werner.

Figure 53B.—Polar Lambert Azimuthal Equal-Area projection, with shorelines, 10° graticule. Central meridian 90° W. Range 90°.



LAMBERT AZIMUTHAL EQUAL-AREA Projection



Figure 53C.—Equatorial Lambert Azimuthal Equal-Area projection, with shorelines, 105° graticule. Central meridian 90° W. Range 90°.



Figure 53D.—Oblique Lambert Azimuthal Equal-Area projection, with shorelines, 10° graticule, central latitude 40° N. Central meridian 90° W. Range 90°.



Figure 53E.—Oblique Lambert Azimuthal Equal-Area projection, with shorelines, 30° graticule, central latitude 40° N. Central meridian 90° W. Range 180° (world map).

AIRY Projection

Classifications

Azimuthal

Minimum error

Nonperspective

Neither conformal nor equal area

Graticule

Polar aspect (fig. 54B):

Meridians: Equally spaced straight lines intersecting at the central pole. Angles between them are the true angles.

Parallels: Nearly but not quite equally spaced circles, centered at the pole, which is a point. Spacing increases slightly away from the center.

Symmetry: About any meridian

Equatorial aspect (fig. 54C):

Meridians: Central meridian is a straight line. Meridians 90° away are circles. Other meridians are complex curves, unequally spaced along the Equator and intersecting at each pole. Spacing increases slightly away from the central meridian.

Parallels: Equator is a straight line. Other parallels are complex curves concave toward the nearest pole. They are unequally spaced along the central meridian, and spacing increases slightly away from the Equator. At the meridian 90° from the central meridian, parallels are equally spaced.

Symmetry: About the central meridian or the Equator

Oblique aspect (fig. 54D):

Meridians: Central meridian is a straight line. Other meridians are complex curves intersecting at each pole.

Parallels: Complex curves unequally spaced along the central meridian; spacing

increases slightly away from the center of projection

Symmetry: About the central meridian

Range

Should be limited to about a hemisphere or less

Scale

Scale increases gradually in all directions with distance from the center, but there is no point at which the scale is correct in all directions.

Distortion

Figure 54A shows distortion for the polar aspect extending 95° or slightly more than one hemisphere. Distortion is moderate throughout the range chosen. No point is completely free of distortion, although the center is free of shape distortion.

Usage

Medium-scale Ordnance Survey map of the United Kingdom

Special feature

The minimum-error azimuthal projection of the region enclosed by a great or small circle of chosen radius from a given center

Origin


Presented by George Biddell Airy (1801–92), British astronomer and geodesist, in 1861

Similar projections

Clarke and James minimum-error perspective azimuthal projections were inspired by Airy's work.

The Azimuthal Equidistant projection (p. 132) closely resembles the Airy, but scale is

Figure 54A.—Polar Airy projection, with Tissot indicatrices, 30° graticule. Range 95°.




constant along radii from the center.

In 1892, A. Breusing proposed the low-error geometric mean of the Stereographic and Lambert Azimuthal Equal-Area projections.

In 1920, A.E. Young of England modified the Breusing (geometric mean) projection to provide a balance between the Stereographic and Azimuthal Equidistant projections, called the Breusing Harmonic projection (fig. 54E).

In 1950, N.A. Urmayev of the Soviet Union proposed two other low-error azimuthal projections.

Figure 54B.—Polar Airy projection, with shorelines, 10° graticule. Central meridian 90° W. Range 90°.



AIRY Projection



Figure 54C.—Equatorial Airy projection, with shorelines, 10° graticule. Central meridian 90° W. Range 90°.



Figure 54D.—Oblique Airy projection, with shorelines, 10° graticule, central latitude 40° N. Central meridian 90° W. Range 90°.




Figure 54E.—Polar Breusing Harmonic projection, with shorelines, 30° graticule. Central meridian 90° W. Range 180° (world map).

TWO-POINT AZIMUTHAL Projection

Classifications

Modified azimuthal
Neither conformal nor equal area

Graticule

Meridians: Straight lines
Parallels: Equator is straight. Other parallels are ellipses, parabolas, or hyperbolas.
Poles: Points (if shown)
Symmetry: Normally none

Scale

Varies considerably. Projection should not be used for measuring distance

Distortion

Direction is correct to all other points from two central points at which there is no distortion.

Other features

Great circles are shown as straight lines.
This projection can be obtained by compressing a Gnomonic projection in the direction parallel to the line joining the two central points.

Usage

Could be used to locate a ship at sea, for example, if the exact location of two radio transmitters and the direction from the ship to the transmitters are known

Origin

Presented by Hans Maurer (1868–1945) of Germany in 1914 and Charles F. Close (1865–1952) of England independently in 1922

Aspects

Typically oblique

Other names

Doubly Azimuthal
Orthodromic
Close
McCaw
Immler


Limiting forms

If the two central points are identical, the Gnomonic projection results.

Similar projections

Gnomonic (p. 116), which can be compressed as described above to produce the Two-Point Azimuthal projection

Figure 55.—Two-Point Azimuthal projection, with shorelines, 10° graticule. Central points are near Washington, D.C. (longitude 77° W., latitude 39° N.) and Honolulu, Hawaii (longitude 158° W., latitude 21.5° N.).



TWO-POINT EQUIDISTANT Projection

Classifications

Modified azimuthal
Neither conformal nor equal area

Graticule

Meridians: Complex curves
Parallels: Complex curves
Poles: Normally points
Symmetry: Normally none

Scale

True along a straight line from either of two central points

Distortion

No points free of distortion

Usage

Map of Asia by the National Geographic Society

Could be used to determine the distance from a ship at sea at a known location from the start and end of a voyage

Origin

Presented by Hans Maurer (1868–1945) of Germany in 1919 and Charles F. Close (1865–1952) of England independently in 1921

Aspects

Typically oblique

Other names

Doubly Equidistant

Limiting forms

If the two central points are identical, the Azimuthal Equidistant projection results.

Similar projections


Azimuthal Equidistant (p. 132), on which distance is correct from only one point

Donald Elliptical, developed by Jay K. Donald of the American Telephone and Telegraph Company in 1956. Used by telephone companies to establish long-distance rates, the Donald is a Two-Point Equidistant projection specially modified for the ellipsoid and confined to the United States and southern Canada.

Chamberlin Trimetric (p. 170), an approximate three-point equidistant projection

Two-Point Azimuthal-Equidistant, presented by Charles F. Close in 1922, has true azimuths from one point and true distances from a second point to all other points.

Figure 56.—Two-Point Equidistant projection, with shorelines, 10° graticule. Central points are near Washington, D.C. (longitude 77° W., latitude 39° N.), and Honolulu, Hawaii (longitude 158° W., latitude 21.5° N.).



MILLER OBLATED STEREOGRAPHIC Projection

Classifications

Modified azimuthal

Conformal

Graticule

Meridians: Complex curves

Parallels: Complex curves

Poles: Points

Symmetry: About the central meridian as applied to the map of Europe and Africa. No symmetry on other prepared maps.

Scale

A line of constant scale bounds an oval region.

Distortion

Scale variations within the oval-shaped region covered are less than variations on standard conformal projections, such as an oblique Mercator or oblique Stereographic projection.

Usage

Used for an area that can be contained within an oval shape to minimize the scale distortions in the area. Applied by O.M. Miller to Africa and Europe in 1953 and to other Eastern Hemisphere regions in conjunction with several non-conformal fill-in projections in 1955. Used by L.P. Lee for the Pacific Ocean in 1974.

Origin

Osborn Maitland Miller (1897–1979) of the American Geographical Society in 1953

Other names

Miller Oblated Stereographic

Oblated Stereographic

Prolated Stereographic

Similar projections

Stereographic projection (p. 120) has lines of constant scale that are concentric circles.




Figure 57.—Miller Oblated Stereographic projection of Europe and Africa, with shorelines, 10° graticule, and lines of constant scale. Scale factors are 0.94, 0.96, 1.00, 1.04, and 1.08 along ovals, beginning with the smallest. Scale factor at center (latitude 18° N., longitude 20° E.) is 0.9245.

WIECHEL Projection

Classifications

Modified azimuthal
Equal area
Pseudoazimuthal

Graticule

Polar aspect:

- Meridians: Semicircles if extended to a world map
- Parallels: Concentric circles
- Poles: Central pole is a point; other pole is a circle
- Symmetry: Radial about the central pole

Scale

Correct along meridians

Distortion

Considerable distortion of shape at the map edge

Usage

A novelty map that, in its polar aspect, has semicircular meridians arranged in a pinwheel

Origin

Presented by H. Wiechel in 1879

Aspects

Normally only polar

Similar projections


Lambert Azimuthal Equal-Area projection (p. 136) has the same spacing of parallels in the polar aspect, but the meridians are straight.

Ginzburg III is a general series of pseudoazimuthal projections developed by G.A. Ginzburg of the Soviet Union in 1952.

The lines of constant distortion are ovals rather than the usual circles of azimuthal projections.

Figure 58.—Polar Wiechel projection. Northern Hemisphere, with shorelines, 10° graticule.

This feature is achieved by the manner in which the lines of constant initial azimuth are curved. The projection series has been used only in the oblique or equatorial aspect.



CRAIG RETROAZIMUTHAL Projection

Classifications

Retroazimuthal
Modified azimuthal
Neither conformal nor equal area

Graticule

Meridians: Equally spaced straight parallel lines
Parallels: Curved
Poles: Curved lines
Symmetry: About the central meridian

Distortion

Extreme distortion of shape and area

Other features

The direction is correct from any location on the map to a central point.
Considerable overlapping when entire sphere is shown

Usage

To show, for example, the correct direction toward Mecca, from any point on the map, by measuring the angle from North. Because the meridians are straight parallel lines, this angle can be readily measured.

Origin

Presented by James Ireland Craig (1868–1952) of the Survey of Egypt in 1909

Other names

Mecca


Similar projections

Hammer Retroazimuthal (p. 152) has curved, unequally spaced meridians.

Hans Maurer of Germany in 1919 and J.E. Jackson of England in 1968 devised retroazimuthal projections characterized by straight, parallel, unequally spaced meridians.

Carl Schoy presented a projection in 1913 having meridians that are straight, equidistant, and parallel but parallels that are curved, so that all great circles passing through the central point are straight lines at their true azimuths from center. It is not retroazimuthal, nor is its distortion only a function of distance from the center, as it is on most azimuthal projections.

Figure 59.—Craig Retroazimuthal projection, centered near St. Louis, Mo. (longitude 90° W., latitude 38.5° N.), with shorelines, 10° graticule.



HAMMER RETROAZIMUTHAL Projection

Classifications

Retroazimuthal
Modified azimuthal
Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight. Other meridians are curved.

Parallels: Curved

Poles: Curved lines

Symmetry: About the central meridian

Distortion

Extreme distortion of area and shape

Other features

The direction from any point to the center of the map is the angle that a straight line connecting the two points makes with a vertical line. This feature is the basis of the term "retroazimuthal." Scimitar-shaped boundary. Considerable overlapping when the entire sphere is shown.

Usage

To determine the direction of a central point from a given location

Origin

Presented by H.H. Ernst von Hammer (1858–1925) of Germany in 1910. Independently presented by Edward A. Reeves (1862–1945) and Arthur R. Hinks (1874–1945) of England in 1929.

Similar projections

Craig Retroazimuthal (p. 150) has straight, equally spaced meridians.





Figure 60.—Hammer Retroazimuthal projection, centered near St. Louis, Mo. (longitude 90° W., latitude 40° N.), with shorelines, 15° graticule. Two hemispheres, one of which appears backwards; they should be superimposed for the full map.



LITTROW Projection

Classifications

Retroazimuthal
Conformal

Graticule

Meridians: Central meridian and meridians 90° away are straight. Others are hyperbolas convex toward the central meridian.

Parallels: Equator is straight. Other parallels are ellipses; the Equator lies along their major axes.

Poles: Not shown

Symmetry: About the central meridian and the Equator

Scale

Varies rapidly

Distortion

Extreme distortion of area

Other features

Cannot show the entire sphere. Shows the correct direction from any point on the map to any point along the central meridian as the angle between a straight line connecting the two points and a vertical straight line parallel to the central meridian.

Usage

By the British Navy in the 19th century to determine directions to a central point from other locations


Origin

Presented by J.J. Littrow of Austria in 1833

Similar projections

Littrow projection is the transverse aspect of a Lagrange projection (p. 180) having certain parameters.

Figure 61.—Litrow projection, one hemisphere, with shorelines,
10° graticule. Central meridian 90° W.



BERGHAUS STAR Projection

Classifications

Equidistant
Composite
Modified azimuthal
Interrupted

Graticule

Meridians: Straight lines, broken at the Equator
Parallels: Concentric circular arcs; North Pole at the center
Poles: Points
Symmetry: Radial about the North Pole

Scale

Correct along meridians in the Northern Hemisphere and along the central meridians of triangular points in the Southern Hemisphere

Distortion

Moderate throughout

Other features

The Northern Hemisphere is a polar Azimuthal

Equidistant projection. The Southern Hemisphere is divided into five equal triangular lobes.

Usage

Largely for artistic map forms. Landforms do not readily fit into the lobes or arms of the star pattern. Used for the logo of the Association of American Geographers.

Origin

Presented by Heinrich Berghaus (1797–1884) of Germany in 1879

Aspects

Polar only


Similar projections

Petermann Star (Germany, 1865) has eight unequal points; parallels are concentric circular arcs.

Jäger Star (Austria, 1865) has eight unequal points; parallels are in straight segments.

Azimuthal Equidistant (p. 132) is the projection for the Northern Hemisphere of the Berghaus Star.

Figure 62.—Berghaus Star projection, with shorelines, 18° graticule. Central meridian 90° W. World map.



AITOFF Projection

Classifications

Modified azimuthal

Neither conformal nor equal area

Graticule

Meridians: Central meridian is a straight line half the length of the Equator. Other meridians are complex curves, equally spaced along the Equator and concave toward the central meridian.

Parallels: Equator is straight. Other parallels are complex curves, equally spaced along the central meridian and concave toward the nearest pole.

Poles: Points

Symmetry: About the Equator and the central meridian

Scale

True along the Equator and the central meridian

Distortion

Moderate shape and area distortion (fig. 63A)


Other features

Elliptical border. An equatorial aspect of one hemisphere of the Azimuthal Equidistant projection, on which horizontal coordinates have been doubled and meridians have been given twice their original longitudes.

Usage

Design inspired H.H. Ernst von Hammer to develop his elliptical projection (p. 160)

Figure 63A.—Aitoff projection, with Tissot indicatrices, 30° graticule.



Origin

David Aitoff (or Aitow) (1854–1933) in 1889

Similar projections


Hammer (Elliptical) (p. 160), equal area and mistakenly called Aitoff in the early 20th century

Briesemeister (p. 162), an oblique modification of the Hammer

Eckert-Greifendorff (p. 166), a modification of the Lambert Azimuthal Equal-Area projection inspired by the Hammer

Winkel Tripel (p. 164) is a combination of the Aitoff and Equidistant Cylindrical projections.

Figure 63B.—Aitoff projection, with shorelines, 15° graticule. Central meridian 90° W.



HAMMER Projection

Classifications

Modified azimuthal

Equal area

Graticule

Meridians: Central meridian is a straight line half the length of the Equator. Other meridians are complex curves, unequally spaced along the Equator and concave toward the central meridian

Parallels: Equator is straight. Other parallels are complex curves, unequally spaced along the central meridian and concave toward the nearest pole.

Poles: Points

Symmetry: About the central meridian and the Equator

Scale

Decreases along the central meridian and the Equator with distance from the center

Distortion

Moderate (fig. 64A). Less shearing action on the outer meridians near the poles than there is on pseudocylindrical projections.

Other features

Elliptical border. An equatorial aspect of one hemisphere of the Lambert Azimuthal Equal-Area projection, on which horizontal coordinates have been doubled and meridians have been given twice their original longitudes.


Usage

Whole-world maps, also in interrupted and condensed forms

Origin

Presented by H.H. Ernst von Hammer (1858–1925) of Germany in 1892

Figure 64A.—Hammer projection, with Tissot indicatrices, 30° graticule.



Aspects

Normal and oblique are both in use.

Other names

Hammer-Aitoff


Mistakenly called Aitoff projection in the early 20th century.

Similar projections

Briesemeister (p. 162), a modified oblique Hammer

Eckert-Greifendorff (p. 166), a further modification of the Lambert Azimuthal Equal-Area projection

Figure 64B.—Hammer projection, with shorelines, 15° graticule. Central meridian 90° W.



BRIESEMEISTER Projection

Classifications

Modified azimuthal
Equal area

Graticule

Meridians: Central meridian is straight. Other meridians are complex curves.

Parallels: Complex curves

Poles: Points

Symmetry: About the central meridian

Scale

True scale at latitudes 3.6° and 86.4° N. at central meridian

Distortion

None at latitudes 3.6° and 86.4° N. at central meridian (fig. 65A)

Other features

An oblique Hammer projection centered at latitude 45° N. and longitude 10° E. and bounded by an ellipse having an axis ratio of 1.75 to 1 instead of 2 to 1, after the vertical coordinates are expanded and the horizontal coordinates are contracted

Usage

Whole-world maps showing continents grouped near the center

Origin

Presented by William A. Briesemeister of the American Geographical Society in 1953

Similar projections

Hammer (Elliptical) (p. 160), the normal aspect of the Briesemeister, but having an axis ratio of 2 to 1

Nordic, by John Bartholomew, is the same as the Briesemeister projection, except that the axis ratio is 2 to 1 and the central meridian is 0°.





Figure 65A.—Briesemeister projection, with Tissot indicatrices, 30° graticule.

Figure 65B.—Briesemeister projection, with shorelines, 15° graticule. Center at 45° N., 10° E.



WINKEL TRIPEL Projection

Classifications

Modified azimuthal

Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight. Other meridians are curved, equally spaced along the Equator and concave toward the central meridian.

Parallels: Equator and the poles are straight. Other parallels are curved, equally spaced along the central meridian and concave toward the nearest pole.

Poles: Straight lines about 0.4 as long as the Equator, depending on the latitude of the standard parallels of the base Equidistant Cylindrical projection

Symmetry: About the central meridian or the Equator

Scale

True along the central meridian. Constant along the Equator.

Distortion

Moderate except near outer meridians in polar regions (fig. 66A)

Other features

Equally spaced central meridian and Equator

Usage

Whole-world maps

Origin

Presented by Oswald Winkel (1873–1953) of Germany in 1921. Obtained by averaging coordinates of Equidistant Cylindrical and Aitoff (not Hammer-Aitoff) projections. Winkel applied the name “Tripel,” normally meaning triple.





Figure 66A.—Winkel Tripel projection, with Tissot indicatrices, 30° graticule. Standard parallels the same as that in figure 66B.

Figure 66B.—Winkel Tripel projection, with shorelines, 15° graticule. Central meridian 90° W. Standard parallels of Cylindrical Equidistant base 40° N. and S.



ECKERT-GREIFENDORFF Projection

Classifications

Modified azimuthal

Equal area

Graticule

Meridians: Central meridian is a straight line 0.46 of the length of the Equator. Other meridians are curved, unequally spaced along the Equator and concave toward the central meridian.

Parallels: Equator is straight. Other parallels are curved, unequally spaced along the central meridian and concave toward the nearest pole.

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

Decreases steadily along the Equator and the central meridian with distance from the center of the projection

Distortion

Moderate near center, but considerable shape distortion near outer meridians

Other features

An equatorial aspect of the Lambert Azimuthal Equal-Area projection for 90° of longitude, on which the horizontal coordinates have been quadrupled and the meridians have been given four times their original longitudes

Usage

Whole-world map

Origin


Max Eckert-Greifendorff (formerly Max Eckert) (1868–1938) of Germany in 1935

Similar projections

Hammer (p. 160), a different modification of the Lambert Azimuthal Equal-Area projection

Quartic Authalic (p. 74) has straight parallels.

Figure 67.—Eckert-Greifendorff projection, with shorelines, 15° graticule. Central meridian 90° W.



WAGNER VII Projection

Classifications

Modified azimuthal
Equal area

Graticule

Meridians: Central meridian is straight and half the length of the Equator. Other meridians are curves, unequally spaced along the Equator and concave toward the central meridian.

Parallels: Equator is straight. Other parallels are curves, unequally spaced along the central meridian and concave toward the nearest pole.

Poles: Curved lines

Symmetry: About the central meridian or the Equator

Scale

Decreases along the central meridian and the Equator with distance from the center of the projection

Distortion

Considerable shape distortion in polar areas (fig. 68A)

Other features

A modification of the Hammer projection; the poles correspond to the 65th parallels on the Hammer, and meridians are repositioned

Usage

World maps, such as climatic maps prepared by the U.S. Department of Commerce




Figure 68A.—Wagner VII projection, with Tissot indicatrices, 30° graticule.

Origin

Presented by Karlheinz Wagner of Germany in 1941


Other names

Hammer-Wagner

Similar projections

Hammer (Elliptical) (p. 160), truncated and otherwise modified for the Wagner VII

Figure 68B.—Wagner VII projection, with shorelines, 15° graticule. Central meridian 90° W.



CHAMBERLIN TRIMETRIC Projection

Classifications

Modified azimuthal
Modified equidistant
Neither conformal nor equal area

Graticule

Meridians: Complex curves
Parallels: Complex curves
Poles: Points
Symmetry: Normally none

Scale

Approximately correct from three selected widely spaced points on map to any other point on map. Cannot be exactly correct.

Distortion

Low distortion throughout map if three source points are properly placed near map limits (fig. 69A)

Usage

Atlas maps of continents by the National Geographic Society and some other atlas publishers

Origin

Presented by Wellman Chamberlin (1908-76) of the National Geographic Society in 1946

Similar projections

Azimuthal Equidistant projection (p. 132); true distance and direction from one point

Two-Point Equidistant projection (p. 146); true distance but not direction from two points





Figure 69A.—Chamberlin Trimetric projection, with Tissot indicatrices, 20° graticule. Source points the same as those in figure 69B.

Figure 69B.—Chamberlin Trimetric projection of North America with national boundaries, 10° graticule. Three source points for this map, as selected by the National Geographic Society: (1) latitude 55° N., longitude 150° W.; (2) latitude 55° N., longitude 35° W.; (3) latitude 10° N., longitude 92°30' W.



TILTED PERSPECTIVE Projection

Classifications

Modified azimuthal
Perspective
Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight; the point of perspective is in its plane, but this meridian may not appear on the map. Other meridians are normally elliptical arcs, but, for certain angles of tilt or centers, they may be straight lines, parabolas, or hyperbolas.

Parallels: One parallel, normally nearer to the pole than is the center of projection (depending on the position of the point of perspective), is a straight line. Other parallels are normally elliptical arcs but, for certain angles of tilt, may be parabolas or hyperbolas.

Poles: Points

Symmetry: Normally none, but there is symmetry about the central meridian if the direction of view is north or south

Range

Less than one hemisphere if the point of perspective is above the surface

Scale

Normally varies widely if a large area is covered

Distortion

Substantial distortion of shape, area, and scale if a large area is covered

Special features

Projection shows the Earth geometrically as it appears when photographed or otherwise imaged onto a plane from any point in space with any pointing of the "camera."

Usage

Pictorial views of the Earth, resembling those seen from space


Origin

Developed primarily during the 20th century to relate to aerial and space photography and photogrammetry

Limiting forms

General Vertical Perspective projection (p. 128)
Orthographic projection (p. 124), if the point of perspective is at infinite distance

Figure 70.—Tilted Perspective projection. Eastern U.S. seaboard, viewed from a point about 160 km above Newburgh, N.Y., 1° graticule. Center of projection latitude 41°30' N., longitude 74° W., tilt 55°, rotation 210°.



BACON GLOBULAR Projection

Classification

Globular

Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight.
Other meridians are circular arcs
equally spaced along the Equator but not
along other parallels.

Parallels: Parallel straight lines equally
spaced on the meridians 90° from center

Poles: Points

Symmetry: About the central meridian or the
Equator

Scale

True along the Equator

Distortion

Extreme area and shape distortion near the
poles

Usage

Early hemispheric maps

Origin


Presented by Roger Bacon (1214–94) of
England, about 1265, but unclear

Similar projections

Glareanus (Loritz) projection of 1527 is
identical.

Apian Globular I hemisphere of 1524 has
straight parallels equally spaced along the
central meridian.

Figure 71.—Bacon Globular projection, with shorelines, 10° graticule.
Central meridian 90° W.



FOURNIER GLOBULAR I Projection

Classification

Globular

Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight. Other meridians are semiellipses equally spaced along the Equator but not along other parallels.

Parallels: Parallel straight lines equally spaced on the meridians 90° from center

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along the central meridian and the Equator

Distortion

Moderate for a hemisphere

Usage

Early whole-world or hemispheric maps

Origin

Presented by Georges Fournier of France in 1643

Similar projections

Nicolosi Globular (p. 176) has circular arcs for meridians.

Mollweide (p. 54) has straight parallels and is used for world maps, but meridians within one hemisphere are the same as those on Fournier I.

Azimuthal Equidistant (equatorial aspect) (p. 134) has complex curves for meridians and parallels.

Figure 72.—Fournier Globular I projection, with shorelines, 10° graticule. Central meridian 90° W.



NICOLOSI GLOBULAR Projection

Classifications

Globular
Polyconic
Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight.
Other meridians are circular arcs, equally spaced along the Equator but not along other parallels.

Parallels: Parallel straight lines equally spaced on the meridians 90° from the center

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along the central meridian and the Equator

Distortion

Moderate for a hemisphere
Considerable area and shape distortion if used beyond the 90th meridians

Usage

Hemispheric maps, common in atlases between 1850 and 1925

Origin

Presented by Giambattista Nicolosi (1610–70) of Rome in 1660

Other names

Arrowsmith
Globular

Similar projections

Van der Grinten IV (p. 204) has parallels of different curvature and is extended to a world map.

Azimuthal Equidistant (equatorial aspect) (p. 134) has complex curves for meridians and parallels.

Fournier Globular I (p. 175) has elliptical arcs for meridians.

Figure 73.—Nicolosi Globular projection, with shorelines, 10° graticule. Central meridian 90° W.



APIAN GLOBULAR I Projection

Classifications

Globular

Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight.
Other meridians are circular arcs equally spaced along the Equator but not along other parallels.

Parallels: Straight lines, equally spaced along the central meridian

Poles: Points

Symmetry: Along the central meridian or the Equator

Scale

True along the central meridian and the Equator

Distortion

Moderate

Usage

Early maps of hemispheres

Origin

Presented by Peter Apian (Petrus Apianus or Peter Bienewitz) (1495–1552) of Saxony in 1524


Similar projections

Glareanus (Loritz) has identical meridians, but straight parallels are equally spaced on the 90th meridians.

Apian Globular II has identical parallels, central meridian, and 90th meridians, but other meridians are equally spaced ellipses.

Ortelius Oval (p. 178) is identical within 90° of the central meridian but is used for a world map.

Figure 74.—Apian Globular I projection, with shorelines, 10° graticule. Central meridian 90° W.



ORTELIUS OVAL Projection

Classifications

Oval

Modified pseudocylindrical

Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight. Other meridians are circular arcs, becoming semicircles between the meridians 90° and 180° from the central meridian. Equally spaced along the Equator but not along other parallels.

Parallels: Straight lines, equally spaced on the central meridian

Poles: Lines half the length of the Equator

Symmetry: Along the central meridian or the Equator

Scale

True along the central meridian and the Equator

Distortion

Moderate

Usage

16th-century whole-world maps

Origin

Battista Agnese (1514–64) used the projection in 1544, but the name comes from Abraham Ortelius (1527–98) of Flanders, who used it in 1570.


Similar projections

Often confused with Eckert III (p. 58), which has elliptic arcs for meridians

Apian Globular I (p. 177) is the same for one hemisphere.

Bordone Oval projection of 1528 has points for poles and near-ellipses or ovals for meridians but is otherwise very similar.

Figure 75.—Ortelius Oval projection, with shorelines, 15° graticule. Central meridian 90° W.



LAGRANGE Projection

Classifications

Conformal
Polyconic

Graticule

Meridians: Central meridian is straight. Other meridians are circular arcs, concave toward the central meridian.

Parallels: One parallel, often the Equator, is straight. Other parallels are circular arcs, concave toward the pole which is on the same side of the straight parallel.

Poles: Points

Symmetry: About the central meridian. Also about the Equator, if the Equator is straight.

Scale

Increases rapidly with distance from the center

Distortion

Great distortion of area and scale when the center is compared with poles or other outer limits (fig. 76A). Conformality fails at the poles.

Other features

The general formula can be used to create a variety of conformal maps.

Usage

Whole-world maps and conformal maps within a circle of smaller portions of the world

Origin

Johann Heinrich Lambert (1728-77) of Alsace presented the world conformally in a circle in 1772. This projection is usually called the Lagrange projection, however, after Joseph Louis Lagrange (1736-1813) of France, who generalized Lambert's concept in 1779.

Similar projections

Van der Grinten I (p. 200) is not conformal.

Special forms

Equatorial or oblique Stereographic projection (fig. 76D, also p. 120)

Littrow projection (p. 154)

Figure 76A.—Lagrange projection of sphere in circle, with Tissot indicatrices, 30° graticule.





Figure 76B.—Lagrange projection of sphere in circle, with shorelines, 15° graticule. Central meridian 90° W.



LAGRANGE Projection

Figure 76C.—Lagrange projection of 0.7 of sphere in circle, with shorelines, 15° graticule. Central meridian 90° W.




Figure 76D.—Equatorial Stereographic projection, with shorelines, 30° graticule. Central meridian 90° W. Special case of Lagrange with hemisphere in circle.



Figure 76E.—Lagrange projection of sphere in circle with latitude 30° N. shown straight, with shorelines, 15° graticule. Central meridian 90° W.

EISENLOHR Projection

Classification

Conformal

Graticule

Meridians: Central meridian is straight. Other meridians are curved and concave toward the central meridian.

Parallels: Equator is straight. Other parallels are curved and concave toward the nearest pole.

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

Increases rapidly with distance from the center. Constant scale around the boundary of a world map.

Distortion

Great distortion of area near the 180th meridians in comparison with the center (fig. 77A).

Other features

Because the boundary of a world map is at a constant scale, the projection provides the minimum overall scale variation for a conformal world map. Like the August Epicycloidal projection (p. 186), the Eisenlohr has no "singular" points at which conformality fails.

Usage

Novelty whole-world maps




Figure 77A.—Eisenlohr projection, with Tissot indicatrices, 30° graticule.

Origin


Presented by Friedrich Eisenlohr of Germany in 1870

Similar projections

August Epicycloidal (p. 186) has a boundary of varying scale.

Van der Grinten IV (p. 204) is not conformal and has circular arcs for meridians and parallels.

Figure 77B.—Eisenlohr projection, with shorelines, 15° graticule. Central meridian 90° W.



AUGUST EPICYCLOIDAL Projection

Classification

Conformal

Graticule

Meridians: Central meridian is straight. Other meridians are curved and concave toward the central meridian.

Parallels: Equator is straight. Other parallels are curved and concave toward the nearest pole.

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

Increases rapidly with distance from the center

Distortion

Great distortion of area toward the 180th meridians in comparison with the center

Other features

Boundary of the world map is a two-cusped epicycloid. Like the Eisenlohr projection (p. 184), this projection has no "singular" point

at which conformality fails. Unlike the Eisenlohr, the map boundary has a varying scale.

Usage

Novelty whole-world maps

Origin

Presented by F. August and G. Bellermann in Germany in 1874

Aspects

Transverse oblique aspect used by Athelstan F. Spilhaus in 1942 to show a map of world oceans

Other names


August

Similar projections

Eisenlohr (p. 184) has a boundary of constant scale.

Van der Grinten IV (p. 204) is not conformal and has circular arcs for meridians and parallels.

Figure 78.—August Epicycloidal projection, with shorelines, 15° graticule. Central meridian 90° W.



GUYOU Projection

Classifications

Conformal

Graticule

Meridians: Central meridian is straight. The 90th meridians are straight but bent at the 45th parallels N. and S. Other meridians are complex curves.

Parallels: Equator is straight. Other parallels are complex curves.

Poles: Points, in the midpoint of opposite sides of a square bounding a hemisphere

Symmetry: About the central meridian or the Equator

Scale

Varies considerably along each meridian and parallel

Distortion

Great distortion of area, especially near the

90th meridian at latitudes 45°N. or S., where conformality fails

Other features

Shows hemisphere conformally in a square and the world in a 2x1 rectangle. Hemispheres can be mosaicked to form a continuous map indefinitely.

Usage

Novelty map of a hemisphere or of the world

Origin

Presented by Émile Guyou of France in 1887


Aspects

A transverse aspect is the Peirce Quincuncial projection (p. 190). An oblique aspect is the Adams projection of a hemisphere in a square (p. 192).

Similar projections

Listed under Aspects above

Figure 79.—Guyou projection, with shorelines, 10° graticule. Central meridian 90° W.



PEIRCE QUINCUNCIAL Projection

Classification

Conformal

Graticule

Meridians: Every 90th meridian is straight but bent at the Equator. Other meridians are complex curves.

Parallels: Equator is a square, bent at the straight meridians. Other parallels are complex curves.

Poles: Points at the center of each hemispherical map

Symmetry: About the central meridian or the Equator

Scale

Varies considerably along each meridian and parallel

Distortion

Great distortion of area, especially near each 90th meridian at the Equator, where conformality fails

Other features

Actually a “star” projection in the normal aspect. Shows a polar hemisphere conformally in a square and the world in a larger square. The first use of elliptic functions for a map projection.

Usage

Novelty whole-world maps

Origin

Presented by Charles Sanders Peirce (1839–1914) of the U.S. Coast and Geodetic Survey in 1879

Aspects

In normal form, the pole is at the center of a

square. In transverse form, placing the pole at the midpoint of one side of a square results in the Guyou projection (p. 188). Placing the pole at the corner of the square results in the Adams projection of a hemisphere in a square (p. 192).


Other names

Quincuncial; name applied by Peirce, because there are normally five parts within a square (the Northern Hemisphere is in the center, and four parts of the Southern Hemisphere are in the corners)

Similar projections

Listed under Aspects above

Figure 80. Peirce Quincuncial projection, with shorelines, 10° graticule. Central meridian 105°W.



ADAMS Projection of a Hemisphere in a Square

Classification

Conformal

Graticule

Meridians: Central meridian is straight. The 90th meridians are straight lines bent at the Equator. Other meridians are curved.

Parallels: Equator is straight. Other parallels are curved.

Poles: Points, in opposite vertices of a square

Symmetry: About the central meridian or the Equator

Scale

Varies along each meridian and each parallel

Distortion

Great distortion of area at the poles or the intersection of the 90th meridians with the Equator. At these four points, conformality fails.

Other features

Hemisphere is enclosed in a square. World can be enclosed in a 2x1 rectangle.

Usage

Novelty map of a hemisphere

Origin

Presented by Oscar Sherman Adams (1874–1962) of the U.S. Coast and Geodetic Survey in 1925


Aspects

One transverse aspect is the Peirce Quincuncial projection (p. 190). Another transverse aspect is a rotation of the original projection. An oblique aspect is the Guyou projection (p. 188).

Similar projections

Listed under Aspects above

Figure 81.—Adams projection of a hemisphere in a square, with shorelines, 10° graticule. Central meridian 90° W.



ADAMS Projection of the World in a Square I

Classification

Conformal

Graticule

Meridians: Central meridian is straight. The 180th meridians are straight lines bent at the latitudes 70°31' N. and S. Other meridians are complex curves.

Parallels: Equator is straight. Other parallels are complex curves.

Poles: Points in midpoints of opposite sides

Symmetry: About the central meridian or the Equator

Scale

Varies along each meridian and each parallel

Distortion

Great area distortion at the higher latitudes. Conformality fails at the poles and at two points on each of the 180th meridians.

Usage

Novel whole-world maps

Origin

Presented by Oscar Sherman Adams (1874–1962) of the U.S. Coast and Geodetic Survey in 1925.

Similar projections

A general formula for representing the whole sphere or a segment of the sphere conformally in a regular polygon of three or more sides can be used to create this map and other conformal maps.




Figure 82.—Adams projection of the world in a square I, with shorelines, 10° graticule. Central meridian 90° W.

ADAMS Projection of the World in a Square II

Classification

Conformal

Graticule

Meridians: Central meridian is straight. The 180th meridians are straight lines bent at the Equator. Other meridians are complex curves.

Parallels: Equator is straight. Other parallels are complex curves.

Poles: Points in opposite vertices

Symmetry: About the central meridian or the Equator

Distortion

Great area distortion near the 180th meridians or at higher latitudes. Conformality fails at the poles and at the intersections of the 180th meridians with the Equator.

Usage

Novel whole-world maps

Origin

Presented by Oscar Sherman Adams (1874–1962) of the U.S. Coast and Geodetic Survey in 1925

Similar projections

See note under Adams projection of the world in a square I (p. 194).




Figure 83.—Adams projection of the world in a square II, with shorelines, 15° graticule. Central meridian 90° W.

LEE Conformal Projection of the World in a Triangle

Classification

Conformal

Graticule

Meridians: Central meridian is straight. The 180th meridians are straight lines bent at a southern latitude. Others are complex curves.

Parallels: Complex curves.

Poles: Points, one in a vertex and the other centered on the opposite side of the triangle

Symmetry: About the central meridian

Scale

Varies along each meridian and parallel

Distortion

Great area distortion near the poles and 180th meridians. Conformality fails at each pole and at one point on each of the 180th meridians.

Usage

Novel whole-world maps

Origin


Laurence Patrick Lee (1913–85) of New Zealand in 1965

Similar projections

See note under Adams projection of the world in a square I (p. 194).

Bernard J.S. Cahill, beginning in 1912, arranged conformal or nonconformal three-sided octants of the world, bounded by the Equator and two meridians 90° apart, in a butterfly arrangement, called the “Butterfly Map.” He promoted these interrupted arrangements for weather maps.

Figure 84.—Lee conformal projection of the world in a triangle, with shorelines, 15° graticule. Central meridian 90° W.



Conformal Projection of the World in an Ellipse

Classification

Conformal

Graticule

Meridians: Central meridian is straight. Other meridians are complex curves.

Parallels: Equator is straight. Other parallels are complex curves.

Poles: Points

Symmetry: About the central meridian and the Equator

Scale

Increases rapidly with distance from the center

Distortion

Great distortion of area near the poles and the 180th meridians. Conformality fails at the poles.

Other features

The shape of the bounding ellipse can be varied. Lee uses an axial ratio of about 1.97 to 1 for convenience in using tabular values.

Usage

Novel whole-world maps

Origin

Presented by Oscar Sherman Adams (1874–1962) of the U.S. Coast and Geodetic Survey in 1925. Corrected by André Gougenheim in 1950 and Laurence Patrick Lee (1913–85) of New Zealand in 1965.


Limiting form

As the modular angle that controls the shape of the ellipse approaches 90°, the map increasingly resembles the Lagrange circular projection (p. 181).

Similar projections

See note under Adams projection of the world in a square I (p. 194).

Figure 85.—Conformal projection of the world in an ellipse, with shorelines, 15° graticule. Central meridian 90° W.



VAN DER GRINTEN I Projection

Classifications

Polyconic

Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight. Other meridians are circular, equally spaced along the Equator and concave toward the central meridian.

Parallels: Equator is straight. Other parallels are circular arcs, concave toward the nearest pole.

Poles: Points

Symmetry: Along the central meridian or the Equator

Scale

True along the Equator. Increases rapidly with distance from the Equator.

Distortion

Great distortion of area near the poles (fig. 86B)

Other features

The world is enclosed in a circle.

Usage

World maps by the U.S. Department of Agriculture, the U.S. Geological Survey, the National Geographic Society, and others

Origin

Presented by Alphons J. van der Grinten (1852-?) of Chicago in 1898. U.S. patent obtained in 1904.

Other names

Van der Grinten (without I)

Similar projections

Lagrange (p. 181) is conformal.

Van der Grinten II (p. 202) has parallels curved to intersect meridians at right angles.

Mercator (p. 10) is conformal and has straight meridians and parallels.

Figure 86A.—Van der Grinten I projection, with Tissot indicatrices, 30° graticule.




Figure 86B.—Van der Grinten I projection, with shorelines, 15° graticule. Central meridian 90° W.



VAN DER GRINTEN II Projection

Classifications

Polyconic

Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight. Other meridians are circular, equally spaced along the Equator and concave toward the central meridian.

Parallels: Equator is straight. Other parallels are circular arcs, concave toward the nearest pole.

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True on the Equator. Increases rapidly with distance from the Equator.

Distortion

Great distortion of area near the poles

Other features

The world is enclosed in a circle.

All meridians intersect parallels at right angles, but projection is not conformal. Spacing of parallels along the central meridian is the same as that for Van der Grinten I.

Usage

World maps, but rarely

Origin

Presented by Alphons J. van der Grinten (1852–?) of Chicago in 1904

Similar projections

Lagrange (p. 181) is conformal.

Van der Grinten I (p. 200) has parallels that do not intersect meridians at right angles.



Figure 87.—Van der Grinten II projection, with shorelines, 15° graticule, central meridian 90° W.

VAN DER GRINTEN III Projection

Classification

Modified pseudocylindrical

Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight. Other meridians are circular, equally spaced along the Equator but unequally spaced along other parallels and concave toward the central meridian.

Parallels: Straight lines

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along the Equator. Increases rapidly with distance from the Equator.

Distortion

Great distortion of area near the poles

Other features

The world is enclosed in a circle.

Spacing of parallels along the central meridian is the same as that for Van der Grinten I.

Usage

World maps, but rarely

Origin

Presented by Alphons J. van der Grinten (1852–?) of Chicago in 1904

Similar projections

Van der Grinten I (p. 200) has curved parallels.

“The Times,” by Moir, has nearly elliptical meridians, but straight parallels are spaced like those of the Gall projection (p. 33).




Figure 88.—Van der Grinten III projection, with shorelines, 15° graticule, central meridian 90° W.

VAN DER GRINTEN IV Projection

Classification

Polyconic

Globular

Neither conformal nor equal area

Graticule

Meridians: Central meridian is straight. Other meridians are circular, equally spaced along the Equator and concave toward the central meridian.

Parallels: Equator is straight. Other parallels are circular, equally spaced on the central meridian and concave toward the nearest pole.

Poles: Points

Symmetry: About the central meridian or the Equator

Scale

True along the central meridian and the Equator

Distortion

Great distortion of area near the 180th meridians at higher latitudes

Other features

The 90th meridians form a complete circle.

Usage

World maps, but rarely

Origin

Presented by Alphons J. van der Grinten (1852–?) of Chicago in 1904

Other names


“Apple Shaped”

Similar projections

Eisenlohr (p. 184) is conformal.

August Epicycloidal (p. 186) is conformal.

Figure 89.—Van der Grinten IV projection, with shorelines, 15° graticule. Central meridian 90° W.



ARMADILLO Projection

Classifications

Orthoapsidal (term coined by originator)
Neither conformal nor equal area

Graticule

Meridians: Central meridian (10° or 15° E.) is straight. Other meridians are elliptical arcs, concave toward the central meridian.
Parallels: Elliptical arcs of the same eccentricity, concave toward the North Pole
Poles: North Pole is semiellipse. South Pole cannot be shown.
Symmetry: About the central meridian

Scale

Gradually decreases with distance from the center

Distortion

Distortion is moderate in central portions.

Other features

An oblique orthographic projection of the world plotted with equidistant meridians and parallels onto a portion of a torus ring (similar to a doughnut). Antarctic region cannot be shown, but the projection was claimed to have "more land in proportion to sea than any other world map." Often plotted with New Zealand, normally hidden from view, appended to Australia as a "pigtail."

Usage

Whole-world maps

Origin

Presented by Erwin J. Raisz (1893–1968) of Harvard University in 1943

Aspects

Oblique is the basic aspect.

Other names

Raisz

Similar projections

Other "orthoapsidal" projections proposed by Raisz in 1943. Raisz coined this term from "orthographic" and "apsidal."

Figure 90.—Armadillo projection, with shorelines, 15° graticule, central meridian 15° E.



GS50 Projection

Classification

Conformal

Graticule

Meridians: Complex curves at slightly varying intervals

Parallels: Complex curves at slightly varying intervals

Poles: Points beyond the useful range of the map

Symmetry: None

Scale

True along an irregular line approximately encompassing the regions of the 50 States of the United States

Varies less than 2 percent from true scale throughout the 50 States and adjacent bodies of water

Distortion

Shape and scale distortion very low for 50-State region (fig. 91B). Greater distortion for regions away from 50 States.

Special features

Portrays irregular region of 50 States at about one-fourth the variation in scale of the best standard projections. Uses tenth-order complex-algebra polynomials to modify the Stereographic projection and is practical only with a computer. Conformality is precise in these cases even though finite polynomial series are used. The coefficients are useful only for this design of a 50-State map.

Origin

Developed by John P. Snyder (1926–) of the U.S. Geological Survey in 1982


Similar projections

Oblated (or Prolated) Stereographic projections by O.M. Miller in 1953 (p. 148) and by L.P. Lee in 1974 are equivalent to third-order complex-algebra polynomial modifications of the oblique Stereographic, applied to continents or the Pacific Ocean.

New Zealand Map Grid presented by W.I. Reilly in 1973 is a sixth-order complex-algebra polynomial modification of the Mercator, applied to New Zealand.


Modified-Stereographic Conformal projection developed by Snyder in 1983 is a sixth-order complex-algebra modification of the oblique Stereographic, applied to Alaska.

Figure 91A.—GS50 projection, with shorelines, 5° graticule.



GS50 Projection

Figure 91B.—GS50 projection, with lines of constant scale.



SPACE OBLIQUE MERCATOR Projection

Classifications

Modified cylindrical
Basically conformal

Graticule

Meridians: Complex curves at slightly varying intervals

Parallels: Complex curves at slightly varying intervals

Poles: Points

Symmetry: Inverse symmetry about each node (crossing of the Equator by satellite groundtrack)

Scale

True along the groundtrack
Varies about 0.01 percent within the normal sensing range of the satellite

Distortion

No distortion along the groundtrack
Distortion is essentially constant along lines of constant distance parallel to the groundtrack.
Conformality is correct within a few parts per million for sensing range.

Special features

Provides continuous conformal mapping of swath sensed by an artificial satellite, such as Landsats 1 through 5, orbiting the Earth; no scale distortion along the groundtrack. The first projection to account for the Earth's rotation with respect to a satellite orbit.

Usage

Format for imagery from mapping satellites Landsats 4 and 5

Origin


Concept presented by Alden P. Colvocoresses (1918–) of the U.S. Geological Survey in 1973.

Mathematical development by John P. Snyder (1926–) in 1977

Similar projections

Oblique Mercator projection (p. 14) would be suitable for a satellite orbiting a nonrotating spherical Earth and is the closest standard approximation for limited intervals involving a rotating Earth.

Figure 92A.—Two orbits of the Space Oblique Mercator projection, shown for Landsat 5, paths 15 (left) and 31 (right). Designed for a narrow band along the groundtrack, which remains true to scale. Note the change in longitude at a given latitude along the groundtrack with successive orbits. 30° graticule. Short, straight lines nearly perpendicular to the groundtrack are sample scan lines, extended 15° from groundtrack.



SPACE OBLIQUE MERCATOR Projection




Figure 92B.—One quadrant of the Space Oblique Mercator projection. An enlargement of part of figure 92A, path 15, beginning at the North Pole, but using 10° graticule.

SATELLITE TRACKING Projections

Classifications

Cylindrical or conic
Neither conformal nor equal area

Graticule

Meridians: Equally spaced straight lines, parallel on cylindrical form and converging toward common point on conic form

Parallels: Unequally spaced straight parallel lines on cylindrical form, closest near the Equator, perpendicular to meridians.

Unequally spaced concentric circular arcs on conic form, centered on the pole of convergence of the meridians, which are therefore radii of the circular arcs. Spacing of parallels increases away from the central latitudes.

Poles: Since satellite groundtracks do not pass over poles, they are not needed on the map. They can be shown in arbitrary locations, however.

Symmetry: About any meridian. On the cylindrical form, also about the Equator.

Scale

On the cylindrical form, true along the Equator or along two parallels equidistant from the Equator

On the conic form, true along one chosen parallel

Distortion

Conformality can be made to occur along two parallels equidistant from the Equator on the cylindrical form and along any two parallels within tracking range on the conic form.

Special feature

Any groundtrack for a given orbiting satellite such as Landsat is shown straight, if the map is designed for that particular satellite. The

groundtrack may or may not have a sharp break at the northern or southern limit of tracking, depending on design.


Origin

Developed by John P. Snyder (1926–) in 1977

Similar projections

Cylindrical form resembles the Mercator (p. 10), and the conic form resembles the Lambert Conformal Conic (p. 95), but spacing of parallels is different. The Mercator and Lambert Conformal Conic are conformal throughout, except at the poles, but the Satellite Tracking projections are only conformal at two latitudes.

Figure 93A.—Cylindrical Satellite Tracking projection (standard parallels 30° N. and S.). Landsats 1, 2, and 3 orbits. Diagonal lines are groundtracks for paths 15, 30, 45, and so on. 15° graticule.



SATELLITE TRACKING Projections





Figure 93B.—Conic Satellite Tracking projection (conformality at parallels 45° and 70° N.). Landsats 1, 2, and 3 orbits. Diagonal lines are groundtracks for paths 15, 30, 45, and so on, and are tangent to the inner circle. 10° graticule.

APPENDIX: FORMULAS FOR PROJECTIONS



APPENDIX: FORMULAS FOR PROJECTIONS

This compilation is abridged in several aspects. Only forward formulas for the sphere are given to compute rectangular coordinates from geographic latitude and longitude. For the inverse formulas and for forward or inverse formulas treating the Earth as an ellipsoid (if available), other references are given below. In addition, the reader is referred to relatively recent literature describing projections in more detail. Older sources are not listed in this volume, but they are frequently given in the literature that is referenced herein.

For a few major projections, the formulas for computing oblique, transverse, or equatorial aspects are given specifically. General transformation formulas are also given for use with any normal aspect, and most general map-plotting computer programs use these or trigonometric identities thereof rather than individual transformation formulas, unless ellipsoidal projections are to be plotted.

On the other hand, because of the importance of computer programs, the formulas below are amplified with conditional equations that should be programmed with the main formulas to avoid errors resulting from dividing by zero and so on.

In these equations, the following symbols apply:

<i>n</i>	cone constant on conic projections, or the ratio of the angle between meridians to the true angle
<i>R</i>	radius of the sphere at the scale of the map
<i>x</i>	rectangular coordinate; distance to the right of the vertical line (<i>Y</i> axis) passing through the origin (if negative, it is distance to the left)
<i>y</i>	rectangular coordinate; distance above the horizontal line (<i>X</i> axis) passing through the origin (if negative, it is distance below)
<i>z</i>	great-circle distance, as an arc of a circle
<i>ln</i>	natural logarithm
Θ	a special parametric variable, or an angle in polar coordinates measured generally counterclockwise from due south in these formulas
λ	longitude east of Greenwich (for longitude west of Greenwich, a minus sign is used)

λ_0	longitude east of Greenwich of the central meridian of the map or of the origin of the rectangular coordinates (for west longitude, a minus sign is used). If ϕ_1 is a pole, λ_0 is the longitude of the meridian extending down on the map from the North Pole or up from the South Pole. On an interrupted projection, λ_0 is the central meridian of each section.
ρ	radius in polar coordinates
ϕ	north latitude (if latitude is south, a minus sign is used)
ϕ_0	middle latitude, or latitude chosen as the origin of rectangular coordinates for a projection, in radians. Its choice does not affect the appearance of the projection.
ϕ_1, ϕ_2	standard parallels of latitude for projections having two standard parallels
ϕ_1 (without ϕ_2)	single standard parallel on cylindrical or conic projections; latitude of central point on azimuthal projections. Its choice normally affects the appearance of the projection.
ϕ_p, λ_p	latitude and longitude, respectively, of transformed pole of oblique, transverse, or equatorial aspect

Angles are to be expressed in radians, but degrees can be equally suitable in some cases, and some numbers are shown as degrees. Radians equal degrees times $\pi/180^\circ$. The \arctan_2 function corresponds to the Fortran ATAN2 function and must therefore be adjusted for quadrant, depending on the signs of both numerator and denominator, if the function is not already programmed. The programmer should frequently apply normal techniques for computing efficiency to minimize computation of trigonometric functions; that is, a given function should be computed only once, and other functions should be derived by using common identities not listed here, since listing these techniques would often make the equations more difficult to follow.

The Newton-Raphson iteration suggested in several cases involves use of the equation

$$\Delta v = -f(v)/[df(v)/dv]$$

After a suitable initial value of v is used, v is repeatedly adjusted with Δv until Δv becomes less than some preset convergence value. For example, equation 31 is transposed to develop a function of Θ , $f(\Theta)$, which should approach zero:

$$f(\Theta) = \Theta + \sin \Theta - (1 + \pi/2) \sin \phi$$

Differentiating, but treating ϕ as a constant,

$$df(\Theta)/d\Theta = 1 + \cos \Theta$$

Then,

$$\Delta\Theta = -[\Theta + \sin \Theta - (1 + \pi/2) \sin \phi]/(1 + \cos \Theta)$$

If an initial value of Θ equal to ϕ is used, convergence is rapid.

CYLINDRICAL PROJECTIONS

Because of the common use of several transverse or oblique aspects of the cylindrical projections, formulas are given separately for them under this heading. They can also be found from the formulas for the normal aspect by using equations 130 and 131.

Mercator

$$x = R(\lambda - \lambda_0)$$

If $\phi = 90^\circ$, $y = \infty$; if $\phi = -90^\circ$, $y = -\infty$. If ϕ is not $\pm 90^\circ$,

$$y = R \ln \tan (\pi/4 + \phi/2)$$

The X axis lies along the Equator, x increasing east; the Y axis lies along the central meridian, y increasing north. For ellipsoidal and inverse formulas and more information, see U.S. Geological Survey (USGS) Professional Paper 1395 (Snyder, 1987, p. 38–47).

Transverse Mercator

Let

$$B = \cos \phi \sin (\lambda - \lambda_0)$$

If $B = 1$, $x = \infty$; if $B = -1$, $x = -\infty$; y is found from equation 5. If B is not ± 1 ,

$$x = \frac{1}{2}R \ln [(1 + B)/(1 - B)]$$

$$y = R \arctan_2 [\tan \phi / \cos (\lambda - \lambda_0)]$$

The X axis lies along the Equator, x increasing east; the Y axis lies along the central meridian, y increasing north. These axes are rotated 90° counterclockwise from those corresponding to the equations for the Oblique Mercator below. For ellipsoidal and inverse formulas and more information, see USGS Professional Paper 1395 (Snyder, 1987, p. 48–64).

Oblique Mercator

$$x = R \arctan_2 \left[\frac{\cos \phi_p \sin \phi - \sin \phi_p \cos \phi \cos (\lambda - \lambda_p)}{\cos \phi \sin (\lambda - \lambda_p)} \right] \quad (6)$$

Let

$$A = \sin \phi_p \sin \phi + \cos \phi_p \cos \phi \cos (\lambda - \lambda_p) \quad (7)$$

(1) If $A = 1$, $y = \infty$; if $A = -1$, $y = -\infty$. If A is not ± 1 ,

$$y = \frac{1}{2} R \ln [(1 + A)/(1 - A)] \quad (8)$$

Note that the X axis lies along the central line, x increasing east. The Y axis crosses the X axis at the actual Equator and a longitude 90° less than λ_p , y increasing north. For ellipsoidal and inverse formulas and more information, see USGS Professional Paper 1395 (Snyder, 1987, p. 66–75).

Lambert Cylindrical Equal-Area

$$x = \text{same as in equation 1}$$

$$y = R \sin \phi \quad (9)$$

For axes, see Mercator. For ellipsoidal and inverse formulas and more information for all aspects, see USGS Professional Paper 1395 (Snyder, 1987, p. 76–85).

Transverse Lambert Cylindrical Equal-Area

$$x = R \cos \phi \sin (\lambda - \lambda_0) \quad (10)$$

$$y = \text{same as in equation 5}$$

For axes, see Transverse Mercator.

**Oblique Lambert
Cylindrical Equal-Area**

$$x = \text{same as in equation 6}$$

Let

$$A = \text{same as in equation 7}$$

$$y = RA \quad (11)$$

For axes, see Oblique Mercator.

**Behrmann Cylindrical
Equal-Area**

$$x = R(\lambda - \lambda_0) \cos 30^\circ \quad (12)$$

$$y = R \sin \phi / \cos 30^\circ \quad (13)$$

For axes, see Mercator.

Plate Carrée

$$x = \text{same as in equation 1}$$

$$y = R\phi \quad (14)$$

For axes, see Mercator.

Equirectangular

$$x = R(\lambda - \lambda_0) \cos \phi_1 \quad (15)$$

$$y = \text{same as in equation 14}$$

For axes, see Mercator.

Cassini

$$x = R \arcsin [\cos \phi \sin (\lambda - \lambda_0)] \quad (16)$$

$$y = \text{same as in equation 5}$$

For axes, see Transverse Mercator. For ellipsoidal and inverse formulas and more information, see USGS Professional Paper 1395 (Snyder, 1987, p. 92–95).

**Oblique Equidistant
Cylindrical**

$$x = \text{same as in equation 6}$$

$$A = \text{same as in equation 7}$$

$$y = R \arcsin A \quad (17)$$

For axes, see Oblique Mercator.

Central Cylindrical

$$x = \text{same as in equation 1}$$

If $\phi = 90^\circ$, $y = \infty$; if $\phi = -90^\circ$, $y = -\infty$. If ϕ is not $\pm 90^\circ$,

$$y = R \tan \phi \quad (18)$$

For axes, see Mercator.

Transverse Central Cylindrical

Calculate B from equation 3. If $B = 1$, $x = \infty$; if $B = -1$, $x = -\infty$. If B is not ± 1 ,

$$x = RB/(1 - B^2)^{1/2} \quad (19)$$

$$y = \text{same as in equation 5}$$

For axes, see Transverse Mercator.

Cylindrical Projections

PSEUDOCYLINDRICAL PROJECTIONS

Cylindrical Projections

Gall (Stereographic)

$$x = R(\lambda - \lambda_0)/2^{\frac{1}{4}} \quad (20)$$

$$y = R(1 + 2^{\frac{1}{4}}/2) \tan \frac{1}{2} \phi \quad (21)$$

For axes, see Mercator.

Miller Cylindrical

$$x = \text{same as in equation 1}$$

$$y = R[\ln \tan (\pi/4 + 0.4\phi)]/0.8 \quad (22)$$

For axes, see Mercator. For more information see the article by Miller (1942) and USGS Professional Paper 1395 (Snyder, 1987, p. 86–89).

Pseudocylindrical Projections

For axes, see Mercator. For transverse and oblique aspects, use the general formulas given in 130 and 131. Snyder (1977) provides original references for most of the formulas and descriptions of several other pseudocylindrical projections.

Sinusoidal

$$x = R(\lambda - \lambda_0) \cos \phi \quad (23)$$

$$y = R\phi \text{ (same as in equation 14)}$$

McBryde-Thomas Flat-Polar

Sinusoidal

$$\Theta/2 + \sin \Theta = (1 + \pi/4) \sin \phi \quad (24)$$

Find Θ for the given ϕ by using a Newton-Raphson iteration of equation 24, in which ϕ is the initial estimate of Θ .

$$x = R[6/(4 + \pi)]^{\frac{1}{4}} (0.5 + \cos \Theta)(\lambda - \lambda_0)/1.5 \quad (25)$$

$$y = R[6/(4 + \pi)]^{\frac{1}{4}} \Theta \quad (26)$$

Eckert V

$$x = R(1 + \cos \phi)(\lambda - \lambda_0)/(2 + \pi)^{\frac{1}{4}} \quad (27)$$

$$y = 2R\phi/(2 + \pi)^{\frac{1}{4}} \quad (28)$$

Winkel I

$$x = R(\lambda - \lambda_0)(\cos \phi_1 + \cos \phi)/2 \quad (29)$$

$$y = R\phi \quad (30)$$

Eckert VI

$$\Theta + \sin \Theta = (1 + \pi/2) \sin \phi \quad (31)$$

Find Θ for a given ϕ by using a Newton-Raphson iteration of equation 31 in which ϕ is the initial estimate of Θ .

$$x = R(1 + \cos \Theta)(\lambda - \lambda_0)/(2 + \pi)^{\frac{1}{4}} \quad (32)$$

$$y = 2R\Theta/(2 + \pi)^{\frac{1}{4}} \quad (33)$$

McBryde S3

For latitudes between $55^{\circ}51'$ N. and S., use equations 23 and 14. For other latitudes, use equations 24, 25, and the following:

$$y = R\{[6/(4 + \pi)]^{\frac{1}{4}} \Theta - 0.069065 \text{ sign } \phi\} \quad (34)$$

Mollweide

$$2\Theta + \sin 2\Theta = \pi \sin \phi \quad (35)$$

Find Θ for the given ϕ by using a Newton-Raphson iteration of equation 35 in which ϕ is the initial estimate of Θ .

$$x = 2(2^{\frac{1}{4}}) R(\lambda - \lambda_0)(\cos \Theta)/\pi \quad (36)$$

$$y = 2^{\frac{1}{4}} R \sin \Theta \quad (37)$$

Eckert III

$$x = 2\{1 + [1 - (2\phi/\pi)^2]^{\frac{1}{4}}\} R(\lambda - \lambda_0)/(4\pi + \pi^2)^{\frac{1}{4}} \quad (38)$$

$$y = 4R\phi/(4\pi + \pi^2)^{\frac{1}{4}} \quad (39)$$

Eckert IV

$$\Theta + \sin \Theta \cos \Theta + 2 \sin \Theta = (4 + \pi)(\sin \phi)/2 \quad (40)$$

Find Θ for the given ϕ by using a Newton-Raphson iteration of equation 40 in which $\phi/2$ is the initial estimate of Θ .

$$x = 2R(\lambda - \lambda_0)(1 + \cos \Theta)/(4\pi + \pi^2)^{\frac{1}{4}} \quad (41)$$

$$y = 2\pi^{\frac{1}{4}}R \sin \Theta/(4 + \pi)^{\frac{1}{4}} \quad (42)$$

Putniš P_{2'}

$$2\Theta + \sin 2\Theta = \{[4\pi + 3(3^{\frac{1}{4}})]/6\} \sin \phi \quad (43)$$

Find Θ for the given ϕ by using a Newton-Raphson iteration of equation 43 in which $\phi/2$ is the initial estimate of Θ .

$$x = 0.86310R(\lambda - \lambda_0) \cos \Theta \quad (44)$$

$$y = 1.56548R \sin \Theta \quad (45)$$

Hatano Asymmetrical Equal-Area

For all latitudes,

$$x = 0.85R(\lambda - \lambda_0) \cos \Theta \quad (46)$$

The parametric angle Θ and y are calculated differently for different latitudes. For latitudes north of the Equator,

$$2\Theta + \sin 2\Theta = 2.67595 \sin \phi \quad (47)$$

$$y = 1.75859R \sin \Theta \quad (48)$$

For latitudes south of the Equator,

Pseudocylindrical Projections

$$2\Theta + \sin 2\Theta = 2.43763 \sin \phi \quad (49)$$

$$y = 1.93052R \sin \Theta \quad (50)$$

Find Θ for the given ϕ in equations 47 and 49 by using a Newton-Raphson iteration in which $\phi/2$ is the initial estimate of Θ .

Goode Homolosine

For latitudes between 40°44' N. and S., use equations 23 and 14. For other latitudes, use equations 35, 36, and the following:

$$y = R[2^{\frac{1}{4}} \sin \Theta - 0.05280 \operatorname{sign} \phi] \quad (51)$$

Boggs Eumorphic

If $\phi = \pm 90^\circ$, then $\Theta = \phi$, $x = 0$, and y is found from equation 53. Otherwise, use equation 35 to find Θ from ϕ . Then,

$$x = 2.00276R(\lambda - \lambda_0)/(\sec \phi + 1.11072 \sec \Theta) \quad (52)$$

$$y = 0.49931R(\phi + 2^{\frac{1}{4}} \sin \Theta) \quad (53)$$

Craster Parabolic

$$x = 3^{\frac{1}{4}}R(\lambda - \lambda_0)[2 \cos(2\phi/3) - 1]/\pi^{\frac{1}{4}} \quad (54)$$

$$y = (3\pi)^{\frac{1}{4}}R \sin(\phi/3) \quad (55)$$

McBryde-Thomas Flat-Polar Parabolic

$$\Theta = \arcsin[(7/3)(\sin \phi)/6^{\frac{1}{4}}] \quad (56)$$

PSEUDOCYLINDRICAL PROJECTIONS

$$x = 6^{\frac{1}{4}}R(\lambda - \lambda_0)[2 \cos(2\theta/3) - 1]/7^{\frac{1}{4}} \quad (57)$$

$$y = 9R[\sin(\theta/3)]/7^{\frac{1}{4}} \quad (58)$$

Putniš P₅

$$x = 1.01346R(\lambda - \lambda_0)[2 - (1 + 12\phi^2/\pi^2)^{\frac{1}{4}}] \quad (64)$$

$$y = 1.01346R\phi \quad (65)$$

Quartic Authalic

$$x = R(\lambda - \lambda_0)(\cos \phi)/\cos(\phi/2) \quad (59)$$

$$y = 2R \sin(\phi/2) \quad (60)$$

Denoyer Semi-Elliptical (approximation)

$$\text{Let } L = |\lambda - \lambda_0|.$$

$$x = R(\lambda - \lambda_0) \cos[(0.95 - L/12 + L^3/600)(0.9\phi + 0.03\phi^5)] \quad (66)$$

$$y = R\phi \text{ (same as in equation 14)}$$

$$\sin(\Theta/2) + \sin \Theta = (1 + 2^{\frac{1}{4}}/2) \sin \phi \quad (61)$$

Find Θ for the given ϕ by using a Newton-Raphson iteration of equation 61 in which ϕ is the initial estimate of Θ .

$$x = R(\lambda - \lambda_0)[1 + 2 \cos \Theta/\cos(\Theta/2)]/[3(2^{\frac{1}{4}}) + 6]^{\frac{1}{4}} \quad (62)$$

$$y = 2(3^{\frac{1}{4}}) R \sin(\Theta/2)/(2 + 2^{\frac{1}{4}})^{\frac{1}{4}} \quad (63)$$

Robinson

A table is used instead of analytical formulas. The coordinates can be used in the following equations for a world map of correct total area (not equal area):

$$x = 0.8487RX(\lambda - \lambda_0) \quad (67)$$

$$y = 1.3523RY$$

$$(68) \quad \text{Eckert I}$$

Pseudocylindrical Projections

where X and Y are interpolated from the following table (Robinson, 1974):

lat ϕ , in degrees	X	Y
90	0.5322	1.0000
85	.5722	.9761
80	.6213	.9394
75	.6732	.8936
70	.7186	.8435
65	.7597	.7903
60	.7986	.7346
55	.8350	.6769
50	.8679	.6176
45	.8962	.5571
40	.9216	.4958
35	.9427	.4340
30	.9600	.3720
25	.9730	.3100
20	.9822	.2480
15	.9900	.1860
10	.9954	.1240
5	.9986	.0620
0	1.0000	.0000

Collignon

$$x = 2R(\lambda - \lambda_0)(1 - \sin \phi)^{\frac{1}{2}} / \pi^{\frac{1}{2}}$$

$$(69)$$

$$y = \pi^{\frac{1}{2}} R [1 - (1 - \sin \phi)^{\frac{1}{2}}]$$

$$(70)$$

Eckert I

$$x = 2[2/(3\pi)]^{\frac{1}{2}} R(\lambda - \lambda_0)(1 - |\phi|/\pi) \quad (71)$$

$$y = 2[2/(3\pi)]^{\frac{1}{2}} R\phi \quad (72)$$

Eckert II

$$x = 2R(\lambda - \lambda_0)(4 - 3 \sin |\phi|)^{\frac{1}{2}} / (6\pi)^{\frac{1}{2}} \quad (73)$$

$$y = (2\pi/3)^{\frac{1}{2}} R [2 - (4 - 3 \sin |\phi|)^{\frac{1}{2}}] \operatorname{sign} \phi \quad (74)$$

Loximuthal

$$x = R(\lambda - \lambda_0)(\phi - \phi_1)/\ln [\tan (\pi/4 + \phi/2)/\tan (\pi/4 + \phi_1/2)] \quad (75)$$

$$y = R(\phi - \phi_1) \quad (76)$$

where ϕ_1 is the central latitude from which, at the central meridian, straight lines are loxodromes of true length and azimuth from the center. However, if $\phi = \phi_1$, equation 75 is replaced by

$$x = R(\lambda - \lambda_0) \cos \phi_1 \quad (77)$$

CONIC PROJECTIONS

For oblique aspects, use general formulas given in equations 130 and 131.

Equidistant Conic

$$n = (\cos \phi_1 - \cos \phi_2)/(\phi_2 - \phi_1) \quad (78)$$

$$\rho = R[(\phi_2 \cos \phi_1 - \phi_1 \cos \phi_2)/(\cos \phi_1 - \cos \phi_2) - \phi] \quad (79)$$

$$\Theta = n(\lambda - \lambda_0) \quad (80)$$

$$x = \rho \sin \Theta \quad (81)$$

$$y = \rho_0 - \rho \cos \Theta \quad (82)$$

where ρ_0 is calculated from equation 79 by using ϕ_0 in place of ϕ . The Y axis lies along the central meridian λ_0 , y increasing north; the X axis crosses the Y axis perpendicularly at (ϕ_0, λ_0) , x increasing east. For ellipsoidal and inverse formulas and more information, see USGS Professional Paper 1395 (Snyder, 1987, p. 111–115).

Lambert Conformal Conic

$$n = \ln(\cos \phi_1 / \cos \phi_2) / \ln[\tan(\pi/4 + \phi_2/2) / \tan(\pi/4 + \phi_1/2)] \quad (83)$$

$$F = \cos \phi_1 \tan^n(\pi/4 + \phi_1/2)/n \quad (84)$$

If $\phi = \pm 90^\circ$ and n has the same sign, $\rho = 0$. If $\phi = \pm 90^\circ$ and n has the opposite sign, $\rho = \infty$. If ϕ is not $\pm 90^\circ$,

$$\rho = RF / \tan^n(\pi/4 + \phi/2) \quad (85)$$

Use also equations 80 through 82, ρ_0 being calculated from equation 85 by using ϕ_0 in place of ϕ . If only one standard parallel is desired, or $\phi_1 = \phi_2$, equation 83 is indeterminate, but $n = \sin \phi_1$. For axes, see Equidistant Conic. For ellipsoidal and inverse formulas and more information, see USGS Professional Paper 1395 (Snyder, 1987, p. 104–110).

Bipolar Oblique Conic Conformal

The formulas are too lengthy to be included here. For formulas and other information, see USGS Professional Paper 1395 (Snyder, 1987, p. 116–123).

Albers Equal-Area Conic

$$n = (\sin \phi_1 + \sin \phi_2)/2 \quad (86)$$

$$C = \cos^2 \phi_1 + 2n \sin \phi_1 \quad (87)$$

$$\rho = R(C - 2n \sin \phi)^{n/2} \quad (88)$$

Use also equations 80 through 82, ρ_0 being calculated from equation 88 by using ϕ_0 in place of ϕ . For axes, see Equidistant Conic. For ellipsoidal and inverse formulas and more information, see USGS Professional Paper 1395 (Snyder, 1987, p. 98–103).

Lambert Equal-Area Conic

$$n = (1 + \sin \phi_1)/2 \quad (89)$$

$$\rho = R[2(1 - \sin \phi)/n]^{n/2} \quad (90)$$

Use also equations 80 through 82, ρ_0 being calculated from equation 90 by using ϕ_0 in place of ϕ . For axes, see Equidistant Conic.

Perspective Conic

$$\phi_0 = (\phi_1 + \phi_2)/2 \quad (91)$$

If ϕ is less than or equal to $(\phi_0 - 90^\circ)$ and ϕ_0 is positive or if ϕ is greater than or equal to $(\phi_0 + 90^\circ)$ and ϕ_0 is negative, the point should be rejected. Otherwise,

$$n = \sin \phi_0 \quad (92)$$

PSEUDOCONIC PROJECTIONS

		Conic Projections
$\rho = R \cos [(\phi_2 - \phi_1)/2][\cot \phi_0 - \tan (\phi - \phi_0)]$	(93)	If ϕ is 0,
Use also equations 80 through 82, ρ_0 being calculated from equation 93 by using ϕ_0 in place of ϕ and thus cancelling the last term. For axes, see Equidistant Conic.		$x = 2RA$ (100) $y = -R\phi_0$ (101)
Polyconic (American)		If ϕ is not 0,
If ϕ is 0,		$E = 2 \arctan (A \sin \phi)$ (102)
$x = R(\lambda - \lambda_0)$ (same as equation 1)		Use also equations 96 and 97, if ϕ is not 0. For axes, see Equidistant Conic.
$y = -R\phi_0$	(94)	Modified Polyconic for International Map of the World Series
If ϕ is not zero,		The formulas are too lengthy to be included here. For formulas and other information, see USGS Professional Paper 1395 (Snyder, 1987, p. 131, 134–137).
$E = (\lambda - \lambda_0) \sin \phi$	(95)	
$x = R \cot \phi \sin E$	(96)	Bonne
$y = R[\phi - \phi_0 + \cot \phi (1 - \cos E)]$	(97)	Pseudoconic Projections
For axes, see Equidistant Conic. For ellipsoidal and inverse formulas and more information, see USGS Professional Paper 1395 (Snyder, 1987, p. 124–133).		$A = \cot \phi_1 + \phi_1 - \phi$ (103)
Rectangular Polyconic		$B = (\lambda - \lambda_0)(\cos \phi)/A$ (104)
If ϕ_1 is 0,		$x = RA \sin B$ (105)
$A = (\lambda - \lambda_0)/2$	(98)	$y = R(\cot \phi_1 - A \cos B)$ (106)
If ϕ_1 is not 0,		The Y axis lies along central meridian λ_0 , y increasing north; the X axis crosses the Y axis perpendicularly at (ϕ_1, λ_0) , x increasing east. For ellipsoidal and inverse formulas and more information, see USGS Professional Paper 1395 (Snyder, 1987, p. 138–140).
$A = \tan [(\lambda - \lambda_0)(\sin \phi_1)/2]/\sin \phi_1$	(99)	Werner
		Use equations 103 through 106, where $\phi_1 = \pi/2$ or 90° , but, if ϕ is also 90° , these equations should not be used, and $x = 0$ and $y = 0$.

AZIMUTHAL PROJECTIONS (PERSPECTIVE)

Gnomonic

North polar aspect:

If ϕ is equal to or less than zero, reject the point. Otherwise,

$$\rho = R \cot \phi \quad (107)$$

$$x = \rho \sin (\lambda - \lambda_0) \quad (108)$$

$$y = -\rho \cos (\lambda - \lambda_0) \quad (109)$$

The origin of the axes is at the North Pole. Meridian λ_0 follows the negative Y axis.

Equatorial aspect:

If $\cos \phi \cos (\lambda - \lambda_0)$ is equal to or less than zero, reject the point. Otherwise,

$$x = R \tan (\lambda - \lambda_0) \quad (110)$$

$$y = R \tan \phi / \cos (\lambda - \lambda_0) \quad (111)$$

The Y axis lies along the central meridian λ_0 , y increasing north; the X axis lies along the Equator, x increasing east.

Oblique aspect (also polar and equatorial aspects):

$$\cos z = \sin \phi_1 \sin \phi + \cos \phi_1 \cos \phi \cos (\lambda - \lambda_0) \quad (112)$$

If $\cos z$ is equal to or less than zero, reject the point. Otherwise,

$$K = \sec z \quad (113)$$

$$x = RK \cos \phi \sin (\lambda - \lambda_0) \quad (114)$$

$$y = RK [\cos \phi_1 \sin \phi - \sin \phi_1 \cos \phi \cos (\lambda - \lambda_0)] \quad (115)$$

The Y axis lies along the central meridian λ_0 , y increasing north; the X axis is perpendicular to the Y axis at (ϕ_1, λ_0) , x increasing east. For inverse formulas and other information, see USGS Professional Paper 1395 (Snyder, 1987, p. 164–168).

Stereographic

North polar aspect:

If ϕ is -90° , reject the point. Otherwise, use equations 108 and 109, but

$$\rho = 2Rk_0 \tan (\pi/4 - \phi/2) \quad (116)$$

where k_0 is the scale factor at the center of the projection.

Equatorial aspect:

$$\cos z = \cos \phi \cos (\lambda - \lambda_0) \quad (117)$$

If $\cos z$ is -1 , reject the point. Otherwise,

$$K = 2k_0/(1 + \cos z) \quad (118)$$

$$x = \text{same as equation 114}$$

$$y = RK \sin \phi \quad (119)$$

Oblique aspect (also polar and equatorial aspects):

Use equations 112, 118, 114, and 115 in order, but, if $\cos z$ is -1 , reject the point. Axes for Gnomonic apply here for all aspects. For inverse and ellipsoidal formulas and other information, see USGS Professional Paper 1395 (Snyder, 1987, p. 154–163).

Orthographic

North polar aspect:

If ϕ is less than zero, reject the point. Otherwise, use equations 108 and 109, but

$$\rho = R \cos \phi \quad (120)$$

Equatorial aspect:

If $\cos \phi \cos (\lambda - \lambda_0)$ is less than zero, reject the point. Otherwise,

x = same as in equation 114, where $K = 1$

$y = R \sin \phi$ (same as in equation 9)

Oblique aspect (also polar and equatorial aspects):

If $\cos z$, determined from equation 112, is less than zero, reject the point. Otherwise,

x = same as in equation 114, where $K = 1$

$y =$ same as in equation 115, where $K = 1$

Axes for Gnomonic apply here for all aspects. For inverse formulas and other information, see USGS Professional Paper 1395 (Snyder, 1987, p. 145–153). A hemisphere for any of the aspects of the Orthographic is bounded by a circle of radius R .

General Vertical Perspective

North polar aspect:

If $\sin \phi$ is less than $1/P$, reject the point. Otherwise, use equations 108 and 109, but

$$\rho = R(P - 1) \cos \phi / (P - \sin \phi) \quad (121)$$

where P is the distance of the point of perspective from the center of the Earth divided by the radius R . If the point is beneath the surface and the center of the Earth, apply a minus sign to P .

Equatorial aspect:

Use equations 117, 114, and 119, but, if $\cos z$ is less than $1/P$, reject the point. Otherwise,

$$K = (P - 1) / (P - \cos z) \quad (122)$$

Oblique aspect (also polar and equatorial aspects):

Use equations 112, 122, 114, and 115 in order, but, if $\cos z$ is less than $1/P$, reject the point. Axes for Gnomonic apply here for all aspects. For inverse and ellipsoidal formulas for either Vertical or Tilted Perspective, see USGS Professional Paper 1395 (Snyder, 1987, p. 169–181).

AZIMUTHAL PROJECTIONS (NONPERSPECTIVE)

Azimuthal Equidistant

North polar aspect:

Use equations 108 and 109, but

$$\rho = R(\pi/2 - \phi) \quad (123)$$

Equatorial aspect:

Use equations 117, 114, and 119, but

$$K = z/\sin z \quad (124)$$

If $\cos z = 1$, equation 124 is indeterminate, but $K = 1$. If $\cos z = -1$, for the point opposite the center, the point is plotted as a bounding circle of radius πR .

Oblique aspect (also polar and equatorial aspects):

Use equations 112, 124 (including notes), 114, and 115 in order. Axes for Gnomonic apply here for all aspects. For inverse and ellipsoidal formulas and other information, see USGS Professional Paper 1395 (Snyder, 1987, p. 191–202).

Lambert Azimuthal Equal-Area

North polar aspect:

Use equations 108 and 109, but

$$\rho = 2R \sin (\pi/4 - \phi/2) \quad (125)$$

Equatorial aspect:

Use equations 117, 114, and 119, but

$$K = [2/(1 + \cos z)]^k \quad (126)$$

If $\cos z = -1$, for the point opposite the center, the point is plotted as a bounding circle of radius $2R$.

Oblique aspect (also polar and equatorial aspects):

Use equations 112, 126 (including notes), 114, and 115 in order. Axes for Gnomonic apply here for all aspects. For inverse and ellipsoidal formulas and other information, see USGS Professional Paper 1395 (Snyder, 1987, p. 182–190).

Airy

North polar aspect:

Use equations 108 and 109, but

$$\rho = 2R [\cot(z/2) \ln \sec(z/2) + \tan(z/2) \cot^2(\beta/2) \ln \sec(\beta/2)] \quad (127)$$

where $z = [(\pi/2) - \phi]$ and β = the range from the center of the region in which distortion is to be minimized, on all aspects. For the north polar aspect, $\beta = (\pi/2 - \phi_b)$, where ϕ_b is the bounding latitude. If $z = 0$, equation 127 is indeterminate, but then $\rho = 0$.

Equatorial aspect:

Use equations 117, 114, and 119, but

$$K = -R\{[\ln((1 + \cos z)/2)]/(1 - \cos z) + 2[\ln \cos(\beta/2)]/[\tan^2(\beta/2)(1 + \cos z)]\} \quad (128)$$

If $z = 0$, equation 128 is indeterminate, but, then

$$K = R\{0.5 - [\ln \cos(\beta/2)]/\tan^2(\beta/2)\} \quad (129)$$

Oblique aspect (also polar and equatorial aspects):

Use equations 112, 128 or 129, 114, and 115 in order. Axes for Gnomonic apply here for all aspects.

General transverse and oblique transformations

These formulas can be used with any of the above normal equations for cylindrical, pseudocylindrical, conic, or other projections to obtain transverse or oblique aspects. They can also be used with polar aspects of azimuthal projections, but the general formulas given above for oblique azimuthal projections are more easily used.

To use these formulas, change ϕ in the normal equations (for example, equations 54 and 55) to ϕ' and $(\lambda - \lambda_0)$ to λ' , and then compute ϕ' and λ' from the following:

$$\phi' = \arcsin [\sin \phi_0' \sin \phi + \cos \phi_0' \cos \phi \sin (\lambda - \lambda_0)] \quad (130)$$

$$\lambda' = \lambda_0' + \arctan_2 \{[\cos \phi \cos (\lambda - \lambda_0)]/[\cos \phi_0' \sin \phi - \sin \phi_0' \cos \phi \sin (\lambda - \lambda_0)]\} \quad (131)$$

But, if $\phi = \phi_0'$ and $(\lambda - \lambda_0) = 90^\circ$ or if $\phi = -\phi_0'$ and $(\lambda - \lambda_0) = -90^\circ$, then equation 131 is indeterminate, and λ' can have any value. Under these conditions, $\phi' = 90^\circ$ and -90° , respectively, and the "point" is a line if the corresponding pole of the basic projections is a line.

In the above formulas, (ϕ_0', λ_0') are the north latitude and longitude east of center, respectively, on the normal projection at which the North Pole ($\phi = 90^\circ$) is to be located ($\phi_0' = 0$ for the transverse aspect, and 90° for the normal aspect).

(ϕ', λ') are the latitude and longitude east of center, respectively, on the normal projection at which point (ϕ, λ) is plotted.

λ_0 is the meridian of the globe 90° less than the meridian λ_p , connecting the actual North Pole to the north pole of the base projection. It is the central meridian of the transverse aspect.

For the transverse aspect, this transformation will cause the X axis, rather than the Y axis, to lie along the central meridian. To reverse the axes, replace y with x and x with $-y$ after all other calculations have been completed.

MODIFIED AZIMUTHAL PROJECTIONS

Two-Point Azimuthal

In the following formulas, the axes of the usual azimuthal projections have been rotated, so that the X axis lies along the two central points A and B (B is more easterly, x increases from A to B) from which the directions to other points are made correct. Points A and B are also equidistant from the Y axis. Azimuth Az_1 and distance d to help determine the center point are

$$Az_1 = \arctan_2 \left[\frac{\cos \phi_B \sin (\lambda_B - \lambda_A)}{\cos \phi_A \sin \phi_B - \sin \phi_A \cos \phi_B \cos (\lambda_B - \lambda_A)} \right] \quad (132)$$

$$d = (\frac{1}{2}) \arccos [\sin \phi_A \sin \phi_B + \cos \phi_A \cos \phi_B \cos (\lambda_B - \lambda_A)] \quad (133)$$

The latitude and longitude (ϕ_1, λ_0) of the point midway between A and B are

$$\phi_1 = \arcsin (\sin \phi_A \cos d + \cos \phi_A \sin d \cos Az_1) \quad (134)$$

$$\lambda_0 = \lambda_A + \arctan_2 [\sin d \sin Az_1 / (\cos \phi_A \cos d - \sin \phi_A \sin d \cos Az_1)] \quad (135)$$

The azimuth Az_2 used to rotate the axes is as follows:

$$Az_2 = \arctan_2 \left[\frac{-\cos \phi_A \sin (\lambda_A - \lambda_0)}{\sin \phi_1 \cos \phi_A (\lambda_A - \lambda_0) - \cos \phi_1 \sin \phi_A} \right] \quad (136)$$

The above Az_1 , Az_2 , d , ϕ_1 , and λ_0 apply to the entire map. For other points (ϕ, λ) , find the nonrotated (x, y) from equations 112 through 115 for the Gnomonic projection, and call those coordinates (x_s, y_s) . Rotate and adjust as follows:

$$x = \cos d (x_s \sin Az_2 + y_s \cos Az_2) \quad (137)$$

$$y = y_s \sin Az_2 - x_s \cos Az_2 \quad (138)$$

Two-Point Equidistant

In the following formulas, the axes of the usual azimuthal projections have been rotated so that the X axis lies along the two

central points A and B (B is more easterly, x increases from A to B) from which the distances to other points are correct. Points A and B are also equidistant from the Y axis.

$$x = R (z_A^2 - z_B^2) / (2z_0) \quad (139)$$

$$y = \pm R [4z_0^2 z_B^2 - (z_0^2 - z_A^2 + z_B^2)^2]^{1/2} / (2z_0) \quad (140)$$

where

$$\cos z_0 = \sin \phi_A \sin \phi_B + \cos \phi_A \cos \phi_B \cos (\lambda_B - \lambda_A) \quad (141)$$

$$\cos z_A = \sin \phi_A \sin \phi + \cos \phi_A \cos \phi \cos (\lambda - \lambda_A) \quad (142)$$

$$\cos z_B = \sin \phi_B \sin \phi + \cos \phi_B \cos \phi \cos (\lambda - \lambda_B) \quad (143)$$

and y normally takes the sign of $[\cos \phi_A \cos \phi_B \sin \phi \sin (\lambda_B - \lambda_A) - \cos \phi_A \sin \phi_B \cos \phi \sin (\lambda - \lambda_A) + \sin \phi_A \cos \phi_B \cos \phi \sin (\lambda - \lambda_B)]$, but, if $(z_0 + z_A + z_B) = 2\pi$, the expression for the sign of y gives zero, and y actually has two values at each point (ϕ, λ) , one using the plus and the other using the minus. A world map is bounded by an ellipse having a semimajor axis $\pi R [1 - z_0 / (2\pi)]$ and a semiminor axis $\pi R (1 - z_0 / \pi)^2$, the two axes coinciding with the X and Y axes, respectively.

Miller Oblated Stereographic

First (x, y) are calculated for the oblique Stereographic projection by using equations 112, 118, 114, and 115 in order, but the coordinates are called (x_s, y_s) . These coordinates are then converted to (x, y) as follows:

$$x = Kx_s [1 - Q (3y_s^2 - x_s^2) / 12] \quad (144)$$

$$y = Ky_s [1 + Q (3x_s^2 - y_s^2) / 12] \quad (145)$$

For the map of Europe and Africa, Miller made $K = 0.9245$ and $Q = 0.2522$, and (ϕ_1, λ_0) in equations 112, 114, and 115 were made lat 18° N., long 20° (later 18°) E. If this projection is to be applied to other applications, K and Q are selected as follows, so that the scales at a selected arc distance d_s from the center along the X axis (left and right) equal the scales at arc distance d_b from the center along the Y axis (up and down):

$$T_a = (1 - \cos d_a)/(1 + \cos d_a) \quad (146)$$

$$T_b = (1 - \cos d_b)/(1 + \cos d_b) \quad (147)$$

$$Q = (T_b - T_a)/(T_a + T_a^2 + T_b + T_b^2) \quad (148)$$

$$K = 1/[(1 + T_a)(1 + QT_a)]^{1/4} \quad (149)$$

The nonconformal fill-in projections used as part of this projection in the Eastern Hemisphere are too complicated to describe here.

Wiechel (north polar aspect only)

$$x = R [\sin(\lambda - \lambda_0) \cos \phi - \cos(\lambda - \lambda_0)(1 - \sin \phi)] \quad (150)$$

$$y = -R [\cos(\lambda - \lambda_0) \cos \phi + \sin(\lambda - \lambda_0)(1 + \sin \phi)] \quad (151)$$

The origin of the axes occurs at the North Pole. The central meridian λ_0 starts from the pole along the negative Y axis but soon deviates because it is curved.

Craig Retroazimuthal

If $(\lambda - \lambda_0) = 0$,

$$x = 0$$

$$y = R (\sin \phi - \cos \phi \tan \phi_1) \quad (152)$$

If $(\lambda - \lambda_0) \neq 0$,

$$x = R (\lambda - \lambda_0) \quad (153)$$

$$y = R (\lambda - \lambda_0)[\sin \phi \cos(\lambda - \lambda_0) - \cos \phi \tan \phi_1]/\sin(\lambda - \lambda_0) \quad (154)$$

The axes pass through (ϕ_1, λ_0) , the Y axis following the central meridian λ_0 and y increasing north. The direction from every point on the map to (ϕ_1, λ_0) is shown correctly with respect to the local meridian.

Hammer Retroazimuthal

Modified Azimuthal Projections

Use equations 112 and 124 (including notes) to calculate K , and then calculate (x, y) from the following alterations to equations 114 and 115:

$$x = RK \cos \phi_1 \sin(\lambda - \lambda_0) \quad (155)$$

$$y = -RK [\sin \phi_1 \cos \phi - \cos \phi_1 \sin \phi \cos(\lambda - \lambda_0)] \quad (156)$$

The axes and role of (ϕ_1, λ_0) are the same as those for the Craig Retroazimuthal.

Littrow

$$x = R \sin(\lambda - \lambda_0)/\cos \phi \quad (157)$$

$$y = R \tan \phi \cos(\lambda - \lambda_0) \quad (158)$$

The X axis lies along the Equator, x increasing east; the Y axis lies along the central meridian, y increasing north.

Berghaus Star

If $\phi \geq 0$ (north latitudes),

$$x = R (\pi/2 - \phi) \sin(\lambda - \lambda_0) \quad (159)$$

$$y = -R (\pi/2 - \phi) \cos(\lambda - \lambda_0) \quad (160)$$

If $\phi < 0$ (south latitudes),

$$y = R \{ -\pi D \sin(\lambda - \lambda_n) + C [(\pi/2 - \phi)^2 (5 - 4 \cos(\lambda - \lambda_n)) - \pi^2 \sin^2(\lambda - \lambda_n)]^{1/2} \}/[5 - 4 \cos(\lambda - \lambda_n)] \quad (161)$$

$$x = -[\pi R \sin(\lambda - \lambda_n) + y D]/C \quad (162)$$

where

$$C = \cos(\lambda - \lambda_0) - 2 \cos(\lambda_n - \lambda_0) \quad (163)$$

MODIFIED AZIMUTHAL PROJECTIONS

$$D = \sin(\lambda - \lambda_0) - 2 \sin(\lambda_n - \lambda_0) \quad (164)$$

but if $C = 0$,

$$x = \pm R (\pi/2 - \phi) \quad (165)$$

taking the sign of $\sin(\lambda_n - \lambda_0)$.

In the above, λ_0 is the central meridian for the Northern Hemisphere and extends along the negative Y axis from the North Pole, which is at the origin. The central meridian of each southern point is λ_n , where n is the number (1 to 5) of the point. Normally, $\lambda_1 = \lambda_0$.

Aitoff

$$D = \arccos \{ \cos \phi \cos [(\lambda - \lambda_0)/2] \} \quad (166)$$

If $D = 0$,

$$x = 0$$

$$y = 0$$

Otherwise,

$$C = \sin \phi / \sin D \quad (167)$$

$$x = \pm 2 RD (1 - C^2)^{\frac{1}{4}} \quad (168)$$

taking the sign of $(\lambda - \lambda_0)$.

$$y = RDC \quad (169)$$

The X axis lies along the Equator, x increasing east; the Y axis lies along the central meridian, y increasing north.

Hammer (Elliptical)

$$W = 0.5$$

$$D = 2/\{1 + \cos \phi \cos [W(\lambda - \lambda_0)]\} \quad (170)$$

$$x = R (D^{\frac{1}{4}}/W) \cos \phi \sin [W(\lambda - \lambda_0)] \quad (171)$$

$$y = RD^{\frac{1}{4}} \sin \phi \quad (172)$$

Axes are same as those for the Aitoff.

Briesemeister

The formulas are the same as those for the Hammer, but x is multiplied and y is divided by $(1.75/2)^{\frac{1}{4}} = 0.93541$, and the projection is recentered at lat 45° N., long 10° E. The equations can be combined as follows:

$$x = R (3.5 D)^{\frac{1}{4}} \cos \phi' \sin (\lambda'/2) \quad (173)$$

$$y = R (2D)^{\frac{1}{4}} \sin \phi' / 1.75^{\frac{1}{4}} \quad (174)$$

where

$$D = 2/[1 + \cos \phi' \cos (\lambda'/2)] \quad (175)$$

$$\phi' = \arcsin \{[\sin \phi - \cos \phi \cos (\lambda - 10^\circ)]/2^{\frac{1}{4}}\} \quad (176)$$

$$\lambda' = \arccos \{[\sin \phi + \cos \phi \cos (\lambda - 10^\circ)]/(2^{\frac{1}{4}} \cos \phi')\} \quad (177)$$

The Y axis coincides with the central meridian 10° E., y increasing north, and the X axis crosses it perpendicularly at lat 45° N.

The world map is bounded by an ellipse having a semimajor axis of $7^{\text{th}}R$ and a semiminor axis of $4R/7^{\text{th}}$.

$$x = 2.66723 RC_0 C_1 \sin [(\lambda - \lambda_0)/3] \quad (183)$$

$$y = 1.24104 RSC_1 \quad (184)$$

Modified Azimuthal Projections

Winkel Tripel

Calculate x and y for the Aitoff projection by using equations 166 through 169 and calling the coordinates x_1 and y_1 , respectively. Then calculate x and y for the Equirectangular projection by using equations 14 and 15, where $\phi_1 = 50.467^\circ$ for Winkel's example or 40° for Bartholomew's usage, and call the coordinates x_2 and y_2 , respectively. Finally, average them:

$$x = (x_1 + x_2)/2 \quad (178)$$

$$y = (y_1 + y_2)/2 \quad (179)$$

The axes are the same as those for the Aitoff projection.

Eckert-Greifendorff

Same formulas and axes as Hammer (equations 170–172), but $W = 0.25$.

Wagner VII

$$S = 0.90631 \sin \phi \quad (180)$$

$$C_0 = (1 - S^2)^{\frac{1}{2}} \quad (181)$$

$$C_1 = (2/\{1 + C_0 \cos [(\lambda - \lambda_0)/3]\})^{\frac{1}{2}} \quad (182)$$

Chamberlin Trimetric

Formulas are too complicated to be provided here. They have been developed (forward and inverse) but not published by George Bynum of the Mobil Corporation.

Tilted Perspective

Calculate x and y for the Vertical Perspective projection by using equations 112, 122, 114, and 115 in order, but, if $\cos z$ is less than $1/P$, reject the point. Call these coordinates x_1 and y_1 , respectively. Then calculate the final (ξ, η) as follows:

$$x = (x_1 \cos \gamma - y_1 \sin \gamma) \cos \omega/A \quad (185)$$

$$y = (y_1 \cos \gamma + x_1 \sin \gamma)/A \quad (186)$$

where

$$A = \{(y_1 \cos \gamma + x_1 \sin \gamma) \sin \omega/[R(P - 1)]\} + \cos \omega \quad (187)$$

γ is the azimuth east of north of the Y axis, the most upward-tilted axis of the plane of projection relative to the tangent plane, and ω is the upward angle of tilt, or the angle between the Y axis and the tangent plane. The X axis lies at the intersection of the tangent and tilted planes. The origin is at (ϕ_1, λ_0) .

MISCELLANEOUS PROJECTIONS

Bacon Globular

$$y = R (\pi/2) \sin \phi \quad (188)$$

$$F = [(\pi/2)^2 / |\lambda - \lambda_0| + |\lambda - \lambda_0|]/2 \quad (189)$$

$$x = \pm R [|\lambda - \lambda_0| - F + (F^2 - y^2/R^2)^{1/2}] \quad (190)$$

taking the sign of $(\lambda - \lambda_0)$. If $\lambda = \lambda_0$, equation 189 is infinite and equation 190 is indeterminate, but $x = 0$. The X axis coincides with the Equator, x increasing east, and the Y axis coincides with the central meridian λ_0 .

Fournier Globular I

If $(\lambda - \lambda_0) = 0$,

$$x = 0$$

$$y = R\phi \quad (14)$$

If $\phi = 0$,

$$x = R (\lambda - \lambda_0) \quad (1)$$

$$y = 0$$

If $|\lambda - \lambda_0| = \pi/2$,

$$x = R (\lambda - \lambda_0) \cos \phi \quad (23)$$

$$y = R (\pi/2) \sin \phi \quad (188)$$

Otherwise,

$$C = \pi^2/4 \quad (191)$$

$$P = |\pi \sin \phi| \quad (192)$$

$$S = (C - \phi^2)/(P - 2|\phi|) \quad (193)$$

$$A = (\lambda - \lambda_0)^2/C - 1 \quad (194)$$

$$y = \pm R (\{S^2 - A[C - PS - (\lambda - \lambda_0)^2]\}^{1/2} - S)/4 \quad (195)$$

$$x = \pm R (\lambda - \lambda_0)[1 - y^2/R^2 C]^{1/2} \quad (196)$$

where x and y take the signs of $(\lambda - \lambda_0)$ and ϕ , respectively. Axes are the same as those for the Bacon Globular.

Nicolosi Globular

If $(\lambda - \lambda_0) = 0$ or $\phi = 0$ or $|\lambda - \lambda_0| = \pi/2$, the limiting formulas for the Fournier I apply. Also, if $|\phi| = \pi/2$,

$$x = 0$$

$$y = R\phi \quad (14)$$

Otherwise,

$$b = \pi/[2(\lambda - \lambda_0)] - 2(\lambda - \lambda_0)/\pi \quad (197)$$

$$c = 2\phi/\pi \quad (198)$$

$$d = (1 - c^2)/(\sin \phi - c) \quad (199)$$

$$M = [b(\sin \phi)/d - b/2]/(1 + b^2/d^2) \quad (200)$$

$$N = [d^2(\sin \phi)/b^2 + d/2]/(1 + d^2/b^2) \quad (201)$$

$$x = (\pi R/2)\{M \pm [M^2 + \cos^2 \phi/(1 + b^2/d^2)]^{1/2}\} \quad (202)$$

$$y = (\pi R/2)\{N \pm [N^2 - (d^2(\sin^2 \phi)/b^2 + d \sin \phi - 1)/(1 + d^2/b^2)]^{1/2}\} \quad (203)$$

in which the plus or minus sign in equation 202 takes the sign of $(\lambda - \lambda_0)$ and the plus or minus sign in equation 203 takes the opposite sign of ϕ . Axes are the same as those for the Bacon Globular.

Apian Globular I

$$y = R\phi \quad (14)$$

To calculate x , use equations 189 and 190 for the Bacon Globular, except that y in equation 190 is taken from equation 14 above. If $\lambda = \lambda_0$, the equations are indeterminate, but $x = 0$. Axes are the same as those for the Bacon.

Ortelius Oval

If $|\lambda - \lambda_0| \leq \pi/2$, use the Apian Globular I formulas in equations 14, 189, and 190. For $|\lambda - \lambda_0| > \pi/2$, use equation 14 for y , but

$$x = \pm R\{[(\pi/2)^2 - \phi^2]^{1/4} + |\lambda - \lambda_0| - \pi/2\} \quad (204)$$

taking the sign of $(\lambda - \lambda_0)$. Axes are the same as those for the Bacon.

Lagrange

If $|\phi| = \pi/2$,

$$x = 0$$

$$y = \pm 2R \quad (205)$$

taking the sign of ϕ . Otherwise, if W is the range in degrees of longitude, divided by 180° , within a bounding circle passing through the poles and ϕ_1 is the parallel shown as a straight line,

$$A_1 = [(1 + \sin \phi_1)/(1 - \sin \phi_1)]^{1/(2W)} \quad (206)$$

$$A = [(1 + \sin \phi)/(1 - \sin \phi)]^{1/(2W)} \quad (207)$$

$$V = A_1 A \quad (208)$$

$$C = (V + 1/V)/2 + \cos [(\lambda - \lambda_0)/W] \quad (209)$$

$$x = 2R \sin [(\lambda - \lambda_0)/W]/C \quad (210)$$

$$y = R(V - 1/V)/C \quad (211)$$

For the usual Lagrange, $W = 2$ and $\phi_1 = 0$. For the equatorial Stereographic, $W = 1$ and $\phi_1 = 0$.

Eisenlohr

$$S_1 = \sin [(\lambda - \lambda_0)/2] \quad (212)$$

$$C_1 = \cos [(\lambda - \lambda_0)/2] \quad (213)$$

$$T = \sin (\phi/2)/[\cos (\phi/2) + (2 \cos \phi)^{1/4} C_1] \quad (214)$$

$$C = [2/(1 + T^2)]^{1/4} \quad (215)$$

$$V = \{[\cos (\phi/2) + ((\cos \phi)/2)^{1/4}(C_1 + S_1)]/[\cos (\phi/2) + ((\cos \phi)/2)^{1/4}(C_1 - S_1)]\}^{1/4} \quad (216)$$

$$x = (3 + 8^{1/4})R [-2 \ln V + C(V - 1/V)] \quad (217)$$

$$y = (3 + 8^{1/4})R [-2 \arctan T + CT(V + 1/V)] \quad (218)$$

Miscellaneous Projections

Axes are the same as those for the Bacon. The factor ahead of R provides a scale factor of 1 at the center.

August Epicycloidal

$$C_1 = [1 - \tan^2(\phi/2)]^{1/2} \quad (219)$$

$$C = 1 + C_1 \cos [(\lambda - \lambda_0)/2] \quad (220)$$

$$x_1 = \sin [(\lambda - \lambda_0)/2]C_1/C \quad (221)$$

$$y_1 = \tan (\phi/2)/C \quad (222)$$

$$x = 4Rx_1(3 + x_1^2 - 3y_1^2)/3 \quad (223)$$

$$y = 4Ry_1(3 + 3x_1^2 - y_1^2)/3 \quad (224)$$

Axes are the same as those for the Bacon. The scale factor is 1 at the center.

Gyou

$$\cos a = [\cos \phi \sin (\lambda - \lambda_0) - \sin \phi]/2^{1/4} \quad (225)$$

$$\cos b = [\cos \phi \sin (\lambda - \lambda_0) + \sin \phi]/2^{1/4} \quad (226)$$

$$\sin m = \pm(1 + \cos a \cos b - \sin a \sin b)^{1/4} \quad (227)$$

$$\sin n = \pm(1 - \cos a \cos b - \sin a \sin b)^{1/4} \quad (228)$$

$$x = R \int_0^{\pi/2} (1 - 0.5 \sin^2 m)^{-1/4} dm \quad (229)$$

$$y = R \int_0^{\pi/2} (1 - 0.5 \sin^2 n)^{-1/4} dn \quad (230)$$

MISCELLANEOUS PROJECTIONS

where m takes the sign of $(\lambda - \lambda_0)$ and n takes the sign of ϕ . If $|\phi| = \pi/2$, then $x = 0$, and $y = \pm 1.85407R$, taking the sign of ϕ . Equations 229 and 230 are elliptical integrals of the first kind and can be computed by means of efficiently converging techniques such as the Burlirsch (1965) algorithm (see p. 239). Axes are the same as those for the Bacon.

Peirce Quincuncial

$$\cos a = \cos \phi [\sin(\lambda - \lambda_0) + \cos(\lambda - \lambda_0)]/2^{\frac{1}{4}} \quad (231)$$

$$\cos b = \cos \phi [\sin(\lambda - \lambda_0) - \cos(\lambda - \lambda_0)]/2^{\frac{1}{4}} \quad (232)$$

These values are then used in equations 227 through 230 for the Guyou. The axes intersect at the pole, the negative Y axis coinciding with the central meridian λ_0 and parallel to a side of the bounding square. In equation 227, m takes the sign of $\sin(\lambda - \lambda_0)$, and, in equation 228, n takes the opposite sign of $\cos(\lambda - \lambda_0)$.

Adams Hemisphere in a Square

$$\cos a = \cos \phi \sin(\lambda - \lambda_0) \quad (233)$$

$$b = \pi/2 - \phi \quad (234)$$

These values are then used in equations 227 through 230 for the Guyou, where m takes the sign of $(\sin \phi + \cos a)$ and n takes the sign of $(\sin \phi - \cos a)$. Axes are tilted 45° with respect to the central meridian and the Equator but are parallel to the sides of the bounding square, x increasing in the quadrant with positive ϕ and $(\lambda - \lambda_0)$.

Adams World in a Square I, poles at opposite vertices

$$\cos a = \cos \phi' \sin[(\lambda - \lambda_0)/2] \quad (235)$$

$$\cos b = \sin \phi' \quad (236)$$

where

$$\sin \phi' = \tan(\phi/2) \quad (237)$$

These values are then used in equations 227 through 230 for the Guyou, where m takes the sign of $(\sin \phi' + \cos a)$ and n takes the sign of $(\sin \phi' - \cos a)$. Axes are the same as those used for the Adams hemisphere in a square.

Adams World in a Square II, poles in centers of sides

$$\cos a = \{\cos \phi' \sin[(\lambda - \lambda_0)/2] - \sin \phi'\}/2^{\frac{1}{4}} \quad (238)$$

$$\cos b = \{\cos \phi' \sin[(\lambda - \lambda_0)/2] + \sin \phi'\}/2^{\frac{1}{4}} \quad (239)$$

where ϕ' is found from equation 237 and a and b are then used in equations 227 through 230. Signs and axes are as described for the Guyou.

Lee Conformal World in a Triangle

The formulas are considered too complicated for listing here. For a discussion, see the monograph by Lee (1976).

Conformal World in an Ellipse

The shape of the bounding ellipse is established by a modular angle ϕ between 0° and 90° . An angle of 23.8958° gives a boundary ellipse in which the Equator is twice the length of the central meridian. If $k_s = \sin \Theta$ and $k_c = \cos \Theta$,

$$K' = \int_0^{\pi/2} (1 - k_c^2 \sin^2 \lambda')^{-\frac{1}{2}} d\lambda' \quad (240)$$

where λ' is a convenient function and K' is the complete elliptic integral of the first kind. For $\Theta = 23.8958^\circ$, $K' = 2.34767$. Then,

$$u_1 = 2(1 - k_c) \cos \phi \quad (241)$$

$$v_1 = (1 + k_c)[1 + \cos \phi \cos(\lambda - \lambda_0)] \quad (242)$$

$$A = k_s^2 [1 - \cos \phi \cos (\lambda - \lambda_0)] \quad (243)$$

$$B = u_1 - v_1 \quad (244)$$

$$C = u_1 + v_1 \quad (245)$$

$$R_1 = A - (A^2 - 4k_c BC)^{1/4} \quad (246)$$

$$\lambda' = -\arcsin [R_1/(2Ck_c)] \quad (247)$$

$$\phi' = \arcsin (\{1 - [R_1/(2B)]^2\}^{1/4}/k_s) \quad (248)$$

$$u_2 = \int_0^{\phi'} (1 - k_s^2 \sin^2 \phi')^{-1/4} d\phi' \quad (249)$$

$$v_2 = K' - \int_0^{\lambda'} (1 - k_c^2 \sin^2 \lambda')^{-1/4} d\lambda' \quad (250)$$

$$u_3 = \exp [\pi u_2/(4K')] \quad (251)$$

$$v_3 = \pi v_2/(4K') \quad (252)$$

$$x = R(u_3 + 1/u_3)(\sin v_3)/2 \quad (253)$$

$$y = R(u_3 - 1/u_3)(\cos v_3)/2 \quad (254)$$

If $(\lambda - \lambda_0) = 0$,

$$x = 0$$

$$y = \pm \pi RB/(1 + C) \quad (257)$$

taking the sign of ϕ . If $|\phi| = \pi/2$,

$$x = 0$$

$$y = \pm \pi R \quad (258)$$

taking the sign of ϕ . Otherwise,

$$A = |\pi/(\lambda - \lambda_0) - (\lambda - \lambda_0)/\pi|/2 \quad (259)$$

$$G = C/(B + C - 1) \quad (260)$$

$$P = G(2/B - 1) \quad (261)$$

$$Q = A^2 + G \quad (262)$$

$$S = P^2 + A^2 \quad (263)$$

$$T = G - P^2 \quad (264)$$

$$x = \pm \pi R \{AT + [A^2 T^2 - S(G^2 - P^2)]^{1/4}\}/S \quad (265)$$

$$y = \pm \pi R \{PQ - A[(A^2 + 1)S - Q^2]^{1/4}\}/S \quad (266)$$

where x and y take the signs of $(\lambda - \lambda_0)$ and ϕ , respectively.

Axes are the same as those for the Bacon Globular.

Equations 249 and 250 are also elliptic integrals of the first kind and, like equations 229, 230, and 240, can be computed by means of efficiently converging techniques such as the Burlirsch (1965) algorithm (see p. 239). Axes are the same as those for the Bacon Globular.

Van der Grinten I

$$B = |2\phi/\pi| \quad (255)$$

$$C = (1 - B^2)^{1/4} \quad (256)$$

If $\phi = 0$,

$$x = R(\lambda - \lambda_0) \quad (1)$$

$$y = 0$$

Van der Grinten II

Use equations 255 and 256. If $(\lambda - \lambda_0) = 0$, use equations for this limitation under Van der Grinten I, including equation 257.

Otherwise, use equation 259 and

$$x_1 = [C(1 + A^2)^{1/4} - AC^2]/(1 + A^2B^2) \quad (267)$$

Miscellaneous Projections

MISCELLANEOUS PROJECTIONS

$$x = \pm \pi R x_1 \quad (268)$$

$$y = \pm \pi R (1 - x_1^2 - 24x_1)^{1/4} \quad (269)$$

where x and y take the signs of $(\lambda - \lambda_0)$ and ϕ , respectively.
Axes are the same as those for the Bacon.

Van der Grinten III

Use equations 255 and 256. If $\phi = 0$ or $(\lambda - \lambda_0) = 0$, use equations for these limitations under Van der Grinten I, including equations 1 and 257. Otherwise, use equation 259 and

$$y_1 = B/(1 + C) \quad (270)$$

$$x = \pm \pi R [(A^2 + 1 - y_1^2)^{1/4} - A] \quad (271)$$

$$y = \pm \pi R y_1 \quad (272)$$

where x and y take the signs of $(\lambda - \lambda_0)$ and ϕ , respectively.
Axes are the same as those for the Bacon.

Van der Grinten IV

If $\phi = 0$,

$$\begin{aligned} x &= R(\lambda - \lambda_0) \\ y &= 0 \end{aligned} \quad (1)$$

If $(\lambda - \lambda_0) = 0$ or $|\phi| = \pi/2$,

$$x = 0$$

$$y = R\phi$$

Otherwise,

$$B = |2\phi/\pi| \quad (255)$$

$$C = (8B - B^4 - 2B^2 - 5)/(2B^3 - 2B^2) \quad (273)$$

$$D = \pm \{[2(\lambda - \lambda_0)/\pi + \pi/(2(\lambda - \lambda_0))]^2 - 4\}^{1/4} \quad (274)$$

taking the sign of $(|\lambda - \lambda_0| - \pi/2)$.

$$\begin{aligned} F &= (B + C)^2(B^2 + C^2D^2 - 1) + (1 - B^2)\{B^2[(B + 3C)^2 \\ &\quad + 4C^2] + 12BC^3 + 4C^4\} \end{aligned} \quad (275)$$

$$x_1 = \{D[(B + C)^2 + C^2 - 1] + 2F^{1/4}\}/[4(B + C)^2 + D^2] \quad (276)$$

$$x = \pm \pi R x_1/2 \quad (277)$$

$$y = \pm (\pi R/2)(1 + D|x_1| - x_1^2)^{1/4} \quad (278)$$

where x and y take the signs of $(\lambda - \lambda_0)$ and ϕ , respectively.
Axes are the same as those for the Bacon.

Armadillo

$$x = R(1 + \cos \phi) \sin [(\lambda - \lambda_0)/2] \quad (279)$$

$$\begin{aligned} y &= R \{(1 + \sin 20^\circ - \cos 20^\circ)/2 + \sin \phi \cos 20^\circ \\ &\quad - (1 + \cos \phi) \sin 20^\circ \cos [(\lambda - \lambda_0)/2]\} \end{aligned} \quad (280)$$

$$\phi_s = -\arctan \{\cos[(\lambda - \lambda_0)/2]/\tan 20^\circ\} \quad (281)$$

where, for a given longitude λ , no latitude ϕ more southerly than ϕ_s should be plotted. The southern part of the map is bounded by a curve plotted at the (x, y) calculated from equations 279 and 280 for a succession of (ϕ, λ) points such that each ϕ is equal to ϕ_s calculated for λ from equation 281. The Y axis coincides with the central meridian, y increasing north. The X axis crosses the central meridian at lat 28.1° N.

GS50

The formulas are too lengthy to be included here. For formulas and other information, see USGS Bulletin 1629 (Snyder, 1985, p. 79–92, 147–51) or USGS Professional Paper 1395 (Snyder, 1987, p. 203–212).

Space Oblique Mercator

The formulas are too lengthy to be included here. For formulas and other information, see USGS Bulletin 1518 (Snyder, 1981a) or USGS Professional Paper 1395 (Snyder, 1987, p. 214–229).

Satellite-Tracking

The formulas are too lengthy to be included here. For formulas and other information, see USGS Professional Paper 1395 (Snyder, 1987, p. 230–238).

Evaluation of elliptic integrals

Equations 229, 230, 240, 249, and 250 can be evaluated by using a rapidly converging algorithm by Bulirsch (1965), applicable to all elliptic integrals of the first kind; that is, to solve

$$F = \int_0^\phi (1 - k_c^2 \sin^2 \phi)^{-1/2} d\phi \quad (282)$$

let $S_p = \sin \phi$, and set convergence values $C_a = 10^{-3}$ and $C_b = 10^{-9}$. Also set a limit $E_p = 10^{-5}$. (k_c is frequently called $\sin \Theta$, where Θ is the modulus). Initially, let

$$h = S_p^2 \quad (283)$$

$$k = (1 - k_c^2)^{1/2} \quad (284)$$

If $h \leq C_b$ (or $\phi = 0$), then $F = 0$. If $h \geq 1.0$ (or $|\phi| = \pi/2$), for the “complete” elliptic integral, use equations 285 through 291, but, if $k \leq E_p$, then $F = \pm\infty$, taking the sign of S_p . Let

$$m = 1.0 \quad (285)$$

$$h = m \quad (286)$$

$$m = k + m \quad (287)$$

If

$$|h - k| > C_a m \quad (288)$$

then let

$$k = (hk)^{1/2} \quad (289)$$

$$m = m/2 \quad (290)$$

and repeat equations 286 through 290 until the condition of equation 288 is no longer met. At this point,

$$F = \pm \pi/m \quad (291)$$

taking the sign of S_p . If $h < 1$ and $h > C_b$ (or $|\phi| < \pi/2$ and $|\phi| > 0$), for the “incomplete” elliptic integral, use equations 293 through 305, but, if $k \leq E_p$,

$$F = \ln [(1 + S_p)/(1 - S_p)]/2 \quad (292)$$

Otherwise, let

$$y = [(1 - h)/h]^{1/2} \quad (293)$$

$$n = 0 \quad (294)$$

$$m = 1.0 \quad (295)$$

$$p = mk \quad (296)$$

$$g = m \quad (297)$$

$$m = k + m \quad (298)$$

$$y = y - (p/y) \quad (299)$$

If $|y| \leq 0$, then let

$$y = C_b k^{1/2} \quad (300)$$

If $|g - k| > C_a g$, then let

$$k = 2p^{1/2} \quad (302)$$

$$n = n + n \quad (303)$$

If $y < 0$, then

$$n = n + 1 \quad (304)$$

and equations 296 through 304 are repeated until the condition of equation 301 is no longer met. At this point, if $y < 0$, then $n = n + 1$. In any case,

$$F = \pm [\arctan(m/y) + \pi n]/m \quad (305)$$

taking the sign of S_p and using the arctan in radians as usual.


The above results are generally accurate to 10 decimal places, if C_a is used as given. For accuracy to 10 places or better in all cases, C_a , C_b , and E_p can be made smaller as required.

Miscellaneous Projections

SELECTED REFERENCES

- Bulirsch, R., 1965, Numerical calculation of elliptic integrals and elliptic functions: *Numerische Mathematik*, v. 7, p. 78–90.
- Deetz, C.H., and Adams, O.S., 1934, Elements of map projection with applications to map and chart construction (4th ed.): U.S. Coast and Geodetic Survey Special Publication 68, 200 p.
- Defense Mapping Agency, 1981, Glossary of mapping, charting, and geodetic terms (4th ed.): Washington, D.C., U.S. Government Printing Office, 204 p.
- Hsu, M.-L., 1981, The role of projections in modern map design, in *Monograph 27: Cartographica*, v. 18, no. 2, p. 151–186.
- Lee, L.P., 1944, The nomenclature and classification of map projections: *Empire Survey Review*, v. 7, p. 190–200.
- , 1976, Conformal projections based on elliptic functions, in *Cartographica Monograph 16*: Toronto, University of Toronto Press, 128 p.
- Maling, D.H., 1960, A review of some Russian map projections: *Empire Survey Review*, v. 15, no. 115, p. 203–215; v. 15, no. 116, p. 255–266; v. 15, no. 117, p. 294–303.
- , 1973, Coordinate systems and map projections: London, George Philip, 255 p.
- Maurer, Hans, 1935, *Ebene Kugelbilder*: Gotha, Germany, Petermanns Mitteilungen, Ergänzungsheft 221, 87 p. [Translated by Peter Ludwig with editing and foreword by William Warntz, 1968, *Harvard papers in theoretical geography*: Cambridge, Mass., Harvard University Laboratory for Computer Graphics and Spatial Analysis, 224 p.]
- Miller, O.M., 1942, Notes on cylindrical world map projections: *Geographical Review*, v. 32, no. 3, p. 424–430.
- Reignier, F., 1957, *Les systèmes de projection et leurs applications*: Paris, Institut Géographique National, v. I (texte), 312 p.; v. II (graphiques), 48 pls.
- Robinson, A.H., 1974, A new map projection: Its development and characteristics: *International Yearbook of Cartography*, v. 14, p. 145–155.
- Robinson, A.H., Sale, R.D., Morrison, J.L., and Muehrcke, P.C., 1984, Elements of cartography (5th ed.): New York, John Wiley, 544 p.
- Royal Society, 1966, *Glossary of technical terms in cartography*: London, 84 p.
- Snyder, J.P., 1977, A comparison of pseudocylindrical map projections: *American Cartographer*, v. 4, no. 1, p. 59–81.
- , 1981, The Space Oblique Mercator—Mathematical development: U.S. Geological Survey Bulletin 1518, 108 p.
- , 1982, Map projections used by the U.S. Geological Survey: U.S. Geological Survey Bulletin 1532, 313 p.
- , 1985, Computer-assisted map projection research: U.S. Geological Survey Bulletin 1629, 157 p.
- , 1987, Map projections: A working manual: U.S. Geological Survey Professional Paper 1395, 383 p.
- Steers, J.A., 1970, An introduction to the study of map projections (15th ed.): London, University of London Press, 294 p.
- Wong, F.K.C., 1965, World map projections in the United States from 1940 to 1960: Syracuse, N.Y., Syracuse University, unpublished M.A. thesis, 193 p.

INDEX



INDEX

[Main references in italics]

A	B
Adams, O.S. ----- 70, 74, 192, 194, 195, 198	B.S.A.M. projection ----- 33
Adams projection of hemisphere in square ----- 188, 190, 192-193, 236	Babinet projection ----- 54
Adams projection of world in square I ----- 194, 236	Bacon Globular projection ----- 174, 234
Adams projection of world in square II ----- 195, 236	Bacon, R. ----- 174
Africa, maps of ----- 37, 148	Baranyi, J. ----- 82
Agnese, B. ----- 178	Bartholomew, J. ----- 54, 56, 92, 113, 114, 162
Airy, G.B. ----- 140	Behrmann Cylindrical Equal-Area projection ----- 16, 19-20, 100, 219
Airy projection ----- 131, 132, 140-143, 228-229	Behrmann, W. ----- 19
Aitoff, D. ----- 158	Bellermann, G. ----- 186
Aitoff projection ----- 158-159, 160, 164, 232	Berghaus, H. ----- 156
Alaska, maps of ----- 208	Berghaus Star projection ----- 132, 156-157, 231-232
Albers, H.C. ----- 100	Bienewitz, P. ----- 177
Albers Equal-Area Conic projection ----- 7, 100-101, 102, 224	Bipolar Oblique Conic Conformal projection ----- 97, 99, 224
American Geographical Society ----- 35, 99, 148, 162	Boggs Eumorphic projection ----- 37, 54, 68-69, 70, 221
American Polyconic projection ----- 106	Boggs, S.W. ----- 68
American Telephone & Telegraph Co. ----- 146	Bomford, G. ----- 33
AMS Lunar projection ----- 131	Bonne, R. ----- 113
Apian(us), P. ----- 177	Bonne projection ----- 92, 112-113, 114, 225
Apian Globular I projection ----- 174, 177, 178, 234	Bordone Oval projection ----- 90, 178
Apian Globular II projection ----- 177	Borneo, maps of ----- 14
"Apple-Shaped" projection ----- 204	Bousfield, G.E. ----- 106
Armadillo projection ----- 206-207, 238	Braun, C. ----- 105
Arrowsmith projection ----- 176	Breusing Geometric projection ----- 132, 140
Association of American Geographers ----- 156	Breusing Harmonic projection ----- 140, 143
Asia, maps of ----- 70, 146	Briesemeister projection ----- 8, 158, 160, 162-163, 232-233
Aspect ----- 2	Briesemeister, W.A. ----- 99, 162
"Atlantis" projection ----- 56	British Isles, maps of ----- 26
Atlantic Ocean, maps of ----- 56	British Navy ----- 154
August Epicycloidal projection ----- 184, 186-187, 204, 235	British Ordnance ----- 92
August, F. ----- 186	Bromley projection ----- 54
August projection ----- 186	Bromley, R.H. ----- 54
Authalic projection <i>See Quartic Authalic projection; equal-area projection</i>	Bulirsch algorithm ----- 236, 237, 239
Azimuthal Equidistant projection ----- 92, 94, 114, 132-135, 140, 146, 156, 170, 175, 176, 228	Bumstead, A.H. ----- 106
Modified ----- 158-159, 170	"Butterfly Map" ----- 196
Azimuthal projection ----- 2, 3, 4, 5, 6, 116-143, 226-229	Bynum, G. ----- 233
Modified ----- 144-173, 230-233	
<i>See also specific projection</i>	
C	
Cahill, B.J.S. ----- 196	
Carte Parallélogrammatique ----- 24	

INDEX

- Cassini de Thury, C.F. 26
Cassini projection 22, 26–27, 28, 219
Central Cylindrical projection 2, 10, 30–32, 105, 219
Central projection 2, 116
Chamberlin Trimetric projection 132, 146, 170–171, 233
Chamberlin, W. 170
Circum-Pacific Map Project 136
Clarke, A.R. 131, 140
Clarke's "Twilight" projection 131
Close, C.F. 144, 146
Close projection 144
Collignon, E. 84
Collignon projection 84–85, 88, 223
Colvocoresses, A.P. 211
Combined projections
 See Composite projection
Complex algebra 2, 120
Composite projection 2, 44, 52, 66, 72, 76, 92, 113, 114, 132, 136, 156
Conformal projection, general 2, 5, 8
 Azimuthal 120–123
 Conic 95–99
 Cylindrical 10–15
 Miscellaneous 180–199, 208–212
 Modified azimuthal 148, 154–155
 See also specific projection
Conformal projection of world in ellipse 198–199, 236–237
Conic projection 2, 4, 6, 7, 92–115, 213–214, 224–225, 229
 As specific projection 92
 See also other specific projections
Conical Orthomorphic projection 97
Conventional aspect 2
Cordiform projection 114
Cossin, J. 37
Craig, J.I. 150
Craig Retroazimuthal projection 150–151, 152, 231
Craster, J.E.E. 70
Craster Parabolic projection 37, 68, 70–71, 72, 221
Cylindric projection 6
 See also Cylindrical projection
- Cylindrical Equal-Area projection 10
 See also Lambert Cylindrical Equal-Area projection
Cylindrical projection 2, 4, 6, 10–36, 213, 218–220, 229
 Modified 33, 211
 See also specific projection
- D**
- Deetz, C.H. 70, 106
De la Hire projection
 See La Hire projection
De l'Isle, N. 92
Denoyer, L.P. 80
Denoyer Semi-Elliptical projection 80–81, 222
Denoyer-Geppert 80
Direct aspect 2
Distortion 2, 6, 8
 See also specific projection
Donald Elliptical projection 146
Donald, J.K. 146
Doubly Azimuthal projection 144
Doubly Equidistant projection 146
- E**
- Eckert (-Greifendorff), M. 46, 50, 58, 60, 86, 88, 166
Eckert-Greifendorff projection 74, 158, 160, 166–167, 233
Eckert projections
 I 84, 86–87, 88, 223
 II 84, 86, 88–89, 223
 III 58–59, 60, 178, 221
 IV 58, 60–61, 221
 V 37, 46–47, 48, 50, 220
 VI 37, 44, 46, 50–51, 220
Edge matching 6–7
Egypt, Survey of 150
Eisenlohr, F. 184
Eisenlohr projection 184–185, 186, 204, 235
Ellipsoid, Earth as 2, 3, 4, 7, 12, 14, 22, 26, 95, 100, 106, 113, 132
Elliptic integrals 236, 237, 239
Elliptical projection 54

INDEX

[Main references in italics]

Equal-area polyconic projections	106	Fournier, G.	175
Equal-area projection, general	2, 5, 8	Fournier Globular I projection	175, 176, 234
Azimuthal	136–139	France, maps of	26, 113
Conic	100–103		
Cylindrical	16–21		
Modified azimuthal	160–163, 166–169		
Pseudoazimuthal	149		
Pseudoconic	112–115		
Pseudocylindrical	37–45, 50–57, 60–77, 84, 88–89		
<i>See also specific projection</i>			
Equatorial aspect	2		
<i>See also specific projection</i>			
Equidistant Conic projection	7, 92–94, 113, 224	Gall Isographic projection	24, 33
Equidistant Cylindrical projection	22–29, 158, 164	Gall, J.	19, 33
<i>See also Plate Carrée projection, Equirectangular projection</i>		Gall Orthographic projection	16, 19, 33
Equidistant Polyconic projection	106	Gall projection	10, 30, 33–34, 35, 203, 220
Equidistant projection, general	2, 5	Modified	33
Azimuthal		Gall Stereographic projection	33
<i>See Azimuthal Equidistant projection</i>			
Conic		Gauss, C.F.	12
<i>See Equidistant Conic projection</i>		Gauss Conformal projection	12
Cylindrical		Gauss-Krüger projection	12
<i>See Equidistant Cylindrical projection</i>		General Vertical Perspective projection	124, 128–131, 172, 227
Equirectangular projection	10, 22, 24–25, 46, 48, 92, 219	Geometric projection	
Equivalent projection		<i>See Perspective projection</i>	
<i>See Equal-area projection</i>		Ginzburg, G.A.	106, 149
Eratosthenes	22	Ginzburg III projection	149
Erdi-Kraus, G.	66	Glareanus projection	174, 177
Euler, L.	92	Globular projection	2, 174–177, 204
Eumorphic projection		As a specific projection	176
<i>See Boggs Eumorphic projection</i>		Glossary	2–4
Eurasia, maps of	106	Gnomonic projection	116
Europe, maps of	148	Gnomonic projection	2, 30, 104, 105, 116–119 130, 144, 226
		Goode Homolosine projection	2, 54, 66–67, 221
		Goode, J.P.	66, 114
		Goode Polar Equal-Area projection	114
		Goode's Atlas	66, 83
		Gougenheim, A.	198
		Graticule	2
		<i>See also specific projection</i>	
		Great Britain, maps of	26, 110, 140
		Great circle	2–3, 116, 120, 124
		Gretschel, H.F.	131
		GS50 projection	208–210, 238
		Guam, maps of	132
		Guyou, E.	188
		Guyou projection	188–189, 190, 192, 235–236

INDEX

H

- Hammer (Elliptical) projection 158, 160–161,
162, 166, 168, 232
Hammer, H.H.E. von 131, 152, 158, 160
Hammer Retroazimuthal projection 150, 152–153, 231
Hammer-Aitoff projection 160, 164
 See also Hammer (Elliptical) projection
Hammer-Wagner projection 168
Harvard University 206
Hassler, F.R. 106
Hatano Asymmetrical Equal-Area projection 64–65,
221
Hatano, M. 64
Hatano Symmetrical Equal-Area projection 64
Herschel, J. 97
Hinks, A.R. 152
Homalographic projection 54
Homolographic projection 54, 66
Homolosine projection 37, 54, 66
 See also Goode Homolosine projection
Hondius, J. 37
Hotine, M. 14
Hotine Oblique Mercator projection 14
Hsu, M.-L., quoted 6
Hyperelliptical projection 54

I

- Imniler projection 144
Indicatrix 3, 8
 See also specific projection
International Map of the World (IMW) 95, 111
Interrupted projection 3, 43, 54, 57, 66, 68, 70,
74, 113, 114, 132, 156

J

- Jackson, J.E. 150
Jäger Star projection 156
James, H. 131, 140

K

- Kamenetskiy, V.V. 33

- Kavrayskiy, V.V. 92, 97
Kavrayskiy VI projection 50
“Kite” projection 113
Krasovskiy, F.N. 100
Krüger, L. 12

L

- Laborde, J. 14
Laborde projection 14
Lagrange, J.L. 180
Lagrange projection 154, 180–183, 198, 200, 202, 235
La Hire, P. de 130
La Hire projection 128, 130–131
Lallemand, C. 111
Lambert Azimuthal Equal-Area projection 100,
101, 102, 114, 132, 136–139, 149, 158, 160, 166, 228
Lambert Conformal Conic projection 7,
95–98, 213, 224
Lambert Cylindrical Equal-Area projection 16–18,
19, 100, 101, 218
Lambert Equal-Area Conic projection 100,
102–103, 224
Lambert, J.H. 12, 16, 19, 97, 102, 136, 180
Landsat 211, 213, 214
Large-scale mapping 3
Latitude, defined 3, 216
Least squares 3
Lee, L.P. 12, 148, 196, 198, 208
 Quoted 5–6
Lee conformal projection of the world
 in triangle 196–197, 236
L'Isle
 See De l'Isle
Littrow, J.J. 154
Littrow projection 154–155, 180, 231
Longitude, defined 3, 216
Lorgna projection 136
Loritz projection 174, 177
“Lotus” projection 113
Loximutthal projection 90–91, 223
Loxodrome 3, 10, 90

INDEX

[Main references in italics]

M	
Madagascar, maps of	14
Maling, D.H., quoted	5, 6
Marinus, Projection of	24
Marinus of Tyre	22, 24
Maurer, H.	106, 131, 144, 146, 150
McBryde, F.W.	44, 52, 72, 76
McBryde S3 projection	52–53, 220
McBryde-Thomas Flat-Polar Parabolic projection	72–73, 221–222
McBryde-Thomas Flat-Polar Quartic projection	74, 75, 76–77, 222
McBryde-Thomas Flat-Polar Sinusoidal projection	44–45, 50, 52, 220
McCaw, G.T.	106
McCaw projection	144
Mecca projection	150
Mendeleev, D.I.	92
Mercator Equal-Area projection	37
Mercator, G.	10, 92
Mercator projection	3, 4, 5, 7, 8, 10–11, 12, 14, 30, 33, 35, 90, 97, 200, 208, 213, 218
<i>See also</i> Oblique Mercator projection, Transverse Mercator projection	
Meridian, defined	3
<i>See also specific projection</i>	
Micronesia, maps of	132
Miller Cylindrical projection	10, 33, 35–36, 220
Miller Oblated Stereographic projection	120, 148, 208, 230–231
Miller, O.M.	35, 99, 148, 208
Miller Prolated Stereographic projection	148, 208
Minimum-error projection	3, 140
Mobil Corp.	233
Modified Gall projection	33
Modified Polyconic projection, for International Map of the World	106, 111, 225
<i>Miscellaneous</i>	106
Modified projections	
<i>See specific types, such as azimuthal and cylindrical</i>	
Modified Stereographic Conformal projection	120, 208

Moir	33, 203
Mollweide, C.B.	54
Mollweide projection	37, 54–57, 64, 66, 68, 175, 220
Moon, maps of the	131
Murdoch, P.	92

N

National Atlas of the United States	100
National Geographic Society	35, 70, 106, 146, 170, 200
Navigation, maps for	3, 10, 90, 116, 144, 146, 154
New Zealand Map Grid	208
Newton-Raphson iteration	216–217, 220, 221, 222
Nicolosi, G.	176
Nicolosi Globular projection	175, 176, 234
Nominal scale	3
Nordic projection	162
Normal aspect	
<i>See Direct aspect; see also specific projection</i>	
North America, maps of	99
Nowicki, A.L.	131

O

Oblated Stereographic projection	148, 208
Oblique aspect	3, 229
<i>See also specific projection</i>	
Oblique Cylindrical Orthomorphic projection	14
Oblique Equidistant Cylindrical projection	
<i>See Oblique Plate Carrée projection</i>	
Oblique Hammer projection	162
Oblique Lambert Conformal Conic projection	98
<i>See also Bipolar Oblique Conic Conformal projection</i>	
Oblique Lambert Cylindrical Equal-Area projection	18, 219
Oblique Mercator projection	14–15, 148, 211, 218
Oblique Sinusoidal projection	37, 41, 42
Oblique Plate Carrée (Oblique Equidistant Cylindrical) projection	28–29, 219
Ordinary Polyconic projection	106
Ordnance Survey	140
Ortelius, A.	178
Ortelius Oval projection	177, 178–179, 235

INDEX

Orthoapsidal projection	3, 206	Postel, G.	132		
Orthodromic projection	144	Postel projection	132		
Orthographic projection	3, 124–127, 130, 172, 226–227	Progressive Military Grid	106		
Orthomorphic projection <i>See</i> Conformal projection		Prolated Stereographic projection	148, 208		
Orthophanic projection	82	Pseudoazimuthal projection	6, 149		
Oval projection	178, 179	Pseudoconic projection	4, 6, 112–115, 225		
Oxford University	33	Pseudocylindric projection	6 <i>See also</i> Pseudocylindrical projection		
P		Pseudocylindrical projection	2, 4, 37–91, 113, 220–223, 229		
Pacific Ocean, maps of	136, 148, 208	Modified	80–81, 178–179, 203		
Parallel of latitude, defined	3 <i>See also</i> specific projection	<i>See also specific projection</i>			
Parameters	3	Ptolemy, C.	92, 113		
Parent, A.	131	Putniņš, R.V.	62, 78		
Peirce, C.S.	190	Putniņš projections			
Peirce Quincuncial projection	188, 190–191, 192, 236	P ₁	62		
Perspective Conic projection	104–105, 224–225	P' ₁	58, 62		
Perspective projection, general	3	P ₂	62		
Azimuthal	116–131, 226–227	P' ₂	60, 62–63, 64, 82, 221		
Conic	104–105, 224–225	P ₃	70		
Cylindrical	16–21, 30–34, 219–220	P' ₃	72		
Modified azimuthal	172–173	P ₄	70		
<i>See also</i> specific projection		P' ₄	72		
Petermann Star projection	156	P ₅	78–79, 222		
Peters, A.	19	P' ₅	78		
Peters projection	16, 19, 21	P ₆	78		
Philbrick, A.K.	66	P' ₆	78		
Planar projection	3				
Planimetric map	3, 12	Q			
Plate Carrée projection	22–23, 24, 26, 28, 46, 92, 94, 219	Quartic Authalic projection	74–75, 76, 166, 222		
Polar aspect	4 <i>See also under</i> specific azimuthal projection	Quincuncial projection	190		
Polyconic projection, type	4, 6, 106–111, 180–183, 200–202, 204–205				
As a specific projection	106–109, 225	R			
Polyhedric projection	84	Raisz, E.J.	206		
		Raisz projection	206		
		Rand McNally & Co.	66, 82		
		Rechteckige Plattcarte, Die	24		
		Rectangular Polyconic projection	106, 107, 110, 225		
		Rectangular projection	24		
		Rectified Skew Orthomorphic projection	14		
		Reeves, E.A.	152		

