

Report

Analysis of cholera surveillance data from the Katanga region, DRC – comparing patterns in 2025 and previous years

Status: week 47 2025

Authors: Paulina Jedynak, Birgit Nikolay

2026-01-15

Table of Contents

Table of Contents.....	1
List of Tables	2
List of Figures	3
1. Summary.....	4
2. Definitions.....	5
3. Context and background	6
4. Objectives	7
5. Methods.....	8
5.1. Data sources, study area and population.....	8
5.2. Analysis of the epidemic indicators in years 2015-2025	10
5.2.1. Descriptive analyses	10
5.2.2. Quantitative analysis methods.....	11
5.3. Prioritized health zones	11
5.4. Cholera epidemic patterns in health zones under the vaccination program	11
5.5. Evaluation of the potential endemicity changes.....	12
5.6. Code	12
6. Results.....	12
6.1. Comparison of indicators for 2025 vs. 2015-2024.....	12
6.1.1. Descriptive analysis	12
6.1.2. Quantitative analysis.....	22
6.2. Characteristics of priority health zones during years 2024-2025	23

6.3.	Cholera epidemic patterns in health zones under the vaccination program	23
6.4.	Evaluation of the potential endemicity changes.....	26
7.	Discussion	31
7.1.	Cholera epidemic dynamics in 2025 and previous years	31
7.2.	Prioritization of the health zones	32
7.3.	Cholera epidemic patterns in health zones under the vaccination program	32
7.4.	Endemicity changes	32
8.	Limitations of the study	33
9.	Conclusions	33
10.	Session info	34
Annex	36

List of Tables

Table 1: Indicators of cholera epidemic used in the present study.	10
Table 2: Cholera epidemic indicators in the DRC's Katanga region in years 2015 - 2025. ..	12
Table 3: Number of cases per year, per province.	13
Table 4: Attack rate (%) per 10,000 inhabitants, per year, per province.	15
Table 5: Number of health zones reporting high (>30) number of cases reported in a week, per year, per province.....	16
Table 6: Number of outbreaks per year, per province.	17
Table 7: Number of deaths per year, per province.	18
Table 8: Case fatality risk (%), per year, per province.	20
Table 9: Number of health zones reporting high (>1%, for weeks with at least 10 cases) CFR per year, per province.....	21
Table 10: Models estimates for different cholera epidemic indicators in the DRC's Katanga region in years 2015 - 2025.	22
Table 11: Cholera epidemic indicators in the DRC's Katanga region in years 2024 - 2025, per prioritized (7 health zones) and non-prioritized health zones (49 health zones).	23
Table 12: Number of days that passed after the vaccination took place (starting from 01/07/2021) until the first instance when at least 10 cases were reported.	25
Table 13: Number of days that passed after the vaccination took place (starting from 01/02/2022) until the first instance when at least 10 cases were reported.	26
Table 14: Number of weeks with zero cases per week, per health zone.	27
Table 15: Number of weeks with less than 10 cases per week, per health zone.	29

List of Figures

Figure 1: A map showing area covered by the ‘preventive’ vaccination program in years 2021-2022 and included in this report.	9
Figure 2: Monthly number of cases in 2025 vs historical median(IQR) from years 2015-2024, per province.	14
<i>Figure 3: Attack rate (%) per 10,000 inhabitants, per year, per province.</i>	<i>15</i>
Figure 4: Number of health zones with high number of cases (>30) reported in a week, per year, per province.	16
<i>Figure 5: Number of outbreaks per year, per province.</i>	<i>17</i>
Figure 6: Monthly number of deaths in 2025 vs historical median(IQR) from years 2015-2024, per province.	19
<i>Figure 7: Number of cases and case fatality risk (calculated if at least 10 cases were reported) per year, per province.</i>	<i>20</i>
Figure 8: Number of health zones with high (>1%, for weeks with at least 10 cases) CFR, per year, per province.	21
Figure 9: Number of cases before and after receiving ‘reactive’ vaccination in April and June 2021.	24
<i>Figure 10: Number of cases before and after ‘preventive’ vaccination in December 2021 and January 2022.</i>	<i>26</i>
Figure 11: Number of weeks with zero cases per week for endemic and non endemic health zones, per year.	27
Figure 12: Number of weeks with less than 10 cases per week for endemic and non endemic health zones per year.	29

1. Summary

Background: Since 2018, MSF-OCP has been regularly engaging in cholera outbreak responses in the Katanga region of the Democratic Republic of Congo (DRC) under the Urgepi project. The actions included resource intense interventions in Lubumbashi city (such as during 2024 and 2025), and other smaller interventions in rural settings. While the intervention area in theory covers all 68 health zones in the Katanga region, in 2024 the project prioritized seven health zones for surveillance and response activities based on the cholera epidemic risk assessed in the years preceding prioritization and including the case fatality and number of outbreaks. However, the surveillance data collected in 2025 (until week 47) indicated potential changes in epidemiological patterns, as in some places the number of cases and case-fatality seemed to have exceeded those recorded in previous years, the epidemic lasted longer than expected, and signs of changes in endemicity were observed. This raised the question of whether the Urgepi cholera strategy was well adapted to these changing patterns.

Aim: To retrospectively describe cholera epidemic patterns in the Katanga region in 2025 and compare it with previous years (2015-2024).

Methods: First, we conducted a descriptive analysis of the DRC's Ministry of Health (MoH) national surveillance data on the 2025 cholera outbreak in the Katanga region and compared it to previous years. We chose several cases- and deaths-related epidemic indicators. Next, we fitted generalized models to assess the overall mean difference between 2025 and previous years in terms of these indicators. Moreover, we compared these indicators for prioritized and non-prioritized health zones to evaluate if the prioritization targets were adequate. We also analyzed cholera vaccination data from 2021-2022 and described how long was the following outbreak-free period. Finally, we evaluated changes in the endemic cholera zones by measuring outbreak-free time in zones defined as endemic or non endemic defined at national and MSF level.

Results: Overall, when compared to previous years, in 2025 we observed a significant increase in most of the studied indicators of cholera epidemic: number of cases and deaths, attack rates (ARs), number of health zones reporting high number of cases or high case fatality risk (CFR), as well as the number of outbreaks and their severity. Moreover, the ARs and percentage of prioritized health zones reporting high number of cases were higher than in the non-prioritized ones, while the CFR and percentage of health zones with elevated CFR were lower among the prioritized zones compared to the non-prioritized ones. One of the prioritized health zones (Mulongo) showed signs of being an endemic area although not classified as one by the local authorities. Finally, the results on the impact of the vaccination

campaigns on the length of the outbreak-free period due to the several limitations of the available data.

Conclusions: Our results suggest that in 2025 there was a significant increase in most of the studied indicators of the cholera epidemic when compared to previous years. Moreover, that health zone prioritization strategy applied in 2024 seemed to efficiently target case-related but not deaths-related epidemic indicators. Finally, one prioritized health zone was identified as health zone with endemic-like transmission, although not classified as such by the MoH. The Urgepi project should remain vigilant to these recent changes in cholera epidemic indicators, and their monitoring should be done to assess if this is a long-term change in patterns or if 2025 was an exceptional year.

2. Definitions

Suspected cholera case definition – the following cholera case definition criteria recommended by the 3rd edition of the technical directive on the *Surveillance Intégrée des Maladies et Riposte* (SIMR, Integrated disease surveillance and response) were applied by the Epidemiological Surveillance Directorate of the DRC's MoH: severe dehydration or death following acute watery diarrhea in a patient over 5 years of age. If there is a cholera epidemic, a case should be suspected in any individual over 5 years of age with acute watery diarrhea, with or without vomiting. The number of reported cases included suspected and confirmed cases.

Prioritized health zones – seven non endemic health zones (Fungurume, Katuba, Kilwa, Kiyambi, Manika, Mukanga, Mulongo) in the Grand Katanga province prioritized within the Urgepi project for cholera surveillance and response activities from 2024 onwards. A health zone was prioritized if it reported ≥ 10 deaths in one year after 2019, case fatality $\geq 3\%$, and two to four cholera outbreaks (i.e., 40 cumulative cases during 4 weeks in a year) in the five preceding years.

Endemic health zones - the DRC's MoH defined cholera endemic health zones as follows: location in lacustrine/fluvial areas, being of a “sanctuary” type (areas where cases persist and where epidemic outbreaks have restarted), and being epidemic-free (meaning reporting zero cases) for not more than 16 weeks per year. The following five health zones located in the Grand Katanga region met the criteria: Kinkondja and Malemba Nkulu (Haut Lomami), Kalemie, Moba, and Nyemba (Tanganyika).

Suspected cholera outbreak – solely for the purpose of the present analysis of the cholera epidemic patterns, an outbreak was defined as an occurrence of at least 10 cases for at least two consecutive weeks. However, for the purpose of evaluating the potential

endemicity changes among the health zones classified as endemic by the DRC's MoH, an outbreak was defined as an occurrence of at least one cholera case.

3. Context and background

Cholera and other diarrheal diseases pose a serious public health threat in the world and in the DRC, which is one of the most affected African countries. This is because of multiple structural, environmental, and social factors including intersection of environmental reservoirs, fragile infrastructure, conflict-driven displacement, rapid urban growth, and limited health and WASH systems. All these converge to sustain endemic transmission and recurrent outbreaks.

Since 2018, Médecins Sans Frontières – Operational Center Paris (MSF-OCP) has been regularly engaging in cholera outbreak responses in the Grand Katanga region of the DRC. Grand Katanga is the southeasternmost DRC region populated by approx. 10 million inhabitants and it plays an important role in the cholera dynamics in the DRC due to high burden of disease and persistent hotspot (with some zones in this region are identified as endemic zones). It is comprised of four provinces (Haut Katanga, Haut Lomami, Lualaba, and Tanganyika) and 68 health zones.

The Urgepi project was initiated by MSF in collaboration with the local MoH mainly to improve prevention and response strategies to measles epidemics. The project may, however, intervene during cholera and other epidemic emergencies. The project includes an operational component (led by MSF) and surveillance and research components (led by Epicentre) and focuses on surveillance strategies (early detection and prioritization of outbreaks in need of interventions), preventive vaccination, laboratory confirmation support (coordination and logistics), and interventions (case management and vaccination activities). The project also includes an operational research component to improve the activities listed above.

While the Urgepi project has a well-defined and previously evaluated measles strategy, the project used to intervene in cholera outbreaks on an *ad-hoc* basis. To improve decision making in response to cholera epidemics, the project attempted to develop the first cholera strategic plan in 2024, as part of which seven non endemic health zones in the Grand Katanga province were prioritized for surveillance and response activities. These health zones were identified based on the cholera epidemic risk assessed in the years preceding prioritization and including the case fatality and number of outbreaks. In the first half of 2025, there were, however, growing concerns about potential changes in cholera outbreak

patterns (i.e. bigger, longer and potentially more severe outbreaks) that may require further adaptation of the project's strategy to this changing context.

In years 2021-2025, several health zones in the Grand Katanga region participated in responsive and preventive vaccination campaigns aiming to constrain cholera transmission. Theoretically, a successful vaccination campaign could significantly decrease or even eradicate new cholera cases in the years following the campaign, so understanding for how long a health zone remains free from cholera outbreaks after the vaccination could help relocating the project resources to other health zones in greater need.

Moreover, cholera endemic health zones are often deprioritized for interventions within the Urgepi project as they cannot be sustained over long periods of time due to limited project resources. Recently, concerns have been growing about the current endemicity classification established by the local MoH, and the need has arisen to identify health zones with long endemic-like periods of transmission and flag them as ineligible for prioritization.

A detailed description of cholera epidemic patterns observed in 2025 compared to previous years was therefore needed to help to assess the pertinence of the current cholera strategy, also in the context of operational limitations, and propose potential improvements for the future.

4. Objectives

Main objective

To describe cholera epidemic patterns in the Katanga region in 2025 and assess differences in these patterns compared to previous years (2015-2024)

Secondary objectives

- To assess if the prioritization scheme of health zones made in 2024 was adequate in terms of various cholera epidemic indicators
- To describe cholera patterns after vaccination campaigns implemented in 2021 and 2022
- To identify health zones with endemic-like periods of cholera transmission

5. Methods

5.1. Data sources, study area and population

The analyses were based on the national surveillance data from the MoH in the DRC that are aggregated and shared on a weekly basis with the Urgepi project since 2018 for surveillance purposes and to evaluate / improve the project's performance (as agreed in an MoU with the 4 provinces of Katanga). There seem to have been issues with the data from 2018, as not a single case or death were reported that year and so it was excluded from the analyses. For the years 2015-2017, data were obtained at national level through the MSF intersectional epidemiologist. Data included the number of cases and deaths reported weekly by health zones. The analysis was limited to 56 health zones that reported at least one cholera case during 2015-2025 (out of a total of 68 health zones) in the four provinces (Haut Katanga, Haut Lomami, Lualaba, Tanganyika) of the Katanga region.

Population estimates by health zones were also provided within the national surveillance data. Due to inconsistent and unexplained population numbers in 2024, the population estimates were imputed for 2024 by averaging population sizes from 2023 and 2025. Other sources of population estimates (i.e. provided by the provinces for MSF operational purposes) were explored; however, the estimates from the surveillance data were the most consistent ones over the years and therefore were retained.

Data on 'reactive' and 'preventive' cholera vaccination campaigns, compiled by the MSF epidemiologist based on local reports, were included in the analysis. Although information of vaccination was available for years 2021-2025, only years 2021 and 2022 were finally analyzed as having the highest number of health zones participating in the vaccination campaigns (21 compared to one, zero, and nine in the following years, respectively). In 2021, 'reactive' vaccination campaigns took place in April (1st dose) and June (2nd dose) in 14 non endemic health zones (Kafubu, Kampemba, Kasenga, Kashobwe, Kenya, Kilwa, Kisanga, Lubumbashi, Lukafu, Mufunga Pweto, Sakania, Sampwe, Tshamilemba, and Vangu), while 'preventive' campaigns took place in December 2021 and January 2022 (one dose only) in seven health zones ([Figure 1](#), Bukama, Kalemie, Kansimba, Kinkondja, Malemba Nkulu, Moba, and Nyemba; all except for Bukama and Kansimba are endemic health zones).




Figure 1: A map showing area covered by the 'preventive' vaccination program in years 2021-2022 and included in this report.

Source: Adapted from materials provided by the Division Provinciale de la Santé Tanganyika.

5.2. Analysis of the epidemic indicators in years 2015-2025

5.2.1. Descriptive analyses

To describe cholera patterns for the years 2015-2025 we used the indicators described in [Table 1](#).

Table 1: Indicators of cholera epidemic used in the present study.

Indicators	What the indicator allows to assess
Cases-related indicators	
Number of cases	Overall disease burden
Attack rate (AR) per 10,000 inhabitants	Number of new cases in a time period, accounting for the population size
Number of health zones reporting high number (over 30) cases per week	The overall extent of the most severe impact of the epidemics
Number of cholera outbreaks	Spatial spread of the epidemic
Deaths-related indicators	
Number of deaths	Severity of the epidemic and potential gaps in access to effective health care and case management
Case fatality risk (CFR)*	Severity of cases and potential gaps in access to effective health care and case management
Number of health zones reporting high CFR* (>1%) for weeks with at least 10 cases reported	Identifies zones with high deaths burden and potential gaps in access to effective health care and case management

*The CFR was calculated only for health zones reporting at least 10 cases per week to avoid large uncertainty in the estimates.

The cholera epidemic indicators were compared across the years for the entire region, per province, or per health zone, where appropriate. Monthly number of cases and deaths in 2025 were plotted against historical median and IQR from years 2015-2024. Yearly CFR was displayed as a line plot, overlaid with a bar plot representing the annual number of cases. All other yearly indicators were represented by bar plots.

5.2.2. Quantitative analysis methods

To quantify the overall mean difference between 2025 and previous years in terms of different epidemic indicators, one generalized additive model (GAM) was fitted per outcome aggregated over a week (number of cases, ARs, number of weeks per health zone per year with over 30 cases, number of deaths, CFR, number of weeks per health zone per year with over high CFR, or number of cholera outbreaks). The explanatory variable was a step change in 2025, where the parametric term was meant to capture the overall mean difference between 2025 vs prior years. Each model was adjusted for both geographic location (province modeled as parametric term and health zones as a random effect) and time (week number within year 2025 and years 2015-2024 modeled as explicit cyclic smoothing to capture short-term seasonal fluctuations). Smoothing parameters were estimated using Restricted Maximum Likelihood method. For count response variables (number of cases, number of deaths, and number of outbreaks), negative binomial models with a log link function were fitted. For proportion response variables (ARs and CFRs), an offset was additionally included (log-population or log-number of cases, respectively) in the model described above. Finally, for binomial outcomes (high number of cases within a week, high CFR within a week), binomial models with logistic link function were fitted. The model estimates and confidence intervals (CIs) were exponentiated and presented as rate ratios (number of cases, number of deaths, and number of outbreaks), incidence rate of cases (AR), fatality risk (CFR), or odds ratios (high number of cases within a week, high CFR within a week) for 2025 vs 2015-2024.

5.3. Prioritized health zones

In 2024, seven non endemic health zones (Fungurume, Katuba, Kilwa, Kiyambi, Manika, Mukanga, and Mulongo) in the Grand Katanga province were prioritized for surveillance and response activities. To describe cholera patterns for the years 2024-2025 in these zones, we chose the same indicators as for the overall analysis.

5.4. Cholera epidemic patterns in health zones under the vaccination program

We calculated the number of days that passed after the end of preventive or responsive vaccination campaigns in 2021 and 2022 (counting from 01/07/2021 and 01/02/2022, respectively) until the start of the first week of the next cholera outbreak (defined as at least 10 cases reported within a week) in the vaccinated health zones.

5.5. Evaluation of the potential endemicity changes

We evaluated which health zones currently classified by the DRC's MoH as non endemic reported zero cases per week for less than 16 weeks in 2025 (concurrent with the MoH endemic health zone definition) or <10 cases per week for less than 24 weeks (a definition more relevant from the MSF operational point of view). This way we could identify the non endemic health zones with prolonged transmission periods. Number of weeks with at least one or 10 cases per week for endemic and non endemic health zones per year were presented on a box plot.

5.6. Code

The analyses were conducted using R v. 4.5.2 and RStudio v. 2025.09.2+418. The code used to generate this report is available in the Epicentre repository:

https://github.com/epicentre-msf/DRC_cholera_surv_report.

6. Results

6.1. Comparison of indicators for 2025 vs. 2015-2024

6.1.1. Descriptive analysis

Table 2: Cholera epidemic indicators in the DRC's Katanga region in years 2015 - 2025.

Year ^a	# cases	AR per 10,000 inhabitants	# HZ with >30 cases ^b	# outbreaks ^c	# deaths	CFR	# HZ with >1% CFR ^d	Population size ^e
2015	3,345	4.5	5	12	50	1.5	7	7,475,707
2016	7,103	9.5	11	23	110	1.5	12	7,475,707
2017	10,008	11.1	13	28	223	2.2	17	9,027,387
2019	11,935	11.6	18	52	275	2.3	26	10,330,225
2020	5,451	7.2	12	31	172	3.2	18	7,593,557
2021	4,284	3.6	8	18	91	2.1	13	11,751,393
2022	6,008	6.3	8	35	133	2.2	12	9,530,757
2023	7,866	8.1	10	28	178	2.3	17	9,704,270
2024	7,679	7.8	18	50	266	3.5	25	9,787,448
2025	19,229	19.5	26	54	397	2.1	36	9,870,626

Abbreviations: AR = attack rate; CFR = case fatality risk; HZ = health zone.

^a 2018 was removed as no cases were reported

^b Number of health zones that reported >30 cases during at least one week during the year

^c An outbreak defined as at least 10 cases for at least two consecutive weeks

^d Number of health zones that reported >1% CFR during at least one week during the year

^e Health zones n = 56, Songa excluded

The main indicators of cholera epidemics over the years 2015-2025 are presented in [Table 2](#). In 2025, the number of cases, attack the rates (ARs), the number of health zones reporting high number of cases (>30), and the number of deaths seemed to be increased compared to previous years. Notably, the case fatality risks (CFRs) were often higher in the past (2020, 2024) than in 2025, however the number of health zones reporting CFRs above 1% seemed to be increased in 2025. The number of outbreaks did not appear to be much higher in 2025 compared to previous years, especially for 2019 and 2024 when the number of outbreaks was similar. However, the number of cases per outbreak was higher in 2025.

6.1.1.1. Cases-related indicators

Number of cases

Table 3: Number of cases per year, per province.

Province ^a	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	721	1,340	131	4,082	3,412	755	343	1,665	4,180	5,856
Haut Lomami	1,226	2,050	4,075	3,218	350	658	2,679	470	1,888	5,351
Lualaba	38	29	45	550	450	167	140	341	369	1,045
Tanganyika	1,360	3,684	5,757	4,085	1,239	2,704	2,846	5,390	1,242	6,977
Total	3,345	7,103	10,008	11,935	5,451	4,284	6,008	7,866	7,679	19,229

^aHealth zones n = 56, Songa excluded; 2018 was removed as 0 cases reported in the entire year

In 2025, there was a major increase in the number of cases, overall and for each province ([Table 3](#)). Apart from 2025, we also observed years with higher number of cases (around 2017-2019) and overall, with more cases in Tanganyika and Haut Lomami than in other provinces. When the population size was considered, the attack rates were similar for all provinces except for Haut Lomami where it was higher ([Table 4](#)).




Figure 2: Monthly number of cases in 2025 vs historical median(IQR) from years 2015-2024, per province.

The increase in cases was particularly pronounced for all provinces in the beginning of 2025, but monthly case numbers since September also exceed the 75th percentile of the previous years in Haut-Lomami, Lualaba and Tanganyika (Figure 2).

Attack rates

Table 4: Attack rate (%) per 10,000 inhabitants, per year, per province.

Province ^a	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	2.3	4.4	0.3	6.8	10.7	1.0	1.1	5.2	13.0	18.2
Haut Lomami	5.3	8.9	13.4	14.3	1.5	2.9	11.6	2.0	8.1	23.2
Lualaba	0.7	0.5	0.8	10.0	8.2	3.0	2.5	6.2	6.7	19.0
Tanganyika	8.8	23.8	37.2	26.4	8.0	17.5	8.2	15.0	3.4	18.3
Total	4.5	9.5	11.1	11.6	7.2	3.6	6.3	8.1	7.8	19.5

^aHealth zones n = 56, Songa excluded; 2018 was removed as 0 cases reported in the entire year




Figure 3: Attack rate (%) per 10,000 inhabitants, per year, per province.

AR seems to be fluctuating over the years but for 2025 they are higher than for previous years for all provinces except for Tanganyika, where for years 2016-2019 the AR was higher than for 2025 (Table 4, Figure 3).

High number of cases

Table 5: Number of health zones reporting high (>30) number of cases reported in a week, per year, per province.

Province ^a	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	1	3	0	6	7	1	1	3	11	9
Haut Lomami	2	4	6	6	1	2	4	0	3	7
Tanganyika	2	4	7	5	3	4	3	6	3	7
Lualaba	0	0	0	1	1	1	0	1	1	3
Total	5	11	13	18	12	8	8	10	18	26

^aHealth zones n = 56, Songa excluded; 2018 was removed as 0 cases reported in the entire year




Figure 4: Number of health zones with high number of cases (>30) reported in a week, per year, per province.

The overall number of health zones that reported over 30 cases during at least one week within each year was higher in 2025 compared to previous years (Table 5). At the province level, this pattern is however only detectable for Lualaba and Haut Lomami (Figure 4). For Haut Lomami, the median number of weeks with over 30 cases was higher than in previous years (data not shown), suggesting longer epidemics with high case numbers.

Number of outbreaks

Table 6: Number of outbreaks per year, per province.

Province ^a	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	2	7	0	16	21	2	3	10	25	23
Haut Lomami	4	6	14	15	2	8	17	2	17	14
Lualaba	0	0	1	6	3	2	1	2	2	4
Tanganyika	6	10	13	15	5	6	14	14	6	13
Total	12	23	28	52	31	18	35	28	50	54

^aHealth zones n = 56, Songa excluded; 2018 was removed as 0 cases reported in the entire year




Figure 5: Number of outbreaks per year, per province.

The overall number of outbreaks was slightly elevated in 2025 compared to previous years (Table 6, Figure 5), and years with similar number of outbreaks happened in the past (2019, 2024). However, considering increased number of cases, attack rates, and elevated number of health zones reporting high number of cases in 2025, although not outnumbering much the previous years, the 2025 outbreaks seemed to be more severe (with more reported cases but generally not lasting longer, except for Haut Lomami).

6.1.1.2. Deaths-related indicators

Number of deaths

Table 7: Number of deaths per year, per province.

Province ^a	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	27	34	8	107	135	23	13	73	197	143
Haut Lomami	9	39	84	87	13	19	67	4	46	105
Lualaba	3	1	3	16	13	11	3	17	5	7
Tanganyika	11	36	128	65	11	38	50	84	18	142
Total	50	110	223	275	172	91	133	178	266	397

^aHealth zones n = 56, Songa excluded; 2018 was removed as 0 cases reported in the entire year

The number of reported deaths largely exceeded previous years in Haut-Lomami and Tanganyika (Table 7). In Haut-Katanga the number of deaths was high but lower than in 2024. Lualaba only reported few deaths.




Figure 6: Monthly number of deaths in 2025 vs historical median(IQR) from years 2015-2024, per province.

Patterns of monthly deaths are not consistent across provinces. In Tanganyika, the number of deaths exceeded the 75th percentile of previous years nearly all year long. In Haut Lomami, the 75th percentile was exceeded in the beginning of the year and then again since October, while for Haut Katanga and Lualaba it was exceeded only during the beginning of 2025 (Figure 6).

Case fatality risk (CFR)

Table 8: Case fatality risk (%), per year, per province.

Province ^a	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	4.4	2.1	6.7	2.2	3.8	2.4	2.9	3.2	4.3	2.2
Haut Lomami	0.9	1.5	1.9	2.8	4.8	2.3	2.3	0.5	2.8	2.0
Lualaba	NaN	NaN	5.9	2.4	1.8	4.2	0.0	4.2	0.8	0.7
Tanganyika	1.0	1.0	2.1	1.5	0.7	1.4	1.6	1.5	1.3	1.9
Total	1.5	1.5	2.2	2.3	3.2	2.1	2.2	2.3	3.5	2.1

^aHealth zones n = 56, Songa excluded; 2018 was removed as 0 cases reported in the entire year
NaN means that less than 10 cases were reported for a province in a given year




Figure 7: Number of cases and case fatality risk (calculated if at least 10 cases were reported) per year, per province.

Although the absolute number of cases seemed to increase in 2025, the CFR did not differ for 2025 compared to previous years (Table 8, Figure 7), as there were pronounced fluctuations for this indicator in the preceding years. Per province comparison showed that


CFR was lower in 2025 compared to several years in the past, for all provinces except for Tanganyika.

High CFR

Table 9: Number of health zones reporting high (>1%, for weeks with at least 10 cases) CFR per year, per province.

Province ^a	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	1	4	1	12	12	2	2	8	14	17
Haut Lomami	3	4	7	6	2	5	5	1	7	8
Lualaba	0	0	1	1	2	1	0	1	1	3
Tanganyika	3	4	8	7	2	5	5	7	3	8
Total	7	12	17	26	18	13	12	17	25	36

^aHealth zones n = 56, Songa excluded; 2018 was removed as 0 cases reported in the entire year



In 2018 0 cases were reported for the entire year
Red dashed line marks median # of health zones with high (>1%) CFR calculated for years 2015-2024

Figure 8: Number of health zones with high (>1%, for weeks with at least 10 cases) CFR, per year, per province.

The overall number of health zones reporting high CFRs was elevated in 2025 compared to previous years (Table 9). The effect was most pronounced for Haut Katanga and Lualaba, while for other provinces comparable numbers of health zones presenting high CFRs were observed in previous years (Figure 8). The median number of weeks with high CFR did not differ for 2025 compared to previous years (data not shown).

6.1.2. Quantitative analysis

Table 10: Models estimates for different cholera epidemic indicators in the DRC's Katanga region in years 2015 - 2025.

Indicator for year 2025	Estimate (95% CI)	p-value
# of cases	RR = 1.7 (1.5; 1.9)	<0.001
AR	IRR = 1.4 (1.3; 1.6)	<0.001
# of HZ with high number of cases	OR = 4.6 (3.7; 5.7)	<0.001
# of outbreaks	RR = 1.9 (1.8; 2.0)	<0.001
# of deaths	RR = 1.9 (1.5; 2.3)	<0.001
CFR	RR = 1.1 (0.9; 1.3)	0.4
# of HZ with high CFR	OR = 1.7 (1.2; 2.2)	<0.001

Abbreviations: AR = attack rate; CFR = case fatality risk; CI = confidence interval; HZ = health zone; IRR = incidence rate ratio; OR = odds ratio; RR = risk ratio.

To model the overall mean difference between 2025 and previous years in terms of different epidemic indicators, one generalized additive model was fitted per outcome. The explanatory variable was a step change in 2025, where the parametric term was meant to capture the overall mean difference between 2025 vs prior years. Each model was adjusted for both geographic location (province modeled as parametric term and health zones as a random effect) and time (week number within year 2025 and years 2015-2024 modeled as explicit cyclic smoothing to capture short-term seasonal fluctuations). Smoothing parameters were estimated using Restricted Maximum Likelihood method. For count response variables (number of cases, number of deaths, and number of outbreaks), negative binomial models with a log link function were fitted. For proportion response variables (AR and CFR), an offset was additionally included (log-population or log-number of cases, respectively) in the model described above. Finally, for binomial outcomes (number of health zones reporting high number of cases or high CFR within a week), binomial models with logistic link function were fitted.

The main indicators of cholera epidemics for year 2025 compared to previous years are presented in Table 10. In 2025, all the indicators apart from the CFR were significantly higher than before. Although the CFR itself was not different, the number of health zones reporting CFR above 1% was increased in 2025. All indicators except for the number of outbreaks were significantly affected by seasonal fluctuations (Annex Figure 1).

6.2. Characteristics of priority health zones during years 2024-2025

The main indicators of cholera epidemics in prioritized vs non-prioritized health zones are presented in [Table 11](#).

Table 11: Cholera epidemic indicators in the DRC's Katanga region in years 2024 - 2025, per prioritized (7 health zones) and non-prioritized health zones (49 health zones).

Year	Prioritized HZs	# cases	AR per 10,000 inhabitants	% HZ with >30 cases ^a	# deaths	Mortality rate per 10,000 inhabitants	CFR ^b	% HZ with >1% CFR	Population size ^c
2024	no	6,377	7.5	32.0	232	0.3	3.6	42.0	8,539,766
	yes	1,302	10.4	29.0	34	0.3	2.6	57.1	1,247,682
2025	no	16,324	18.9	44.0	353	0.4	2.2	64.0	8,616,728
	yes	2,905	23.2	57.0	44	0.4	1.5	57.1	1,253,898

Abbreviations: AR = attack rate; CFR = case fatality risk; HZ = health zone.

^a Number of health zones that reported >30 cases during at least one week within a year

^b Number of health zones that reported >1% CFR during at least one week within a year

^c Health zones n = 56, Songa excluded

In 2024, compared to non-prioritized zones, the prioritized zones showed higher ARs (7.5 vs 10.4%) as well as lower CFRs (3.6 vs 2.6), but much higher percentage of health zones with elevated CFR (42.0 vs 57.1%). In 2025, the ARs and percentage of prioritized health zones reporting high number of cases were higher than in the non-prioritized ones (18.9 vs 23.2% and 44 vs. 57%, respectively). The CFR and percentage of health zones with elevated CFR were lower among the prioritized zones compared to the non-prioritized ones (CFR: 2.2 vs 1.5 and health zones with >1% CFR: 64.0 vs 57.2%, respectively).

6.3. Cholera epidemic patterns in health zones under the vaccination program

In April and June 2021, 14 non endemic health zones were provided with two 'reactive' vaccination doses. The median number of days after vaccination to the first occurrence of at least 10 cases per week was 917.5 days [Q1: 583; Q3: 940] (mean 772 days, [Table 12](#), [Figure 9](#)), and the longest outbreak-free period was reported for Tshamilemba (1306 days), the shortest for Kilwa (235 days).




Figure 9: Number of cases before and after receiving 'reactive' vaccination in April and June 2021.

Table 12: Number of days that passed after the vaccination took place (starting from 01/07/2021) until the first instance when at least 10 cases were reported.

Health zone	# days to a week with >10 cases
Kafubu	1,047
Kampemba	935
Kasenga	578
Kashobwe	242
Kenya	935
Kilwa	235
Kisanga	599
Lubumbashi	1,054
Lukafu	914
Mufunga Sampwe	487
Pweto	613
Sakania	942
Tshamilemba	1,306
Vangu	921

In December of the same year and January 2022, ‘preventive’ vaccination campaigns were launched in five endemic and two non endemic health zones. The number of days to the first occurrence of at least 10 cases was immediate for all vaccinated health zones except for Kansimba (non endemic zone) and Malemba Nkulu which reported >10 cases after 356 and 251 days, respectively ([Figure 10](#), [Table 13](#)).




Figure 10: Number of cases before and after 'preventive' vaccination in December 2021 and January 2022.

Table 13: Number of days that passed after the vaccination took place (starting from 01/02/2022) until the first instance when at least 10 cases were reported.

Health zone	# days to a week with >10 cases
Bukama*	6
Kalemie	6
Kansimba*	356
Kinkondja	6
Malemba Nkulu	251
Moba	6
Nyemba	6

*Non-endemic health zones.

6.4. Evaluation of the potential endemicity changes

To evaluate whether the DRC's MoH classification of endemic and non endemic zones was reflected in the number of reported cases, we calculated the number of weeks reporting zero cases per week for endemic and non endemic zones across years.




Figure 11: Number of weeks with zero cases per week for endemic and non endemic health zones, per year.

For most of the years, the median number of weeks reporting zero cases was very distinct in endemic/ non endemic health zones (except for 2020 and 2021 when the numbers were similar, Figure 11). In 2025, the difference was very clear however there were a few non endemic health zones (Bukama, Kabondo Dianda, Kabalo, Kansimba, Manono) that reported zero cases for less than 15 and 16 weeks, which was the 75th percentile of number of weeks reporting zero cases for the endemic zones and the MoH's threshold for non endemic zones, respectively (Table 14).

Table 14: Number of weeks with zero cases per week, per health zone.

Province	Health zone	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	Kafubu	45	42	47	41	42	47	46	46	18	26
Haut Katanga	Kamalondo	47	37	45	28	25	47	46	47	36	30
Haut Katanga	Kambove	44	39	47	46	43	39	46	47	44	35
Haut Katanga	Kampemba	47	24	41	5	14	47	46	36	14	16
Haut Katanga	Kapolowe	43	44	46	31	24	44	47	44	37	37
Haut Katanga	Kasenga	47	47	47	46	35	47	47	32	47	47
Haut Katanga	Kashobwe	46	47	47	46	37	46	43	30	47	47
Haut Katanga	Katuba	47	31	42	15	14	47	46	31	21	20
Haut Katanga	Kenya	46	31	42	13	15	44	47	36	17	22
Haut Katanga	Kikula	38	28	47	29	42	38	41	45	34	30
Haut Katanga	Kilela Balanda	47	46	47	46	47	47	47	47	42	47

Province	Health zone	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	Kilwa	44	34	37	25	32	45	37	34	47	43
Haut Katanga	Kipushi	47	46	46	33	44	47	47	46	29	27
Haut Katanga	Kisanga	47	31	41	21	16	45	47	30	18	21
Haut Katanga	Kowe	47	43	46	41	45	47	47	47	42	34
Haut Katanga	Likasi	43	30	47	44	47	47	41	47	41	40
Haut Katanga	Lubumbashi	47	35	45	18	18	47	45	45	22	25
Haut Katanga	Lukafu	47	46	47	44	16	43	47	38	32	36
Haut Katanga	Mitwaba	46	46	47	46	43	43	47	43	46	46
Haut Katanga	Mufunga Sampwe	25	42	46	11	14	37	41	23	40	37
Haut Katanga	Mumbunda	46	36	39	24	33	47	46	38	26	24
Haut Katanga	Panda	44	41	47	46	46	47	46	47	43	45
Haut Katanga	Pweto	47	47	46	20	44	46	47	23	38	46
Haut Katanga	Rwashi	47	29	46	30	29	46	45	46	31	24
Haut Katanga	Sakania	47	47	47	44	46	47	47	45	34	40
Haut Katanga	Tshamilemba	46	40	47	33	32	47	47	47	34	26
Haut Katanga	Vangu	47	44	47	26	44	47	46	47	33	31
Haut Lomami	Bukama	20	16	13	9	37	37	6	16	8	9
Haut Lomami	Butumba	28	24	28	26	41	38	28	37	28	26
Haut Lomami	Kabondo Dianda	29	18	27	23	29	31	33	42	26	9
Haut Lomami	Kabongo	47	46	47	47	47	47	47	47	47	47
Haut Lomami	Kaniama	47	47	47	47	47	47	47	47	46	45
Haut Lomami	Kayamba	47	47	47	47	47	47	47	47	47	46
Haut Lomami	Kinkondja	8	11	12	7	24	33	26	32	21	23
Haut Lomami	Kitenge	47	41	41	36	47	47	47	47	47	46
Haut Lomami	Lwamba	46	37	32	41	47	47	40	47	35	31
Haut Lomami	Malemba Nkulu	15	18	13	12	25	36	19	31	8	15
Haut Lomami	Mukanga	32	36	23	18	40	45	6	31	12	22
Haut Lomami	Mulongo	33	47	33	26	41	40	16	32	7	17
Lualaba	Bunkeya	47	47	47	30	31	46	42	38	34	29
Lualaba	Dilala	47	45	47	47	40	47	47	46	46	46
Lualaba	Fungurume	46	43	42	23	34	35	39	41	29	27
Lualaba	Kanzenze	47	47	47	37	47	47	47	47	46	47
Lualaba	Lualaba	47	47	47	46	42	47	43	44	40	38
Lualaba	Lubudi	37	40	47	25	40	47	47	47	47	45
Lualaba	Manika	47	47	47	35	39	47	44	44	39	47
Tanganyika	Ankoro	40	47	28	30	47	46	37	28	40	26
Tanganyika	Kabalo	42	30	22	26	37	39	33	27	40	14
Tanganyika	Kalemie	1	0	1	0	3	6	1	0	1	0
Tanganyika	Kansimba	44	34	33	29	20	35	4	13	16	10
Tanganyika	Kiyambi	47	47	43	29	47	47	47	23	45	43
Tanganyika	Kongolo	46	41	25	25	45	39	36	34	43	22
Tanganyika	Manono	37	46	28	20	35	46	33	35	35	14
Tanganyika	Mbulala	46	45	38	37	46	47	47	47	47	46
Tanganyika	Moba	15	10	22	4	20	36	1	6	21	1
Tanganyika	Nyemba	3	2	1	0	4	12	0	0	0	0
Tanganyika	Nyunzu	45	46	46	38	47	41	46	42	42	41

In pink, prioritized health zones


In blue, endemic health zones

In red, potential outlier among non endemic zones in 2025 (0 cases reported for <16 weeks)

In bold, potential outliers in non endemic zones in 2025 (0 cases reported for <15 weeks, which was Q75 for endemic zones in 2025)

In 2018 there were 0 cases reported for the entire year

To evaluate if there are health zones with prolonged endemic-like transmission that should be deprioritized for MSF activities, we also calculated the number of weeks with less than 10 cases reported (a less conservative threshold that is more relevant from the MSF's operational point of view).



Health zones n = 56, Songa excluded
In 2018 there were 0 cases reported for the entire year
Dashed line marks 24 weeks threshold

Figure 12: Number of weeks with less than 10 cases per week for endemic and non endemic health zones per year.

For some years, the median number of weeks reporting <10 cases was similar in endemic/ non endemic health zones (2015, 2020, 2021, and 2024, Figure 12). In 2025, the difference was very clear, however there was one non endemic health zone (prioritized health zone Mulongo) that reported <10 for less than 24 weeks, Table 15. One more non endemic zone (Kisanga) reported <10 cases for <27 weeks, which was the 75th percentile for the endemic zones, Table 15.

Table 15: Number of weeks with less than 10 cases per week, per health zone.

Province	Health zone	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	Kafubu	47	47	47	47	45	47	47	47	27	43
Haut Katanga	Kamalondo	47	47	47	47	44	47	47	47	45	42
Haut Katanga	Kambove	47	45	47	47	47	41	47	47	47	47

Province	Health zone	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Haut Katanga	Kampemba	47	37	47	33	32	47	47	47	34	27
Haut Katanga	Kapolowe	47	47	47	47	43	47	47	47	45	45
Haut Katanga	Kasenga	47	47	47	47	40	47	47	37	47	47
Haut Katanga	Kashobwe	47	47	47	47	43	47	45	43	47	47
Haut Katanga	Katuba	47	47	47	28	36	47	47	44	38	29
Haut Katanga	Kenya	47	47	47	39	42	47	47	47	42	32
Haut Katanga	Kikula	47	35	47	40	45	47	45	47	40	33
Haut Katanga	Kilela Balanda	47	47	47	47	47	47	47	47	46	47
Haut Katanga	Kilwa	47	38	47	31	43	47	43	43	47	47
Haut Katanga	Kipushi	47	47	47	45	47	47	47	47	42	43
Haut Katanga	Kisanga	47	47	47	36	47	47	47	37	32	26
Haut Katanga	Kowe	47	46	47	47	47	47	47	47	47	47
Haut Katanga	Likasi	47	43	47	46	47	47	46	47	47	46
Haut Katanga	Lubumbashi	47	47	47	43	45	47	47	47	39	41
Haut Katanga	Lukafu	47	47	47	47	32	47	47	47	37	43
Haut Katanga	Mitwaba	47	47	47	47	47	44	47	46	47	47
Haut Katanga	Mufunga Sampwe	37	47	47	34	21	47	45	39	47	40
Haut Katanga	Mumbunda	47	47	46	40	46	47	47	47	44	34
Haut Katanga	Panda	47	47	47	47	47	47	47	47	47	47
Haut Katanga	Pweto	47	47	47	42	47	47	47	37	47	47
Haut Katanga	Rwashi	47	46	47	46	43	47	47	47	47	42
Haut Katanga	Sakania	47	47	47	45	47	47	47	47	42	44
Haut Katanga	Tshamilemba	47	47	47	46	47	47	47	47	47	38
Haut Katanga	Vangu	47	47	47	47	47	47	47	47	39	43
Haut Lomami	Bukama	37	32	36	29	47	45	22	38	33	37
Haut Lomami	Butumba	43	45	39	40	47	43	40	47	43	32
Haut Lomami	Kabondo Dianda	38	26	46	39	44	44	45	47	41	33
Haut Lomami	Kabongo	47	47	47	47	47	47	47	47	47	47
Haut Lomami	Kaniama	47	47	47	47	47	47	47	47	47	47
Haut Lomami	Kayamba	47	47	47	47	47	47	47	47	47	47
Haut Lomami	Kinkondja	37	28	30	18	45	42	44	47	39	32
Haut Lomami	Kitenge	47	47	47	47	47	47	47	47	47	47
Haut Lomami	Lwamba	47	47	41	47	47	47	47	47	46	40
Haut Lomami	Malemba Nkulu	44	45	25	27	45	40	43	47	43	27
Haut Lomami	Mukanga	47	47	38	38	47	47	25	46	26	34
Haut Lomami	Mulongo	47	47	33	41	47	43	30	41	29	19
Lualaba	Bunkeya	47	47	47	46	42	47	46	43	45	37
Lualaba	Dilala	47	47	47	47	47	47	47	47	47	47
Lualaba	Fungurume	47	47	45	38	46	42	45	41	38	31
Lualaba	Kanzenze	47	47	47	45	47	47	47	47	47	47
Lualaba	Lualaba	47	47	47	47	47	47	46	47	47	43
Lualaba	Lubudi	47	47	47	46	44	47	47	47	47	46
Lualaba	Manika	47	47	47	41	45	47	47	47	46	47
Tanganyika	Ankoro	47	47	32	43	47	47	43	47	47	39
Tanganyika	Kabalo	47	42	33	43	47	45	47	41	47	37
Tanganyika	Kalemie	30	20	1	9	30	32	22	9	34	0
Tanganyika	Kansimba	47	47	46	46	44	40	45	39	46	37
Tanganyika	Kiyambi	47	47	47	37	47	47	47	32	47	47
Tanganyika	Kongolo	47	47	37	34	47	46	47	39	47	39
Tanganyika	Manono	47	47	35	45	47	47	47	46	47	42
Tanganyika	Mbulala	47	47	47	47	47	47	47	47	47	47

Province	Health zone	2015	2016	2017	2019	2020	2021	2022	2023	2024	2025
Tanganyika	Moba	45	28	29	21	38	42	24	26	45	26
Tanganyika	Nyemba	35	23	8	24	40	30	12	4	31	3
Tanganyika	Nyunzu	47	47	47	46	47	47	47	45	47	47

In pink, prioritized health zones

In blue, endemic health zones

In red, potential outlier among non endemic zones in 2025 (<10 cases reported for <24 weeks)

In bold, potential outliers in non endemic zones in 2025 (<10 cases reported for <27 weeks, which was Q75 for endemic zones in 2025)

In 2018 there were 0 cases reported for the entire year

7. Discussion

7.1. Cholera epidemic dynamics in 2025 and previous years

Overall, in 2025 we observed a significant increase of all the studied indicators of cholera epidemic (except for CFR), compared to previous years.

As for the cases-related indicators, in 2025 we observed an increased total number of cases and the elevated number of weeks with high number of cases. When the population size was considered, the increase of ARs for 2025 compared to previous years followed a similar pattern for all provinces, suggesting that all provinces were similarly affected by the cholera epidemic. Finally, we observed a slight overall increase in the number of cholera outbreaks in 2025 compared to previous years, however on the province level the number of outbreaks was not exceptionally higher compared to previous years. Nevertheless, the reported outbreaks seemed to be bigger in size, with high number of reported cases but generally not lasting longer (except for Haut Lomami).

As for the deaths-related indicators, in 2025 we observed increased mortality rates that followed a similar pattern for all provinces, suggesting that the entire region was similarly affected in this regard. The CFR remained stable over the years and the observed increase in the overall number of deaths was related to more cases reported, which may suggest that cholera outbreaks were not more severe in 2025 compared to previous years, but may also be related to the access to care or case management. On the other hand, the number of weeks with high CFR was significantly elevated, which may suggest a higher mortality burden for the most affected health zones for extended periods of time, possibly due to higher number of cases, unequal access to health care facilities, etc.

For most indicators (except for the number of outbreaks and total number of deaths for years 2015-2024), the seasonal differences significantly impacted the estimators. Both the number of cases and deaths, as well as their associated indicators, were elevated towards the end of the year and during the first half of the following one, which coincides with the

rainy season (April-October). On the other hand, CFR was higher in the second half of the year.

It should be noted that the present analysis implies that surveillance performance was stable over time, however its quality and accuracy may have improved since 2015, hence affecting the trends observed in our study. Nevertheless, this would have been a gradual change over time and not just a recent improvement (i.e., during 2025).

7.2. Prioritization of the health zones

In 2025, ARs and number of health zones reporting high number of cases were elevated compared to non-prioritized zones. On the other hand, the CFR and number of health zones reporting high CFR were lower in prioritized zones, in the contrary to the prioritization goals. This finding raises doubts about the effectiveness of the prioritization approach in identifying health zones at risk of high CFR, and historic high CFR may not be a good predictor of a high CFR during future cholera epidemics.

7.3. Cholera epidemic patterns in health zones under the vaccination program

There was no clear effect of any vaccination scheme on the number of cholera cases. For the ‘reactive’ vaccination, the time to the next outbreak varied greatly – from eight months to over three years. Moreover, such outbreak-free periods were observed in the years before the vaccination in 2021-2022 as well as in the non-vaccinated zones. As for the ‘preventive’ vaccination in the mainly cholera endemic health zones, the data suggests that it was not effective in cholera transmission mitigation in four out of six studied health zones. Overall, due to several limitations of the data on the vaccination against cholera, we could not draw a conclusion about the length of the outbreak-free periods following vaccination in the Grand Katanga region.

7.4. Endemicity changes

In 2025 the DRC’s MoH classification of endemic and non endemic zones and regarding the elevated number of weeks reporting >0 cases by the non endemic zones, held only to some extent. For 10% of the non endemic health zones, zero cases were reported for less than 16 weeks of 2025. For the less conservative and more operationally relevant threshold (<10 cases per week for at least 24 weeks per year), one zone (Mulongo that met the MoH’s criteria of being a non endemic zone), surpassed it in 2025. Therefore, our results suggest

that at least Mulongo should be flagged as health zone with endemic-like cholera transmission pattern, although not classified as such by the MoH.

8. Limitations of the study

Due to significant discrepancies in absolute population numbers and hence their likely inaccuracy, the indicators affected by the population sizes need to be interpreted with caution. The lack of surveillance data for 2018 may also bias the epidemiological indicators estimated for years 2015-2024.

The complex structure of the DRC's surveillance system, which requires information to be transmitted from health areas to health zones and ultimately to the provincial level, may lead to information loss, resulting in underreporting of cholera cases and deaths.

Moreover, these data are collected in health care structures that participate in the surveillance system, therefore cases occurring in the community are not captured by surveillance. Additionally, several private entities providing health care services, such as those common in the city of Lubumbashi, may not report to the general surveillance system. The observed spatial distribution of cases may be biased by the fact that access of patients to healthcare facilities may vary between health zones due to geographic and socio-economic factors as well as due to unequal distribution of private health care structures in the region.

The effects of the vaccination against cholera were difficult to estimate due to a high variability of outbreaks across time and geographic zones – the same health zone may show different epidemic pattern depending on the year, and in the same year there can be significant differences between geographical locations in cholera epidemic patterns. Moreover, very few health zones were covered by the vaccination program in 2021-2022 and the coverage varied among them, especially for the 'preventive' campaign. All these factors might have impeded the evaluation of the effects of the vaccination and complicated the comparisons between periods before and after the vaccination campaign regarding its effects on the number of cholera cases.

9. Conclusions

In conclusion, while epidemics seem to have been bigger in size in 2025, no changes in severity of disease were observed. The Urgepi project should remain vigilant to these recent changes in case numbers, and monitoring of these indicators should be done to assess if this is a long-term change in patterns or if 2025 was an exceptional year in terms

of case load. Our results suggest that health zone prioritization strategy applied in 2024 was efficiently targeting the ARs but not CFRs; an alternative prioritization approach (e.g. based on locally available cholera response capacities) could be considered and tested. Finally, Mulongo health zone should be flagged to the Urgepi project as health zone with endemic-like transmission, although not classified as such by the DRC's MoH.

10. Session info

```
R version 4.5.1 (2025-06-13 ucrt)
Platform: x86_64-w64-mingw32/x64
Running under: Windows 11 x64 (build 26200)

Matrix products: default
  LAPACK version 3.12.1

locale:
[1] LC_COLLATE=English_United States.utf8
[2] LC_CTYPE=English_United States.utf8
[3] LC_MONETARY=English_United States.utf8
[4] LC_NUMERIC=C
[5] LC_TIME=English_United States.utf8

time zone: Europe/Paris
tzcode source: internal

attached base packages:
[1] grid      stats  graphics grDevices utils  datasets methods
[8] base

other attached packages:
[1] viridis_0.6.5   viridisLite_0.4.2 lubridate_1.9.4 forcats_1.0.1
[5] stringr_1.6.0   readr_2.1.6      tibble_3.3.0    tidyverse_2.0.0
[9] yardstick_1.3.2 workflowsets_1.1.1 workflows_1.3.0 tune_2.0.1
[13] tidyr_1.3.1     tailor_0.1.0     rsample_1.3.1   recipes_1.3.1
[17] purrr_1.2.0     parsnip_1.3.3    modeldata_1.5.1 infer_1.0.9
[21] dplyr_1.1.4     dials_1.4.2      scales_1.4.0    broom_1.0.10
[25] tidymodels_1.4.1 rio_1.2.4         patchwork_1.3.2 MASS_7.3-65
[29] janitor_2.2.1   ISOweek_0.6-2    here_1.0.2      gratia_0.11.1
[33] ggpubr_0.6.2    gghighlight_0.5.0 ggh4x_0.3.1     ggplot2_4.0.1
[37] gamm4_0.2-7     mgcv_1.9-4       nlme_3.1-168    lme4_1.1-37
[41] Matrix_1.7-4    flextable_0.9.10


loaded via a namespace (and not attached):
[1] RColorBrewer_1.1-3 rstudioapi_0.17.1 jsonlite_2.0.0
[4] magrittr_2.0.4      farver_2.1.2      nloptr_2.2.1
[7] rmarkdown_2.30      ragg_1.5.0        vctrs_0.6.5
[10] minqa_1.2.8         askpass_1.2.1     rstatix_0.7.3
[13] htmltools_0.5.8.1  cellranger_1.1.0  Formula_1.2-5
[16] parallelly_1.45.1  uuid_1.2-1        lifecycle_1.0.4
```

```

[19] pkgconfig_2.0.3    R6_2.6.1          fastmap_1.2.0
[22] rbibutils_2.4      future_1.68.0     snakecase_0.11.1
[25] digest_0.6.39     furr_0.3.1        rprojroot_2.1.1
[28] textshaping_1.0.4  labeling_0.4.3    timechange_0.3.0
[31] abind_1.4-8        compiler_4.5.1     fontquiver_0.2.1
[34] withr_3.0.2        S7_0.2.1          backports_1.5.0
[37] carData_3.0-5      R.utils_2.13.0    ggsignif_0.6.4
[40] lava_1.8.2         openssl_2.3.4     tools_4.5.1
[43] otel_0.2.0         zip_2.3.3          future.apply_1.20.0
[46] nnet_7.3-20        R.oo_1.27.1        glue_1.8.0
[49] generics_0.1.4     gtable_0.3.6      tzdb_0.5.0
[52] R.methodsS3_1.8.2  class_7.3-23      hms_1.1.4
[55] data.table_1.17.8  xml2_1.5.0         car_3.1-3
[58] pillar_1.11.1      lhs_1.2.0          splines_4.5.1
[61] lattice_0.22-7     survival_3.8-3     tidyselect_1.2.1
[64] fontLiberation_0.1.0 knitr_1.50         gridExtra_2.3
[67] fontBitstreamVera_0.1.1 reformulas_0.4.2  xfun_0.54
[70] hardhat_1.4.2      timeDate_4051.111 stringi_1.8.7
[73] DiceDesign_1.10    yaml_2.3.10        pacman_0.5.1
[76] boot_1.3-32        ggokabeito_0.1.0   evaluate_1.0.5
[79] codetools_0.2-20   officer_0.7.1      gdtools_0.4.4
[82] cli_3.6.5          rpart_4.1.24       tweedie_2.3.5
[85] systemfonts_1.3.1  Rdpack_2.6.4       mirai_2.5.2
[88] readxl_1.4.5       Rcpp_1.1.0         globals_0.18.0
[91] nanonext_1.7.2     parallel_4.5.1     gower_1.0.2
[94] mvnfast_0.2.8      GPfit_1.0-9        listenv_0.10.0
[97] ipred_0.9-15       prodlim_2025.04.28 rlang_1.1.6
[100] cowplot_1.2.0

```

Annex



Shaded area: 95% CI
Lines represent trends captured for weeks 1:47 for years 2015–2024 and 2025

Annex Figure 1: Differences in the predicted values of cholera epidemic indicators (2025 vs 2015–2024). For number of outbreaks, the model found no evidence of seasonal structure across weeks.

Abbreviations: AR = attack rate; CFR = case fatality risk; HZ = health zone.

To model the overall mean difference between 2025 and previous years in terms of different epidemic indicators, one generalized additive model was fitted per outcome. The explanatory variable was a step change in 2025, where the parametric term was meant to capture the overall mean difference between 2025 vs prior years. Each model was adjusted for both geographic location (province modeled as parametric term and health zones as a random effect) and time (week number within year 2025 and years 2015-2024 modeled as explicit cyclic smoothing to capture short-term seasonal fluctuations). Smoothing parameters were estimated using Restricted Maximum Likelihood method. For count response variables (number of cases, number of deaths, and number of outbreaks), negative binomial models with a log link function were fitted. For proportion response variables (AR and CFR), an offset was additionally included (log-population or log-number of cases, respectively) in the model described above. Finally, for binomial outcomes (number of health zones reporting high number of cases or high CFR within a week), binomial models with logistic link function were fitted.