

Nipah Virus, Malaysia and Singapore, 1997–1999

Overview

This dataset records weekly case counts of Nipah virus encephalitis during the 1997–1999 outbreak in Malaysia and Singapore. The outbreak resulted in 257 human cases, primarily among pig farmers. Transmission occurred via a zoonotic spillover pathway from fruit bats to pigs, and then from pigs to humans.

Source

Pulliam et al. (2011).

Data

The dataset contains 49 rows and 5 columns of weekly case counts by region:

- date: week of report
- perak: cases in Perak state
- negeri_sembilan: cases in Negeri Sembilan state
- selangor: cases in Selangor state
- singapore: cases in Singapore

```
using Outbreaks
using DataFrames
df = Outbreaks.nipah_malaysia()
first(df, 6)
```

	date	perak	negeri_sembilan	selangor	singapore
	Date	Int64	Int64	Int64	Int64
1	1997-01-04	0	0	0	0
2	1997-01-11	1	0	0	0
3	1997-01-18	0	0	0	0
4	1997-01-25	1	0	0	0
5	1997-02-01	1	0	0	0
6	1997-02-08	0	0	0	0

Summary statistics

```
describe(df)
```

	variable	mean	min	median	max	nmissing	eltype
	Symbol	Union...	Any	Any	Any	Int64	DataType
1	date		1997-01-04	1997-08-16	1999-04-10	0	Date
2	perak	0.734694	0	0.0	4	0	Int64
3	negeri_sembilan	4.18367	0	0.0	49	0	Int64
4	selangor	0.102041	0	0.0	3	0	Int64
5	singapore	0.22449	0	0.0	11	0	Int64