

Andrew Gard - equitable.equations@gmail.com



Discrete Bivariate Distributions

Suppose we have two discrete random variables X and Y . Often, we're interested in their **joint distribution**, that is, the probabilities of different combinations of values.

Suppose we have two discrete random variables X and Y . Often, we're interested in their **joint distribution**, that is, the probabilities of different combinations of values.

The probability $P(X = x, Y = y)$, where x and y are particular values, is called the **joint probability mass function**, often abbreviated $f(x, y)$.

Example. Suppose the random variables X and Y have the following joint probability distribution.

$f(x, y)$	y			
x	1	2	3	4
1	-	0.12	0.10	0.09
2	-	0.18	0.07	-
3	0.02	0.14	0.11	-
4	0.05	0.12	-	-

Example. Suppose the random variables X and Y have the following joint probability distribution.

$f(x, y)$	y			
x	1	2	3	4
1	-	0.12	0.10	0.09
2	-	0.18	0.07	-
3	0.02	0.14	0.11	-
4	0.05	0.12	-	-

- $P(x = 2 \text{ and } y = 3)$

Example. Suppose the random variables X and Y have the following joint probability distribution.

$f(x, y)$	y			
x	1	2	3	4
1	-	0.12	0.10	0.09
2	-	0.18	0.07	-
3	0.02	0.14	0.11	-
4	0.05	0.12	-	-

- $P(x = 2 \text{ and } y = 3) = f(2, 3) = 0.07$.

Example. Suppose the random variables X and Y have the following joint probability distribution.

$f(x, y)$	y			
x	1	2	3	4
1	-	0.12	0.10	0.09
2	-	0.18	0.07	-
3	0.02	0.14	0.11	-
4	0.05	0.12	-	-

- $P(x = 2 \text{ and } y = 3) = f(2, 3) = 0.07$.
- $P(x = 2 \text{ or } y = 3)$

Example. Suppose the random variables X and Y have the following joint probability distribution.

$f(x, y)$	y			
x	1	2	3	4
1	-	0.12	0.10	0.09
2	-	0.18	0.07	-
3	0.02	0.14	0.11	-
4	0.05	0.12	-	-

- $P(X = 2 \text{ and } Y = 3) = f(2, 3) = 0.07$.
- $P(X = 2 \text{ or } Y = 3) = f(2, 2) + f(2, 3) + f(1, 3) + f(3, 3)$

Example. Suppose the random variables X and Y have the following joint probability distribution.

$f(x, y)$	y			
x	1	2	3	4
1	-	0.12	0.10	0.09
2	-	0.18	0.07	-
3	0.02	0.14	0.11	-
4	0.05	0.12	-	-

- $P(X = 2 \text{ and } Y = 3) = f(2, 3) = 0.07$.
- $P(X = 2 \text{ or } Y = 3) = f(2, 2) + f(2, 3) + f(1, 3) + f(3, 3)$
 $= 0.18 + 0.07 + 0.10 + 0.11 = 0.46$.

For any such joint probability mass function $f(x, y)$, three things are true.

For any such joint probability mass function $f(x, y)$, three things are true.

- $0 \leq f(x, y) \leq 1$ for any $(x, y) \in S$ where S is the sample space of all possible pairs (x, y) .

For any such joint probability mass function $f(x, y)$, three things are true.

- $0 \leq f(x, y) \leq 1$ for any $(x, y) \in S$ where S is the sample space of all possible pairs (x, y) .
- $\sum_{(x,y) \in S} f(x, y) = 1$.

For any such joint probability mass function $f(x, y)$, three things are true.

- $0 \leq f(x, y) \leq 1$ for any $(x, y) \in S$ where S is the sample space of all possible pairs (x, y) .
- $\sum_{(x,y) \in S} f(x, y) = 1$.
- For any event $A \subset S$,

$$P[(X, Y) \in A] = \sum_{(x,y) \in A} f(x, y)$$

Naturally, we can still consider the distribution of each random variable separately.

Naturally, we can still consider the distribution of each random variable separately. We call the corresponding pmfs the **marginal probability mass functions**.

Naturally, we can still consider the distribution of each random variable separately. We call the corresponding pmfs the **marginal probability mass functions**. In symbols,

$$f_X(x) = P(X = x) = \sum_y f(x, y), \quad x \in S_x$$

$$f_Y(y) = P(Y = y) = \sum_x f(x, y), \quad y \in S_y$$

Here S_x and S_y are the sample spaces (or *supports*) of X and Y , respectively.

It's natural to add marginal probabilities to tables like the one in the first example.

$f(x, y)$	y				
x	1	2	3	4	$f_X(x)$
1	-	0.12	0.10	0.09	0.31
2	-	0.18	0.07	-	0.25
3	0.02	0.14	0.11	-	0.27
4	0.05	0.12	-	-	0.17
$f_Y(y)$	0.07	0.56	0.28	0.09	

It's natural to add marginal probabilities to tables like the one in the first example.

$f(x, y)$	y				
x	1	2	3	4	$f_X(x)$
1	-	0.12	0.10	0.09	0.31
2	-	0.18	0.07	-	0.25
3	0.02	0.14	0.11	-	0.27
4	0.05	0.12	-	-	0.17
$f_Y(y)$	0.07	0.56	0.28	0.09	

Note that $\sum_x f_X(x) = \sum_y f_Y(y) = 1$.

Two random variables X and Y are said to be **independent** if and only if

$$f(x, y) = f_X(x) \cdot f_Y(y)$$

for every $x \in S_x$ and $y \in S_y$.

Two random variables X and Y are said to be **independent** if and only if

$$f(x, y) = f_X(x) \cdot f_Y(y)$$

for every $x \in S_x$ and $y \in S_y$. If this equality fails for even one (x, y) pair, we say the random variables are **dependent**.

The random variables in the previous example are not independent.

$f(x, y)$	y				
x	1	2	3	4	$f_X(x)$
1	-	0.12	0.10	0.09	0.31
2	-	0.18	0.07	-	0.25
3	0.02	0.14	0.11	-	0.27
4	0.05	0.12	-	-	0.17
$f_Y(y)$	0.07	0.56	0.28	0.09	

The random variables in the previous example are not independent.

$f(x, y)$	y				
x	1	2	3	4	$f_X(x)$
1	-	0.12	0.10	0.09	0.31
2	-	0.18	0.07	-	0.25
3	0.02	0.14	0.11	-	0.27
4	0.05	0.12	-	-	0.17
$f_Y(y)$	0.07	0.56	0.28	0.09	

For instance, $f(1, 2) = 0.12$ while $f_X(1) \cdot f_Y(2) = (0.31)(0.56) = 0.1736$.

Example. Suppose X and Y have the following joint pmf.

$$f(x, y) = Cx^2y, \quad (x, y) \in \{(1, 1), (1, 2), (2, 2), (3, 1), (3, 2)\}$$

First find C , then compute the marginal probability mass functions $f_X(x)$ and $f_Y(y)$. Finally, determine whether X and Y are independent or not.

Example. Suppose X and Y have the following joint pmf.

$$f(x, y) = Cx^2y, \quad (x, y) \in \{(1, 1), (1, 2), (2, 2), (3, 1), (3, 2)\}$$

First find C , then compute the marginal probability mass functions $f_X(x)$ and $f_Y(y)$. Finally, determine whether X and Y are independent or not.

The probabilities must all sum to one, that is,

$$1 = f(1, 1) + f(1, 2) + f(2, 2) + f(3, 1) + f(3, 2)$$

Example. Suppose X and Y have the following joint pmf.

$$f(x, y) = Cx^2y, \quad (x, y) \in \{(1, 1), (1, 2), (2, 2), (3, 1), (3, 2)\}$$

First find C , then compute the marginal probability mass functions $f_X(x)$ and $f_Y(y)$. Finally, determine whether X and Y are independent or not.

The probabilities must all sum to one, that is,

$$\begin{aligned} 1 &= f(1, 1) + f(1, 2) + f(2, 2) + f(3, 1) + f(3, 2) \\ &= C + 2C + 8C + 9C + 18C = 38C \end{aligned}$$

Example. Suppose X and Y have the following joint pmf.

$$f(x, y) = Cx^2y, \quad (x, y) \in \{(1, 1), (1, 2), (2, 2), (3, 1), (3, 2)\}$$

First find C , then compute the marginal probability mass functions $f_X(x)$ and $f_Y(y)$. Finally, determine whether X and Y are independent or not.

The probabilities must all sum to one, that is,

$$\begin{aligned} 1 &= f(1, 1) + f(1, 2) + f(2, 2) + f(3, 1) + f(3, 2) \\ &= C + 2C + 8C + 9C + 18C = 38C \end{aligned}$$

$$C = 1/38$$

$$f(x, y) = \frac{1}{38}x^2y$$

We've determined that

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

We've determined that

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

So

$$f_X(1) = \sum_y f(1, y) = f(1, 1) + f(1, 2)$$

We've determined that

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

So

$$f_X(1) = \sum_y f(1, y) = f(1, 1) + f(1, 2) = \frac{1}{38} + \frac{2}{38} = \frac{3}{38}$$

We've determined that

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

So

$$f_X(1) = \sum_y f(1, y) = f(1, 1) + f(1, 2) = \frac{1}{38} + \frac{2}{38} = \frac{3}{38}$$

$$f_X(2) = \sum_y f(2, y) = f(2, 2)$$

We've determined that

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

So

$$f_X(1) = \sum_y f(1, y) = f(1, 1) + f(1, 2) = \frac{1}{38} + \frac{2}{38} = \frac{3}{38}$$

$$f_X(2) = \sum_y f(2, y) = f(2, 2) = \frac{8}{38}$$

We've determined that

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

So

$$f_X(1) = \sum_y f(1, y) = f(1, 1) + f(1, 2) = \frac{1}{38} + \frac{2}{38} = \frac{3}{38}$$

$$f_X(2) = \sum_y f(2, y) = f(2, 2) = \frac{8}{38}$$

$$f_X(3) = \sum_y f(3, y) = f(3, 1) + f(3, 2)$$

We've determined that

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

So

$$f_X(1) = \sum_y f(1, y) = f(1, 1) + f(1, 2) = \frac{1}{38} + \frac{2}{38} = \frac{3}{38}$$

$$f_X(2) = \sum_y f(2, y) = f(2, 2) = \frac{8}{38}$$

$$f_X(3) = \sum_y f(3, y) = f(3, 1) + f(3, 2) = \frac{9}{38} + \frac{18}{38} = \frac{27}{38}$$

We've determined that

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

So

$$f_X(1) = \sum_y f(1, y) = f(1, 1) + f(1, 2) = \frac{1}{38} + \frac{2}{38} = \frac{3}{38}$$

$$f_X(2) = \sum_y f(2, y) = f(2, 2) = \frac{8}{38}$$

$$f_X(3) = \sum_y f(3, y) = f(3, 1) + f(3, 2) = \frac{9}{38} + \frac{18}{38} = \frac{27}{38}$$

Notice that $\sum_x f_X(x) = \frac{3}{38} + \frac{8}{38} + \frac{27}{38} = 1.$

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

Similarly,

$$f_Y(1) = \sum_x f(x, 1) = f(1, 1) + f(3, 1) = \frac{1}{38} + \frac{9}{38} = \frac{10}{38}$$

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

Similarly,

$$f_Y(1) = \sum_x f(x, 1) = f(1, 1) + f(3, 1) = \frac{1}{38} + \frac{9}{38} = \frac{10}{38}$$

$$\begin{aligned} f_Y(2) &= \sum_x f(x, 2) = f(1, 2) + f(2, 2) + f(3, 2) = \frac{2}{38} + \frac{8}{38} + \frac{18}{38} \\ &= \frac{28}{38} \end{aligned}$$

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

Similarly,

$$f_Y(1) = \sum_x f(x, 1) = f(1, 1) + f(3, 1) = \frac{1}{38} + \frac{9}{38} = \frac{10}{38}$$

$$\begin{aligned} f_Y(2) &= \sum_x f(x, 2) = f(1, 2) + f(2, 2) + f(3, 2) = \frac{2}{38} + \frac{8}{38} + \frac{18}{38} \\ &= \frac{28}{38} \end{aligned}$$

Again, the marginal probabilities sum to one.

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

We've determined that

$$f_X(x) = \begin{cases} 3/38 & \text{if } x = 1 \\ 8/38 & \text{if } x = 2 \\ 27/38 & \text{if } x = 3 \end{cases}, \quad f_Y(y) = \begin{cases} 10/38 & \text{if } y = 1 \\ 28/38 & \text{if } y = 2 \end{cases}.$$

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

We've determined that

$$f_X(x) = \begin{cases} 3/38 & \text{if } x = 1 \\ 8/38 & \text{if } x = 2 \\ 27/38 & \text{if } x = 3 \end{cases}, \quad f_Y(y) = \begin{cases} 10/38 & \text{if } y = 1 \\ 28/38 & \text{if } y = 2 \end{cases}.$$

If X and Y are independent, then $f(x, y) = f_X(x)f_Y(y)$ for all (x, y) .

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

We've determined that

$$f_X(x) = \begin{cases} 3/38 & \text{if } x = 1 \\ 8/38 & \text{if } x = 2 \\ 27/38 & \text{if } x = 3 \end{cases}, \quad f_Y(y) = \begin{cases} 10/38 & \text{if } y = 1 \\ 28/38 & \text{if } y = 2 \end{cases}.$$

If X and Y are independent, then $f(x, y) = f_X(x)f_Y(y)$ for all (x, y) . However, when $X = 1$ and $Y = 1$,

$$f_X(1)f_Y(1) = \frac{3}{38} \cdot \frac{10}{38} = \frac{30}{1444} \approx 0.021$$

$$f(1, 1) = \frac{1}{38} \approx 0.026$$

$$f(x, y) = \frac{1}{38}x^2y, \quad (x, y) = (1, 1), (1, 2), (2, 2), (3, 1), (3, 2)$$

We've determined that

$$f_X(x) = \begin{cases} 3/38 & \text{if } x = 1 \\ 8/38 & \text{if } x = 2 \\ 27/38 & \text{if } x = 3 \end{cases}, \quad f_Y(y) = \begin{cases} 10/38 & \text{if } y = 1 \\ 28/38 & \text{if } y = 2 \end{cases}.$$

If X and Y are independent, then $f(x, y) = f_X(x)f_Y(y)$ for all (x, y) . However, when $X = 1$ and $Y = 1$,

$$f_X(1)f_Y(1) = \frac{3}{38} \cdot \frac{10}{38} = \frac{30}{1444} \approx 0.021$$

$$f(1, 1) = \frac{1}{38} \approx 0.026$$

Since $f(1, 1) \neq f_X(1)f_Y(1)$, the random variables are not independent.