

Step 2: PCA m=2, varimax rotation**The FACTOR Procedure**

Input Data Type	Raw Data
Number of Records Read	572
Number of Records Used	572
N for Significance Tests	572

Correlations								
	palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic
palmitic	1.00000	0.83560	-0.17039	-0.83734	0.46068	0.31933	0.22830	0.50195
palmitoleic	0.83560	1.00000	-0.22219	-0.85244	0.62163	0.09311	0.08548	0.41635
stearic	-0.17039	-0.22219	1.00000	0.11360	-0.19782	0.01892	-0.04098	0.14038
oleic	-0.83734	-0.85244	0.11360	1.00000	-0.85032	-0.21817	-0.31996	-0.42415
linoleic	0.46068	0.62163	-0.19782	-0.85032	1.00000	-0.05744	0.21097	0.08904
linolenic	0.31933	0.09311	0.01892	-0.21817	-0.05744	1.00000	0.62024	0.57832
arachidic	0.22830	0.08548	-0.04098	-0.31996	0.21097	0.62024	1.00000	0.32866
eicosenoic	0.50195	0.41635	0.14038	-0.42415	0.08904	0.57832	0.32866	1.00000

Step 2: PCA m=2, varimax rotation**The FACTOR Procedure****Initial Factor Method: Principal Components**

Partial Correlations Controlling all other Variables								
	palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic
palmitic	1.00000	-0.49371	-0.81558	-0.96675	-0.96050	-0.30224	-0.51243	-0.50208
palmitoleic	-0.49371	1.00000	-0.62247	-0.65857	-0.60200	-0.38686	-0.46754	-0.16313
stearic	-0.81558	-0.62247	1.00000	-0.84270	-0.83486	-0.36214	-0.47383	-0.31466
oleic	-0.96675	-0.65857	-0.84270	1.00000	-0.99043	-0.37359	-0.54207	-0.53022
linoleic	-0.96050	-0.60200	-0.83486	-0.99043	1.00000	-0.39491	-0.49354	-0.54421
linolenic	-0.30224	-0.38686	-0.36214	-0.37359	-0.39491	1.00000	0.22546	0.15585
arachidic	-0.51243	-0.46754	-0.47383	-0.54207	-0.49354	0.22546	1.00000	-0.28058
eicosenoic	-0.50208	-0.16313	-0.31466	-0.53022	-0.54421	0.15585	-0.28058	1.00000

Kaiser's Measure of Sampling Adequacy: Overall MSA = 0.36626306								
palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic	
0.37770058	0.52941582	0.04962441	0.39785809	0.27911482	0.54256492	0.33795395	0.50405344	

Step 2: PCA m=2, varimax rotation**The FACTOR Procedure****Initial Factor Method: Principal Components****Prior Communality Estimates: ONE**

Eigenvalues of the Correlation Matrix: Total = 8 Average = 1				
	Eigenvalue	Difference	Proportion	Cumulative
1	3.72141001	1.95561249	0.4652	0.4652
2	1.76579752	0.74944209	0.2207	0.6859
3	1.01635543	0.22345660	0.1270	0.8129
4	0.79289883	0.45908117	0.0991	0.9121

Eigenvalues of the Correlation Matrix: Total = 8 Average = 1				
	Eigenvalue	Difference	Proportion	Cumulative
5	0.33381767	0.08499900	0.0417	0.9538
6	0.24881867	0.12999856	0.0311	0.9849
7	0.11882011	0.11673835	0.0149	0.9997
8	0.00208176		0.0003	1.0000

2 factors will be retained by the NFACTOR criterion.

Factor Pattern		
	Factor1	Factor2
palmitic	0.88882	-0.06589
palmitoleic	0.86853	-0.32013
stearic	-0.19030	0.34334
oleic	-0.95331	0.21083
linoleic	0.70546	-0.45632
linolenic	0.42245	0.80373
arachidic	0.44042	0.59425
eicosenoic	0.60162	0.53787

Variance Explained by Each Factor	
Factor1	Factor2
3.7214100	1.7657975

Final Community Estimates: Total = 5.487208							
palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic
0.79433932	0.85682226	0.15409583	0.95325274	0.70590405	0.82444057	0.54709786	0.65125490

Residual Correlations With Uniqueness on the Diagonal								
	palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic
palmitic	0.20566	0.04255	0.02137	0.02388	-0.19641	-0.00320	-0.12400	0.00266
palmitoleic	0.04255	0.14318	0.05300	0.04303	-0.13717	-0.01650	-0.10680	0.06601
stearic	0.02137	0.05300	0.84590	-0.14020	0.09310	-0.17665	-0.16120	0.07019
oleic	0.02388	0.04303	-0.14020	0.04675	-0.08159	0.01510	-0.02539	0.03599
linoleic	-0.19641	-0.13717	0.09310	-0.08159	0.29410	0.01130	0.17144	-0.08994
linolenic	-0.00320	-0.01650	-0.17665	0.01510	0.01130	0.17556	-0.04343	-0.10814
arachidic	-0.12400	-0.10680	-0.16120	-0.02539	0.17144	-0.04343	0.45290	-0.25593
eicosenoic	0.00266	0.06601	0.07019	0.03599	-0.08994	-0.10814	-0.25593	0.34875

Root Mean Square Off-Diagonal Residuals: Overall = 0.10560557							
palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic
0.09008464	0.07683248	0.11570032	0.06644227	0.12554076	0.08055349	0.14647815	0.11702773

Partial Correlations Controlling Factors								
	palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic
palmitic	1.00000	0.24795	0.05123	0.24351	-0.79862	-0.01682	-0.40630	0.00993
palmitoleic	0.24795	1.00000	0.15230	0.52600	-0.66845	-0.10408	-0.41941	0.29541
stearic	0.05123	0.15230	1.00000	-0.70503	0.18666	-0.45839	-0.26043	0.12923
oleic	0.24351	0.52600	-0.70503	1.00000	-0.69582	0.16666	-0.17452	0.28184
linoleic	-0.79862	-0.66845	0.18666	-0.69582	1.00000	0.04973	0.46975	-0.28082
linolenic	-0.01682	-0.10408	-0.45839	0.16666	0.04973	1.00000	-0.15402	-0.43703

Partial Correlations Controlling Factors								
	palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic
arachidic	-0.40630	-0.41941	-0.26043	-0.17452	0.46975	-0.15402	1.00000	-0.64397
eicosenoic	0.00993	0.29541	0.12923	0.28184	-0.28082	-0.43703	-0.64397	1.00000

Root Mean Square Off-Diagonal Partials: Overall = 0.39311690							
palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic
0.36383980	0.39318741	0.34895384	0.45589257	0.52175748	0.25806937	0.39611518	0.35214584

Step 2: PCA m=2, varimax rotation

The FACTOR Procedure
Rotation Method: Varimax

Orthogonal Transformation Matrix		
	1	2
1	0.86214	0.50666
2	-0.50666	0.86214

Rotated Factor Pattern		
	Factor1	Factor2
palmitic	0.79967	0.39353
palmitoleic	0.91099	0.16406
stearic	-0.33802	0.19959
oleic	-0.92871	-0.30124
linoleic	0.83941	-0.03598
linolenic	-0.04301	0.90697
arachidic	0.07862	0.73547
eicosenoic	0.24617	0.76854

Variance Explained by Each Factor		
	Factor1	Factor2
	3.2193882	2.2678193

Final Community Estimates: Total = 5.487208							
palmitic	palmitoleic	stearic	oleic	linoleic	linolenic	arachidic	eicosenoic
0.79433932	0.85682226	0.15409583	0.95325274	0.70590405	0.82444057	0.54709786	0.65125490

Step 2: PCA m=2, varimax rotation

The FACTOR Procedure
Rotation Method: Varimax



