# Causal Boundary and its relations with the conformal boundary

#### Jónatan Herrera Fernández

(joint work with Jose Luis Flores and Miguel Sánchez, in preparation)

July-2009





# **INDEX**

#### Introduction

- Physical motivation.
  - Study of singularities.
  - Asymptotic properties of fields.
  - Global properties of spacetimes.

- Mathematical motivation.
  - Classical problem
  - Interest of the mathematical tools involved

- Physical motivation.
  - Study of singularities.
  - Asymptotic properties of fields.
  - Global properties of spacetimes.

- Mathematical motivation.

- Physical motivation.
  - Study of singularities.
  - Asymptotic properties of fields.
  - Global properties of spacetimes.

- Mathematical motivation.

- Physical motivation.
  - Study of singularities.
  - Asymptotic properties of fields.
  - Global properties of spacetimes.

- Mathematical motivation.

- Physical motivation.
  - Study of singularities.
  - Asymptotic properties of fields.
  - Global properties of spacetimes.

- Mathematical motivation.
  - Classical problem.

- Physical motivation.
  - Study of singularities.
  - Asymptotic properties of fields.
  - Global properties of spacetimes.

- Mathematical motivation.
  - Classical problem.

- Physical motivation.
  - Study of singularities.
  - Asymptotic properties of fields.
  - Global properties of spacetimes.

- Mathematical motivation.
  - Classical problem.
  - Interest of the mathematical tools involved.

- Any timelike curve (future and past) has an endpoint.
- Some causal structures are preserved.

- Multiple solutions:

- Any timelike curve (future and past) has an endpoint.
- Some causal structures are preserved.

- Multiple solutions:

- Any timelike curve (future and past) has an endpoint.
- Some causal structures are preserved.

- Multiple solutions:

- Any timelike curve (future and past) has an endpoint.
- Some causal structures are preserved.

- Multiple solutions:
  - A-boundary.

  - Geroch boundary.
  - Conformal boundary. (Most common in physics)
  - Causal Boundary.

- Any timelike curve (future and past) has an endpoint.
- Some causal structures are preserved.

- Multiple solutions:
  - A-boundary.

  - Geroch boundary.
  - Conformal boundary. (Most common in physics)
  - Causal Boundary.

- Any timelike curve (future and past) has an endpoint.
- Some causal structures are preserved.

- Multiple solutions:
  - A-boundary.
  - B-boundary.

  - Conformal boundary. (Most common in physics)
  - Causal Boundary.

- Any timelike curve (future and past) has an endpoint.
- Some causal structures are preserved.

- Multiple solutions:
  - A-boundary.
  - B-boundary.
  - Geroch boundary.
  - Conformal boundary. (Most common in physics)
  - Causal Boundary.

- Any timelike curve (future and past) has an endpoint.
- Some causal structures are preserved.

- Multiple solutions:
  - A-boundary.
  - B-boundary.
  - Geroch boundary.
  - Conformal boundary. (Most common in physics)
  - Causal Boundary.

- Any timelike curve (future and past) has an endpoint.
- Some causal structures are preserved.

- Multiple solutions:
  - A-boundary.
  - B-boundary.
  - Geroch boundary.
  - Conformal boundary. (Most common in physics)
  - Causal Boundary.

# **INDEX**

#### Causal Boundary

- Applicable to "strongly causal" spacetimes.
- Construction:

Applicable to "strongly causal" spacetimes.

CONFORMAL BOUNDARY

- Construction:
  - For any inextendible future or past timelike curves, attach

  - The future (past) causal boundary is the set of future (past)
  - The (total) causal boundary is defined as the union of the

- Applicable to "strongly causal" spacetimes.
- Construction:
  - For any inextendible future or past timelike curves, attach an ideal point.
  - Two inextendible future (past) timelike curves are attached the same ideal point if they have the same past (resp. future).
  - The future (past) causal boundary is the set of future (past) ideal points.
  - The (total) causal boundary is defined as the union of the future and past causal boundaries under "certain identifications".


- Applicable to "strongly causal" spacetimes.
- Construction:
  - For any inextendible future or past timelike curves, attach an ideal point.
  - Two inextendible future (past) timelike curves are attached the same ideal point if they have the same past (resp. future).
  - The future (past) causal boundary is the set of future (past) ideal points.
  - The (total) causal boundary is defined as the union of the future and past causal boundaries under "certain identifications".


- Applicable to "strongly causal" spacetimes.
- Construction:
  - For any inextendible future or past timelike curves, attach an ideal point.
  - Two inextendible future (past) timelike curves are attached the same ideal point if they have the same past (resp. future).
  - The future (past) causal boundary is the set of future (past) ideal points.
  - The (total) causal boundary is defined as the union of the

- Applicable to "strongly causal" spacetimes.
- Construction:
  - For any inextendible future or past timelike curves, attach an ideal point.
  - Two inextendible future (past) timelike curves are attached the same ideal point if they have the same past (resp. future).
  - The future (past) causal boundary is the set of future (past) ideal points.
  - The (total) causal boundary is defined as the union of the future and past causal boundaries under "certain identifications".

- Past Set:  $P \subset V$  such that  $I^{-}[P] = P$ .
- IP: Past set which cannot be expressed by the union of two
- PIP: IP,  $P \subset V$  such that  $P = I^-(p)$  for  $p \in V$ .
- TIP: IP,  $P \subset V$  such that  $P \neq I^{-}(p) \ \forall \ p \in V \iff \exists$  an
- We have the following identifications:


$$\hat{\partial}(V) \equiv TIPs, V \equiv PIPs, \hat{V} \equiv IPs$$






- Past Set:  $P \subset V$  such that  $I^{-}[P] = P$ .
- IP: Past set which cannot be expressed by the union of two proper past sets.
- PIP: IP,  $P \subset V$  such that  $P = I^-(p)$  for  $p \in V$ .
- TIP: IP,  $P \subset V$  such that  $P \neq I^{-}(p) \ \forall \ p \in V \iff \exists$  an
- We have the following identifications:


$$\hat{\partial}(V) \equiv TIPs, V \equiv PIPs, \hat{V} \equiv IPs$$






- Past Set:  $P \subset V$  such that  $I^{-}[P] = P$ .
- IP: Past set which cannot be expressed by the union of two proper past sets.
- PIP: IP,  $P \subset V$  such that  $P = I^-(p)$  for  $p \in V$ .
- TIP: IP,  $P \subset V$  such that  $P \neq I^{-}(p) \ \forall \ p \in V \iff \exists$  an
- We have the following identifications:


$$\hat{\partial}(V) \equiv TIPs, V \equiv PIPs, \hat{V} \equiv IPs$$






- Past Set:  $P \subset V$  such that  $I^{-}[P] = P$ .
- IP: Past set which cannot be expressed by the union of two proper past sets.
- PIP: IP,  $P \subset V$  such that  $P = I^-(p)$  for  $p \in V$ .
- TIP: IP,  $P \subset V$  such that  $P \neq I^{-}(p) \forall p \in V \iff \exists$  an
- We have the following identifications:


$$\hat{\partial}(V) \equiv TIPs, V \equiv PIPs, \hat{V} \equiv IPs$$






- Past Set:  $P \subset V$  such that  $I^{-}[P] = P$ .
- IP: Past set which cannot be expressed by the union of two proper past sets.
- PIP: IP,  $P \subset V$  such that  $P = I^-(p)$  for  $p \in V$ .
- TIP: IP,  $P \subset V$  such that  $P \neq I^{-}(p) \ \forall \ p \in V \iff \exists$  an inextendible timelike curve  $\gamma$ ,  $I^{-}[\gamma] = P$ .
- We have the following identifications:


$$\hat{\partial}(V) \equiv TIPs, V \equiv PIPs, \hat{V} \equiv IPs$$






- Past Set:  $P \subset V$  such that  $I^{-}[P] = P$ .
- IP: Past set which cannot be expressed by the union of two proper past sets.
- PIP: IP,  $P \subset V$  such that  $P = I^-(p)$  for  $p \in V$ .
- TIP: IP,  $P \subset V$  such that  $P \neq I^{-}(p) \ \forall \ p \in V \iff \exists$  an inextendible timelike curve  $\gamma$ ,  $I^{-}[\gamma] = P$ .
- We have the following identifications:

$$\hat{\partial}(V) \equiv TIPs, V \equiv PIPs, \hat{V} \equiv IPs$$







# Past Causal Boundary

- Future Set:  $F \subset V$  such that  $I^+[F] = F$ .
- IF: Future set which cannot be expressed by the union of two proper past sets.
- PIF: IP,  $F \subset V$  such that  $F = I^+(p)$  for  $p \in V$ .
- TIF: IP,  $F \subset V$  such that  $F \neq I^+(p)$   $\forall p \in V$ .

$$\check{\partial}(\mathit{V}) \equiv \mathit{TIFs}, \mathit{V} \equiv \mathit{PIFs}, \check{\mathit{V}} \equiv \mathit{IFs}$$





 In a first approach, we can consider that the (total) causal boundary defined by:

$$\partial(V) = \hat{\partial}(V) \cup \check{\partial}(V).$$

In our example, this definition provides:



We need to introduce some non-trivial "identifications".

 In a first approach, we can consider that the (total) causal boundary defined by:

$$\partial(V) = \hat{\partial}(V) \cup \check{\partial}(V).$$

In our example, this definition provides:



We need to introduce some non-trivial "identifications".

 We will follow an approach based on the Marolf-Ross and Flores approach.

## Definition (Szabados relation)

$$P \sim_{S} F$$
 if 
$$\begin{cases} P \text{ is a maximal IP in } \downarrow F := I^{-}[\{q \in V : q \ll p, \forall p \in F\}] \\ F \text{ is a maximal IF in } \uparrow P := I^{+}[\{p \in V : q \ll p, \forall q \in P\}] \end{cases}$$

$$\partial(V) := \{ (P, F) \in \hat{\partial}(V) \times \check{\partial}(V) : P \sim_{S} F \}$$

$$V \equiv \{(I^{-}(p), I^{+}(p)) : p \in V\}$$

 We will follow an approach based on the Marolf-Ross and Flores approach.

#### Definition (Szabados relation)

$$P \sim_S F$$
 if 
$$\begin{cases} P \text{ is a maximal IP in } \downarrow F := I^-[\{q \in V : q \ll p, \forall p \in F\}] \\ F \text{ is a maximal IF in } \uparrow P := I^+[\{p \in V : q \ll p, \forall q \in P\}] \end{cases}$$


#### Definition (Total Causal Boundary)

$$\partial(V) := \{(P, F) \in \hat{\partial}(V) \times \check{\partial}(V) : P \sim_{\mathcal{S}} F\}$$


If P(F) is not S-related with nobody, then  $(P, \emptyset) \in \partial(V)$  $((\emptyset, F) \in \partial(V)).$ 

$$V \equiv \{(I^{-}(p), I^{+}(p)) : p \in V\}$$

### (Total) Causal Boundary




•  $(P, F) \ll (P', F')$  if, and only if,  $F \cap P' \neq \emptyset$ .



## *Topology*

• Limit-Operator: $(P, F) \in L(\sigma)$ , with  $\sigma = \{(P_n, F_n)\}_n \subset V$  if:  $\begin{cases}
P \in \hat{L}(P_n) := \{P' \in \hat{V} : P' \subset LI(P_n) \text{ maximal in } LS(P_n)\} \\
F \in \check{L}(F_n) := \{F' \in \check{V} : P' \subset LI(F_n) \text{ maximal in } LS(F_n)\}
\end{cases}$ 




• The *closed sets* of  $\overline{V}$  are the subsets  $C \subset \overline{V}$  such that  $L(\sigma) \in C$  for all sequence  $\sigma \subset C$ .

#### **INDEX**

Conformal Boundary


## Asymptotically-flat spacetimes

 Generalize the structure of Minkowski spacetime seen inside the Einstein spacetime.



The concept of conformal boundary appear naturally!

 Generalize the structure of Minkowski spacetime seen inside the Einstein spacetime.



The concept of conformal boundary appear naturally!

## Conformal Boundary

- Consider a conformal embedding  $i: V \to V_0$ . (Envelopment)
- Define:
- Topology: Induced from  $V_0$ .
- Causality:  $x \ll_i y$ ,  $\exists \gamma : [a, b] \rightarrow i(V)$  future-directed

## Conformal Boundary

• Consider a conformal embedding  $i: V \to V_0$ . (Envelopment)

CONFORMAL BOUNDARY

- Define:
  - $\overline{i}(V)$  (Conformal Completion).
  - $\partial_i V = i(V) \setminus V$  (Conformal Boundary).
  - $\partial_i^* V$  set of point in  $\partial_i V$  reachable by timelike curves in V.
- Topology: Induced from  $V_0$ .
- Causality:  $x \ll_i y$ ,  $\exists \gamma : [a, b] \rightarrow i(V)$  future-directed

- Consider a conformal embedding  $i: V \to V_0$ . (Envelopment)
- Define:
  - i(V) (Conformal Completion).
  - $\partial_i V = i(V) \setminus V$  (Conformal Boundary).
  - $\partial_i^* V$  set of point in  $\partial_i V$  reachable by timelike curves in V.
- Topology: Induced from  $V_0$ .
- Causality:  $x \ll_i y$ ,  $\exists \gamma : [a, b] \rightarrow i(V)$  future-directed

- Consider a conformal embedding  $i: V \to V_0$ . (Envelopment)
- Define:
  - i(V) (Conformal Completion).
  - $\partial_i V = i(V) \setminus V$  (Conformal Boundary).
  - $\partial_i^* V$  set of point in  $\partial_i V$  reachable by timelike curves in V.
- Topology: Induced from  $V_0$ .
- Causality:  $x \ll_i y$ ,  $\exists \gamma : [a, b] \rightarrow i(V)$  future-directed

- Consider a conformal embedding i : V → V<sub>0</sub>. (Envelopment)
- Define:
  - i(V) (Conformal Completion).
  - $\partial_i V = i(V) \setminus V$  (Conformal Boundary).
  - $\partial_i^* V$  set of point in  $\partial_i V$  reachable by timelike curves in V.
- Topology: Induced from  $V_0$ .
- Causality:  $x \ll_i y$ ,  $\exists \gamma : [a, b] \rightarrow \overline{i(V)}$  future-directed timelike curve with  $\gamma((a, b)) \subset V$ ,  $\gamma(a) = x$ ,  $\gamma(b) = y$

- Consider a conformal embedding  $i: V \to V_0$ . (Envelopment)
- Define:
  - i(V) (Conformal Completion).
  - $\partial_i V = i(V) \setminus V$  (Conformal Boundary).
  - $\partial_i^* V$  set of point in  $\partial_i V$  reachable by timelike curves in V.
- Topology: Induced from  $V_0$ .
- Causality:  $x \ll_i y$ ,  $\exists \gamma : [a, b] \rightarrow i(V)$  future-directed

#### Conformal Boundary

- Consider a conformal embedding  $i: V \to V_0$ . (Envelopment)
- Define:
  - i(V) (Conformal Completion).
  - $\partial_i V = i(V) \setminus V$  (Conformal Boundary).
  - $\partial_i^* V$  set of point in  $\partial_i V$  reachable by timelike curves in V.
- Topology: Induced from  $V_0$ .
- Causality:  $x \ll_i y$ ,  $\exists \gamma : [a, b] \rightarrow \overline{i(V)}$  future-directed timelike curve with  $\gamma((a, b)) \subset V$ ,  $\gamma(a) = x$ ,  $\gamma(b) = v$ .


#### **INDEX**

Relations between conformal and causal boundary

Causal Boundary	Conformal Boundary
Conformal invariant	No conformal invariant
Uniqueness	Dependence on the conformal embedding
Systematic	No systematic
Lack of physical spacetimes with known causal boundary	Most common boundary in physics


Due to the bad behaviour of the conformal boundary, we cannot expect in general any possible relation:

As a point set



Due to the bad behaviour of the conformal boundary, we cannot expect in general any possible relation:

- As a point set
- As a topological structure



- There exists conformal embeddings where conformal and causal boundary coincide.
- How can we know when the conformal embedding is good
- Under certain technical conditions for the conformal

- There exists conformal embeddings where conformal and causal boundary coincide.
- How can we know when the conformal embedding is good enough?
- Under certain technical conditions for the conformal

- There exists conformal embeddings where conformal and causal boundary coincide.
- How can we know when the conformal embedding is good enough?
- Under certain technical conditions for the conformal boundary, it is possible to compare them.

Consider a future (resp. past) timelike curve  $\gamma: [a, b] \to i(M)$ .

#### Deformally timelike

 $\gamma$  is deformally timelike if there exist a neighbourhood  $\gamma(b) \in U = U_0 \cap i(M)$  (where  $U_0$  is an open set of  $M_0$ ) such that  $w \in U, \gamma(a) \ll w$ .

Consider a future (resp. past) timelike curve  $\gamma: [a, b] \to i(M)$ .

#### Deformally timelike


 $\gamma$  is deformally timelike if there exist a neighbourhood  $\gamma(b) \in U = U_0 \cap i(M)$  (where  $U_0$  is an open set of  $M_0$ ) such that  $w \in U, \gamma(a) \ll w$ .

#### Transitively timelike


 $\gamma$  is *transitively timelike* if there exist a neighbourhood  $\gamma(b) \in V = V_0 \cap i(M)$  such that for all  $w \in V$  with  $\gamma(b) \leq w$ , then  $\gamma(a) \ll_i w$ .

CONFORMAL BOUNDARY

#### Deformally timelike



#### Transitively timelike



#### Regular Border Curve

 $\gamma$  is a regular border curve if it is deformally timelike and transitively timelike.

#### Regular Border Curve

 $\gamma$  is a regular border curve if it is deformally timelike and transitively timelike.

#### Chronologically complete

An envelopment  $i: V \to V_0$  is chronologically complete if all the timelike curves contained in i(V) has an endpoint in  $V_0$ .

#### Regular Border Curve

 $\gamma$  is a regular border curve if it is deformally timelike and transitively timelike .

#### Chronologically complete

An envelopment  $i: V \to V_0$  is chronologically complete if all the timelike curves contained in i(V) has an endpoint in  $V_0$ .

#### Regular accesible


An envelopment  $i: V \to V_0$  is regular accesible if all TIPs and TIFs in V can be defined by regular border curves.

#### Projection

$$\pi: \partial V \rightarrow \partial_i^* V$$
  
 $(P,F) \rightarrow \pi(P,F) = X$ 

where *x* is the endpoint of a timelike curve such that:

- $I^-[\gamma] = P$  if  $\gamma$  is future-directed.
- $I^+[\gamma] = F$  if  $\gamma$  is past-directed.



#### Theorem


Consider  $i: V \rightarrow V_0$  an envelopment chronologically complete and regular accesible. Then, the following statements hold:

- π is well defined.
- The extension  $\pi: \overline{V} \to \overline{V}_i^*$  is an homeomorphism and
- is a chronological isomorphism.

## **Applications**

Thanks to this study, it is straightforward to compute the causal boundary in the following cases:

Minkowski spacetime.



## Applications


Thanks to this study, it is straightforward to compute the causal boundary in the following cases:

- Minkowski spacetime.
- Asymptotically flat spacetimes.

## **Applications**

Thanks to this study, it is straightforward to compute the causal boundary in the following cases:

- Minkowski spacetime.
- Asymptotically flat spacetimes.
- Schwarzschild spacetime...



# Thank you for your attention.