

# Height Estimates for $r$ -Mean Curvature Spacelike Hypersurfaces in Product Spaces

A.Gervasio Colares  
( Universidade Federal do Ceará, Brazil )

Joint work with H. F. de Lima  
in *General Relativity and Gravitation*, v.40 (2008), 2131-2147

V International Meeting on Lorentzian Geometry  
Martina Franca, Italy

8-11 July, 2009

- Consider an  $n$ -dimensional Riemannian manifold  $M$ , and let  $\mathbb{R}$  be the 1-dimensional Euclidean space.

# Lorentzian Products

- Consider an  $n$ -dimensional Riemannian manifold  $M$ , and let  $\mathbb{R}$  be the 1-dimensional Euclidean space.
- We will denote by  $-\mathbb{R} \times M$  the  $(n + 1)$ -dimensional product manifold  $\mathbb{R} \times M$  endowed with the Lorentzian metric

# Lorentzian Products

- Consider an  $n$ -dimensional Riemannian manifold  $M$ , and let  $\mathbb{R}$  be the 1-dimensional Euclidean space.
- We will denote by  $-\mathbb{R} \times M$  the  $(n + 1)$ -dimensional product manifold  $\mathbb{R} \times M$  endowed with the Lorentzian metric

$$\langle \cdot, \cdot \rangle = -\pi_{\mathbb{R}}^* dt^2 + \pi_M^* \langle \cdot, \cdot \rangle_M,$$

# Lorentzian Products

- Consider an  $n$ -dimensional Riemannian manifold  $M$ , and let  $\mathbb{R}$  be the 1-dimensional Euclidean space.
- We will denote by  $-\mathbb{R} \times M$  the  $(n + 1)$ -dimensional product manifold  $\mathbb{R} \times M$  endowed with the Lorentzian metric

$$\langle \cdot, \cdot \rangle = -\pi_{\mathbb{R}}^* dt^2 + \pi_M^* \langle \cdot, \cdot \rangle_M,$$

- where  $\pi_{\mathbb{R}}$  and  $\pi_M$  denote the canonical projections of  $-\mathbb{R} \times M$  onto each factor and  $\langle \cdot, \cdot \rangle_M$  stands for the Riemannian metric on  $M$ .

# Lorentzian Products

- Consider an  $n$ -dimensional Riemannian manifold  $M$ , and let  $\mathbb{R}$  be the 1-dimensional Euclidean space.
- We will denote by  $-\mathbb{R} \times M$  the  $(n + 1)$ -dimensional product manifold  $\mathbb{R} \times M$  endowed with the Lorentzian metric

$$\langle \cdot, \cdot \rangle = -\pi_{\mathbb{R}}^* dt^2 + \pi_M^* \langle \cdot, \cdot \rangle_M,$$

- where  $\pi_{\mathbb{R}}$  and  $\pi_M$  denote the canonical projections of  $-\mathbb{R} \times M$  onto each factor and  $\langle \cdot, \cdot \rangle_M$  stands for the Riemannian metric on  $M$ .
- In particular, when the Riemannian factor  $M$  is the Euclidean space  $\mathbb{R}^n$  then  $-\mathbb{R} \times \mathbb{R}^n$  is the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$ .

# Lorentzian Products

- Consider a smooth immersion

# Lorentzian Products

- Consider a smooth immersion

$$\Psi : \Sigma \rightarrow -\mathbb{R} \times M$$



- Consider a smooth immersion

$$\Psi : \Sigma \rightarrow -\mathbb{R} \times M$$

of a  $n$ -dimensional connected manifold  $\Sigma$  into  $-\mathbb{R} \times M$ , and assume that the induced metric via  $\Psi$  is a Riemannian metric on  $\Sigma$ ; that is,  $\Sigma$  is a spacelike hypersurface.

# Lorentzian Products

- Consider a smooth immersion

$$\Psi : \Sigma \rightarrow -\mathbb{R} \times M$$

of a  $n$ -dimensional connected manifold  $\Sigma$  into  $-\mathbb{R} \times M$ , and assume that the induced metric via  $\Psi$  is a Riemannian metric on  $\Sigma$ ; that is,  $\Sigma$  is a spacelike hypersurface.

- Since  $\partial_t = (\partial/\partial t)_{(t,x)}$ ,  $(t, x) \in -\mathbb{R} \times M$ , is a unitary timelike vector field globally defined on the ambient, then there exists a unique unitary timelike normal field  $N$  globally defined on  $\Sigma$  which is in the same time-orientation as  $\partial_t$ , so that

# Lorentzian Products

- Consider a smooth immersion

$$\Psi : \Sigma \rightarrow -\mathbb{R} \times M$$

of a  $n$ -dimensional connected manifold  $\Sigma$  into  $-\mathbb{R} \times M$ , and assume that the induced metric via  $\Psi$  is a Riemannian metric on  $\Sigma$ ; that is,  $\Sigma$  is a spacelike hypersurface.

- Since  $\partial_t = (\partial/\partial t)_{(t,x)}$ ,  $(t, x) \in -\mathbb{R} \times M$ , is a unitary timelike vector field globally defined on the ambient, then there exists a unique unitary timelike normal field  $N$  globally defined on  $\Sigma$  which is in the same time-orientation as  $\partial_t$ , so that

$$\langle N, \partial_t \rangle \leq -1 < 0 \text{ on } \Sigma.$$

# Lorentzian Products

- Consider a smooth immersion

$$\Psi : \Sigma \rightarrow -\mathbb{R} \times M$$

of a  $n$ -dimensional connected manifold  $\Sigma$  into  $-\mathbb{R} \times M$ , and assume that the induced metric via  $\Psi$  is a Riemannian metric on  $\Sigma$ ; that is,  $\Sigma$  is a spacelike hypersurface.

- Since  $\partial_t = (\partial/\partial t)_{(t,x)}$ ,  $(t, x) \in -\mathbb{R} \times M$ , is a unitary timelike vector field globally defined on the ambient, then there exists a unique unitary timelike normal field  $N$  globally defined on  $\Sigma$  which is in the same time-orientation as  $\partial_t$ , so that

$$\langle N, \partial_t \rangle \leq -1 < 0 \text{ on } \Sigma.$$

- We will refer to that normal field  $N$  as the future-pointing Gauss map of the hypersurface. Its opposite will be referred as the past-pointing Gauss map of  $\Sigma$ .

# Higher order mean curvatures

- Let  $A$  be the shape operator of  $\Sigma$  with respect to either the future or the past-pointing Gauss map  $N$ . It is a self-adjoint linear operator on each tangent space  $T_p\Sigma$  and its eigenvalues  $k_1(p), \dots, k_n(p)$  are the principal curvatures of the hypersurface. Associated to the shape operator there are  $n$  algebraic invariants given by

# Higher order mean curvatures

- Let  $A$  be the shape operator of  $\Sigma$  with respect to either the future or the past-pointing Gauss map  $N$ . It is a self-adjoint linear operator on each tangent space  $T_p\Sigma$  and its eigenvalues  $k_1(p), \dots, k_n(p)$  are the principal curvatures of the hypersurface. Associated to the shape operator there are  $n$  algebraic invariants given by

$$S_k(p) = \sigma_k(k_1(p), \dots, k_n(p)), \quad 1 \leq k \leq n,$$

# Higher order mean curvatures

- Let  $A$  be the shape operator of  $\Sigma$  with respect to either the future or the past-pointing Gauss map  $N$ . It is a self-adjoint linear operator on each tangent space  $T_p\Sigma$  and its eigenvalues  $k_1(p), \dots, k_n(p)$  are the principal curvatures of the hypersurface. Associated to the shape operator there are  $n$  algebraic invariants given by

$$S_k(p) = \sigma_k(k_1(p), \dots, k_n(p)), \quad 1 \leq k \leq n,$$

where  $\sigma_k : \mathbb{R}^n \rightarrow \mathbb{R}$  is the elementary symmetric function in  $\mathbb{R}$  given by

# Higher order mean curvatures

- Let  $A$  be the shape operator of  $\Sigma$  with respect to either the future or the past-pointing Gauss map  $N$ . It is a self-adjoint linear operator on each tangent space  $T_p\Sigma$  and its eigenvalues  $k_1(p), \dots, k_n(p)$  are the principal curvatures of the hypersurface. Associated to the shape operator there are  $n$  algebraic invariants given by

$$S_k(p) = \sigma_k(k_1(p), \dots, k_n(p)), \quad 1 \leq k \leq n,$$

where  $\sigma_k : \mathbb{R}^n \rightarrow \mathbb{R}$  is the elementary symmetric function in  $\mathbb{R}$  given by

$$\sigma_k(x_1 \cdots x_n) = \sum_{i_1 < \dots < i_k} x_{i_1} \cdots x_{i_k}$$



# Higher order mean curvatures

- The  $k$ th-mean curvature  $H_k$  of the hypersurface is then defined by

# Higher order mean curvatures

- The  $k$ th-mean curvature  $H_k$  of the hypersurface is then defined by

$$\binom{n}{k} H_k = (-1)^k S_k = \sigma_k(-k_1, \dots, -k_n),$$

# Higher order mean curvatures

- The  $k$ th-mean curvature  $H_k$  of the hypersurface is then defined by

$$\binom{n}{k} H_k = (-1)^k S_k = \sigma_k(-k_1, \dots, -k_n),$$

for every  $0 \leq k \leq n$ .

# Higher order mean curvatures

- The  $k$ th-mean curvature  $H_k$  of the hypersurface is then defined by

$$\binom{n}{k} H_k = (-1)^k S_k = \sigma_k(-k_1, \dots, -k_n),$$

for every  $0 \leq k \leq n$ .

- When  $k = 1$ ,

# Higher order mean curvatures

- The  $k$ th-mean curvature  $H_k$  of the hypersurface is then defined by

$$\binom{n}{k} H_k = (-1)^k S_k = \sigma_k(-k_1, \dots, -k_n),$$

for every  $0 \leq k \leq n$ .

- When  $k = 1$ ,

$$H_1 = -1/n \sum k_i = -1/n \operatorname{trace}(A) = H,$$

# Higher order mean curvatures

- The  $k$ th-mean curvature  $H_k$  of the hypersurface is then defined by

$$\binom{n}{k} H_k = (-1)^k S_k = \sigma_k(-k_1, \dots, -k_n),$$

for every  $0 \leq k \leq n$ .

- When  $k = 1$ ,

$$H_1 = -1/n \sum k_i = -1/n \operatorname{trace}(A) = H,$$

the mean curvature of  $\Sigma$ . By the choice of the sign  $(-1)^k$ , the mean curvature vector  $\mathbf{H}$  is given by  $\mathbf{H} = HN$ . Therefore,  $H(p) > 0$  at a point  $p \in \Sigma$  if and only if  $\mathbf{H}(p)$  is in the same time-orientation as  $N$ .

# Ellipticity and positive definiteness

- The Newton transformations  $P_k$  are defined inductively from  $A$  by  $P_0 = I$  and

# Ellipticity and positive definiteness

- The Newton transformations  $P_k$  are defined inductively from  $A$  by  $P_0 = I$  and

$$P_k = \binom{n}{k} H_k I + A \cdot P_{k-1}, \quad k = 1, \dots, n,$$



# Ellipticity and positive definiteness

- The Newton transformations  $P_k$  are defined inductively from  $A$  by  $P_0 = I$  and

$$P_k = \binom{n}{k} H_k I + A \cdot P_{k-1}, \quad k = 1, \dots, n,$$

and  $I$  the identity.

# Ellipticity and positive definiteness

- The Newton transformations  $P_k$  are defined inductively from  $A$  by  $P_0 = I$  and

$$P_k = \binom{n}{k} H_k I + A \cdot P_{k-1}, \quad k = 1, \dots, n,$$

and  $I$  the identity.

- Then  $P_k(p)$  is a self-adjoint linear operator on the tangent space  $T_p \Sigma$  which commutes with  $A(p)$  and  $A(p)$  and  $P_k(p)$  can be simultaneously diagonalized.

# Ellipticity and positive definiteness

- The Newton transformations  $P_k$  are defined inductively from  $A$  by  $P_0 = I$  and

$$P_k = \binom{n}{k} H_k I + A \cdot P_{k-1}, \quad k = 1, \dots, n,$$

and  $I$  the identity.

- Then  $P_k(p)$  is a self-adjoint linear operator on the tangent space  $T_p \Sigma$  which commutes with  $A(p)$  and  $A(p)$  and  $P_k(p)$  can be simultaneously diagonalized.
- Let  $\nabla$  be the Levi-Civita connection of  $\Sigma$ . To each  $P_k$ , we associate the second order linear differential operator

# Ellipticity and positive definiteness

- The Newton transformations  $P_k$  are defined inductively from  $A$  by  $P_0 = I$  and

$$P_k = \binom{n}{k} H_k I + A \cdot P_{k-1}, \quad k = 1, \dots, n,$$

and  $I$  the identity.

- Then  $P_k(p)$  is a self-adjoint linear operator on the tangent space  $T_p \Sigma$  which commutes with  $A(p)$  and  $A(p)$  and  $P_k(p)$  can be simultaneously diagonalized.
- Let  $\nabla$  be the Levi-Civita connection of  $\Sigma$ . To each  $P_k$ , we associate the second order linear differential operator

$$L_k(f) = \text{trace}(P_k \circ \nabla^2 f).$$

# Ellipticity and positive definiteness

- There  $\nabla^2 f$  denotes the self-adjoint linear operator metrically equivalent to the hessian of  $f$  : for all tangent  $X, Y$ ,

# Ellipticity and positive definiteness

- There  $\nabla^2 f$  denotes the self-adjoint linear operator metrically equivalent to the hessian of  $f$  : for all tangent  $X, Y$ ,

$$\langle \nabla^2 f(X), Y \rangle = \langle \nabla_X(\nabla f), Y \rangle .$$

# Ellipticity and positive definiteness

- There  $\nabla^2 f$  denotes the self-adjoint linear operator metrically equivalent to the hessian of  $f$  : for all tangent  $X, Y$ ,

$$\langle \nabla^2 f(X), Y \rangle = \langle \nabla_X(\nabla f), Y \rangle .$$

- Observe that  $P_k$  commutes with  $\nabla^2 f$  and

# Ellipticity and positive definiteness

- There  $\nabla^2 f$  denotes the self-adjoint linear operator metrically equivalent to the hessian of  $f$  : for all tangent  $X, Y$ ,

$$\langle \nabla^2 f(X), Y \rangle = \langle \nabla_X(\nabla f), Y \rangle .$$

- Observe that  $P_k$  commutes with  $\nabla^2 f$  and

$$\operatorname{div}(P_k(\nabla f)) = \langle \operatorname{div} P_k, \nabla f \rangle + L_k(f),$$



# Ellipticity and positive definiteness

- There  $\nabla^2 f$  denotes the self-adjoint linear operator metrically equivalent to the hessian of  $f$  : for all tangent  $X, Y$ ,

$$\langle \nabla^2 f(X), Y \rangle = \langle \nabla_X(\nabla f), Y \rangle .$$

- Observe that  $P_k$  commutes with  $\nabla^2 f$  and

$$\operatorname{div}(P_k(\nabla f)) = \langle \operatorname{div} P_k, \nabla f \rangle + L_k(f),$$

where, for an orthonormal basis  $E_1, \dots, E_n$ ,

# Ellipticity and positive definiteness

- There  $\nabla^2 f$  denotes the self-adjoint linear operator metrically equivalent to the hessian of  $f$  : for all tangent  $X, Y$ ,

$$\langle \nabla^2 f(X), Y \rangle = \langle \nabla_X(\nabla f), Y \rangle .$$

- Observe that  $P_k$  commutes with  $\nabla^2 f$  and

$$\operatorname{div}(P_k(\nabla f)) = \langle \operatorname{div} P_k, \nabla f \rangle + L_k(f),$$

where, for an orthonormal basis  $E_1, \dots, E_n$ ,

$$\operatorname{div} P_k := \operatorname{trace}(\nabla P_k) = \sum_{i=1}^n (\nabla_{E_i} P_k) E_i.$$

# Ellipticity and positive definiteness

- There  $\nabla^2 f$  denotes the self-adjoint linear operator metrically equivalent to the hessian of  $f$  : for all tangent  $X, Y$ ,

$$\langle \nabla^2 f(X), Y \rangle = \langle \nabla_X(\nabla f), Y \rangle .$$

- Observe that  $P_k$  commutes with  $\nabla^2 f$  and

$$\operatorname{div}(P_k(\nabla f)) = \langle \operatorname{div} P_k, \nabla f \rangle + L_k(f),$$

where, for an orthonormal basis  $E_1, \dots, E_n$ ,

$$\operatorname{div} P_k := \operatorname{trace}(\nabla P_k) = \sum_{i=1}^n (\nabla_{E_i} P_k) E_i.$$

- Then  $L_k(f) = \operatorname{div}(P_k(\nabla f)) \Leftrightarrow \operatorname{div}(P_k) = 0$ .

# Ellipticity and positive definiteness

- By an elliptic point in a spacelike hypersurface  $\Sigma$  we mean a point of  $\Sigma$  where all the principal curvatures are negative, with respect to an appropriate choice of the Gauss map  $N$ .

# Ellipticity and positive definiteness

- By an elliptic point in a spacelike hypersurface  $\Sigma$  we mean a point of  $\Sigma$  where all the principal curvatures are negative, with respect to an appropriate choice of the Gauss map  $N$ .

## Lemma 1

Let  $\Sigma$  be a spacelike hypersurface immersed into a Lorentzian product space. If there exists an elliptic point of  $\Sigma$ , with respect to an appropriate choice of the Gauss map  $N$ , and  $H_{k+1} > 0$  on  $\Sigma$ , for  $1 \leq k \leq n-1$ , then for all  $1 \leq j \leq k$  the operator  $L_j$  is elliptic or, equivalently,  $P_j$  is positive definite ( for that appropriate choice of the Gauss map if  $j$  is odd )

# Ellipticity and positive definiteness

- By an elliptic point in a spacelike hypersurface  $\Sigma$  we mean a point of  $\Sigma$  where all the principal curvatures are negative, with respect to an appropriate choice of the Gauss map  $N$ .

## Lemma 1

Let  $\Sigma$  be a spacelike hypersurface immersed into a Lorentzian product space. If there exists an elliptic point of  $\Sigma$ , with respect to an appropriate choice of the Gauss map  $N$ , and  $H_{k+1} > 0$  on  $\Sigma$ , for  $1 \leq k \leq n-1$ , then for all  $1 \leq j \leq k$  the operator  $L_j$  is elliptic or, equivalently,  $P_j$  is positive definite ( for that appropriate choice of the Gauss map if  $j$  is odd )

- The proof follows from that of Cheng-Rosenberg in An. Ac. Br. Ci., 2005 or Barbosa-Colares in Ann. Gl. Ann. and Geom., 1997, considering that in our case, and by our sign convention in the definition of the  $j$ -th mean curvatures, we have

# Ellipticity and positive definiteness

- By an elliptic point in a spacelike hypersurface  $\Sigma$  we mean a point of  $\Sigma$  where all the principal curvatures are negative, with respect to an appropriate choice of the Gauss map  $N$ .

## Lemma 1

Let  $\Sigma$  be a spacelike hypersurface immersed into a Lorentzian product space. If there exists an elliptic point of  $\Sigma$ , with respect to an appropriate choice of the Gauss map  $N$ , and  $H_{k+1} > 0$  on  $\Sigma$ , for  $1 \leq k \leq n-1$ , then for all  $1 \leq j \leq k$  the operator  $L_j$  is elliptic or, equivalently,  $P_j$  is positive definite ( for that appropriate choice of the Gauss map if  $j$  is odd )

- The proof follows from that of Cheng-Rosenberg in An. Ac. Br. Ci., 2005 or Barbosa-Colares in Ann. Gl. Ann. and Geom., 1997, considering that in our case, and by our sign convention in the definition of the  $j$ -th mean curvatures, we have

$$\binom{n}{k} H_j = \sigma_j(-k_1, \dots, -k_n) = (-1)^j S_j.$$

# Height and support functions

- In this setting, we consider two particular functions naturally attached to a spacelike hypersurface  $\Sigma$  immersed into a Lorentzian product space  $-\mathbb{R} \times \mathbb{R}^n$ : the vertical height function  $h = (\pi_{\mathbb{R}})|_{\Sigma}$  and the support function  $\eta = \langle N, \partial_t \rangle$ , where  $N$  denotes the Gauss map of  $\Sigma^n$  and  $\partial_t$  is the coordinate vector field induced on  $-\mathbb{R} \times M^n$ .
- The following lemma corresponds to the analytical framework that we will use to obtain our main result.



# Height and support functions

## Lemma 2

Let  $\Sigma^n$  be an immersed spacelike hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$ , with Gauss map  $N$ . For every  $r = 0, \dots, n - 1$  we have:

# Height and support functions

## Lemma 2

Let  $\Sigma^n$  be an immersed spacelike hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$ , with Gauss map  $N$ . For every  $r = 0, \dots, n-1$  we have:

(a)  $L_r h = -(r+1) \binom{n}{r+1} H_{r+1} \eta;$

# Height and support functions

## Lemma 2

Let  $\Sigma^n$  be an immersed spacelike hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$ , with Gauss map  $N$ . For every  $r = 0, \dots, n-1$  we have:

- (a)  $L_r h = -(r+1) \binom{n}{r+1} H_{r+1} \eta$ ;
- (b)  $\Delta \eta = n \langle \nabla H, \partial_t \rangle + (|A|^2 + \text{Ric}_M((\pi_M)_* N, (\pi_M)_* N)) \eta$ , where  $\text{Ric}_M$  denotes the Ricci tensor of  $M^n$ .

Moreover, if  $M^n$  has constant sectional curvature  $\kappa_M$ , then

# Height and support functions

## Lemma 2

Let  $\Sigma^n$  be an immersed spacelike hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$ , with Gauss map  $N$ . For every  $r = 0, \dots, n-1$  we have:

- (a)  $L_r h = -(r+1) \binom{n}{r+1} H_{r+1} \eta$ ;
- (b)  $\Delta \eta = n \langle \nabla H, \partial_t \rangle + (|A|^2 + \text{Ric}_M((\pi_M)_* N, (\pi_M)_* N)) \eta$ , where  $\text{Ric}_M$  denotes the Ricci tensor of  $M^n$ .

Moreover, if  $M^n$  has constant sectional curvature  $\kappa_M$ , then

$$L_r \eta = \binom{n}{r+1} \langle \nabla H_{r+1}, \partial_t \rangle + \text{tr}(A^2 \circ P_r) \eta$$

# Height and support functions

## Lemma 2

Let  $\Sigma^n$  be an immersed spacelike hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$ , with Gauss map  $N$ . For every  $r = 0, \dots, n-1$  we have:

- (a)  $L_r h = -(r+1) \binom{n}{r+1} H_{r+1} \eta$ ;
- (b)  $\Delta \eta = n \langle \nabla H, \partial_t \rangle + (|A|^2 + \text{Ric}_M((\pi_M)_* N, (\pi_M)_* N)) \eta$ , where  $\text{Ric}_M$  denotes the Ricci tensor of  $M^n$ .

Moreover, if  $M^n$  has constant sectional curvature  $\kappa_M$ , then

$$\begin{aligned} L_r \eta &= \binom{n}{r+1} \langle \nabla H_{r+1}, \partial_t \rangle + \text{tr}(A^2 \circ P_r) \eta \\ &= +\kappa_M \left( (r+1) \binom{n}{r+1} H_r |\nabla h|^2 - \langle P_r \nabla h, \nabla h \rangle \right) \eta. \end{aligned}$$

# Height and support functions

## Remark

The formulae collected in the above lemma are the Lorentzian versions of the ones obtained by X. Cheng and H. Rosenberg. We also note that L.J. Alías jointly with A. G. Colares obtained a generalization of these formulae in the context of the Generalized Robertson- Walker spacetimes. Moreover, A.L. Albuje and L.J. Alías obtained the corresponding formulae for the Laplacian of the height and support functions of a space-like surface immersed in a 3-dimensional Lorentzian product space.

Now, we are in the position to state and prove our main result.

## Theorem

Let  $\Sigma^n$  be a compact immersed spacelike hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$  whose Riemannian fiber  $M^n$  has nonnegative constant sectional curvature  $\kappa_M$ . Suppose that  $\Sigma^n$  has positive constant  $r$ -mean curvature  $H_r$ , for some  $1 \leq r \leq n$ , and that its boundary  $\partial\Sigma^n$  is contained in the slice  $\{0\} \times M^n$ . Then, the vertical height of  $\Sigma^n$  satisfies the inequality

## Theorem

Let  $\Sigma^n$  be a compact immersed spacelike hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$  whose Riemannian fiber  $M^n$  has nonnegative constant sectional curvature  $\kappa_M$ . Suppose that  $\Sigma^n$  has positive constant  $r$ -mean curvature  $H_r$ , for some  $1 \leq r \leq n$ , and that its boundary  $\partial\Sigma^n$  is contained in the slice  $\{0\} \times M^n$ . Then, the vertical height of  $\Sigma^n$  satisfies the inequality

$$|h| \leq \frac{C - 1}{H_r^{1/r}}$$



## Theorem

Let  $\Sigma^n$  be a compact immersed spacelike hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$  whose Riemannian fiber  $M^n$  has nonnegative constant sectional curvature  $\kappa_M$ . Suppose that  $\Sigma^n$  has positive constant  $r$ -mean curvature  $H_r$ , for some  $1 \leq r \leq n$ , and that its boundary  $\partial\Sigma^n$  is contained in the slice  $\{0\} \times M^n$ . Then, the vertical height of  $\Sigma^n$  satisfies the inequality

$$|h| \leq \frac{C - 1}{H_r^{1/r}}$$

where  $C = \max_{\partial\Sigma} |\eta|$ . Moreover, in the case  $r = 1$  one can replace the condition on the sectional curvature of  $M^n$  by that of the Ricci curvature of  $M^n$  being nonnegative.

# Main theorem and proof

Proof.

# Main theorem and proof

## Proof.

Suppose, for example, that  $N$  is in the same time-orientation of  $\partial_t$  (i.e.,  $\langle N, \partial_t \rangle \leq -1$ ). At a lowest point, all the principal curvatures have the same sign. Since we assume that  $H_r > 0$ , we know that at this point all the principal curvatures are negative and hence we can apply Lemma 1 to obtain that  $L_{r-1}$  is elliptic and  $H_j$  are positive,  $1 \leq j \leq r-1$ .

# Main theorem and proof

## Proof.

Suppose, for example, that  $N$  is in the same time-orientation of  $\partial_t$  (i.e.,  $\langle N, \partial_t \rangle \leq -1$ ). At a lowest point, all the principal curvatures have the same sign. Since we assume that  $H_r > 0$ , we know that at this point all the principal curvatures are negative and hence we can apply Lemma 1 to obtain that  $L_{r-1}$  is elliptic and  $H_j$  are positive,  $1 \leq j \leq r-1$ .

Now, we define on  $\Sigma$  the function

# Main theorem and proof

## Proof.

Suppose, for example, that  $N$  is in the same time-orientation of  $\partial_t$  (i.e.,  $\langle N, \partial_t \rangle \leq -1$ ). At a lowest point, all the principal curvatures have the same sign. Since we assume that  $H_r > 0$ , we know that at this point all the principal curvatures are negative and hence we can apply Lemma 1 to obtain that  $L_{r-1}$  is elliptic and  $H_j$  are positive,  $1 \leq j \leq r-1$ .

Now, we define on  $\Sigma$  the function

$$\varphi = ch - \eta,$$

# Main theorem and proof

## Proof.

Suppose, for example, that  $N$  is in the same time-orientation of  $\partial_t$  (i.e.,  $\langle N, \partial_t \rangle \leq -1$ ). At a lowest point, all the principal curvatures have the same sign. Since we assume that  $H_r > 0$ , we know that at this point all the principal curvatures are negative and hence we can apply Lemma 1 to obtain that  $L_{r-1}$  is elliptic and  $H_j$  are positive,  $1 \leq j \leq r-1$ .

Now, we define on  $\Sigma$  the function

$$\varphi = ch - \eta,$$

where  $c$  is a negative constant. We have that  $\varphi|_{\partial\Sigma} \leq C$ , where  $C = \max_{\partial\Sigma} |\eta|$ .

# Main theorem and proof

## Proof ( cont. )

On the other hand, since  $P_{r-1}$  is positive definite, we have that

# Main theorem and proof

## Proof ( cont. )

On the other hand, since  $P_{r-1}$  is positive definite, we have that

$$\langle P_{r-1} \nabla h, \nabla h \rangle \leq \text{tr}(P_{r-1}) |\nabla h|^2 = r \binom{n}{r} H_{r-1} |\nabla h|^2.$$



# Main theorem and proof

## Proof ( cont. )

On the other hand, since  $P_{r-1}$  is positive definite, we have that

$$\langle P_{r-1} \nabla h, \nabla h \rangle \leq \text{tr}(P_{r-1}) |\nabla h|^2 = r \binom{n}{r} H_{r-1} |\nabla h|^2.$$

Thus, from Lemma 2 and using the assumption that the Riemannian fiber  $M^n$  has nonnegative constant sectional curvature  $\kappa_M$ , we obtain that

# Main theorem and proof

## Proof ( cont. )

On the other hand, since  $P_{r-1}$  is positive definite, we have that

$$\langle P_{r-1} \nabla h, \nabla h \rangle \leq \text{tr}(P_{r-1}) |\nabla h|^2 = r \binom{n}{r} H_{r-1} |\nabla h|^2.$$

Thus, from Lemma 2 and using the assumption that the Riemannian fiber  $M^n$  has nonnegative constant sectional curvature  $\kappa_M$ , we obtain that

$$L_{r-1} \varphi = -\langle N, \partial_t \rangle \left( r \binom{n}{r} c H_r + \text{tr}(A^2 \circ P_{r-1}) \right)$$

# Main theorem and proof

## Proof ( cont. )

On the other hand, since  $P_{r-1}$  is positive definite, we have that

$$\langle P_{r-1} \nabla h, \nabla h \rangle \leq \text{tr}(P_{r-1}) |\nabla h|^2 = r \binom{n}{r} H_{r-1} |\nabla h|^2.$$

Thus, from Lemma 2 and using the assumption that the Riemannian fiber  $M^n$  has nonnegative constant sectional curvature  $\kappa_M$ , we obtain that

$$\begin{aligned} L_{r-1} \varphi = & -\langle N, \partial_t \rangle \left( r \binom{n}{r} c H_r + \text{tr}(A^2 \circ P_{r-1}) \right) \\ & - \kappa_M \left( r \binom{n}{r} H_{r-1} |\nabla h|^2 - \langle P_{r-1} \nabla h, \nabla h \rangle \right) \eta \end{aligned}$$

# Main theorem and proof

## Proof (cont.)

On the other hand, since  $P_{r-1}$  is positive definite, we have that

$$\langle P_{r-1} \nabla h, \nabla h \rangle \leq \text{tr}(P_{r-1}) |\nabla h|^2 = r \binom{n}{r} H_{r-1} |\nabla h|^2.$$

Thus, from Lemma 2 and using the assumption that the Riemannian fiber  $M^n$  has nonnegative constant sectional curvature  $\kappa_M$ , we obtain that

$$\begin{aligned} L_{r-1} \varphi &= -\langle N, \partial_t \rangle \left( r \binom{n}{r} c H_r + \text{tr}(A^2 \circ P_{r-1}) \right) \\ &\quad - \kappa_M \left( r \binom{n}{r} H_{r-1} |\nabla h|^2 - \langle P_{r-1} \nabla h, \nabla h \rangle \right) \eta \\ &\geq -\langle N, \partial_t \rangle \left( r \binom{n}{r} c H_r + \text{tr}(A^2 \circ P_{r-1}) \right). \end{aligned}$$

# Main theorem and proof

## Proof ( cont. )

Furthermore, we have in the case  $r = 1$  (by the Cauchy-Schwarz inequality)

# Main theorem and proof

## Proof ( cont. )

Furthermore, we have in the case  $r = 1$  (by the Cauchy-Schwarz inequality)

$$H H_r - H_{r+1} = H^2 - H_2 \geq 0.$$

# Main theorem and proof

## Proof ( cont. )

Furthermore, we have in the case  $r = 1$  (by the Cauchy-Schwarz inequality)

$$H H_r - H_{r+1} = H^2 - H_2 \geq 0.$$

In the case  $r > 1$ , we know from Garding inequalities that

# Main theorem and proof

## Proof ( cont. )

Furthermore, we have in the case  $r = 1$  (by the Cauchy-Schwarz inequality)

$$H H_r - H_{r+1} = H^2 - H_2 \geq 0.$$

In the case  $r > 1$ , we know from Garding inequalities that

$$H_{r-1} \geq H_r^{(r-1)/r} > 0,$$



# Main theorem and proof

## Proof ( cont. )

Furthermore, we have in the case  $r = 1$  (by the Cauchy-Schwarz inequality)

$$H H_r - H_{r+1} = H^2 - H_2 \geq 0.$$

In the case  $r > 1$ , we know from Garding inequalities that

$$H_{r-1} \geq H_r^{(r-1)/r} > 0,$$

and also that

# Main theorem and proof

## Proof ( cont. )

Furthermore, we have in the case  $r = 1$  (by the Cauchy-Schwarz inequality)

$$H H_r - H_{r+1} = H^2 - H_2 \geq 0.$$

In the case  $r > 1$ , we know from Garding inequalities that

$$H_{r-1} \geq H_r^{(r-1)/r} > 0,$$

and also that

$$H \geq H_{r-1}^{1/(r-1)}.$$

# Main theorem and proof

## Proof ( cont. )

Furthermore, we have in the case  $r = 1$  (by the Cauchy-Schwarz inequality)

$$H H_r - H_{r+1} = H^2 - H_2 \geq 0.$$

In the case  $r > 1$ , we know from Garding inequalities that

$$H_{r-1} \geq H_r^{(r-1)/r} > 0,$$

and also that

$$H \geq H_{r-1}^{1/(r-1)}.$$

Moreover, we have also that  $H_r^2 - H_{r-1}H_{r+1} \geq 0$ . Thus,

# Main theorem and proof

## Proof ( cont. )

Furthermore, we have in the case  $r = 1$  (by the Cauchy-Schwarz inequality)

$$H H_r - H_{r+1} = H^2 - H_2 \geq 0.$$

In the case  $r > 1$ , we know from Garding inequalities that

$$H_{r-1} \geq H_r^{(r-1)/r} > 0,$$

and also that

$$H \geq H_{r-1}^{1/(r-1)}.$$

Moreover, we have also that  $H_r^2 - H_{r-1}H_{r+1} \geq 0$ . Thus,

$$H_{r+1} \leq \frac{H_r^2}{H_{r-1}}.$$

# Main theorem and proof

## Proof ( cont. )

Then, from these above inequalities, we obtain that

# Main theorem and proof

## Proof ( cont. )

Then, from these above inequalities, we obtain that

$$H H_r - H_{r+1} \geq \frac{H_r}{H_{r-1}} (H H_{r-1} - H_r) \geq \frac{H_r}{H_{r-1}} (H H_{r-1} - H_{r-1}^{r/(r-1)})$$

# Main theorem and proof

## Proof ( cont. )

Then, from these above inequalities, we obtain that

$$\begin{aligned} H H_r - H_{r+1} &\geq \frac{H_r}{H_{r-1}} (H H_{r-1} - H_r) \geq \frac{H_r}{H_{r-1}} (H H_{r-1} - H_{r-1}^{r/(r-1)}) \\ &= H_r (H - H_{r-1}^{1/(r-1)}) \geq 0. \end{aligned}$$

# Main theorem and proof

## Proof ( cont. )

Then, from these above inequalities, we obtain that

$$\begin{aligned} H H_r - H_{r+1} &\geq \frac{H_r}{H_{r-1}} (H H_{r-1} - H_r) \geq \frac{H_r}{H_{r-1}} (H H_{r-1} - H_{r-1}^{r/(r-1)}) \\ &= H_r (H - H_{r-1}^{1/(r-1)}) \geq 0. \end{aligned}$$

Therefore,



# Main theorem and proof

## Proof ( cont. )

Then, from these above inequalities, we obtain that

$$\begin{aligned} H H_r - H_{r+1} &\geq \frac{H_r}{H_{r-1}} (H H_{r-1} - H_r) \geq \frac{H_r}{H_{r-1}} (H H_{r-1} - H_{r-1}^{r/(r-1)}) \\ &= H_r (H - H_{r-1}^{1/(r-1)}) \geq 0. \end{aligned}$$

Therefore,

$$\text{tr}(A^2 \circ P_{r-1}) = \binom{n}{r} (n H H_r - (n-r) H_{r+1}) \geq r \binom{n}{r} H_r^{(r+1)/r}.$$

# Main theorem and proof

## Proof ( cont. )

Consequently, by taking

# Main theorem and proof

## Proof ( cont. )

Consequently, by taking

$$c = -H_r^{1/r}$$

# Main theorem and proof

## Proof ( cont. )

Consequently, by taking

$$c = -H_r^{1/r}$$

in the definition of the function  $\varphi$ , we get that  $L_{r-1}\varphi \geq 0$  on  $\Sigma$ . Then, we conclude from the maximum principle that  $\varphi \leq C$  on  $\Sigma$ . Therefore,

# Main theorem and proof

## Proof ( cont. )

Consequently, by taking

$$c = -H_r^{1/r}$$

in the definition of the function  $\varphi$ , we get that  $L_{r-1}\varphi \geq 0$  on  $\Sigma$ . Then, we conclude from the maximum principle that  $\varphi \leq C$  on  $\Sigma$ . Therefore,

$$0 \geq h \geq \frac{1 - C}{H_r^{1/r}}.$$

# Main theorem and proof

## Proof ( cont. )

Consequently, by taking

$$c = -H_r^{1/r}$$

in the definition of the function  $\varphi$ , we get that  $L_{r-1}\varphi \geq 0$  on  $\Sigma$ . Then, we conclude from the maximum principle that  $\varphi \leq C$  on  $\Sigma$ . Therefore,

$$0 \geq h \geq \frac{1 - C}{H_r^{1/r}}.$$

In the case of  $N$  is in the opposite time-orientation of  $\partial_t$  (i.e.,  $\langle N, \partial_t \rangle \geq 1$ ), the proof carries in a similar way and we conclude that

# Main theorem and proof

## Proof ( cont. )

Consequently, by taking

$$c = -H_r^{1/r}$$

in the definition of the function  $\varphi$ , we get that  $L_{r-1}\varphi \geq 0$  on  $\Sigma$ . Then, we conclude from the maximum principle that  $\varphi \leq C$  on  $\Sigma$ . Therefore,

$$0 \geq h \geq \frac{1 - C}{H_r^{1/r}}.$$

In the case of  $N$  is in the opposite time-orientation of  $\partial_t$  (i.e.,  $\langle N, \partial_t \rangle \geq 1$ ), the proof carries in a similar way and we conclude that

$$0 \leq h \leq \frac{C - 1}{H_r^{1/r}}.$$

# Main theorem and proof

Proof ( cont. )

Finally, since



# Main theorem and proof

## Proof ( cont. )

Finally, since

$$\Delta\eta = \eta\langle\Delta H, \partial_t\rangle + (|A|^2 + \text{Ric}_M((\pi_M)*N, (\pi_M)*N))\eta,$$

# Main theorem and proof

## Proof ( cont. )

Finally, since

$$\Delta\eta = \eta\langle\Delta H, \partial_t\rangle + (|A|^2 + \text{Ric}_M((\pi_M) * N, (\pi_M) * N))\eta,$$

we note that in the case  $r = 1$  one can replace the condition on the sectional curvature of  $M^n$  by that of the Ricci curvature of  $M^n$  being nonnegative.

# Sharpness of the estimate

## Remark

By considering the hyperbolic caps of the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$ , we show that our estimate is sharp.

# Sharpness of the estimate

## Remark

By considering the hyperbolic caps of the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$ , we show that our estimate is sharp.

Fixed a positive constant  $\lambda$ , we easily verify that the hyperbolic cap

# Sharpness of the estimate

## Remark

By considering the hyperbolic caps of the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$ , we show that our estimate is sharp.

Fixed a positive constant  $\lambda$ , we easily verify that the hyperbolic cap

$$\Sigma_{\lambda}^n = \left\{ x \in \mathbb{L}^{n+1}; \langle x, x \rangle = -\lambda^2, \lambda \leq x_1 \leq \sqrt{1 + \lambda^2} \right\}$$

# Sharpness of the estimate

## Remark

By considering the hyperbolic caps of the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$ , we show that our estimate is sharp.

Fixed a positive constant  $\lambda$ , we easily verify that the hyperbolic cap

$$\Sigma_{\lambda}^n = \left\{ x \in \mathbb{L}^{n+1}; \langle x, x \rangle = -\lambda^2, \lambda \leq x_1 \leq \sqrt{1 + \lambda^2} \right\}$$

is an example of spacelike hypersurface of the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$  which has positive constant  $r$ -mean curvature

# Sharpness of the estimate

## Remark

By considering the hyperbolic caps of the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$ , we show that our estimate is sharp.

Fixed a positive constant  $\lambda$ , we easily verify that the hyperbolic cap

$$\Sigma_{\lambda}^n = \left\{ x \in \mathbb{L}^{n+1}; \langle x, x \rangle = -\lambda^2, \lambda \leq x_1 \leq \sqrt{1 + \lambda^2} \right\}$$

is an example of spacelike hypersurface of the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$  which has positive constant  $r$ -mean curvature

$$H_r = \frac{1}{\lambda^r},$$

# Sharpness of the estimate

## Remark

By considering the hyperbolic caps of the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$ , we show that our estimate is sharp.

Fixed a positive constant  $\lambda$ , we easily verify that the hyperbolic cap

$$\Sigma_\lambda^n = \left\{ x \in \mathbb{L}^{n+1}; \langle x, x \rangle = -\lambda^2, \lambda \leq x_1 \leq \sqrt{1 + \lambda^2} \right\}$$

is an example of spacelike hypersurface of the Lorentz-Minkowski space  $\mathbb{L}^{n+1}$  which has positive constant  $r$ -mean curvature

$$H_r = \frac{1}{\lambda^r},$$

for each  $1 \leq r \leq n$  (if we choose the Gauss map  $N$  in the same time-orientation of  $e_1$ , for the case  $r$  odd).



# Sharpness of the estimate

## Remark (cont.)

We also easily verify that the hyperbolic image of  $\Sigma_\lambda^n$  is contained in the geodesic ball of center  $e_1 \in \mathbb{H}^n$  and radius

# Sharpness of the estimate

## Remark (cont.)

We also easily verify that the hyperbolic image of  $\Sigma_\lambda^n$  is contained in the geodesic ball of center  $e_1 \in \mathbb{H}^n$  and radius

$$\rho = \cosh^{-1} \sqrt{1 + \frac{1}{\lambda^2}}.$$

# Sharpness of the estimate

## Remark (cont.)

We also easily verify that the hyperbolic image of  $\Sigma_\lambda^n$  is contained in the geodesic ball of center  $e_1 \in \mathbb{H}^n$  and radius

$$\rho = \cosh^{-1} \sqrt{1 + \frac{1}{\lambda^2}}.$$

Therefore, since the vertical height of such hyperbolic cap is

# Sharpness of the estimate

## Remark (cont.)

We also easily verify that the hyperbolic image of  $\Sigma_\lambda^n$  is contained in the geodesic ball of center  $e_1 \in \mathbb{H}^n$  and radius

$$\rho = \cosh^{-1} \sqrt{1 + \frac{1}{\lambda^2}}.$$

Therefore, since the vertical height of such hyperbolic cap is

$$h = \sqrt{1 + \lambda^2} - \lambda = \frac{\cosh \rho - 1}{H_r^{1/r}},$$

# Sharpness of the estimate

## Remark (cont.)

We also easily verify that the hyperbolic image of  $\Sigma_\lambda^n$  is contained in the geodesic ball of center  $e_1 \in \mathbb{H}^n$  and radius

$$\rho = \cosh^{-1} \sqrt{1 + \frac{1}{\lambda^2}}.$$

Therefore, since the vertical height of such hyperbolic cap is

$$h = \sqrt{1 + \lambda^2} - \lambda = \frac{\cosh \rho - 1}{H_r^{1/r}},$$

we conclude that our estimate for the vertical height function is sharp.

# An application

- For what follows, we observe that a complete immersed spacelike hypersurface  $\Sigma^n$  of a Lorentzian product space  $-\mathbb{R} \times M^{n+1}$ , with one end, can be regarded as

# An application

- For what follows, we observe that a complete immersed spacelike hypersurface  $\Sigma^n$  of a Lorentzian product space  $-\mathbb{R} \times M^{n+1}$ , with one end, can be regarded as

$$\Sigma^n = \Sigma_t^n \cup \mathcal{C},$$

# An application

- For what follows, we observe that a complete immersed spacelike hypersurface  $\Sigma^n$  of a Lorentzian product space  $-\mathbb{R} \times M^{n+1}$ , with one end, can be regarded as

$$\Sigma^n = \Sigma_t^n \cup \mathcal{C},$$

where  $\Sigma_t^n$  is a compact hypersurface with boundary contained into a slice  $M_t = \{t\} \times M^n$  and  $\mathcal{C}^n$  is diffeomorphic to the cylinder  $[t, \infty) \times \mathbb{S}^{n-1}$ .



# An application

- For what follows, we observe that a complete immersed spacelike hypersurface  $\Sigma^n$  of a Lorentzian product space  $-\mathbb{R} \times M^{n+1}$ , with one end, can be regarded as

$$\Sigma^n = \Sigma_t^n \cup \mathcal{C},$$

where  $\Sigma_t^n$  is a compact hypersurface with boundary contained into a slice  $M_t = \{t\} \times M^n$  and  $\mathcal{C}^n$  is diffeomorphic to the cylinder  $[t, \infty) \times \mathbb{S}^{n-1}$ .

- Given a complete space-like hypersurface with one end,  $\Sigma^n = \Sigma_t^n \cup \mathcal{C}$ , we say that the end of  $\Sigma^n$  is *divergent* if, considering  $\mathcal{C}^n$  with cylindrical coordinates  $p = (s, q) \in [t, \infty) \times \mathbb{S}^{n-1}$ , we have that

# An application

- For what follows, we observe that a complete immersed spacelike hypersurface  $\Sigma^n$  of a Lorentzian product space  $-\mathbb{R} \times M^{n+1}$ , with one end, can be regarded as

$$\Sigma^n = \Sigma_t^n \cup \mathcal{C},$$

where  $\Sigma_t^n$  is a compact hypersurface with boundary contained into a slice  $M_t = \{t\} \times M^n$  and  $\mathcal{C}^n$  is diffeomorphic to the cylinder  $[t, \infty) \times \mathbb{S}^{n-1}$ .

- Given a complete space-like hypersurface with one end,  $\Sigma^n = \Sigma_t^n \cup \mathcal{C}$ , we say that the end of  $\Sigma^n$  is *divergent* if, considering  $\mathcal{C}^n$  with cylindrical coordinates  $p = (s, q) \in [t, \infty) \times \mathbb{S}^{n-1}$ , we have that

$$\lim_{s \rightarrow \infty} h(p) = \infty,$$

# An application

- For what follows, we observe that a complete immersed spacelike hypersurface  $\Sigma^n$  of a Lorentzian product space  $-\mathbb{R} \times M^{n+1}$ , with one end, can be regarded as

$$\Sigma^n = \Sigma_t^n \cup \mathcal{C},$$

where  $\Sigma_t^n$  is a compact hypersurface with boundary contained into a slice  $M_t = \{t\} \times M^n$  and  $\mathcal{C}^n$  is diffeomorphic to the cylinder  $[t, \infty) \times \mathbb{S}^{n-1}$ .

- Given a complete space-like hypersurface with one end,  $\Sigma^n = \Sigma_t^n \cup \mathcal{C}$ , we say that the end of  $\Sigma^n$  is *divergent* if, considering  $\mathcal{C}^n$  with cylindrical coordinates  $p = (s, q) \in [t, \infty) \times \mathbb{S}^{n-1}$ , we have that

$$\lim_{s \rightarrow \infty} h(p) = \infty,$$

where  $h$  denotes the vertical height function of  $\mathcal{C}^n$ .

## Theorem

Let  $\Sigma^n$  be a complete immersed space-like hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$ , with one end. Suppose that one of the following conditions is satisfied:

## Theorem

Let  $\Sigma^n$  be a complete immersed space-like hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$ , with one end. Suppose that one of the following conditions is satisfied:

- (a) The Riemannian fiber  $M^n$  has nonnegative Ricci curvature  $\text{Ric}_M$  and  $\Sigma^n$  has positive constant mean curvature  $H$ .

## Theorem

Let  $\Sigma^n$  be a complete immersed space-like hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$ , with one end. Suppose that one of the following conditions is satisfied:

- (a) The Riemannian fiber  $M^n$  has nonnegative Ricci curvature  $\text{Ric}_M$  and  $\Sigma^n$  has positive constant mean curvature  $H$ .
- (b) The Riemannian fiber  $M^n$  has nonnegative constant sectional curvature  $\kappa_M$  and  $\Sigma^n$  has positive constant  $r$ -mean curvature  $H_r$ , for some  $1 \leq r \leq n$ .

## Theorem

Let  $\Sigma^n$  be a complete immersed space-like hypersurface of a Lorentzian product space  $-\mathbb{R} \times M^n$ , with one end. Suppose that one of the following conditions is satisfied:

- (a) The Riemannian fiber  $M^n$  has nonnegative Ricci curvature  $\text{Ric}_M$  and  $\Sigma^n$  has positive constant mean curvature  $H$ .
- (b) The Riemannian fiber  $M^n$  has nonnegative constant sectional curvature  $\kappa_M$  and  $\Sigma^n$  has positive constant  $r$ -mean curvature  $H_r$ , for some  $1 \leq r \leq n$ .

If the support function  $\eta$  of  $\Sigma^n$  is bounded, then its end is not divergent.

# Height Estimates for $r$ -Mean Curvature Spacelike Hypersurfaces in Product Spaces

A.Gervasio Colares  
( Universidade Federal do Ceará, Brazil )

Joint work with H. F. de Lima  
in *General Relativity and Gravitation*, v.40 (2008), 2131-2147

V International Meeting on Lorentzian Geometry  
Martina Franca, Italy

Thanks