

# Deformations of $2k$ -Einstein structures

Levi Lopes de Lima

Department of Mathematics  
Federal University of Ceará - Fortaleza/Brazil

Martina Franca - July, 2009

Joint work with Newton Santos (UFPI/Brazil).

# Joint work with Newton Santos (UFPI/Brazil).

⇒ Here are the main results:

# Joint work with Newton Santos (UFPI/Brazil).

⇒ Here are the main results:

- The space of infinitesimal deformations of  $2k$ -Einstein structures is finite dimensional at compact non-euclidean space forms.

# Joint work with Newton Santos (UFPI/Brazil).

⇒ Here are the main results:

- The space of infinitesimal deformations of  $2k$ -Einstein structures is finite dimensional at compact non-euclidean space forms.
- Compact spherical space forms are rigid as  $2k$ -Einstein structures.

# Joint work with Newton Santos (UFPI/Brazil).

⇒ Here are the main results:

- The space of infinitesimal deformations of  $2k$ -Einstein structures is finite dimensional at compact non-euclidean space forms.
- Compact spherical space forms are rigid as  $2k$ -Einstein structures.
- Gauss-Bonnet curvatures can be prescribed on certain compact space forms.

## Joint work with Newton Santos (UFPI/Brazil).

⇒ Here are the main results:

- The space of infinitesimal deformations of  $2k$ -Einstein structures is finite dimensional at compact non-euclidean space forms.
- Compact spherical space forms are rigid as  $2k$ -Einstein structures.
- Gauss-Bonnet curvatures can be prescribed on certain compact space forms.

## Joint work with Newton Santos (UFPI/Brazil).

⇒ Here are the main results:

- The space of infinitesimal deformations of  $2k$ -Einstein structures is finite dimensional at compact non-euclidean space forms.
- Compact spherical space forms are rigid as  $2k$ -Einstein structures.
- Gauss-Bonnet curvatures can be prescribed on certain compact space forms.



# The standard mathematical motivation for studying the Einstein tensor

# The standard mathematical motivation for studying the Einstein tensor

If  $S_g$  is the *scalar curvature* of a metric  $g$  on  $X$ , consider the *Hilbert-Einstein functional*  $\mathcal{A} : \mathcal{M}(X) \rightarrow \mathbb{R}$ :

$$\mathcal{A}(g) = \int_X S_g \nu_g.$$

# The standard mathematical motivation for studying the Einstein tensor

If  $S_g$  is the *scalar curvature* of a metric  $g$  on  $X$ , consider the *Hilbert-Einstein functional*  $\mathcal{A} : \mathcal{M}(X) \rightarrow \mathbb{R}$ :

$$\mathcal{A}(g) = \int_X S_g \nu_g.$$

Then,

- $g$  is critical for  $\mathcal{A} \Leftrightarrow \mathcal{T}_g = 0$ , where

$$\mathcal{T}_g = \text{Ric}_g - \frac{1}{2}S_g g$$

is the *Einstein tensor* of  $g$ .

- Actually it is proved that  $-\mathcal{T}$  is the  $L^2$ -gradient of  $\mathcal{A}$ . In particular,  $\mathcal{T}_g = 0$  is divergence free for *any*  $g$ !

# The standard mathematical motivation for studying the Einstein tensor

If  $\mathcal{S}_g$  is the *scalar curvature* of a metric  $g$  on  $X$ , consider the *Hilbert-Einstein functional*  $\mathcal{A} : \mathcal{M}(X) \rightarrow \mathbb{R}$ :

$$\mathcal{A}(g) = \int_X \mathcal{S}_g \nu_g.$$

Then,

- $g$  is critical for  $\mathcal{A} \Leftrightarrow \mathcal{T}_g = 0$ , where

$$\mathcal{T}_g = \text{Ric}_g - \frac{1}{2}\mathcal{S}_g g$$

is the *Einstein tensor* of  $g$ .

- Actually it is proved that  $-\mathcal{T}$  is the  $L^2$ -gradient of  $\mathcal{A}$ . In particular,  $\mathcal{T}_g = 0$  is divergence free for *any*  $g$ !
- Moreover,  $g$  is critical for  $\mathcal{A}|_{\mathcal{M}_1(X)} \Leftrightarrow \text{Ric}_g = \lambda g$  ( $g$  is Einstein).

# The standard mathematical motivation for studying the Einstein tensor

If  $S_g$  is the *scalar curvature* of a metric  $g$  on  $X$ , consider the *Hilbert-Einstein functional*  $\mathcal{A} : \mathcal{M}(X) \rightarrow \mathbb{R}$ :

$$\mathcal{A}(g) = \int_X S_g \nu_g.$$

Then,

- $g$  is critical for  $\mathcal{A} \Leftrightarrow \mathcal{T}_g = 0$ , where

$$\mathcal{T}_g = \text{Ric}_g - \frac{1}{2}S_g g$$

is the *Einstein tensor* of  $g$ .

- Actually it is proved that  $-\mathcal{T}$  is the  $L^2$ -gradient of  $\mathcal{A}$ . In particular,  $\mathcal{T}_g = 0$  is divergence free for *any*  $g$ !
- Moreover,  $g$  is critical for  $\mathcal{A}|_{\mathcal{M}_1(X)} \Leftrightarrow \text{Ric}_g = \lambda g$  ( $g$  is Einstein).

# Another mathematical motivation: an old result by Vermeil, Weyl and Cartan

# Another mathematical motivation: an old result by Vermeil, Weyl and Cartan

## Theorem

Let  $T = T_{ij}$  be a twice covariant tensor on a Riemannian (or Lorentzian) manifold  $(X, g)$  such that:

- 1  $T_{ij} = T_{ji}$ ;
- 2  $\nabla_i T_j^i = 0$ ;
- 3 Locally,  $T = \sum_{|\alpha|=2} A_\alpha(g, \partial g) \partial^\alpha g + B(g, \partial g)$ .

# Another mathematical motivation: an old result by Vermeil, Weyl and Cartan

## Theorem

Let  $T = T_{ij}$  be a twice covariant tensor on a Riemannian (or Lorentzian) manifold  $(X, g)$  such that:

- 1  $T_{ij} = T_{ji}$ ;
- 2  $\nabla_i T_j^i = 0$ ;
- 3 Locally,  $T = \sum_{|\alpha|=2} A_\alpha(g, \partial g) \partial^\alpha g + B(g, \partial g)$ .

Then,

$$T_{ij} = ag_{ij} + bT_{ij},$$



# Another mathematical motivation: an old result by Vermeil, Weyl and Cartan

## Theorem

Let  $T = T_{ij}$  be a twice covariant tensor on a Riemannian (or Lorentzian) manifold  $(X, g)$  such that:

- 1  $T_{ij} = T_{ji}$ ;
- 2  $\nabla_i T_j^i = 0$ ;
- 3 Locally,  $T = \sum_{|\alpha|=2} A_\alpha(g, \partial g) \partial^\alpha g + B(g, \partial g)$ .

Then,

$$T_{ij} = ag_{ij} + bT_{ij},$$

where

$$T_{ij} = \text{Ric}_{ij} - \frac{S}{2}g_{ij}, \quad S = g^{ij}\text{Ric}_{ij}.$$

# Another mathematical motivation: an old result by Vermeil, Weyl and Cartan

## Theorem

Let  $T = T_{ij}$  be a twice covariant tensor on a Riemannian (or Lorentzian) manifold  $(X, g)$  such that:

- 1  $T_{ij} = T_{ji}$ ;
- 2  $\nabla_i T_j^i = 0$ ;
- 3 Locally,  $T = \sum_{|\alpha|=2} A_\alpha(g, \partial g) \partial^\alpha g + B(g, \partial g)$ .

Then,

$$T_{ij} = ag_{ij} + bT_{ij},$$

where

$$T_{ij} = \text{Ric}_{ij} - \frac{S}{2}g_{ij}, \quad S = g^{ij}\text{Ric}_{ij}.$$

The main point here is:

# The main point here is:

There are generalized Ricci tensors  $\text{Ric}_g^{(2k)}$  and scalar curvatures  $\mathcal{S}_g^{(2k)}$  (and hence of the generalized Einstein tensors!) such that:

- $\text{Ric}_g^{(2)} = \text{Ric}_g$  and  $\mathcal{S}_g^{(2)} = \mathcal{S}_g/2$ ;
- The condition  $\text{Ric}_g^{(2k)} = \lambda g$  defines  $2k$ -Einstein metrics;
- The variational picture above extends to this more general setting.

# The main point here is:

There are generalized Ricci tensors  $\text{Ric}_g^{(2k)}$  and scalar curvatures  $\mathcal{S}_g^{(2k)}$  (and hence of the generalized Einstein tensors!) such that:

- $\text{Ric}_g^{(2)} = \text{Ric}_g$  and  $\mathcal{S}_g^{(2)} = \mathcal{S}_g/2$ ;
- The condition  $\text{Ric}_g^{(2k)} = \lambda g$  defines  $2k$ -Einstein metrics;
- The variational picture above extends to this more general setting.

# The main point here is:

There are generalized Ricci tensors  $\text{Ric}_g^{(2k)}$  and scalar curvatures  $\mathcal{S}_g^{(2k)}$  (and hence of the generalized Einstein tensors!) such that:

- $\text{Ric}_g^{(2)} = \text{Ric}_g$  and  $\mathcal{S}_g^{(2)} = \mathcal{S}_g/2$ ;
- The condition  $\text{Ric}_g^{(2k)} = \lambda g$  defines  $2k$ -Einstein metrics;
- The variational picture above extends to this more general setting.

# A not so old theorem by Lovelock

# A not so old theorem by Lovelock

Replace the third condition in the VWC theorem by only requiring that

$$T = T(g, \partial g, \partial^2 g)$$



# A not so old theorem by Lovelock

Replace the third condition in the VWC theorem by only requiring that

$$T = T(g, \partial g, \partial^2 g)$$

Then:

$$T = \sum_{k \leq [n/2]} a_k T_g^{(2k)}.$$

# A not so old theorem by Lovelock

Replace the third condition in the VWC theorem by only requiring that

$$T = T(g, \partial g, \partial^2 g)$$

Then:

$$T = \sum_{k \leq [n/2]} a_k \mathcal{T}_g^{(2k)}.$$

- Here,  $\mathcal{T}_g^{(0)} = g$ ,  $\mathcal{T}_g^{(2)} = \mathcal{T}_g$ , but who are the Lovelock tensors  $\mathcal{T}_g^{(2k)}$ ,  $k \geq 2$ ?

# A not so old theorem by Lovelock

Replace the third condition in the VWC theorem by only requiring that

$$T = T(g, \partial g, \partial^2 g)$$

Then:

$$T = \sum_{k \leq [n/2]} a_k \mathcal{T}_g^{(2k)}.$$

- Here,  $\mathcal{T}_g^{(0)} = g$ ,  $\mathcal{T}_g^{(2)} = \mathcal{T}_g$ , but who are the Lovelock tensors  $\mathcal{T}_g^{(2k)}$ ,  $k \geq 2$ ?

# Sectional curvature of $2k$ -planes (Thorpe)

# Sectional curvature of $2k$ -planes (Thorpe)

## Definition

If  $\mathfrak{p}$  is a tangent  $2k$ -plane to  $(X, g)$ , its *sectional curvature* is

$$\sigma^{(2k)}(\mathfrak{p}) = c_k \sum \delta_{j_1 j_2 \dots j_{2k-1} j_{2k}}^{i_1 i_2 \dots i_{2k-1} i_{2k}} R_{i_1 i_2}^{j_1 j_2} \dots R_{i_{2k-1} i_{2k}}^{j_{2k-1} j_{2k}}.$$

# Sectional curvature of $2k$ -planes (Thorpe)

## Definition

If  $\mathfrak{p}$  is a tangent  $2k$ -plane to  $(X, g)$ , its *sectional curvature* is

$$\sigma^{(2k)}(\mathfrak{p}) = c_k \sum \delta_{j_1 j_2 \dots j_{2k-1} j_{2k}}^{i_1 i_2 \dots i_{2k-1} i_{2k}} R_{i_1 i_2}^{j_1 j_2} \dots R_{i_{2k-1} i_{2k}}^{j_{2k-1} j_{2k}}.$$

Here,  $R_{ij}^{kl} = g(R_g(e_i, e_j)e_l, e_k)$  and  $\{e_i\}$  is an orthonormal basis of  $\mathfrak{p}$ .

# Sectional curvature of $2k$ -planes (Thorpe)

## Definition

If  $\mathfrak{p}$  is a tangent  $2k$ -plane to  $(X, g)$ , its *sectional curvature* is

$$\sigma^{(2k)}(\mathfrak{p}) = c_k \sum \delta_{j_1 j_2 \dots j_{2k-1} j_{2k}}^{i_1 i_2 \dots i_{2k-1} i_{2k}} R_{i_1 i_2}^{j_1 j_2} \dots R_{i_{2k-1} i_{2k}}^{j_{2k-1} j_{2k}}.$$

Here,  $R_{ij}^{kl} = g(R_g(e_i, e_j)e_l, e_k)$  and  $\{e_i\}$  is an orthonormal basis of  $\mathfrak{p}$ .

- If  $k = 1$  then  $\sigma^{(2)}(\mathfrak{p})$  is the standard sectional curvature.

# Sectional curvature of $2k$ -planes (Thorpe)

## Definition

If  $\mathfrak{p}$  is a tangent  $2k$ -plane to  $(X, g)$ , its *sectional curvature* is

$$\sigma^{(2k)}(\mathfrak{p}) = c_k \sum \delta_{j_1 j_2 \dots j_{2k-1} j_{2k}}^{i_1 i_2 \dots i_{2k-1} i_{2k}} R_{i_1 i_2}^{j_1 j_2} \dots R_{i_{2k-1} i_{2k}}^{j_{2k-1} j_{2k}}.$$

Here,  $R_{ij}^{kl} = g(R_g(e_i, e_j)e_l, e_k)$  and  $\{e_i\}$  is an orthonormal basis of  $\mathfrak{p}$ .

- If  $k = 1$  then  $\sigma^{(2)}(\mathfrak{p})$  is the standard sectional curvature.
- If  $n = 2k$  then  $\sigma^{(n)}(x) = \mathcal{S}^{(n)}(T_x X)$  is the Gauss-Bonnet integrand.



# Sectional curvature of $2k$ -planes (Thorpe)

## Definition

If  $\mathfrak{p}$  is a tangent  $2k$ -plane to  $(X, g)$ , its *sectional curvature* is

$$\sigma^{(2k)}(\mathfrak{p}) = c_k \sum \delta_{j_1 j_2 \dots j_{2k-1} j_{2k}}^{i_1 i_2 \dots i_{2k-1} i_{2k}} R_{i_1 i_2}^{j_1 j_2} \dots R_{i_{2k-1} i_{2k}}^{j_{2k-1} j_{2k}}.$$

Here,  $R_{ij}^{kl} = g(R_g(e_i, e_j)e_l, e_k)$  and  $\{e_i\}$  is an orthonormal basis of  $\mathfrak{p}$ .

- If  $k = 1$  then  $\sigma^{(2)}(\mathfrak{p})$  is the standard sectional curvature.
- If  $n = 2k$  then  $\sigma^{(n)}(x) = \mathcal{S}^{(n)}(T_x X)$  is the Gauss-Bonnet integrand.
- In general,  $\sigma^{(2k)}(\mathfrak{p})$  is the Gauss-Bonnet integrand of  $\exp(\mathfrak{p})$  computed at  $\exp(0)$ .

# $2k$ -Ricci tensor and $2k$ -Gauss-Bonnet curvature

# $2k$ -Ricci tensor and $2k$ -Gauss-Bonnet curvature

## Definition

Se  $v \in T_x X$ ,  $|v| = 1$ , and  $\{e_1, \dots, e_{n-1}, e_n = v\} \subset T_x X$  is orthonormal, the  $2k$ -Ricci tensor is given by

$$\text{Ric}_g^{(2k)}(v, v) = \sum \sigma^{(2k)}(e_{i_1} \wedge \dots \wedge e_{i_{2k-1}} \wedge v).$$

# $2k$ -Ricci tensor and $2k$ -Gauss-Bonnet curvature

## Definition

Se  $v \in T_x X$ ,  $|v| = 1$ , and  $\{e_1, \dots, e_{n-1}, e_n = v\} \subset T_x X$  is orthonormal, the  $2k$ -Ricci tensor is given by

$$\text{Ric}_g^{(2k)}(v, v) = \sum \sigma^{(2k)}(e_{i_1} \wedge \dots \wedge e_{i_{2k-1}} \wedge v).$$

## Definition

The  $2k$ -Gauss-Bonnet curvature of  $(X, g)$  is

$$S_g^{(2k)} = \frac{1}{(2k)!} \text{tr}_g \text{Ric}_g^{(2k)}.$$

Locally,

$$S^{(2k)} = d_k \sum \delta_{j_1 j_2 \dots j_{2k-1} j_{2k}}^{i_1 i_2 \dots i_{2k-1} i_{2k}} R_{i_1 i_2}^{j_1 j_2} \dots R_{i_{2k-1} i_{2k}}^{j_{2k-1} j_{2k}}.$$

# $2k$ -Ricci tensor and $2k$ -Gauss-Bonnet curvature

## Definition

Se  $v \in T_x X$ ,  $|v| = 1$ , and  $\{e_1, \dots, e_{n-1}, e_n = v\} \subset T_x X$  is orthonormal, the  $2k$ -Ricci tensor is given by

$$\text{Ric}_g^{(2k)}(v, v) = \sum \sigma^{(2k)}(e_{i_1} \wedge \dots \wedge e_{i_{2k-1}} \wedge v).$$

## Definition

The  $2k$ -Gauss-Bonnet curvature of  $(X, g)$  is

$$\mathcal{S}_g^{(2k)} = \frac{1}{(2k)!} \text{tr}_g \text{Ric}_g^{(2k)}.$$

Locally,

$$\mathcal{S}^{(2k)} = d_k \sum \delta_{j_1 j_2 \dots j_{2k-1} j_{2k}}^{i_1 i_2 \dots i_{2k-1} i_{2k}} R_{i_1 i_2}^{j_1 j_2} \dots R_{i_{2k-1} i_{2k}}^{j_{2k-1} j_{2k}}.$$

# The variational picture I

# The variational picture I

## Definition

The  $2k$ -Lovelock tensor of  $(X, g)$  is

$$\mathcal{T}_g^{(2k)} = \frac{\text{Ric}_g^{(2k)}}{(2k-1)!} - \mathcal{S}_g^{(2k)} g.$$

# The variational picture I

## Definition

The  $2k$ -Lovelock tensor of  $(X, g)$  is

$$\mathcal{T}_g^{(2k)} = \frac{\text{Ric}_g^{(2k)}}{(2k-1)!} - \mathcal{S}_g^{(2k)} g.$$

If  $\mathcal{A}^{(2k)} : \mathcal{M}(X) \rightarrow \mathbb{R}$  is the *Lovelock functional*

$$\mathcal{A}^{(2k)}(g) = \int_X \mathcal{S}_g^{(2k)} \nu_g,$$



# The variational picture I

## Definition

The  $2k$ -Lovelock tensor of  $(X, g)$  is

$$\mathcal{T}_g^{(2k)} = \frac{\text{Ric}_g^{(2k)}}{(2k-1)!} - \mathcal{S}_g^{(2k)} g.$$

If  $\mathcal{A}^{(2k)} : \mathcal{M}(X) \rightarrow \mathbb{R}$  is the *Lovelock functional*

$$\mathcal{A}^{(2k)}(g) = \int_X \mathcal{S}_g^{(2k)} \nu_g,$$

then

- $g$  is critical for  $\mathcal{A}^{(2k)} \Leftrightarrow \mathcal{T}_g^{(2k)} = 0$ . In fact,  $-\mathcal{T}_g^{(2k)}$  is the  $L^2$ -gradient of  $\mathcal{A}^{(2k)}$  and hence  $\text{div } \mathcal{T}_g^{(2k)} = 0$ .

# The variational picture I

## Definition

The  $2k$ -Lovelock tensor of  $(X, g)$  is

$$\mathcal{T}_g^{(2k)} = \frac{\text{Ric}_g^{(2k)}}{(2k-1)!} - \mathcal{S}_g^{(2k)} g.$$

If  $\mathcal{A}^{(2k)} : \mathcal{M}(X) \rightarrow \mathbb{R}$  is the *Lovelock functional*

$$\mathcal{A}^{(2k)}(g) = \int_X \mathcal{S}_g^{(2k)} \nu_g,$$

then

- $g$  is critical for  $\mathcal{A}^{(2k)} \Leftrightarrow \mathcal{T}_g^{(2k)} = 0$ . In fact,  $-\mathcal{T}_g^{(2k)}$  is the  $L^2$ -gradient of  $\mathcal{A}^{(2k)}$  and hence  $\text{div } \mathcal{T}_g^{(2k)} = 0$ .

# The variational picture II: $2k$ -Einstein metrics

# The variational picture II: $2k$ -Einstein metrics

More importantly, we have

- $g$  is critical for  $\mathcal{A}^{(2k)}|_{\mathcal{M}_1(X)} \Leftrightarrow \text{Ric}_g^{(2k)} = \lambda_g g$  ( $g$  is  $2k$ -Einstein, by definition).

# The variational picture II: $2k$ -Einstein metrics

More importantly, we have

- $g$  is critical for  $\mathcal{A}^{(2k)}|_{\mathcal{M}_1(X)} \Leftrightarrow \operatorname{Ric}_g^{(2k)} = \lambda_g g$  ( $g$  is  $2k$ -Einstein, by definition).
- If  $2k = n$  then any compact manifold is  $2k$ -Einstein by Gauss-Bonnet! Thus we may assume  $n > 2k$ .

## The variational picture II: $2k$ -Einstein metrics

More importantly, we have

- $g$  is critical for  $\mathcal{A}^{(2k)}|_{\mathcal{M}_1(X)} \Leftrightarrow \text{Ric}_g^{(2k)} = \lambda_g g$  ( $g$  is  $2k$ -Einstein, by definition).
- If  $2k = n$  then any compact manifold is  $2k$ -Einstein by Gauss-Bonnet! Thus we may assume  $n > 2k$ .

Moreover, in this case, the fact that  $\mathcal{T}^{(2k)}$  is divergence free implies that

$$\lambda_g = \frac{(2k)!}{n} S_g^{(2k)}$$

# The variational picture II: $2k$ -Einstein metrics

More importantly, we have

- $g$  is critical for  $\mathcal{A}^{(2k)}|_{\mathcal{M}_1(X)} \Leftrightarrow \text{Ric}_g^{(2k)} = \lambda_g g$  ( $g$  is  $2k$ -Einstein, by definition).
- If  $2k = n$  then any compact manifold is  $2k$ -Einstein by Gauss-Bonnet! Thus we may assume  $n > 2k$ .

Moreover, in this case, the fact that  $\mathcal{T}^{(2k)}$  is divergence free implies that

$$\lambda_g = \frac{(2k)!}{n} \mathcal{S}_g^{(2k)}$$

is a constant. In particular,  $\mathcal{S}_g^{(2k)}$  is a constant as well.

# A few examples



# A few examples

- Space forms;

# A few examples

- Space forms;
- Isotropically irreducible homogeneous spaces (this includes all irreducible symmetric spaces);

# A few examples

- Space forms;
- Isotropically irreducible homogeneous spaces (this includes all irreducible symmetric spaces);
- If  $M^4$  is Ricci-flat but not flat and  $N$  is flat then  $M \times N$  is Einstein but not 4-Einstein.

# A few examples

- Space forms;
- Isotropically irreducible homogeneous spaces (this includes all irreducible symmetric spaces);
- If  $M^4$  is Ricci-flat but not flat and  $N$  is flat then  $M \times N$  is Einstein but not 4-Einstein.
- If  $M^n$  is *any* Riemannian manifold and  $N^m$  is flat then  $M \times N$  is  $2k$ -Einstein for  $2k > n$  (for  $\lambda = 0$ ).

# A few examples

- Space forms;
- Isotropically irreducible homogeneous spaces (this includes all irreducible symmetric spaces);
- If  $M^4$  is Ricci-flat but not flat and  $N$  is flat then  $M \times N$  is Einstein but not 4-Einstein.
- If  $M^n$  is *any* Riemannian manifold and  $N^m$  is flat then  $M \times N$  is  $2k$ -Einstein for  $2k > n$  (for  $\lambda = 0$ ).

# The moduli space of $2k$ -Einstein structures

# The moduli space of $2k$ -Einstein structures

Let  $X$  be a *closed* manifold and

- $D(X)$  the diffeomorphism group of  $X$ ;
- $E^{(2k)}(X)$  the space of  $2k$ -Einstein metrics on  $X$ .

# The moduli space of $2k$ -Einstein structures

Let  $X$  be a *closed* manifold and

- $D(X)$  the diffeomorphism group of  $X$ ;
- $E^{(2k)}(X)$  the space of  $2k$ -Einstein metrics on  $X$ .

and consider the standard action

$$\begin{array}{ccccc} (\mathbb{R}^+ \times D(X)) & \times & \mathcal{M}(X) & \rightarrow & \mathcal{M}(X) \\ & & \cup & & \\ (\mathbb{R}^+ \times D(X)) & \times & E^{(2k)}(X) & \rightarrow & E^{(2k)}(X) \end{array}$$



# The moduli space of $2k$ -Einstein structures

Let  $X$  be a *closed* manifold and

- $D(X)$  the diffeomorphism group of  $X$ ;
- $E^{(2k)}(X)$  the space of  $2k$ -Einstein metrics on  $X$ .

and consider the standard action

$$\begin{array}{ccccc} (\mathbb{R}^+ \times D(X)) & \times & \mathcal{M}(X) & \rightarrow & \mathcal{M}(X) \\ & & \cup & & \\ (\mathbb{R}^+ \times D(X)) & \times & E^{(2k)}(X) & \rightarrow & E^{(2k)}(X) \end{array}$$

## Definition

The *moduli space of  $2k$ -Einstein structures on  $X$*  is

$$\mathcal{E}^{(2k)}(X) = \frac{E^{(2k)}(X)}{\mathbb{R}^+ \times D(X)}.$$

The quotient map is represented by  $g \mapsto [g]$ .

# Looking at $\mathcal{E}^{(2k)}(X)$ infinitesimally I

# Looking at $\mathcal{E}^{(2k)}(X)$ infinitesimally I

Let  $t \in (-\epsilon, \epsilon) \mapsto [g_t] \in \mathcal{E}^{(2k)}(X)$  a one-parameter family of  $2k$ -Einstein structures. Here,  $g_t \in E^{(2k)}(X)$ , with  $g_0 = g$ . Thus,  $[g_t]$  is a *deformation* of  $[g]$ .

# Looking at $\mathcal{E}^{(2k)}(X)$ infinitesimally I

Let  $t \in (-\epsilon, \epsilon) \mapsto [g_t] \in \mathcal{E}^{(2k)}(X)$  a one-parameter family of  $2k$ -Einstein structures. Here,  $g_t \in E^{(2k)}(X)$ , with  $g_0 = g$ . Thus,  $[g_t]$  is a *deformation* of  $[g]$ .

- What constraints  $h = dg_t/dt|_{t=0} \in S^2(X)$  should satisfy?

# Looking at $\mathcal{E}^{(2k)}(X)$ infinitesimally I

Let  $t \in (-\epsilon, \epsilon) \mapsto [g_t] \in \mathcal{E}^{(2k)}(X)$  a one-parameter family of  $2k$ -Einstein structures. Here,  $g_t \in E^{(2k)}(X)$ , with  $g_0 = g$ . Thus,  $[g_t]$  is a *deformation* of  $[g]$ .

- What constraints  $h = dg_t/dt|_{t=0} \in S^2(X)$  should satisfy?

One has:

$$\boxed{\int_X \operatorname{tr}_g h \nu_g = 0} \quad (\text{scale invariance}) \quad (I)$$

# Looking at $\mathcal{E}^{(2k)}(X)$ infinitesimally I

Let  $t \in (-\epsilon, \epsilon) \mapsto [g_t] \in \mathcal{E}^{(2k)}(X)$  a one-parameter family of  $2k$ -Einstein structures. Here,  $g_t \in E^{(2k)}(X)$ , with  $g_0 = g$ . Thus,  $[g_t]$  is a *deformation* of  $[g]$ .

- What constraints  $h = dg_t/dt|_{t=0} \in S^2(X)$  should satisfy?

One has:

$$\boxed{\int_X \operatorname{tr}_g h \nu_g = 0} \quad (\text{scale invariance}) \quad (\text{I})$$

# Looking at $\mathcal{E}^{(2k)}(X)$ infinitesimally II (Berger-Ebin)

## Looking at $\mathcal{E}^{(2k)}(X)$ infinitesimally II (Berger-Ebin)

We should also discard *trivial* deformations satisfying  $g_t = \varphi_t^* g$ ,  $\varphi_t \in D(X)$ . Equivalently,  $g_t$  remains in the  $D(X)$ -orbit  $\mathcal{O}(g)$  of  $g$ .



# Looking at $\mathcal{E}^{(2k)}(X)$ infinitesimally II (Berger-Ebin)

We should also discard *trivial* deformations satisfying  $g_t = \varphi_t^* g$ ,  $\varphi_t \in D(X)$ . Equivalently,  $g_t$  remains in the  $D(X)$ -orbit  $\mathcal{O}(g)$  of  $g$ .

## Theorem

If  $\delta_g^*$  is the  $L^2$ -adjoint of the divergence operator  $\delta_g : S^2(X) \rightarrow \mathcal{D}^1(X)$  then  $T_g \mathcal{O}(g) = \text{Im } \delta_g^*$ . Moreover,

$$S^2(X) = \text{Im } \delta_g^* \oplus \ker \delta_g.$$

Thus,

$$\boxed{\delta_g h = 0} \quad (\text{diffeomorphism invariance}) \quad (\text{II})$$

## Looking at $\mathcal{E}^{(2k)}(X)$ infinitesimally II (Berger-Ebin)

We should also discard *trivial* deformations satisfying  $g_t = \varphi_t^* g$ ,  $\varphi_t \in D(X)$ . Equivalently,  $g_t$  remains in the  $D(X)$ -orbit  $\mathcal{O}(g)$  of  $g$ .

### Theorem

If  $\delta_g^*$  is the  $L^2$ -adjoint of the divergence operator  $\delta_g : S^2(X) \rightarrow \mathcal{D}^1(X)$  then  $T_g \mathcal{O}(g) = \text{Im } \delta_g^*$ . Moreover,

$$S^2(X) = \text{Im } \delta_g^* \oplus \ker \delta_g.$$

Thus,

$$\boxed{\delta_g h = 0} \quad (\text{diffeomorphism invariance}) \quad (\text{II})$$

# Looking at $\mathcal{E}^{(2k)}$ infinitesimally III

# Looking at $\mathcal{E}^{(2k)}$ infinitesimally III

We also have

$$\mathrm{Ric}_{g_t}^{(2k)} = \lambda_{g_t} g_t,$$

# Looking at $\mathcal{E}^{(2k)}$ infinitesimally III

We also have

$$\mathrm{Ric}_{g_t}^{(2k)} = \lambda_{g_t} g_t,$$

so that

$$\dot{\mathrm{Ric}}_g^{(2k)} h = \dot{\lambda}_g h g + \lambda_g h.$$

# Looking at $\mathcal{E}^{(2k)}$ infinitesimally III

We also have

$$\mathrm{Ric}_{g_t}^{(2k)} = \lambda_{g_t} g_t,$$

so that

$$\dot{\mathrm{Ric}}_g^{(2k)} h = \dot{\lambda}_g h g + \lambda_g h.$$

But

$$\lambda_{g_t} = \frac{(2k)!}{n} \mathcal{S}_{g_t}^{(2k)} = \frac{(2k)!}{n} \mathcal{A}^{(2k)}(g_t) \implies \dot{\lambda}_g h = 0,$$

# Looking at $\mathcal{E}^{(2k)}$ infinitesimally III

We also have

$$\mathrm{Ric}_{g_t}^{(2k)} = \lambda_{g_t} g_t,$$

so that

$$\dot{\mathrm{Ric}}_g^{(2k)} h = \dot{\lambda}_g h g + \lambda_g h.$$

But

$$\lambda_{g_t} = \frac{(2k)!}{n} \mathcal{S}_{g_t}^{(2k)} = \frac{(2k)!}{n} \mathcal{A}^{(2k)}(g_t) \implies \dot{\lambda}_g h = 0,$$

and we must also have

$$\dot{\mathrm{Ric}}_g^{(2k)} h = \lambda_g h \quad (\text{III})$$

# Looking at $\mathcal{E}^{(2k)}$ infinitesimally III

We also have

$$\mathrm{Ric}_{g_t}^{(2k)} = \lambda_{g_t} g_t,$$

so that

$$\dot{\mathrm{Ric}}_g^{(2k)} h = \dot{\lambda}_g h g + \lambda_g h.$$

But

$$\lambda_{g_t} = \frac{(2k)!}{n} \mathcal{S}_{g_t}^{(2k)} = \frac{(2k)!}{n} \mathcal{A}^{(2k)}(g_t) \implies \dot{\lambda}_g h = 0,$$

and we must also have

$$\boxed{\dot{\mathrm{Ric}}_g^{(2k)} h = \lambda_g h} \quad (\text{III})$$



# (Infinitesimal) rigidity

# (Infinitesimal) rigidity

## Definition

If  $\mathcal{I}_g = \text{tr}_g^{-1}(0) \cap \delta_g^{-1}(0)$ , the *space of infinitesimal deformations* of  $[g]$  is

$$\varepsilon_{[g]}^{(2k)} \doteq \ker \mathcal{L}_g^{(2k)}|_{\mathcal{I}_g}, \quad \mathcal{L}_g^{(2k)} = \dot{\text{Ric}}_g^{(2k)} - \lambda_g.$$

# (Infinitesimal) rigidity

## Definition

If  $\mathcal{I}_g = \text{tr}_g^{-1}(0) \cap \delta_g^{-1}(0)$ , the *space of infinitesimal deformations* of  $[g]$  is

$$\varepsilon_{[g]}^{(2k)} \doteq \ker \mathcal{L}_g^{(2k)}|_{\mathcal{I}_g}, \quad \mathcal{L}_g^{(2k)} = \dot{\text{Ric}}_g^{(2k)} - \lambda_g.$$

- In particular,  $\dim \varepsilon_{[g]}^{(2k)} < +\infty$  if  $\mathcal{L}_g^{(2k)}|_{\mathcal{I}_g}$  is elliptic.

# (Infinitesimal) rigidity

## Definition

If  $\mathcal{I}_g = \text{tr}_g^{-1}(0) \cap \delta_g^{-1}(0)$ , the *space of infinitesimal deformations* of  $[g]$  is

$$\varepsilon_{[g]}^{(2k)} \doteq \ker \mathcal{L}_g^{(2k)}|_{\mathcal{I}_g}, \quad \mathcal{L}_g^{(2k)} = \dot{\text{Ric}}_g^{(2k)} - \lambda_g.$$

- In particular,  $\dim \varepsilon_{[g]}^{(2k)} < +\infty$  if  $\mathcal{L}_g^{(2k)}|_{\mathcal{I}_g}$  is elliptic.

## Definition

$[g]$  is *infinitesimally rigid* if  $\varepsilon_{[g]}^{(2k)}$  is trivial.

# (Infinitesimal) rigidity

## Definition

If  $\mathcal{I}_g = \text{tr}_g^{-1}(0) \cap \delta_g^{-1}(0)$ , the *space of infinitesimal deformations* of  $[g]$  is

$$\varepsilon_{[g]}^{(2k)} \doteq \ker \mathcal{L}_g^{(2k)}|_{\mathcal{I}_g}, \quad \mathcal{L}_g^{(2k)} = \dot{\text{Ric}}_g^{(2k)} - \lambda_g.$$

- In particular,  $\dim \varepsilon_{[g]}^{(2k)} < +\infty$  if  $\mathcal{L}_g^{(2k)}|_{\mathcal{I}_g}$  is elliptic.

## Definition

$[g]$  is *infinitesimally rigid* if  $\varepsilon_{[g]}^{(2k)}$  is trivial.

## Definition

A  $2k$ -Einstein structure is *rigid* if it is isolated in  $\mathcal{E}^{2k}(X)$ .

# The Einstein case (Berger-Ebin)

# The Einstein case (Berger-Ebin)

## Theorem

$\dim \varepsilon_{[g]}^{(2)} < +\infty$  for any *Einstein metric*.

# The Einstein case (Berger-Ebin)

## Theorem

$\dim \varepsilon_{[g]}^{(2)} < +\infty$  for any *Einstein metric*.

In this case,

$$\dot{\text{Ric}}_g^{(2)} h = \frac{1}{2} \Delta_L h - \mathcal{L}_{\omega_h^*} g,$$



# The Einstein case (Berger-Ebin)

## Theorem

$\dim \varepsilon_{[g]}^{(2)} < +\infty$  for any *Einstein metric*.

In this case,

$$\dot{\text{Ric}}_g^{(2)} h = \frac{1}{2} \Delta_L h - \mathcal{L}_{\omega_h^*} g,$$

where  $\Delta_L$  is the Lichnerowicz Laplacian and

$$\omega_h = \delta_g h + \frac{1}{2} \text{tr}_g h = 0.$$

# The Einstein case (Berger-Ebin)

## Theorem

$\dim \varepsilon_{[g]}^{(2)} < +\infty$  for any Einstein metric.

In this case,

$$\dot{\text{Ric}}_g^{(2)} h = \frac{1}{2} \Delta_L h - \mathcal{L}_{\omega_h^*} g,$$

where  $\Delta_L$  is the Lichnerowicz Laplacian and

$$\omega_h = \delta_g h + \frac{1}{2} \text{tr}_g h = 0.$$

- However,  $M^n \times T^4$  satisfies  $\text{Ric}_g^{(4)} = 0$ , for any metric on  $M^n$ .  
Thus,  $\dim \varepsilon_{[g]}^{(4)} = +\infty$  in this case.

# The Einstein case (Berger-Ebin)

## Theorem

$\dim \varepsilon_{[g]}^{(2)} < +\infty$  for any Einstein metric.

In this case,

$$\dot{\text{Ric}}_g^{(2)} h = \frac{1}{2} \Delta_L h - \mathcal{L}_{\omega_h^*} g,$$

where  $\Delta_L$  is the Lichnerowicz Laplacian and

$$\omega_h = \delta_g h + \frac{1}{2} \text{tr}_g h = 0.$$

- However,  $M^n \times T^4$  satisfies  $\text{Ric}_g^{(4)} = 0$ , for any metric on  $M^n$ .  
Thus,  $\dim \varepsilon_{[g]}^{(4)} = +\infty$  in this case.

# Our results

# Our results

## Theorem

*If  $4 \leq 2k < n$  then  $\dim \varepsilon_{[g]}^{(2k)} < +\infty$  at nonflat space forms. Moreover, spherical space forms are infinitesimally rigid.*

# Our results

## Theorem

*If  $4 \leq 2k < n$  then  $\dim \varepsilon_{[g]}^{(2k)} < +\infty$  at nonflat space forms. Moreover, spherical space forms are infinitesimally rigid.*

## Theorem

*Spherical space forms are rigid as  $2k$ -Einstein structures.*

# Our results

## Theorem

*If  $4 \leq 2k < n$  then  $\dim \varepsilon_{[g]}^{(2k)} < +\infty$  at nonflat space forms. Moreover, spherical space forms are infinitesimally rigid.*

## Theorem

*Spherical space forms are rigid as  $2k$ -Einstein structures.*

# The formalism of double forms (Nomizu-Kulkarni)



# The formalism of double forms (Nomizu-Kulkarni)

Let

- $\mathcal{D}^p(X)$  differential  $p$ -forms on  $X$ ;
- $\mathcal{D}^{(p,q)} = \mathcal{D}^p(X) \otimes \mathcal{D}^q(X)$  differential  $(p, q)$ -forms on  $X$ ;
- $\mathcal{C}^p(X) \subset \mathcal{D}^{(p,p)}(X)$  symmetric  $(p, p)$ -forms on  $X$ .
- $\mathcal{D}(X) = \bigoplus_{p,q} \mathcal{D}^{(p,q)}(X)$  algebra of double forms on  $X$ .

# The formalism of double forms (Nomizu-Kulkarni)

Let

- $\mathcal{D}^p(X)$  differential  $p$ -forms on  $X$ ;
- $\mathcal{D}^{(p,q)} = \mathcal{D}^p(X) \otimes \mathcal{D}^q(X)$  differential  $(p, q)$ -forms on  $X$ ;
- $\mathcal{C}^p(X) \subset \mathcal{D}^{(p,p)}(X)$  symmetric  $(p, p)$ -forms on  $X$ .
- $\mathcal{D}(X) = \bigoplus_{p,q} \mathcal{D}^{(p,q)}(X)$  algebra of double forms on  $X$ .

For example,  $g \in \mathcal{M}(X) \implies g \in \mathcal{C}^1(X)$  and  $R_g \in \mathcal{C}^2(X)$ .

# The formalism of double forms (Nomizu-Kulkarni)

Let

- $\mathcal{D}^p(X)$  differential  $p$ -forms on  $X$ ;
- $\mathcal{D}^{(p,q)} = \mathcal{D}^p(X) \otimes \mathcal{D}^q(X)$  differential  $(p, q)$ -forms on  $X$ ;
- $\mathcal{C}^p(X) \subset \mathcal{D}^{(p,p)}(X)$  symmetric  $(p, p)$ -forms on  $X$ .
- $\mathcal{D}(X) = \bigoplus_{p,q} \mathcal{D}^{(p,q)}(X)$  algebra of double forms on  $X$ .

For example,  $g \in \mathcal{M}(X) \implies g \in \mathcal{C}^1(X)$  and  $R_g \in \mathcal{C}^2(X)$ .

Taking product with  $g$  defines a map

$$g : \mathcal{D}^{(p-1,q-1)}(X) \rightarrow \mathcal{D}^{(p,q)}(X)$$

whose (pointwise) adjoint is the *contraction map*  $c_g$ .

# The formalism of double forms (Nomizu-Kulkarni)

Let

- $\mathcal{D}^p(X)$  differential  $p$ -forms on  $X$ ;
- $\mathcal{D}^{(p,q)} = \mathcal{D}^p(X) \otimes \mathcal{D}^q(X)$  differential  $(p, q)$ -forms on  $X$ ;
- $\mathcal{C}^p(X) \subset \mathcal{D}^{(p,p)}(X)$  symmetric  $(p, p)$ -forms on  $X$ .
- $\mathcal{D}(X) = \bigoplus_{p,q} \mathcal{D}^{(p,q)}(X)$  algebra of double forms on  $X$ .

For example,  $g \in \mathcal{M}(X) \implies g \in \mathcal{C}^1(X)$  and  $R_g \in \mathcal{C}^2(X)$ .

Taking product with  $g$  defines a map

$$g : \mathcal{D}^{(p-1,q-1)}(X) \rightarrow \mathcal{D}^{(p,q)}(X)$$

whose (poinwise) adjoint is the *contraction map*  $c_g$ .

# Recovering the invariants (Labbi)

# Recovering the invariants (Labbi)

In this notation,

$$\mathrm{Ric}_g^{(2k)} = c_g^{2k-1} R_g^k, \quad S_g^{(2k)} = \frac{1}{(2k)!} c_g^{2k} R_g^k$$

# Recovering the invariants (Labbi)

In this notation,

$$\operatorname{Ric}_g^{(2k)} = c_g^{2k-1} R_g^k, \quad S_g^{(2k)} = \frac{1}{(2k)!} c_g^{2k} R_g^k$$

so that

$$\dot{\operatorname{Ric}}_g^{(2k)} h = k c_g^{2k-1} R_g^{k-1} \dot{R}_g h + (2k-1) c_g^{2k-2} \dot{c}_g h R_g^k,$$

# Recovering the invariants (Labbi)

In this notation,

$$\operatorname{Ric}_g^{(2k)} = c_g^{2k-1} R_g^k, \quad \mathcal{S}_g^{(2k)} = \frac{1}{(2k)!} c_g^{2k} R_g^k$$

so that

$$\dot{\operatorname{Ric}}_g^{(2k)} h = k c_g^{2k-1} R_g^{k-1} \dot{R}_g h + (2k-1) c_g^{2k-2} \dot{c}_g h R_g^k,$$

- Thus, the symbol of  $\mathcal{L}_g^{(2k)}$  is determined by the first term in the righthand side above. If  $k = 1$  the symbol is first order in  $g$ . On the other hand, if  $k \geq 2$ , the symbol depends on  $R$ !



# Recovering the invariants (Labbi)

In this notation,

$$\operatorname{Ric}_g^{(2k)} = c_g^{2k-1} R_g^k, \quad \mathcal{S}_g^{(2k)} = \frac{1}{(2k)!} c_g^{2k} R_g^k$$

so that

$$\dot{\operatorname{Ric}}_g^{(2k)} h = k c_g^{2k-1} R_g^{k-1} \dot{R}_g h + (2k-1) c_g^{2k-2} \dot{c}_g h R_g^k,$$

- Thus, the symbol of  $\mathcal{L}_g^{(2k)}$  is determined by the first term in the righthand side above. If  $k = 1$  the symbol is first order in  $g$ . On the other hand, if  $k \geq 2$ , the symbol depends on  $R$ !

# The proofs (sketch)

# The proofs (sketch)

If  $g = g_\mu$  has constant sectional curvature  $\mu$  then  $R_g = \frac{\mu}{2}g^2$  and a computation shows that

$$\mathcal{L}_g^{(2k)} \Big|_{\mathcal{I}_g} = (n - 2k) C_{n,k} \mu^{k-1} \left( \frac{1}{2} \nabla^* \nabla + \mu \right).$$

## The proofs (sketch)

If  $g = g_\mu$  has constant sectional curvature  $\mu$  then  $R_g = \frac{\mu}{2}g^2$  and a computation shows that

$$\mathcal{L}_g^{(2k)} \Big|_{\mathcal{I}_g} = (n - 2k) C_{n,k} \mu^{k-1} \left( \frac{1}{2} \nabla^* \nabla + \mu \right).$$

Since  $\nabla^* \nabla$ , the Bochner Laplacian, is elliptic and nonnegative, this proves our first theorem. The second theorem follows from a local structure theorem that implies in particular that infinitesimal rigidity implies rigidity.

## The proofs (sketch)

If  $g = g_\mu$  has constant sectional curvature  $\mu$  then  $R_g = \frac{\mu}{2}g^2$  and a computation shows that

$$\mathcal{L}_g^{(2k)} \Big|_{\mathcal{I}_g} = (n - 2k) C_{n,k} \mu^{k-1} \left( \frac{1}{2} \nabla^* \nabla + \mu \right).$$

Since  $\nabla^* \nabla$ , the Bochner Laplacian, is elliptic and nonnegative, this proves our first theorem. The second theorem follows from a local structure theorem that implies in particular that infinitesimal rigidity implies rigidity.

Another result: prescribing  $\mathcal{S}^{(2k)}$

## Another result: prescribing $\mathcal{S}^{(2k)}$

Recall: there are well-known *topological* obstructions to the existence of metrics with  $\mathcal{S}^{(2)} > 0$ . Nevertheless, the following theorem has been established (Kazdan-Warner):

## Another result: prescribing $\mathcal{S}^{(2k)}$

Recall: there are well-known *topological* obstructions to the existence of metrics with  $\mathcal{S}^{(2)} > 0$ . Nevertheless, the following theorem has been established (Kazdan-Warner):

### Theorem

*Any function that is negative somewhere (on  $X$  compact) is the scalar curvature function of some metric.*



## Another result: prescribing $\mathcal{S}^{(2k)}$

Recall: there are well-known *topological* obstructions to the existence of metrics with  $\mathcal{S}^{(2)} > 0$ . Nevertheless, the following theorem has been established (Kazdan-Warner):

### Theorem

*Any function that is negative somewhere (on  $X$  compact) is the scalar curvature function of some metric.*

### Theorem

*Let  $(X, g_\mu)$  be a compact non-flat space form not isometric to the round sphere and let  $f$  be a function on  $X$  that has the same sign as  $\mathcal{S}_{g_\mu}^{(2k)} \approx \mu^k$  somewhere. Then  $f$  is the  $2k$ -Gauss-Bonnet curvature of some metric on  $X$ .*