

Partially conformal vector fields

Oscar Palmas (Joint work with A. G. Colares)

V International Meeting on Lorentzian Geometry
2009

Abstract

We define the partially conformal vector fields and give a characterization of Lorentzian manifolds which admit this kind of fields as doubly warped products with two 1-dimensional factors.

These fields can be associated to foliations by $(n - 1)$ -umbilical hypersurfaces. As a particular case in Lorentzian space forms we have foliations by rotation hypersurfaces.

A natural question in this context is whether a given $(n - 1)$ -umbilical hypersurface is a leaf of such a foliation. We also give partial answers to this question.

[Introduction](#)[Definition and
example](#)[Basic properties](#)[Relation with
warped products](#)[Conditions for a
hypersurface to be
 \$\(n - 1\)\$ -umbilical](#)[Some references](#)

Abstract

We define the partially conformal vector fields and give a characterization of Lorentzian manifolds which admit this kind of fields as doubly warped products with two 1-dimensional factors.

These fields can be associated to foliations by $(n - 1)$ -umbilical hypersurfaces. As a particular case in Lorentzian space forms we have foliations by rotation hypersurfaces.

A natural question in this context is whether a given $(n - 1)$ -umbilical hypersurface is a leaf of such a foliation. We also give partial answers to this question.

[Introduction](#)[Definition and
example](#)[Basic properties](#)[Relation with
warped products](#)[Conditions for a
hypersurface to be
 \$\(n - 1\)\$ -umbilical](#)[Some references](#)

Abstract

We define the partially conformal vector fields and give a characterization of Lorentzian manifolds which admit this kind of fields as doubly warped products with two 1-dimensional factors.

These fields can be associated to foliations by $(n - 1)$ -umbilical hypersurfaces. As a particular case in Lorentzian space forms we have foliations by rotation hypersurfaces.

A natural question in this context is whether a given $(n - 1)$ -umbilical hypersurface is a leaf of such a foliation. We also give partial answers to this question.

[Introduction](#)[Definition and
example](#)[Basic properties](#)[Relation with
warped products](#)[Conditions for a
hypersurface to be
 \$\(n - 1\)\$ -umbilical](#)[Some references](#)

General aim

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

To establish some uniqueness theorems for constant mean curvature (or constant r -th mean curvature H_r) hypersurfaces of some spacetimes...

at least when the ambient spacetime has a lot of these hypersurfaces.

General aim

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

To establish some uniqueness theorems for constant mean curvature (or constant r -th mean curvature H_r) hypersurfaces of some spacetimes...
at least when the ambient spacetime has a lot of these hypersurfaces.

Previous works

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

As we may see, our general aim has a close relation with some characterization of totally umbilical hypersurfaces in spacetimes. In particular, we recall that

Alías-Romero-Sánchez and Montiel considered this problem in spacetimes satisfying the timelike convergence condition (Ricci curvature non negative on timelike [and null] directions) as well as those satisfying the weaker null convergence condition (Ricci curvature ≥ 0 along null directions) and CMC hypersurfaces.

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

As we may see, our general aim has a close relation with some characterization of totally umbilical hypersurfaces in spacetimes. In particular, we recall that

Alías-Romero-Sánchez and Montiel considered this problem in spacetimes satisfying the timelike convergence condition (Ricci curvature non negative on timelike [and null] directions) as well as those satisfying the weaker null convergence condition (Ricci curvature ≥ 0 along null directions) and CMC hypersurfaces.

Previous works

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

They studied the class of ambient spacetimes \bar{M}_1^{n+1} admitting a *closed conformal* timelike vector field K , which satisfies

$$\bar{\nabla}_u K = \phi u, \quad u \in TM,$$

for some real function ϕ .

Our way of generalization

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

As it is well-known, already in Riemannian space forms there are many hypersurfaces with constant mean curvature (or constant H_r) which are not totally umbilical; for example, rotational hypersurfaces in space forms.

In fact, Lorentzian space forms also admit plenty of rotational hypersurfaces, which are examples of $(n - 1)$ -umbilical hypersurfaces.

So our aim is to extend these results to this setting, namely, to a class of ambient spacetimes admitting a foliation by $(n - 1)$ -umbilical hypersurfaces.

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n - 1)$ -umbilical

Some references

Our way of generalization

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

As it is well-known, already in Riemannian space forms there are many hypersurfaces with constant mean curvature (or constant H_r) which are not totally umbilical; for example, rotational hypersurfaces in space forms.

In fact, Lorentzian space forms also admit plenty of rotational hypersurfaces, which are examples of $(n - 1)$ -umbilical hypersurfaces.

So our aim is to extend these results to this setting, namely, to a class of ambient spacetimes admitting a foliation by $(n - 1)$ -umbilical hypersurfaces.

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n - 1)$ -umbilical

Some references

Our way of generalization

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

As it is well-known, already in Riemannian space forms there are many hypersurfaces with constant mean curvature (or constant H_r) which are not totally umbilical; for example, rotational hypersurfaces in space forms.

In fact, Lorentzian space forms also admit plenty of rotational hypersurfaces, which are examples of $(n - 1)$ -umbilical hypersurfaces.

So our aim is to extend these results to this setting, namely, to a class of ambient spacetimes admitting a foliation by $(n - 1)$ -umbilical hypersurfaces.

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n - 1)$ -umbilical

Some references

Our questions

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

In particular, we attacked the following questions:

- ▶ Given a spacetime \bar{M}_1^{n+1} , under which conditions it may be foliated by $(n - 1)$ -umbilical hypersurfaces?
- ▶ Given such a foliation, which conditions imply that a hypersurface with CMC (or constant H_r) is $(n - 1)$ -umbilical or, in particular, a leaf of the aforementioned foliation?

Our questions

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

In particular, we attacked the following questions:

- ▶ Given a spacetime \bar{M}_1^{n+1} , under which conditions it may be foliated by $(n - 1)$ -umbilical hypersurfaces?
- ▶ Given such a foliation, which conditions imply that a hypersurface with CMC (or constant H_r) is $(n - 1)$ -umbilical or, in particular, a leaf of the aforementioned foliation?

Our questions

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

In particular, we attacked the following questions:

- ▶ Given a spacetime \bar{M}_1^{n+1} , under which conditions it may be foliated by $(n - 1)$ -umbilical hypersurfaces?
- ▶ Given such a foliation, which conditions imply that a hypersurface with CMC (or constant H_r) is $(n - 1)$ -umbilical or, in particular, a leaf of the aforementioned foliation?

An example

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

First we give an example of foliation of (open sets of) the de Sitter space. Recall that the scalar product in \mathbb{R}_1^{n+2} is given by

$$\langle u, v \rangle = -u_0 v_0 + u_1 v_1 + \cdots + u_{n+1} v_{n+1}$$


and that the de Sitter space is defined as

$$\mathbb{S}_1^{n+1} = \{ u \in \mathbb{R}_1^{n+2} \mid |u|^2 = 1 \}$$

Hyperbolic cylinders

$$M = \{p \in \mathbb{S}_1^{n+1} \mid -x_0^2 + x_1^2 = -\sinh^2 r\},$$

where $r > 0$ and $n > 2$. Varying r we obtain a foliation, shown in the figure.



Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

Closed partially conformal vector fields

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

Definition

Let $(\bar{M}_1^{n+1}, \bar{\nabla})$ be a spacetime and K be a timelike vector field defined on \bar{M} . We say that K is *closed, partially conformal* in \bar{M} if and only if there is a vector field $W \in \mathfrak{X}(\bar{M})$ everywhere orthogonal to K such that

$$\bar{\nabla}_X K = \phi X \text{ for } X \in W^\perp \quad \text{and} \quad \bar{\nabla}_W K = \psi W$$

for some functions $\phi, \psi : \bar{M} \rightarrow \mathbb{R}$. W is called the vector field *associated* to K .

Example in \mathbb{S}_1^{n+1}

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

To obtain the foliation by hyperbolic cylinders, we take the unit vector field N normal to each of these hypersurfaces

$$N(p) = \frac{1}{\sinh \rho \cosh \rho} (x_0, \dots, x_{n-1}, 0, 0) + (\tanh \rho) p$$

and prove that, if e_0, \dots, e_{n+1} is the canonical basis of \mathbb{R}_1^{n+2} , then

$$\nabla_{e_0+e_i} N = (\coth \rho)(e_0 + e_i), \quad i = 1, \dots, n-1,$$

and

$$\nabla_{e_n-e_{n+1}} N = (\tanh \rho)(e_n - e_{n+1}).$$

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n-1$)-umbilical

Some references

Example in \mathbb{S}_1^{n+1}

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

To obtain the foliation by hyperbolic cylinders, we take the unit vector field N normal to each of these hypersurfaces

$$N(p) = \frac{1}{\sinh \rho \cosh \rho} (x_0, \dots, x_{n-1}, 0, 0) + (\tanh \rho) p$$

and prove that, if e_0, \dots, e_{n+1} is the canonical basis of \mathbb{R}_1^{n+2} , then

$$\nabla_{e_0+e_i} N = (\coth \rho)(e_0 + e_i), \quad i = 1, \dots, n-1,$$

and

$$\nabla_{e_n-e_{n+1}} N = (\tanh \rho)(e_n - e_{n+1}).$$

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n-1$)-umbilical

Some references

Basic properties

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Take

\bar{M}_1^{n+1} : a spacetime possessing a closed partially conformal timelike vector field K ,

and

K^\perp : the distribution defined by taking the orthogonal complement of K .

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

Theorem

K^\perp is involutive and each leaf of the corresponding foliation \mathcal{K}^\perp is a $(n - 1)$ -umbilical hypersurface with $n - 1$ equal and *constant* principal curvatures.

Basic properties

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Take

\bar{M}_1^{n+1} : a spacetime possessing a closed partially
conformal timelike vector field K ,

and

K^\perp : the distribution defined by taking the orthogonal
complement of K .

Theorem

K^\perp is involutive and each leaf of the corresponding
foliation \mathcal{K}^\perp is a $(n - 1)$ -umbilical hypersurface with $n - 1$
equal and **constant** principal curvatures.

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n - 1)$ -umbilical

Some references

Relation with warped products

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Note in the examples that these open subsets may be written as a warped product of the form $-I \times_f F \times J$, where I, J are intervals.

Theorem

Let \bar{M}_1^{n+1} be a spacetime.

- ▶ If $\bar{M} = -I \times_{f_1} F \times_{f_2} J$, then $K = f_1 \partial_0$ is a closed partially conformal timelike vector field defined on \bar{M} (with $W = \frac{1}{f_2} \partial_{n+1}$).
- ▶ If \bar{M} admits a closed partially conformal timelike vector field K and the vector field W is parallel, then \bar{M} can be expressed locally as $-I \times_f F \times J$.

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

Relation with warped products

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Note in the examples that these open subsets may be written as a warped product of the form $-I \times_f F \times J$, where I, J are intervals.

Theorem

Let \bar{M}_1^{n+1} be a spacetime.

- ▶ If $\bar{M} = -I \times_{f_1} F \times_{f_2} J$, then $K = f_1 \partial_0$ is a closed partially conformal timelike vector field defined on \bar{M} (with $W = \frac{1}{f_2} \partial_{n+1}$).
- ▶ If \bar{M} admits a closed partially conformal timelike vector field K and the vector field W is parallel, then \bar{M} can be expressed locally as $-I \times_f F \times J$.

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

Relation with warped products

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Note in the examples that these open subsets may be written as a warped product of the form $-I \times_f F \times J$, where I, J are intervals.

Theorem

Let \bar{M}_1^{n+1} be a spacetime.

- ▶ If $\bar{M} = -I \times_{f_1} F \times_{f_2} J$, then $K = f_1 \partial_0$ is a closed partially conformal timelike vector field defined on \bar{M} (with $W = \frac{1}{f_2} \partial_{n+1}$).
- ▶ If \bar{M} admits a closed partially conformal timelike vector field K and the vector field W is parallel, then \bar{M} can be expressed locally as $-I \times_f F \times J$.

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

Relation with warped products

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Note in the examples that these open subsets may be written as a warped product of the form $-I \times_f F \times J$, where I, J are intervals.

Theorem

Let \bar{M}_1^{n+1} be a spacetime.

- ▶ If $\bar{M} = -I \times_{f_1} F \times_{f_2} J$, then $K = f_1 \partial_0$ is a closed partially conformal timelike vector field defined on \bar{M} (with $W = \frac{1}{f_2} \partial_{n+1}$).
- ▶ If \bar{M} admits a closed partially conformal timelike vector field K and the vector field W is parallel, then \bar{M} can be expressed locally as $-I \times_f F \times J$.

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

Is a given hypersurface a leaf?

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n - 1)$ -umbilical

Some references

Finally, we give conditions for a hypersurface with constant r -th mean curvature H_r to be a leaf of the foliation determined by a closed partially conformal vector field, thus being $(n - 1)$ -umbilical.

Our Theorem, first part

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

Theorem

*Let $\bar{M}^{n+1} = -I \times_f F \times J$ be a spacetime with F compact.
Suppose that the following strong null convergence
condition holds:*

$$K_F > \sup(ff'' - f'^2),$$

*where K_F stands for the (not necessarily constant)
sectional curvature of F .*

Our Theorem, first part, continued

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Let $M \subset \bar{M}$ be a spacelike hypersurface contained in a region $-[t_1, t_2] \times_f F \times J$ where f' does not vanish, and satisfying the following conditions:

1. M is everywhere transverse to K ;
2. Denoting $\bar{M}_s = -I \times_f F \times \{s\}$ and $M_s = M \cap \bar{M}_s$, we have that M_s is compact and has constant r -th mean curvature H_r in \bar{M}_s for $1 \leq r \leq n$.
3. If \tilde{N} denotes a unit timelike vector field everywhere normal to M , there are constants a, b such that $\tilde{N} = aK + bW$.

Then M is $(n-1)$ -umbilical in \bar{M} .

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n-1)$ -umbilical

Some references

Our Theorem, first part, continued

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Let $M \subset \bar{M}$ be a spacelike hypersurface contained in a region $-[t_1, t_2] \times_f F \times J$ where f' does not vanish, and satisfying the following conditions:

1. M is everywhere transverse to K ;
2. Denoting $\bar{M}_s = -I \times_f F \times \{s\}$ and $M_s = M \cap \bar{M}_s$, we have that M_s is compact and has constant r -th mean curvature H_r in \bar{M}_s for $1 \leq r \leq n$.
3. If \tilde{N} denotes a unit timelike vector field everywhere normal to M , there are constants a, b such that $\tilde{N} = aK + bW$.

Then M is $(n-1)$ -umbilical in \bar{M} .

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n-1)$ -umbilical

Some references

Our Theorem, first part, continued

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Let $M \subset \bar{M}$ be a spacelike hypersurface contained in a region $-[t_1, t_2] \times_f F \times J$ where f' does not vanish, and satisfying the following conditions:

1. M is everywhere transverse to K ;
2. Denoting $\bar{M}_s = -I \times_f F \times \{s\}$ and $M_s = M \cap \bar{M}_s$, we have that M_s is compact and has constant r -th mean curvature H_r in \bar{M}_s for $1 \leq r \leq n$.
3. If \tilde{N} denotes a unit timelike vector field everywhere normal to M , there are constants a, b such that $\tilde{N} = aK + bW$.

Then M is $(n-1)$ -umbilical in \bar{M} .

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n-1)$ -umbilical

Some references

Our Theorem, first part, continued

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Let $M \subset \bar{M}$ be a spacelike hypersurface contained in a region $-[t_1, t_2] \times_f F \times J$ where f' does not vanish, and satisfying the following conditions:

1. M is everywhere transverse to K ;
2. Denoting $\bar{M}_s = -I \times_f F \times \{s\}$ and $M_s = M \cap \bar{M}_s$, we have that M_s is compact and has constant r -th mean curvature H_r in \bar{M}_s for $1 \leq r \leq n$.
3. If \tilde{N} denotes a unit timelike vector field everywhere normal to M , there are constants a, b such that $\tilde{N} = aK + bW$.

Then M is $(n-1)$ -umbilical in \bar{M} .

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n-1)$ -umbilical

Some references

Our Theorem, second part

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n - 1)$ -umbilical

Some references

Moreover, if

4. M has constant r -th mean curvature H_r , $1 \leq r \leq n$;
5. The leaves of the foliation determined by K have constant r -th mean curvature; and
6. There is a s_0 and a point p of M_{s_0} such that $\tilde{N}(p) = K(p)$ and locally, M lies above (or below) the leaf of \mathcal{K}^\perp passing through p ,

Then M is a **leaf** of the foliation determined by K .

Our Theorem, second part

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n - 1)$ -umbilical

Some references

Moreover, if

4. M has constant r -th mean curvature H_r , $1 \leq r \leq n$;
5. The leaves of the foliation determined by K have constant r -th mean curvature; and
6. There is a s_0 and a point p of M_{s_0} such that $\tilde{N}(p) = K(p)$ and locally, M lies above (or below) the leaf of \mathcal{K}^\perp passing through p ,

Then M is **a leaf** of the foliation determined by K .

Our Theorem, second part

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
($n - 1$)-umbilical

Some references

Moreover, if

4. M has constant r -th mean curvature H_r , $1 \leq r \leq n$;
5. The leaves of the foliation determined by K have constant r -th mean curvature; and
6. There is a s_0 and a point p of M_{s_0} such that $\tilde{N}(p) = K(p)$ and locally, M lies above (or below) the leaf of \mathcal{K}^\perp passing through p ,

Then M is a leaf of the foliation determined by K .

Our Theorem, second part

Partially conformal
vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Introduction

Definition and
example

Basic properties

Relation with
warped products

Conditions for a
hypersurface to be
 $(n - 1)$ -umbilical

Some references

Moreover, if







4. M has constant r -th mean curvature H_r , $1 \leq r \leq n$;
5. The leaves of the foliation determined by K have constant r -th mean curvature; and
6. There is a s_0 and a point p of M_{s_0} such that $\tilde{N}(p) = K(p)$ and locally, M lies above (or below) the leaf of \mathcal{K}^\perp passing through p ,

Then M is **a leaf** of the foliation determined by K .

Partially conformal vector fields

Oscar Palmas
(Joint work with
A. G. Colares)

Some references

-  Alías, L. J. and A. G. Colares, Math. Proc. of the Cambridge Phil. Soc. **143** (3), 2007, 703–729.
-  Alías, L. J., A. Romero and M. Sánchez, Gen. Relat. Grav. **27**, 1995, 71–84.
-  Alías, L. J., A. Romero and M. Sánchez, Tôhoku Math. J. **49**, 1997, 337–345.
-  do Carmo, M. P. and M. Dajczer, Trans. Amer. Math. Soc. **277** (2), 1983, 685–709.
-  Fontenele, F. and S. L. Silva, Illinois Jour. of Math. **45** (1), 2001, 213–228.
-  Montiel, S., Math. Ann. **314**, 1999, 529–553.

Partially conformal vector fields

Oscar Palmas (Joint work with A. G. Colares)

V International Meeting on Lorentzian Geometry
2009

Grazie!