

Surfaces in exotic AdS spaces

Jorge H. S. Lira

Universidade Federal do Ceará
Fortaleza - Brazil

V International Meeting on Lorentzian Geometry
July 2009

Joint work in progress with Luis Alías (Universidad de Murcia, Spain) and J. Hinojosa (UFRPE, Brazil)

It concerns surfaces immersed into three-dimensional homogeneous Lorentzian geometries.

In this sense, it is worth to mention some related contributions by I. Taimanov, B. Daniel, F. Mercuri, P. Piccione and collaborators.

Joint work in progress with Luis Alías (Universidad de Murcia, Spain) and J. Hinojosa (UFRPE, Brazil)

It concerns surfaces immersed into three-dimensional homogeneous Lorentzian geometries.

In this sense, it is worth to mention some related contributions by I. Taimanov, B. Daniel, F. Mercuri, P. Piccione and collaborators.

Joint work in progress with Luis Alías (Universidad de Murcia, Spain) and J. Hinojosa (UFRPE, Brazil)

It concerns surfaces immersed into three-dimensional homogeneous Lorentzian geometries.

In this sense, it is worth to mention some related contributions by I. Taimanov, B. Daniel, F. Mercuri, P. Piccione and collaborators.

Joint work in progress with Luis Alías (Universidad de Murcia, Spain) and J. Hinojosa (UFRPE, Brazil)

It concerns surfaces immersed into three-dimensional homogeneous Lorentzian geometries.

In this sense, it is worth to mention some related contributions by I. Taimanov, B. Daniel, F. Mercuri, P. Piccione and collaborators.

AdS spaces

Anti de Sitter 3-space (AdS) is the universal cover of the hyperquadric M in \mathbb{R}^4 defined by

AdS spaces

Anti de Sitter 3-space (AdS) is the universal cover of the hyperquadric M in \mathbb{R}^4 defined by

$$x^2 + y^2 - u^2 - v^2 = 1$$

AdS spaces

Anti de Sitter 3-space (AdS) is the universal cover of the hyperquadric M in \mathbb{R}^4 defined by

$$x^2 + y^2 - u^2 - v^2 = 1$$

We identify

AdS spaces

Anti de Sitter 3-space (AdS) is the universal cover of the hyperquadric M in \mathbb{R}^4 defined by

$$x^2 + y^2 - u^2 - v^2 = 1$$

We identify

$$M \sim SU_{1,1} = \left\{ Z = \begin{pmatrix} z & w \\ \bar{w} & \bar{z} \end{pmatrix} \in GL(2, \mathbb{C}) : |z|^2 - |w|^2 = 1 \right\}$$

AdS spaces

Anti de Sitter 3-space (AdS) is the universal cover of the hyperquadric M in \mathbb{R}^4 defined by

$$x^2 + y^2 - u^2 - v^2 = 1$$

We identify

$$M \sim SU_{1,1} = \left\{ Z = \begin{pmatrix} z & w \\ \bar{w} & \bar{z} \end{pmatrix} \in GL(2, \mathbb{C}) : |z|^2 - |w|^2 = 1 \right\}$$

through the correspondence

AdS spaces

Anti de Sitter 3-space (AdS) is the universal cover of the hyperquadric M in \mathbb{R}^4 defined by

$$x^2 + y^2 - u^2 - v^2 = 1$$

We identify

$$M \sim SU_{1,1} = \left\{ Z = \begin{pmatrix} z & w \\ \bar{w} & \bar{z} \end{pmatrix} \in GL(2, \mathbb{C}) : |z|^2 - |w|^2 = 1 \right\}$$

through the correspondence

$$(x, y, u, v) \in \mathbb{R}^4 \mapsto (z = x + iy, w = u + iv) \in \mathbb{C}^2.$$

AdS spaces

Anti de Sitter 3-space (AdS) is the universal cover of the hyperquadric M in \mathbb{R}^4 defined by

$$x^2 + y^2 - u^2 - v^2 = 1$$

We identify

$$M \sim SU_{1,1} = \left\{ Z = \begin{pmatrix} z & w \\ \bar{w} & \bar{z} \end{pmatrix} \in GL(2, \mathbb{C}) : |z|^2 - |w|^2 = 1 \right\}$$

through the correspondence

$$(x, y, u, v) \in \mathbb{R}^4 \mapsto (z = x + iy, w = u + iv) \in \mathbb{C}^2.$$

Left invariant frames

The Lie algebra $\mathfrak{su}_{1,1}$ of $SU_{1,1}$ is spanned by the vectors

Left invariant frames

The Lie algebra $\mathfrak{su}_{1,1}$ of $SU_{1,1}$ is spanned by the vectors

$$\varsigma_1 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad \varsigma_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \varsigma_3 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \quad (1)$$

Left invariant frames

The Lie algebra $\mathfrak{su}_{1,1}$ of $SU_{1,1}$ is spanned by the vectors

$$\varsigma_1 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad \varsigma_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \varsigma_3 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \quad (1)$$

which satisfy split-quaternionic relations

Left invariant frames

The Lie algebra $\mathfrak{su}_{1,1}$ of $SU_{1,1}$ is spanned by the vectors

$$\varsigma_1 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad \varsigma_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \varsigma_3 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \quad (1)$$

which satisfy split-quaternionic relations

$$[\varsigma_1, \varsigma_2] = 2\varsigma_3, \quad [\varsigma_2, \varsigma_3] = -2\varsigma_1, \quad [\varsigma_3, \varsigma_1] = 2\varsigma_2. \quad (2)$$

Left invariant frames

The Lie algebra $\mathfrak{su}_{1,1}$ of $SU_{1,1}$ is spanned by the vectors

$$\varsigma_1 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad \varsigma_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \varsigma_3 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \quad (1)$$

which satisfy split-quaternionic relations

$$[\varsigma_1, \varsigma_2] = 2\varsigma_3, \quad [\varsigma_2, \varsigma_3] = -2\varsigma_1, \quad [\varsigma_3, \varsigma_1] = 2\varsigma_2. \quad (2)$$

Their left-invariant translations in $SU_{1,1}$ are

Left invariant frames

The Lie algebra $\mathfrak{su}_{1,1}$ of $SU_{1,1}$ is spanned by the vectors

$$\varsigma_1 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad \varsigma_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \varsigma_3 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \quad (1)$$

which satisfy split-quaternionic relations

$$[\varsigma_1, \varsigma_2] = 2\varsigma_3, \quad [\varsigma_2, \varsigma_3] = -2\varsigma_1, \quad [\varsigma_3, \varsigma_1] = 2\varsigma_2. \quad (2)$$

Their left-invariant translations in $SU_{1,1}$ are

$$Z \in SU_{1,1} \mapsto Z_{\varsigma_a} \in T_Z SU_{1,1}, \quad a = 1, 2, 3. \quad (3)$$

Why exotics?

Given $\tau \in \mathbb{R}^+$, we fix a Lorentzian metric in $\mathfrak{su}_{1,1}$ so that

Why exotics?

Given $\tau \in \mathbb{R}^+$, we fix a Lorentzian metric in $\mathfrak{su}_{1,1}$ so that

$$\langle \varsigma_1, \varsigma_1 \rangle = -\tau^2, \quad \langle \varsigma_2, \varsigma_2 \rangle = \langle \varsigma_3, \varsigma_3 \rangle = 1, \quad \langle \varsigma_i, \varsigma_j \rangle = 0, \quad \text{if } i \neq j. \quad (4)$$

Why exotics?

Given $\tau \in \mathbb{R}^+$, we fix a Lorentzian metric in $\mathfrak{su}_{1,1}$ so that

$$\langle \varsigma_1, \varsigma_1 \rangle = -\tau^2, \quad \langle \varsigma_2, \varsigma_2 \rangle = \langle \varsigma_3, \varsigma_3 \rangle = 1, \quad \langle \varsigma_i, \varsigma_j \rangle = 0, \quad \text{if } i \neq j. \quad (4)$$

We then define a left-invariant Lorentzian metric $g_\tau = \langle \cdot, \cdot \rangle$ in $SU_{1,1}$ by

Why exotics?

Given $\tau \in \mathbb{R}^+$, we fix a Lorentzian metric in $\mathfrak{su}_{1,1}$ so that

$$\langle \varsigma_1, \varsigma_1 \rangle = -\tau^2, \quad \langle \varsigma_2, \varsigma_2 \rangle = \langle \varsigma_3, \varsigma_3 \rangle = 1, \quad \langle \varsigma_i, \varsigma_j \rangle = 0, \quad \text{if } i \neq j. \quad (4)$$

We then define a left-invariant Lorentzian metric $g_\tau = \langle \cdot, \cdot \rangle$ in $SU_{1,1}$ by

$$\langle Z_{\varsigma_a}, Z_{\varsigma_b} \rangle = \langle \varsigma_a, \varsigma_b \rangle, \quad (5)$$

Why exotics?

Given $\tau \in \mathbb{R}^+$, we fix a Lorentzian metric in $\mathfrak{su}_{1,1}$ so that

$$\langle \varsigma_1, \varsigma_1 \rangle = -\tau^2, \quad \langle \varsigma_2, \varsigma_2 \rangle = \langle \varsigma_3, \varsigma_3 \rangle = 1, \quad \langle \varsigma_i, \varsigma_j \rangle = 0, \quad \text{if } i \neq j. \quad (4)$$

We then define a left-invariant Lorentzian metric $g_\tau = \langle \cdot, \cdot \rangle$ in $SU_{1,1}$ by

$$\langle Z_{\varsigma_a}, Z_{\varsigma_b} \rangle = \langle \varsigma_a, \varsigma_b \rangle, \quad (5)$$

what implies that the left-invariant frame

Why exotics?

Given $\tau \in \mathbb{R}^+$, we fix a Lorentzian metric in $\mathfrak{su}_{1,1}$ so that

$$\langle \varsigma_1, \varsigma_1 \rangle = -\tau^2, \quad \langle \varsigma_2, \varsigma_2 \rangle = \langle \varsigma_3, \varsigma_3 \rangle = 1, \quad \langle \varsigma_i, \varsigma_j \rangle = 0, \quad \text{if } i \neq j. \quad (4)$$

We then define a left-invariant Lorentzian metric $g_\tau = \langle \cdot, \cdot \rangle$ in $SU_{1,1}$ by

$$\langle Z_{\varsigma_a}, Z_{\varsigma_b} \rangle = \langle \varsigma_a, \varsigma_b \rangle, \quad (5)$$

what implies that the left-invariant frame

$$E_1 = \frac{1}{\tau} Z_{\varsigma_1}, \quad E_2 = Z_{\varsigma_2}, \quad E_3 = Z_{\varsigma_3} \quad (6)$$

is g_τ -orthonormal.

$(2 + 1)$ gravity

The curvature tensor of (M, g_τ) is completely determined by its Ricci contraction whose components are

$(2 + 1)$ gravity

The curvature tensor of (M, g_τ) is completely determined by its Ricci contraction whose components are

$$\text{Ric}(E_1, E_1) = -2\tau^2,$$

$(2 + 1)$ gravity

The curvature tensor of (M, g_τ) is completely determined by its Ricci contraction whose components are

$$\text{Ric}(E_1, E_1) = -2\tau^2,$$

$$\text{Ric}(E_2, E_2) = 4 - 2\tau^2,$$

$(2 + 1)$ gravity

The curvature tensor of (M, g_τ) is completely determined by its Ricci contraction whose components are

$$\text{Ric}(E_1, E_1) = -2\tau^2,$$

$$\text{Ric}(E_2, E_2) = 4 - 2\tau^2,$$

$$\text{Ric}(E_3, E_3) = 4 - 2\tau^2,$$

$(2 + 1)$ gravity

The curvature tensor of (M, g_τ) is completely determined by its Ricci contraction whose components are

$$\text{Ric}(E_1, E_1) = -2\tau^2,$$

$$\text{Ric}(E_2, E_2) = 4 - 2\tau^2,$$

$$\text{Ric}(E_3, E_3) = 4 - 2\tau^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0.$$

$(2 + 1)$ gravity

The curvature tensor of (M, g_τ) is completely determined by its Ricci contraction whose components are

$$\text{Ric}(E_1, E_1) = -2\tau^2,$$

$$\text{Ric}(E_2, E_2) = 4 - 2\tau^2,$$

$$\text{Ric}(E_3, E_3) = 4 - 2\tau^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0.$$

Thus, the scalar curvature is given by

$(2 + 1)$ gravity

The curvature tensor of (M, g_τ) is completely determined by its Ricci contraction whose components are

$$\text{Ric}(E_1, E_1) = -2\tau^2,$$

$$\text{Ric}(E_2, E_2) = 4 - 2\tau^2,$$

$$\text{Ric}(E_3, E_3) = 4 - 2\tau^2,$$

$$\text{Ric}(E_1, E_2) = \text{Ric}(E_1, E_3) = \text{Ric}(E_2, E_3) = 0.$$

Thus, the scalar curvature is given by

$$R = 8 - 2\tau^2. \tag{7}$$



It follows that g_τ satisfies the $(2 + 1)$ -dimensional Einstein's equation

It follows that g_τ satisfies the $(2 + 1)$ -dimensional Einstein's equation

$$\text{Ric} - \frac{R}{2} g_\tau = -\tau^2 g_\tau + 4(1 - \tau^2) T, \quad (8)$$

It follows that g_τ satisfies the $(2 + 1)$ -dimensional Einstein's equation

$$Ric - \frac{R}{2} g_\tau = -\tau^2 g_\tau + 4(1 - \tau^2) T, \quad (8)$$

where the energy-momentum tensor T is given by

It follows that g_τ satisfies the $(2 + 1)$ -dimensional Einstein's equation

$$\text{Ric} - \frac{R}{2} g_\tau = -\tau^2 g_\tau + 4(1 - \tau^2) T, \quad (8)$$

where the energy-momentum tensor T is given by

$$T = \chi \otimes \chi, \quad \chi = E_1^b. \quad (9)$$

It follows that g_τ satisfies the $(2 + 1)$ -dimensional Einstein's equation

$$\text{Ric} - \frac{R}{2} g_\tau = -\tau^2 g_\tau + 4(1 - \tau^2) T, \quad (8)$$

where the energy-momentum tensor T is given by

$$T = \chi \otimes \chi, \quad \chi = E_1^\flat. \quad (9)$$

Since E_1 is a Killing vector field, T is divergence-free.

It follows that g_τ satisfies the $(2 + 1)$ -dimensional Einstein's equation

$$\text{Ric} - \frac{R}{2} g_\tau = -\tau^2 g_\tau + 4(1 - \tau^2) T, \quad (8)$$

where the energy-momentum tensor T is given by

$$T = \chi \otimes \chi, \quad \chi = E_1^\flat. \quad (9)$$

Since E_1 is a Killing vector field, T is divergence-free.

More precisely, we have

It follows that g_τ satisfies the $(2 + 1)$ -dimensional Einstein's equation

$$\text{Ric} - \frac{R}{2} g_\tau = -\tau^2 g_\tau + 4(1 - \tau^2) T, \quad (8)$$

where the energy-momentum tensor T is given by

$$T = \chi \otimes \chi, \quad \chi = E_1^\flat. \quad (9)$$

Since E_1 is a Killing vector field, T is divergence-free.

More precisely, we have

$$\nabla_V E_1 = \tau V \times E_1, \quad V \in \Gamma(TM). \quad (10)$$

It follows that g_τ satisfies the $(2 + 1)$ -dimensional Einstein's equation

$$\text{Ric} - \frac{R}{2} g_\tau = -\tau^2 g_\tau + 4(1 - \tau^2) T, \quad (8)$$

where the energy-momentum tensor T is given by

$$T = \chi \otimes \chi, \quad \chi = E_1^\flat. \quad (9)$$

Since E_1 is a Killing vector field, T is divergence-free.

More precisely, we have

$$\nabla_V E_1 = \tau V \times E_1, \quad V \in \Gamma(TM). \quad (10)$$

Thus, (8) is the Euler-Lagrange equation for extremals of the action

$$\frac{\delta}{\delta g} \int_M (R - 2\tau^2) dV_g + 2(1 - \tau^2) \frac{\delta}{\delta g} \int_M |\chi|^2 dV_g = 0. \quad (11)$$

Thus, (8) is the Euler-Lagrange equation for extremals of the action

$$\frac{\delta}{\delta g} \int_M (R - 2\tau^2) dV_g + 2(1 - \tau^2) \frac{\delta}{\delta g} \int_M |\chi|^2 dV_g = 0. \quad (11)$$

Equation (10) says that the distribution spanned by E_2, E_3 is non-integrable.

Thus, (8) is the Euler-Lagrange equation for extremals of the action

$$\frac{\delta}{\delta g} \int_M (R - 2\tau^2) dV_g + 2(1 - \tau^2) \frac{\delta}{\delta g} \int_M |\chi|^2 dV_g = 0. \quad (11)$$

Equation (10) says that the distribution spanned by E_2, E_3 is non-integrable.

This means that g_τ is a stationary non-static metric in M . We describe that distribution in terms of the Hopf fibration $\pi : SU_{1,1} \rightarrow \mathbb{H}^2$.

Hopf fibration

Hopf fibration

When $\tau = 1$, the Lie group $SU_{1,1}$ acts isometrically in $\mathfrak{su}_{1,1} = \mathbb{L}^3$ by its adjoint action

$$\mathrm{Ad}(\Phi) \cdot E = \Phi E \Phi^{-1}, \quad E \in \mathfrak{su}_{1,1}. \quad (12)$$

Hopf fibration

When $\tau = 1$, the Lie group $SU_{1,1}$ acts isometrically in $\mathfrak{su}_{1,1} = \mathbb{L}^3$ by its adjoint action

$$\mathrm{Ad}(\Phi) \cdot E = \Phi E \Phi^{-1}, \quad E \in \mathfrak{su}_{1,1}. \quad (12)$$

The stabilizer of this action at $\varsigma_1 \in \mathbb{H}^2$ is the group \mathbb{S}^1 defined by

$$\mathbb{S}^1 = \{u 1 + v \varsigma_1 : u^2 + v^2 = 1\}.$$

Hopf fibration

When $\tau = 1$, the Lie group $SU_{1,1}$ acts isometrically in $\mathfrak{su}_{1,1} = \mathbb{L}^3$ by its adjoint action

$$\mathrm{Ad}(\Phi) \cdot E = \Phi E \Phi^{-1}, \quad E \in \mathfrak{su}_{1,1}. \quad (12)$$

The stabilizer of this action at $\varsigma_1 \in \mathbb{H}^2$ is the group \mathbb{S}^1 defined by

$$\mathbb{S}^1 = \{u \, 1 + v \varsigma_1 : u^2 + v^2 = 1\}.$$

The Lie subalgebra is denoted by \mathfrak{l}

An algebraic complement of \mathfrak{l} is given by $\mathfrak{p} = \text{span}\{\varsigma_2, \varsigma_3\}$.

An algebraic complement of \mathfrak{l} is given by $\mathfrak{p} = \text{span}\{\varsigma_2, \varsigma_3\}$. Thus,

$$\mathfrak{su}_{1,1} = \mathfrak{l} \oplus \mathfrak{p}.$$

An algebraic complement of \mathfrak{l} is given by $\mathfrak{p} = \text{span}\{\varsigma_2, \varsigma_3\}$. Thus,

$$\mathfrak{su}_{1,1} = \mathfrak{l} \oplus \mathfrak{p}.$$

These are eigenspaces for the involution $\sigma : \mathfrak{su}_{1,1} \rightarrow \mathfrak{su}_{1,1}$ defined as

An algebraic complement of \mathfrak{l} is given by $\mathfrak{p} = \text{span}\{\varsigma_2, \varsigma_3\}$. Thus,

$$\mathfrak{su}_{1,1} = \mathfrak{l} \oplus \mathfrak{p}.$$

These are eigenspaces for the involution $\sigma : \mathfrak{su}_{1,1} \rightarrow \mathfrak{su}_{1,1}$ defined as

$$\sigma(E) = \text{Ad}(\varsigma_1) \cdot E$$

An algebraic complement of \mathfrak{l} is given by $\mathfrak{p} = \text{span}\{\varsigma_2, \varsigma_3\}$. Thus,

$$\mathfrak{su}_{1,1} = \mathfrak{l} \oplus \mathfrak{p}.$$

These are eigenspaces for the involution $\sigma : \mathfrak{su}_{1,1} \rightarrow \mathfrak{su}_{1,1}$ defined as

$$\sigma(E) = \text{Ad}(\varsigma_1) \cdot E$$

and extended to the whole group by

$$\sigma(Z) = \text{Ad}(E_1) \cdot Z.$$

An algebraic complement of \mathfrak{l} is given by $\mathfrak{p} = \text{span}\{\varsigma_2, \varsigma_3\}$. Thus,

$$\mathfrak{su}_{1,1} = \mathfrak{l} \oplus \mathfrak{p}.$$

These are eigenspaces for the involution $\sigma : \mathfrak{su}_{1,1} \rightarrow \mathfrak{su}_{1,1}$ defined as

$$\sigma(E) = \text{Ad}(\varsigma_1) \cdot E$$

and extended to the whole group by

$$\sigma(Z) = \text{Ad}(E_1) \cdot Z.$$

Then

$$Z = \begin{pmatrix} z & w \\ \bar{w} & \bar{z} \end{pmatrix} \mapsto \sigma(Z) = \begin{pmatrix} z & -w \\ -\bar{w} & \bar{z} \end{pmatrix}$$

Thusm we define $\pi : SU_{1,1} \rightarrow \mathbb{H}^2$ by

$$\pi(Z) = \text{Ad}(Z) \cdot \varsigma_1 \tag{13}$$

Thusm we define $\pi : SU_{1,1} \rightarrow \mathbb{H}^2$ by

$$\pi(Z) = \text{Ad}(Z) \cdot \varsigma_1 \tag{13}$$

Identifying points in the same orbits is equivalent to define the relation

Thusm we define $\pi : SU_{1,1} \rightarrow \mathbb{H}^2$ by

$$\pi(Z) = \text{Ad}(Z) \cdot \varsigma_1 \tag{13}$$

Identifying points in the same orbits is equivalent to define the relation

$$(z, w) \sim e^{i\theta}(z, w).$$

Some useful references (for $\tau = 1$) are

Some useful references (for $\tau = 1$) are

- M. Bañados, M. Henneaux, C. Teitelboim and J. Zanelli, *Geometry of the 2 + 1 black hole*, Phys. Rev. D **48** 4 (1993), 1506–1525.

Some useful references (for $\tau = 1$) are

- M. Bañados, M. Henneaux, C. Teitelboim and J. Zanelli, *Geometry of the 2 + 1 black hole*, Phys. Rev. D **48** 4 (1993), 1506–1525.
- P. Valtancoli, *2 + 1 gravity on Riemann surfaces with conformal gauge*, Class. Quantum Grav. **14** (1997), 1795–1809.

Surfaces in AdS spaces

Now, let Σ be a Riemannian simply-connected surface

Surfaces in AdS spaces

Now, let Σ be a Riemannian simply-connected surface and let $X : \Sigma \rightarrow M$ be an isometric immersion.

Surfaces in AdS spaces

Now, let Σ be a Riemannian simply-connected surface and let $X : \Sigma \rightarrow M$ be an isometric immersion.

In terms of local conformal parameters $z = u + iv$ on Σ the induced metric is given by

Surfaces in AdS spaces

Now, let Σ be a Riemannian simply-connected surface and let $X : \Sigma \rightarrow M$ be an isometric immersion.

In terms of local conformal parameters $z = u + iv$ on Σ the induced metric is given by

$$\langle X_z, X_z \rangle = \langle X_{\bar{z}}, X_{\bar{z}} \rangle = 0, \quad \langle X_z, X_{\bar{z}} \rangle = \frac{1}{2} e^{2\omega}. \quad (14)$$

Surfaces in AdS spaces

Now, let Σ be a Riemannian simply-connected surface and let $X : \Sigma \rightarrow M$ be an isometric immersion.

In terms of local conformal parameters $z = u + iv$ on Σ the induced metric is given by

$$\langle X_z, X_z \rangle = \langle X_{\bar{z}}, X_{\bar{z}} \rangle = 0, \quad \langle X_z, X_{\bar{z}} \rangle = \frac{1}{2} e^{2\omega}. \quad (14)$$

The coefficients of the complexified second fundamental form are

Surfaces in AdS spaces

Now, let Σ be a Riemannian simply-connected surface and let $X : \Sigma \rightarrow M$ be an isometric immersion.

In terms of local conformal parameters $z = u + iv$ on Σ the induced metric is given by

$$\langle X_z, X_z \rangle = \langle X_{\bar{z}}, X_{\bar{z}} \rangle = 0, \quad \langle X_z, X_{\bar{z}} \rangle = \frac{1}{2}e^{2\omega}. \quad (14)$$

The coefficients of the complexified second fundamental form are

$$\langle \nabla_{X_z} X_z, N \rangle = \frac{q}{2}, \quad \langle \nabla_{X_z} X_{\bar{z}}, N \rangle = -\frac{1}{2}He^{2\omega}, \quad \langle \nabla_{X_{\bar{z}}} X_{\bar{z}}, N \rangle = \frac{\bar{q}}{2}, \quad (15)$$

Surfaces in AdS spaces

Now, let Σ be a Riemannian simply-connected surface and let $X : \Sigma \rightarrow M$ be an isometric immersion.

In terms of local conformal parameters $z = u + iv$ on Σ the induced metric is given by

$$\langle X_z, X_z \rangle = \langle X_{\bar{z}}, X_{\bar{z}} \rangle = 0, \quad \langle X_z, X_{\bar{z}} \rangle = \frac{1}{2}e^{2\omega}. \quad (14)$$

The coefficients of the complexified second fundamental form are

$$\langle \nabla_{X_z} X_z, N \rangle = \frac{q}{2}, \quad \langle \nabla_{X_z} X_{\bar{z}}, N \rangle = -\frac{1}{2}He^{2\omega}, \quad \langle \nabla_{X_{\bar{z}}} X_{\bar{z}}, N \rangle = \frac{\bar{q}}{2}, \quad (15)$$

where H and q are the mean curvature and the Hopf differential of the immersion.

The Gauss-Weingarten equations are written in terms of the complex adapted frame $\mathcal{F} = (X_z, X_{\bar{z}}, N)$ as

The Gauss-Weingarten equations are written in terms of the complex adapted frame $\mathcal{F} = (X_z, X_{\bar{z}}, N)$ as

$$\nabla_{X_z} X_z = 2\omega_z X_z + \frac{\epsilon}{2} q N, \quad (16)$$

The Gauss-Weingarten equations are written in terms of the complex adapted frame $\mathcal{F} = (X_z, X_{\bar{z}}, N)$ as

$$\nabla_{X_z} X_z = 2\omega_z X_z + \frac{\epsilon}{2} q N, \quad (16)$$

$$\nabla_{X_z} X_{\bar{z}} = \nabla_{X_{\bar{z}}} X_{\bar{z}} = \frac{\epsilon}{2} H e^{2\omega} N, \quad (17)$$

The Gauss-Weingarten equations are written in terms of the complex adapted frame $\mathcal{F} = (X_z, X_{\bar{z}}, N)$ as

$$\nabla_{X_z} X_z = 2\omega_z X_z + \frac{\epsilon}{2} q N, \quad (16)$$

$$\nabla_{X_z} X_{\bar{z}} = \nabla_{X_{\bar{z}}} X_z = \frac{\epsilon}{2} H e^{2\omega} N, \quad (17)$$

$$\nabla_{X_{\bar{z}}} X_{\bar{z}} = 2\omega_{\bar{z}} X_{\bar{z}} + \frac{\epsilon}{2} \bar{q} N. \quad (18)$$

The Gauss-Weingarten equations are written in terms of the complex adapted frame $\mathcal{F} = (X_z, X_{\bar{z}}, N)$ as

$$\nabla_{X_z} X_z = 2\omega_z X_z + \frac{\epsilon}{2} q N, \quad (16)$$

$$\nabla_{X_z} X_{\bar{z}} = \nabla_{X_{\bar{z}}} X_z = \frac{\epsilon}{2} H e^{2\omega} N, \quad (17)$$

$$\nabla_{X_{\bar{z}}} X_{\bar{z}} = 2\omega_{\bar{z}} X_{\bar{z}} + \frac{\epsilon}{2} \bar{q} N. \quad (18)$$

We define complex-valued functions $Z^a(z, \bar{z})$, $a = 1, 2, 3$, so that

$$X_z = Z^1 E_1|_X + Z^2 E_2|_X + Z^3 E_3|_X \quad (19)$$

We define complex-valued functions $Z^a(z, \bar{z})$, $a = 1, 2, 3$, so that

$$X_z = Z^1 E_1|_X + Z^2 E_2|_X + Z^3 E_3|_X \quad (19)$$

That is,

We define complex-valued functions $Z^a(z, \bar{z})$, $a = 1, 2, 3$, so that

$$X_z = Z^1 E_1|_X + Z^2 E_2|_X + Z^3 E_3|_X \quad (19)$$

That is,

$$X^{-1}X_z = Z^1 e_1 + Z^2 e_2 + Z^3 e_3 \in \mathfrak{su}_{1,1}. \quad (20)$$

We define complex-valued functions $Z^a(z, \bar{z})$, $a = 1, 2, 3$, so that

$$X_z = Z^1 E_1|_X + Z^2 E_2|_X + Z^3 E_3|_X \quad (19)$$

That is,

$$X^{-1}X_z = Z^1 e_1 + Z^2 e_2 + Z^3 e_3 \in \mathfrak{su}_{1,1}. \quad (20)$$

Hence, (14) gives

$$-(Z^1)^2 + (Z^2)^2 + (Z^3)^2 = 0, \quad (21)$$

We define complex-valued functions $Z^a(z, \bar{z})$, $a = 1, 2, 3$, so that

$$X_z = Z^1 E_1|_X + Z^2 E_2|_X + Z^3 E_3|_X \quad (19)$$

That is,

$$X^{-1}X_z = Z^1 e_1 + Z^2 e_2 + Z^3 e_3 \in \mathfrak{su}_{1,1}. \quad (20)$$

Hence, (14) gives

$$-(Z^1)^2 + (Z^2)^2 + (Z^3)^2 = 0, \quad (21)$$

$$-|Z^1|^2 + |Z^2|^2 + |Z^3|^2 = \frac{1}{2}e^\omega. \quad (22)$$

We define complex-valued functions $Z^a(z, \bar{z})$, $a = 1, 2, 3$, so that

$$X_z = Z^1 E_1|_X + Z^2 E_2|_X + Z^3 E_3|_X \quad (19)$$

That is,

$$X^{-1} X_z = Z^1 e_1 + Z^2 e_2 + Z^3 e_3 \in \mathfrak{su}_{1,1}. \quad (20)$$

Hence, (14) gives

$$-(Z^1)^2 + (Z^2)^2 + (Z^3)^2 = 0, \quad (21)$$

$$-|Z^1|^2 + |Z^2|^2 + |Z^3|^2 = \frac{1}{2} e^\omega. \quad (22)$$

In terms of Z^a , the Gauss-Weingarten equations become a first-order system of **nine** complex-scalar equations.

Thus, if we define in Σ a $\mathfrak{su}_{1,1}$ -valued 1-form α by

Thus, if we define in Σ a $\mathfrak{su}_{1,1}$ -valued 1-form α by

$$\gamma = e_a \otimes (Z^a dz + \bar{Z}^a d\bar{z}), \quad (23)$$

Thus, if we define in Σ a $\mathfrak{su}_{1,1}$ -valued 1-form α by

$$\Upsilon = e_a \otimes (Z^a dz + \bar{Z}^a d\bar{z}), \quad (23)$$

where $Z^a = Z^a(z, \bar{z})$ satisfy the Gauss-Weingarten system,

Thus, if we define in Σ a $\mathfrak{su}_{1,1}$ -valued 1-form α by

$$\Upsilon = e_a \otimes (Z^a dz + \bar{Z}^a d\bar{z}), \quad (23)$$

where $Z^a = Z^a(z, \bar{z})$ satisfy the Gauss-Weingarten system, then there exists an isometric immersion $X : \Sigma \rightarrow M$ with prescribed extrinsic curvature

Thus, if we define in Σ a $\mathfrak{su}_{1,1}$ -valued 1-form α by

$$\Upsilon = e_a \otimes (Z^a dz + \bar{Z}^a d\bar{z}), \quad (23)$$

where $Z^a = Z^a(z, \bar{z})$ satisfy the Gauss-Weingarten system, then there exists an isometric immersion $X : \Sigma \rightarrow M$ with prescribed extrinsic curvature

$$II = \frac{q}{2} dz^2 - \frac{H}{2} dzd\bar{z} + \frac{\bar{q}}{2} d\bar{z}^2. \quad (24)$$

Thus, if we define in Σ a $\mathfrak{su}_{1,1}$ -valued 1-form α by

$$\Upsilon = e_a \otimes (Z^a dz + \bar{Z}^a d\bar{z}), \quad (23)$$

where $Z^a = Z^a(z, \bar{z})$ satisfy the Gauss-Weingarten system, then there exists an isometric immersion $X : \Sigma \rightarrow M$ with prescribed extrinsic curvature

$$II = \frac{q}{2} dz^2 - \frac{H}{2} dzd\bar{z} + \frac{\bar{q}}{2} d\bar{z}^2. \quad (24)$$

and induced metric

Thus, if we define in Σ a $\mathfrak{su}_{1,1}$ -valued 1-form α by

$$\Upsilon = e_a \otimes (Z^a dz + \bar{Z}^a d\bar{z}), \quad (23)$$

where $Z^a = Z^a(z, \bar{z})$ satisfy the Gauss-Weingarten system, then there exists an isometric immersion $X : \Sigma \rightarrow M$ with prescribed extrinsic curvature

$$II = \frac{q}{2} dz^2 - \frac{H}{2} dzd\bar{z} + \frac{\bar{q}}{2} d\bar{z}^2. \quad (24)$$

and induced metric

$$I = e^{2\omega} |dz|^2. \quad (25)$$

The immersion X is a Darboux's primitive for Υ , that is,

The immersion X is a Darboux's primitive for Υ , that is,

$$X^{-1}dX = \Upsilon \tag{26}$$

and it is locally given as an integral leaf of the distribution

$$(p, Z) \in \Sigma \times SU_{1,1} \mapsto \ker \Xi_{(p,Z)}, \tag{27}$$

The immersion X is a Darboux's primitive for Υ , that is,

$$X^{-1}\mathrm{d}X = \Upsilon \tag{26}$$

and it is locally given as an integral leaf of the distribution

$$(p, Z) \in \Sigma \times SU_{1,1} \mapsto \ker \Xi_{(p,Z)}, \tag{27}$$

where Ξ is a $\mathfrak{su}_{1,1}$ -valued 1-form defined in $\Sigma \times SU_{1,1}$ by

The immersion X is a Darboux's primitive for Υ , that is,

$$X^{-1}dX = \Upsilon \tag{26}$$

and it is locally given as an integral leaf of the distribution

$$(p, Z) \in \Sigma \times SU_{1,1} \mapsto \ker \Xi_{(p,Z)}, \tag{27}$$

where Ξ is a $\mathfrak{su}_{1,1}$ -valued 1-form defined in $\Sigma \times SU_{1,1}$ by

$$\Xi = \Omega - \Upsilon, \tag{28}$$

The immersion X is a Darboux's primitive for Υ , that is,

$$X^{-1}dX = \Upsilon \tag{26}$$

and it is locally given as an integral leaf of the distribution

$$(p, Z) \in \Sigma \times SU_{1,1} \mapsto \ker \Xi_{(p,Z)}, \tag{27}$$

where Ξ is a $\mathfrak{su}_{1,1}$ -valued 1-form defined in $\Sigma \times SU_{1,1}$ by

$$\Xi = \Omega - \Upsilon, \tag{28}$$

Ω being the Maurer-Cartan form in $SU_{1,1}$.

Generalized Weierstrass representation

Generalized Weierstrass representation

We then define a pair of functions ψ_1 and ψ_2 by

Generalized Weierstrass representation

We then define a pair of functions ψ_1 and ψ_2 by

$$\psi_1^2 = Z^2 + iZ^3 \quad \psi_2^2 = \bar{Z}^2 + i\bar{Z}^3, \quad (29)$$

Generalized Weierstrass representation

We then define a pair of functions ψ_1 and ψ_2 by

$$\psi_1^2 = Z^2 + iZ^3 \quad \psi_2^2 = \bar{Z}^2 + i\bar{Z}^3, \quad (29)$$

so that

$$Z^1 = \psi_1 \bar{\psi}_2, \quad Z^2 = \frac{1}{2}(\bar{\psi}_2^2 + \psi_1^2), \quad Z^3 = \frac{i}{2}(\bar{\psi}_2^2 - \psi_1^2). \quad (30)$$

Generalized Weierstrass representation

We then define a pair of functions ψ_1 and ψ_2 by

$$\psi_1^2 = Z^2 + iZ^3 \quad \psi_2^2 = \bar{Z}^2 + i\bar{Z}^3, \quad (29)$$

so that

$$Z^1 = \psi_1 \bar{\psi}_2, \quad Z^2 = \frac{1}{2}(\bar{\psi}_2^2 + \psi_1^2), \quad Z^3 = \frac{i}{2}(\bar{\psi}_2^2 - \psi_1^2). \quad (30)$$

Given an isometric equation $X : \Sigma \rightarrow M$ with prescribed extrinsic curvature II , these functions satisfy the system

Given an isometric equation $X : \Sigma \rightarrow M$ with prescribed extrinsic curvature II , these functions satisfy the system

$$\partial_{\bar{z}}\psi_1 = V\psi_2, \quad \partial_z\psi_2 = -U\psi_1, \quad (31)$$

Given an isometric equation $X : \Sigma \rightarrow M$ with prescribed extrinsic curvature II , these functions satisfy the system

$$\partial_{\bar{z}}\psi_1 = V\psi_2, \quad \partial_z\psi_2 = -U\psi_1, \quad (31)$$

where

Given an isometric equation $X : \Sigma \rightarrow M$ with prescribed extrinsic curvature II , these functions satisfy the system

$$\partial_{\bar{z}}\psi_1 = V\psi_2, \quad \partial_z\psi_2 = -U\psi_1, \quad (31)$$

where

$$U = -\frac{1}{2}\left(H - \frac{i}{\tau}\right)(|\psi_2|^2 - |\psi_1|^2) - \frac{i}{2}\left(\tau - \frac{1}{\tau}\right)(|\psi_2|^2 + |\psi_1|^2) \quad (32)$$

Given an isometric equation $X : \Sigma \rightarrow M$ with prescribed extrinsic curvature II , these functions satisfy the system

$$\partial_{\bar{z}}\psi_1 = V\psi_2, \quad \partial_z\psi_2 = -U\psi_1, \quad (31)$$

where

$$U = -\frac{1}{2}\left(H - \frac{i}{\tau}\right)(|\psi_2|^2 - |\psi_1|^2) - \frac{i}{2}\left(\tau - \frac{1}{\tau}\right)(|\psi_2|^2 + |\psi_1|^2) \quad (32)$$

$$V = \frac{1}{2}\left(H + \frac{i}{\tau}\right)(|\psi_2|^2 - |\psi_1|^2) + \frac{i}{2}\left(\tau - \frac{1}{\tau}\right)(|\psi_2|^2 + |\psi_1|^2). \quad (33)$$

Theorem

Let $\psi = (\psi_1\sqrt{dz}, \psi_2\sqrt{dz})$ be a field of spinors in Σ satisfying the Dirac-type equation $\mathbf{D}\psi = 0$, where

Theorem

Let $\psi = (\psi_1\sqrt{dz}, \psi_2\sqrt{d\bar{z}})$ be a field of spinors in Σ satisfying the Dirac-type equation $\mathbf{D}\psi = 0$, where

$$\mathbf{D} = \begin{pmatrix} 0 & \partial_z \\ -\partial_{\bar{z}} & 0 \end{pmatrix} + \begin{pmatrix} U & 0 \\ 0 & V \end{pmatrix}. \quad (34)$$

Theorem

Let $\psi = (\psi_1\sqrt{dz}, \psi_2\sqrt{d\bar{z}})$ be a field of spinors in Σ satisfying the Dirac-type equation $\mathbf{D}\psi = 0$, where

$$\mathbf{D} = \begin{pmatrix} 0 & \partial_z \\ -\partial_{\bar{z}} & 0 \end{pmatrix} + \begin{pmatrix} U & 0 \\ 0 & V \end{pmatrix}. \quad (34)$$

If ψ satisfies the first order system

$$\partial_z \psi_1 = \omega_z \psi_1 + i\left(\tau - \frac{1}{\tau}\right) \psi_1^2 \bar{\psi}_2 - e^{-\omega} \frac{q}{2} \psi_2, \quad (35)$$

$$\partial_z \bar{\psi}_2 = \omega_z \bar{\psi}_2 - i\left(\tau - \frac{1}{\tau}\right) \psi_1 \bar{\psi}_2^2 - e^{-\omega} \frac{q}{2} \bar{\psi}_1, \quad (36)$$

Theorem

Let $\psi = (\psi_1\sqrt{dz}, \psi_2\sqrt{d\bar{z}})$ be a field of spinors in Σ satisfying the Dirac-type equation $\mathbf{D}\psi = 0$, where

$$\mathbf{D} = \begin{pmatrix} 0 & \partial_z \\ -\partial_{\bar{z}} & 0 \end{pmatrix} + \begin{pmatrix} U & 0 \\ 0 & V \end{pmatrix}. \quad (34)$$

If ψ satisfies the first order system

$$\partial_z \psi_1 = \omega_z \psi_1 + i\left(\tau - \frac{1}{\tau}\right) \psi_1^2 \bar{\psi}_2 - e^{-\omega} \frac{q}{2} \psi_2, \quad (35)$$

$$\partial_z \bar{\psi}_2 = \omega_z \bar{\psi}_2 - i\left(\tau - \frac{1}{\tau}\right) \psi_1 \bar{\psi}_2^2 - e^{-\omega} \frac{q}{2} \bar{\psi}_1, \quad (36)$$

then there exists an immersion $X : \Sigma \rightarrow (M, g_\tau)$ with induced metric I , mean curvature H and Hopf differential qdz^2 .



In order to obtain compatibility conditions for the system of **four** first-order equations in Theorem just above, we define

In order to obtain compatibility conditions for the system of **four** first-order equations in Theorem just above, we define

$$E_1 = \underbrace{\alpha}_{\text{lapse}} N + \underbrace{\beta}_{\text{shift}}, \quad (37)$$

In order to obtain compatibility conditions for the system of **four** first-order equations in Theorem just above, we define

$$E_1 = \underbrace{\alpha}_{\text{lapse}} N + \underbrace{\beta}_{\text{shift}}, \quad (37)$$

that is,

$$\beta := -2e^{-2\omega}(\bar{\psi}_1\psi_2X_z + \psi_1\bar{\psi}_2X_{\bar{z}}) \quad (38)$$

In order to obtain compatibility conditions for the system of **four** first-order equations in Theorem just above, we define

$$E_1 = \underbrace{\alpha}_{\text{lapse}} N + \underbrace{\beta}_{\text{shift}}, \quad (37)$$

that is,

$$\beta := -2e^{-2\omega}(\bar{\psi}_1\psi_2X_z + \psi_1\bar{\psi}_2X_{\bar{z}}) \quad (38)$$

$$\alpha := -e^{-\omega}(|\psi_1|^2 + |\psi_2|^2) \quad (39)$$

In order to obtain compatibility conditions for the system of **four** first-order equations in Theorem just above, we define

$$E_1 = \underbrace{\alpha}_{\text{lapse}} N + \underbrace{\beta}_{\text{shift}}, \quad (37)$$

that is,

$$\beta := -2e^{-2\omega}(\bar{\psi}_1\psi_2X_z + \psi_1\bar{\psi}_2X_{\bar{z}}) \quad (38)$$

$$\alpha := -e^{-\omega}(|\psi_1|^2 + |\psi_2|^2) \quad (39)$$

Let J and ∇^Σ be the complex structure and Riemannian connection in Σ associated to I . We refer to the equations

$$K_{\text{int}} - K_{\text{ext}} = 4\tau\left(\tau - \frac{1}{\tau}\right)\alpha^2 + \tau^2 \quad (40)$$

Let J and ∇^Σ be the complex structure and Riemannian connection in Σ associated to I . We refer to the equations

$$K_{\text{int}} - K_{\text{ext}} = 4\tau\left(\tau - \frac{1}{\tau}\right)\alpha^2 + \tau^2 \quad (40)$$

$$(\nabla_V^\Sigma K)W - (\nabla_W^\Sigma K)V = 4\alpha\tau\left(\tau - \frac{1}{\tau}\right)I(JV, W)J\beta \quad (41)$$

Let J and ∇^Σ be the complex structure and Riemannian connection in Σ associated to I . We refer to the equations

$$K_{\text{int}} - K_{\text{ext}} = 4\tau\left(\tau - \frac{1}{\tau}\right)\alpha^2 + \tau^2 \quad (40)$$

$$(\nabla_V^\Sigma K)W - (\nabla_W^\Sigma K)V = 4\alpha\tau\left(\tau - \frac{1}{\tau}\right)I(JV, W)J\beta \quad (41)$$

$$\alpha KV + \nabla_V \beta = -\alpha\tau JV \quad (42)$$

Let J and ∇^Σ be the complex structure and Riemannian connection in Σ associated to I . We refer to the equations

$$K_{\text{int}} - K_{\text{ext}} = 4\tau\left(\tau - \frac{1}{\tau}\right)\alpha^2 + \tau^2 \quad (40)$$

$$(\nabla_V^\Sigma K)W - (\nabla_W^\Sigma K)V = 4\alpha\tau\left(\tau - \frac{1}{\tau}\right)I(JV, W)J\beta \quad (41)$$

$$\alpha KV + \nabla_V \beta = -\alpha\tau JV \quad (42)$$

$$d\alpha(V) + II(V, \beta) = \tau I(JV, \beta). \quad (43)$$

Let J and ∇^Σ be the complex structure and Riemannian connection in Σ associated to I . We refer to the equations

$$K_{\text{int}} - K_{\text{ext}} = 4\tau\left(\tau - \frac{1}{\tau}\right)\alpha^2 + \tau^2 \quad (40)$$

$$(\nabla_V^\Sigma K)W - (\nabla_W^\Sigma K)V = 4\alpha\tau\left(\tau - \frac{1}{\tau}\right)I(JV, W)J\beta \quad (41)$$

$$\alpha KV + \nabla_V \beta = -\alpha\tau JV \quad (42)$$

$$d\alpha(V) + II(V, \beta) = \tau I(JV, \beta). \quad (43)$$

as compatibility equations for I , II , α and β .

Let J and ∇^Σ be the complex structure and Riemannian connection in Σ associated to I . We refer to the equations

$$K_{\text{int}} - K_{\text{ext}} = 4\tau\left(\tau - \frac{1}{\tau}\right)\alpha^2 + \tau^2 \quad (40)$$

$$(\nabla_V^\Sigma K)W - (\nabla_W^\Sigma K)V = 4\alpha\tau\left(\tau - \frac{1}{\tau}\right)I(JV, W)J\beta \quad (41)$$

$$\alpha KV + \nabla_V \beta = -\alpha\tau JV \quad (42)$$

$$d\alpha(V) + II(V, \beta) = \tau I(JV, \beta). \quad (43)$$

as compatibility equations for I , II , α and β . Here, K is the Weingarten map $I^{-1}II$.

Theorem

Let (Σ, I) be a simply-connected Riemannian surface, $II \in \Gamma(T^*\Sigma \odot T^*\Sigma)$, $\beta \in \Gamma(T\Sigma)$ and $\alpha \in C^\infty(\Sigma, \mathbb{R})$ satisfying the compatibility equations above. Then, there exists an isometric immersion $X : \Sigma \rightarrow M$ such that

$$I(V, W) = \langle X_* V, X_* W \rangle, \quad (44)$$

$$II(V, W) = \langle \nabla_{X_* V} X_* W, N \rangle, \quad (45)$$

where $V, W \in \Gamma(T\Sigma)$ and $p \in \Sigma \mapsto N|_{X(p)} \in T_{X(p)}M$ is a normal unit vector field along X given by

$$E_1|_{X(p)} = \alpha(p)N|_{X(p)} + \beta(p). \quad (46)$$

Initial value problem

Maximal surfaces

From now on, we assume that $H = 0$.

Maximal surfaces

From now on, we assume that $H = 0$.

Maximal surfaces

From now on, we assume that $H = 0$.

The quadric

$$\mathbb{H}^2 = \{a e_1 + b e_2 + c e_3 \in \mathfrak{su}_{1,1} : -a^2 + b^2 + c^2 = 1\}$$

Maximal surfaces

From now on, we assume that $H = 0$.

The quadric

$$\mathbb{H}^2 = \{a e_1 + b e_2 + c e_3 \in \mathfrak{su}_{1,1} : -a^2 + b^2 + c^2 = 1\}$$

is the hyperboloid model in $\mathbb{L}^3 = \mathfrak{su}_{1,1}$ for the hyperbolic plane.

Maximal surfaces

From now on, we assume that $H = 0$.

The quadric

$$\mathbb{H}^2 = \{a e_1 + b e_2 + c e_3 \in \mathfrak{su}_{1,1} : -a^2 + b^2 + c^2 = 1\}$$

is the hyperboloid model in $\mathbb{L}^3 = \mathfrak{su}_{1,1}$ for the hyperbolic plane.

Denoting by \mathbb{D} the unit disc in \mathbb{C} , the stereographic projection $\pi : \mathbb{H}^2 \rightarrow \mathbb{D}$ is defined by

Maximal surfaces

From now on, we assume that $H = 0$.

The quadric

$$\mathbb{H}^2 = \{a e_1 + b e_2 + c e_3 \in \mathfrak{su}_{1,1} : -a^2 + b^2 + c^2 = 1\}$$

is the hyperboloid model in $\mathbb{L}^3 = \mathfrak{su}_{1,1}$ for the hyperbolic plane.

Denoting by \mathbb{D} the unit disc in \mathbb{C} , the stereographic projection $\pi : \mathbb{H}^2 \rightarrow \mathbb{D}$ is defined by

$$\pi(a, b, c) = \frac{b + ic}{1 + a}. \quad (47)$$

Given an isometric immersion $X : \Sigma \rightarrow M$, its Gauss map $g : \Sigma \rightarrow \mathbb{D}$ is defined by

Given an isometric immersion $X : \Sigma \rightarrow M$, its **Gauss map** $g : \Sigma \rightarrow \mathbb{D}$ is defined by

$$g(p) = \pi(X^{-1}N(p)), \quad p \in \Sigma. \quad (48)$$

Given an isometric immersion $X : \Sigma \rightarrow M$, its **Gauss map** $g : \Sigma \rightarrow \mathbb{D}$ is defined by

$$g(p) = \pi(X^{-1}N(p)), \quad p \in \Sigma. \quad (48)$$

In terms of spinors, we have

$$g = \frac{\psi_1}{\psi_2}. \quad (49)$$

Given an isometric immersion $X : \Sigma \rightarrow M$, its **Gauss map** $g : \Sigma \rightarrow \mathbb{D}$ is defined by

$$g(p) = \pi(X^{-1}N(p)), \quad p \in \Sigma. \quad (48)$$

In terms of spinors, we have

$$g = \frac{\psi_1}{\bar{\psi}_2}. \quad (49)$$

and denoting $f = \bar{\psi}_2^2$, we may recover the results above in terms of f and g

The novelty is the following one

Proposition

If the immersion $X : \Sigma \rightarrow (SU_{1,1}, g_\tau)$ is maximal, then it holds that

$$g_{z\bar{z}} + 2 \frac{2\varsigma^2 - (1 + |g|^2)}{(1 + |g|^2)^2 - 4\varsigma^2 |g|^2} \bar{g} g_z g_{\bar{z}} = 0, \quad (50)$$

where $\varsigma = 1/\tau$.

In order to state a converse for this Proposition, we denote

In order to state a converse for this Proposition, we denote

$$\mathbb{D}_\tau = \begin{cases} \{w \in \mathbb{C} : |w| < \tau_0\}, & 0 < \tau \leq 1 \\ \{w \in \mathbb{C} : |w| < 1\}, & 1 < \tau. \end{cases} \quad (51)$$

where $\tau_0 = \frac{1}{\tau}(1 - \sqrt{1 - \tau^2})$ when $\tau \in (0, 1]$.

In order to state a converse for this Proposition, we denote

$$\mathbb{D}_\tau = \begin{cases} \{w \in \mathbb{C} : |w| < \tau_0\}, & 0 < \tau \leq 1 \\ \{w \in \mathbb{C} : |w| < 1\}, & 1 < \tau. \end{cases} \quad (51)$$

where $\tau_0 = \frac{1}{\tau}(1 - \sqrt{1 - \tau^2})$ when $\tau \in (0, 1]$.

We then define the complete Riemannian metric in \mathbb{D}_τ

In order to state a converse for this Proposition, we denote

$$\mathbb{D}_\tau = \begin{cases} \{w \in \mathbb{C} : |w| < \tau_0\}, & 0 < \tau \leq 1 \\ \{w \in \mathbb{C} : |w| < 1\}, & 1 < \tau. \end{cases} \quad (51)$$

where $\tau_0 = \frac{1}{\tau}(1 - \sqrt{1 - \tau^2})$ when $\tau \in (0, 1]$.

We then define the complete Riemannian metric in \mathbb{D}_τ

$$ds_\tau^2 = \frac{|dw|^2}{(1 + |w|^2)^2 - 4s^2|w|^2}, \quad w \in \mathbb{D}_\tau. \quad (52)$$

Theorem

The Gauss map $g : \Sigma \rightarrow (\mathbb{D}_\tau, ds_\tau^2)$ of a maximal immersion $X : \Sigma \rightarrow (SU_{1,1}, g_\tau)$ is a harmonic map.

Theorem

The Gauss map $g : \Sigma \rightarrow (\mathbb{D}_\tau, ds_\tau^2)$ of a maximal immersion $X : \Sigma \rightarrow (SU_{1,1}, g_\tau)$ is a harmonic map.

We observe that if g is harmonic, all maps $g^\vartheta = e^{i\vartheta}g$, $\vartheta \in \mathbb{R}$, are also harmonic.

Theorem

The Gauss map $g : \Sigma \rightarrow (\mathbb{D}_\tau, ds_\tau^2)$ of a maximal immersion $X : \Sigma \rightarrow (SU_{1,1}, g_\tau)$ is a harmonic map.

We observe that if g is harmonic, all maps $g^\vartheta = e^{i\vartheta}g$, $\vartheta \in \mathbb{R}$, are also harmonic.

Theorem

The quadratic differential

$$\begin{aligned} Q &= (\psi_1 \partial_z \bar{\psi}_2 - \bar{\psi}_2 \partial_z \psi_1) dz^2 \\ &= \left(\frac{q}{2} - 2i\left(\tau - \frac{1}{\tau}\right) f^2 g^2 \right) dz^2 \end{aligned} \tag{53}$$

is holomorphic when $H = 0$.

The next result states that we may determine a maximal immersion of Σ into $SU_{1,1}$ starting with a harmonic map g with target in \mathbb{D}_τ .

The next result states that we may determine a maximal immersion of Σ into $SU_{1,1}$ starting with a harmonic map g with target in \mathbb{D}_τ .

Theorem

Let Σ be a simply connected Riemannian surface. Let $g : \Sigma \rightarrow (\mathbb{D}_\tau, ds_\tau^2)$ be a harmonic never holomorphic map. Given $z_0 \in \Sigma$ and $Z_0 \in SU_{1,1}$, there exists a unique conformal maximal immersion $X : \Sigma \rightarrow SU_{1,1}$ whose Gauss map is g and such that $X(z_0) = Z_0$.

The next result states that we may determine a maximal immersion of Σ into $SU_{1,1}$ starting with a harmonic map g with target in \mathbb{D}_τ .

Theorem

Let Σ be a simply connected Riemannian surface. Let $g : \Sigma \rightarrow (\mathbb{D}_\tau, ds_\tau^2)$ be a harmonic never holomorphic map. Given $z_0 \in \Sigma$ and $Z_0 \in SU_{1,1}$, there exists a unique conformal maximal immersion $X : \Sigma \rightarrow SU_{1,1}$ whose Gauss map is g and such that $X(z_0) = Z_0$.

The next result states that we may determine a maximal immersion of Σ into $SU_{1,1}$ starting with a harmonic map g with target in \mathbb{D}_τ .

Theorem

Let Σ be a simply connected Riemannian surface. Let $g : \Sigma \rightarrow (\mathbb{D}_\tau, ds_\tau^2)$ be a harmonic never holomorphic map. Given $z_0 \in \Sigma$ and $Z_0 \in SU_{1,1}$, there exists a unique conformal maximal immersion $X : \Sigma \rightarrow SU_{1,1}$ whose Gauss map is g and such that $X(z_0) = Z_0$.

The immersion obtained in the theorem above satisfies

The immersion obtained in the theorem above satisfies

$$\begin{aligned}
 -i\tau X^{-1}X_z &= \frac{2g\bar{g}_z}{(1+|g|^2)^2 - 4\varsigma^2|g|^2} \hat{e}_1 \\
 &+ \frac{(1+g^2)\bar{g}_z}{(1+|g|^2)^2 - 4\varsigma^2|g|^2} \hat{e}_2 \\
 &+ \frac{(1-g^2)\bar{g}_z}{(1+|g|^2)^2 - 4\varsigma^2|g|^2} \hat{e}_3.
 \end{aligned} \tag{54}$$

The immersion obtained in the theorem above satisfies

$$\begin{aligned}
 -i\tau X^{-1}X_z &= \frac{2g\bar{g}_z}{(1+|g|^2)^2 - 4\varsigma^2|g|^2} \hat{e}_1 \\
 &+ \frac{(1+g^2)\bar{g}_z}{(1+|g|^2)^2 - 4\varsigma^2|g|^2} \hat{e}_2 \\
 &+ \frac{(1-g^2)\bar{g}_z}{(1+|g|^2)^2 - 4\varsigma^2|g|^2} \hat{e}_3.
 \end{aligned} \tag{54}$$

Moreover, there exists a family of maximal immersions $X_\vartheta : \Sigma \rightarrow SU_{1,1}$, $\vartheta \in \mathbb{R}$, represented by the harmonic maps $g^\vartheta = e^{i\vartheta}g$, such that $X_0 = X$.

Some examples

Some examples

- minimal discs.

Some examples

- minimal discs.
- geodesics, including the radial ones.

Some examples

- minimal discs.
- geodesics, including the radial ones.
- solutions by separation of variables \rightarrow “helicoids”

Some examples

- minimal discs.
- geodesics, including the radial ones.
- solutions by separation of variables \rightarrow “helicoids”
- associated families

Some examples

- minimal discs.
- geodesics, including the radial ones.
- solutions by separation of variables \rightarrow “helicoids”
- associated families

Questions

- correspondence with CMC surfaces in some space with fibration over S^2_1 .

Some examples

- minimal discs.
- geodesics, including the radial ones.
- solutions by separation of variables \rightarrow “helicoids”
- associated families

Questions

- correspondence with CMC surfaces in some space with fibration over \mathbb{S}_1^2 .
- more intricate examples.

Further developments ($\tau = 1$)

Given a map $X : \Sigma \rightarrow SU_{1,1} \subset \mathbb{R}^{2,2}$ we lift a frame along X to

Further developments ($\tau = 1$)

Given a map $X : \Sigma \rightarrow SU_{1,1} \subset \mathbb{R}^{2,2}$ we lift a frame along X to

$$(\Phi_-, \Phi_+) \in SU_{1,1} \times SU_{1,1}$$

Further developments ($\tau = 1$)

Given a map $X : \Sigma \rightarrow SU_{1,1} \subset \mathbb{R}^{2,2}$ we lift a frame along X to

$$(\Phi_-, \Phi_+) \in SU_{1,1} \times SU_{1,1}$$

and define the $\mathfrak{su}_{1,1} \oplus \mathfrak{su}_{1,1}$ -valued form

Further developments ($\tau = 1$)

Given a map $X : \Sigma \rightarrow SU_{1,1} \subset \mathbb{R}^{2,2}$ we lift a frame along X to

$$(\Phi_-, \Phi_+) \in SU_{1,1} \times SU_{1,1}$$

and define the $\mathfrak{su}_{1,1} \oplus \mathfrak{su}_{1,1}$ -valued form

$$(\alpha_-, \alpha_+) = (\Phi_-^{-1} d\Phi_-, \Phi_+^{-1} d\Phi_+).$$

Further developments ($\tau = 1$)

Given a map $X : \Sigma \rightarrow SU_{1,1} \subset \mathbb{R}^{2,2}$ we lift a frame along X to

$$(\Phi_-, \Phi_+) \in SU_{1,1} \times SU_{1,1}$$

and define the $\mathfrak{su}_{1,1} \oplus \mathfrak{su}_{1,1}$ -valued form

$$(\alpha_-, \alpha_+) = (\Phi_-^{-1} d\Phi_-, \Phi_+^{-1} d\Phi_+).$$

Using a suitable decomposition $\mathfrak{su}_{1,1} \oplus \mathfrak{su}_{1,1} = \mathfrak{d} \oplus \mathfrak{m}$ we introduce a parameter $\lambda \in \mathbb{S}^1$ by

Further developments ($\tau = 1$)

Given a map $X : \Sigma \rightarrow SU_{1,1} \subset \mathbb{R}^{2,2}$ we lift a frame along X to

$$(\Phi_-, \Phi_+) \in SU_{1,1} \times SU_{1,1}$$

and define the $\mathfrak{su}_{1,1} \oplus \mathfrak{su}_{1,1}$ -valued form

$$(\alpha_-, \alpha_+) = (\Phi_-^{-1} d\Phi_-, \Phi_+^{-1} d\Phi_+).$$

Using a suitable decomposition $\mathfrak{su}_{1,1} \oplus \mathfrak{su}_{1,1} = \mathfrak{d} \oplus \mathfrak{m}$ we introduce a parameter $\lambda \in \mathbb{S}^1$ by

$$\alpha_{\pm}(\lambda) = \alpha_{\pm\mathfrak{d}} + \lambda^{-1} \alpha'_{\pm\mathfrak{m}} + \lambda \alpha''_{\pm\mathfrak{m}}, \quad \lambda \in \mathbb{S}^1. \quad (55)$$

It happens that

$$d\alpha(\lambda) + \frac{1}{2}[\alpha(\lambda), \alpha(\lambda)] = 0,$$

It happens that

$$d\alpha(\lambda) + \frac{1}{2}[\alpha(\lambda), \alpha(\lambda)] = 0,$$

and then given a solution

It happens that

$$d\alpha(\lambda) + \frac{1}{2}[\alpha(\lambda), \alpha(\lambda)] = 0,$$

and then given a solution

$$\Phi(\lambda) : \Sigma \rightarrow \mathrm{SU}(1,1) \times \mathrm{SU}(1,1), \quad \lambda \in \mathbb{S}^1$$

of the Maurer-Cartan equation

It happens that

$$d\alpha(\lambda) + \frac{1}{2}[\alpha(\lambda), \alpha(\lambda)] = 0,$$

and then given a solution

$$\Phi(\lambda) : \Sigma \rightarrow \mathrm{SU}(1,1) \times \mathrm{SU}(1,1), \quad \lambda \in \mathbb{S}^1$$

of the Maurer-Cartan equation

$$\Phi^{-1}d\Phi = \alpha(\lambda),$$

with components $\Phi = (\Phi_-, \Phi_+)$

It happens that

$$d\alpha(\lambda) + \frac{1}{2}[\alpha(\lambda), \alpha(\lambda)] = 0,$$

and then given a solution

$$\Phi(\lambda) : \Sigma \rightarrow \mathrm{SU}(1,1) \times \mathrm{SU}(1,1), \quad \lambda \in \mathbb{S}^1$$

of the Maurer-Cartan equation

$$\Phi^{-1}d\Phi = \alpha(\lambda),$$

with components $\Phi = (\Phi_-, \Phi_+)$ we define a family of maximal immersions $X_\lambda : \Sigma \rightarrow \mathrm{SU}_{1,1}$ by

$$X_\lambda = \Phi^+(\lambda)\Phi_-^{-1}(\lambda), \quad \lambda \in \mathbb{S}^1.$$

An isometric immersion $X : \Sigma \rightarrow SU_{1,1}$ with CMC H gives rise to a form $\vartheta \in \Lambda^1(\Sigma, \mathfrak{su}_{1,1})$ satisfying

An isometric immersion $X : \Sigma \rightarrow SU_{1,1}$ with CMC H gives rise to a form $\vartheta \in \Lambda^1(\Sigma, \mathfrak{su}_{1,1})$ satisfying

$$d * \vartheta = \frac{H}{2} [\vartheta, \vartheta]. \quad (56)$$

An isometric immersion $X : \Sigma \rightarrow SU_{1,1}$ with CMC H gives rise to a form $\vartheta \in \Lambda^1(\Sigma, \mathfrak{su}_{1,1})$ satisfying

$$d * \vartheta = \frac{H}{2} [\vartheta, \vartheta]. \quad (56)$$

$$d\vartheta = -\frac{1}{2} [\vartheta, \vartheta]. \quad (57)$$

Flat Lorentz surfaces

Theorem (–, F. Vitório)

Let Σ be a simply-connected flat Lorentz surface. An isometric immersion $X : \Sigma \rightarrow SL(2, \mathbb{R})$ whose mean curvature satisfies $H \geq 1$ may be represented by

$$X = F_1 F_2^T,$$

where $F_i : \Sigma \rightarrow SL(2, \mathbb{R})$, $i = 1, 2$, are given by

$$dF_1 = F_1 \begin{pmatrix} 0 & h \\ 1 & 0 \end{pmatrix} du \quad \text{and} \quad dF_2 = F_2 \begin{pmatrix} 0 & 1 \\ g & 0 \end{pmatrix} dv$$

for any Lorentz-holomorphic function h and Lorentz-antiholomorphic function g defined in Σ in terms of a null coordinate system (u, v)

Thank you!