

Einstein-like Walker manifolds and generalized locally symmetric spaces

Sandra Gavino Fernández



Department of Geometry and Topology
University of Santiago de Compostela

July 11th, 2009

Work in progress with Miguel Brozos Vázquez

Aim: To investigate some generalizations of locally symmetric spaces.

Different approaches:

- 1 Einstein-like manifolds (Algebraic conditions)
- 2 \mathcal{C} -spaces and \mathfrak{P} -spaces (Geometry of geodesics)
- 3 \mathfrak{D} -spaces and \mathfrak{T} -spaces (Geometry of unit circles)

Aim: To investigate some generalizations of locally symmetric spaces.

Different approaches:

- 1 Einstein-like manifolds (Algebraic conditions)
- 2 \mathcal{C} -spaces and \mathfrak{P} -spaces (Geometry of geodesics)
- 3 \mathfrak{D} -spaces and \mathfrak{T} -spaces (Geometry of unit circles)

Aim: To investigate some generalizations of locally symmetric spaces.

Different approaches:

- 1 Einstein-like manifolds (Algebraic conditions)
- 2 \mathcal{C} -spaces and \mathfrak{P} -spaces (Geometry of geodesics)
- 3 \mathfrak{D} -spaces and \mathfrak{T} -spaces (Geometry of unit circles)

Aim: To investigate some generalizations of locally symmetric spaces.

Different approaches:

- 1 Einstein-like manifolds (Algebraic conditions)
- 2 \mathcal{C} -spaces and \mathfrak{P} -spaces (Geometry of geodesics)
- 3 \mathfrak{D} -spaces and \mathfrak{T} -spaces (Geometry of unit circles)

Aim: To investigate some generalizations of locally symmetric spaces.

Different approaches:

- 1 Einstein-like manifolds (Algebraic conditions)
- 2 \mathfrak{C} -spaces and \mathfrak{P} -spaces (Geometry of geodesics)
- 3 \mathfrak{D} -spaces and \mathfrak{T} -spaces (Geometry of unit circles)

Aim: To investigate some generalizations of locally symmetric spaces.

Different approaches:

- 1 Einstein-like manifolds (Algebraic conditions)
- 2 \mathfrak{C} -spaces and \mathfrak{P} -spaces (Geometry of geodesics)
- 3 \mathfrak{D} -spaces and \mathfrak{T} -spaces (Geometry of unit circles)

Aim: To investigate some generalizations of locally symmetric spaces.

Different approaches:

- 1 Einstein-like manifolds (Algebraic conditions)
- 2 \mathfrak{C} -spaces and \mathfrak{P} -spaces (Geometry of geodesics)
- 3 \mathfrak{D} -spaces and \mathfrak{T} -spaces (Geometry of unit circles)

Three-dimensional Walker metrics

Definition

A manifold (\mathcal{M}, g) is **Walker** if it admits a null parallel distribution \mathcal{D} of dimension r .

Definition

A manifold (\mathcal{M}, g) is **Walker** if it admits a null parallel distribution \mathcal{D} of dimension r .

- In dimension 3 there exists coordinates such that:

$$g_f = \begin{pmatrix} 0 & 0 & 1 \\ 0 & \varepsilon & 0 \\ 1 & 0 & f \end{pmatrix} \text{ where } f = f(t, x, y) \text{ and } \varepsilon = \pm 1.$$

Definition

A manifold (\mathcal{M}, g) is **Walker** if it admits a null parallel distribution \mathcal{D} of dimension r .

- In dimension 3 there exists coordinates such that:

$$g_f = \begin{pmatrix} 0 & 0 & 1 \\ 0 & \varepsilon & 0 \\ 1 & 0 & f \end{pmatrix} \text{ where } f = f(t, x, y) \text{ and } \varepsilon = \pm 1.$$

- \mathcal{D} is locally generated by ∂_t .

Definition

A manifold (\mathcal{M}, g) is **Walker** if it admits a null parallel distribution \mathcal{D} of dimension r .

- In dimension 3 there exists coordinates such that:

$$g_f = \begin{pmatrix} 0 & 0 & 1 \\ 0 & \varepsilon & 0 \\ 1 & 0 & f \end{pmatrix} \text{ where } f = f(t, x, y) \text{ and } \varepsilon = \pm 1.$$

- \mathcal{D} is locally generated by ∂_t .
- (\mathcal{M}, g) is strictly Walker if \mathcal{D} admits a null parallel vector field.

Definition

A manifold (\mathcal{M}, g) is **Walker** if it admits a null parallel distribution \mathcal{D} of dimension r .

- In dimension 3 there exists coordinates such that:

$$g_f = \begin{pmatrix} 0 & 0 & 1 \\ 0 & \varepsilon & 0 \\ 1 & 0 & f \end{pmatrix} \text{ where } f = f(t, x, y) \text{ and } \varepsilon = \pm 1.$$

- \mathcal{D} is locally generated by ∂_t .
- (\mathcal{M}, g) is strictly Walker if \mathcal{D} admits a null parallel vector field.

$$- > (\mathcal{M}, g) \text{ is strictly Walker} \Leftrightarrow f = f(x, y)$$

Definition

A manifold (\mathcal{M}, g) is **Walker** if it admits a null parallel distribution \mathcal{D} of dimension r .

- In dimension 3 there exists coordinates such that:

$$g_f = \begin{pmatrix} 0 & 0 & 1 \\ 0 & \varepsilon & 0 \\ 1 & 0 & f \end{pmatrix} \text{ where } f = f(t, x, y) \text{ and } \varepsilon = \pm 1.$$

- \mathcal{D} is locally generated by ∂_t .
- (\mathcal{M}, g) is strictly Walker if \mathcal{D} admits a null parallel vector field.
 - $>$ (\mathcal{M}, g) is strictly Walker $\Leftrightarrow f = f(x, y)$
 - $>$ (\mathcal{M}, g) is strictly Walker $\Leftrightarrow Ric^2 = 0$

- 1 Algebraic conditions
- 2 Geometry of the Jacobi operator
- 3 Geometry of the skew-symmetric curvature operator

- 1 Algebraic conditions
- 2 Geometry of the Jacobi operator
- 3 Geometry of the skew-symmetric curvature operator

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension n .

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension n .

① ∇ denotes the **Levi-Civita connection**.

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension n .

- ① ∇ denotes the Levi-Civita connection.
- ② $R(x, y) = [\nabla_x, \nabla_y] - \nabla_{[x, y]}$ is the **curvature** operator.

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension n .

- ① ∇ denotes the Levi-Civita connection.
- ② $R(x, y) = [\nabla_x, \nabla_y] - \nabla_{[x, y]}$ is the curvature operator.

Ricci tensor

$$\rho(x, y) = \sum_i \varepsilon_i R(e_i, x, y, e_i)$$

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension n .

- ① ∇ denotes the Levi-Civita connection.
- ② $R(x, y) = [\nabla_x, \nabla_y] - \nabla_{[x, y]}$ is the curvature operator.

Ricci tensor

$$\rho(x, y) = \sum_i \varepsilon_i R(e_i, x, y, e_i)$$

Scalar curvature

$$\tau = \sum_i \varepsilon_i \rho(e_i, e_i)$$

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension n .

- ① ∇ denotes the Levi-Civita connection.
- ② $R(x, y) = [\nabla_x, \nabla_y] - \nabla_{[x, y]}$ is the curvature operator.

Ricci tensor

$$\rho(x, y) = \sum_i \varepsilon_i R(e_i, x, y, e_i)$$

Scalar curvature

$$\tau = \sum_i \varepsilon_i \rho(e_i, e_i)$$

Ricci operator

$g(\rho(x), y) := \rho(x, y)$ for a unit x and y .

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension n .

- ① ∇ denotes the Levi-Civita connection.
- ② $R(x, y) = [\nabla_x, \nabla_y] - \nabla_{[x, y]}$ is the curvature operator.

Ricci tensor

$$\rho(x, y) = \sum_i \varepsilon_i R(e_i, x, y, e_i)$$

Scalar curvature

$$\tau = \sum_i \varepsilon_i \rho(e_i, e_i)$$

Ricci operator

$$g(\rho(x), y) := \rho(x, y) \text{ for a unit } x \text{ and } y.$$

Skew-symmetric operator

$$g(R(x, y)z, w) = R(x, y, z, w) \text{ for } x, y \in T_p \mathcal{M}.$$

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension n .

- ① ∇ denotes the Levi-Civita connection.
- ② $R(x, y) = [\nabla_x, \nabla_y] - \nabla_{[x, y]}$ is the curvature operator.

Ricci tensor

$$\rho(x, y) = \sum_i \varepsilon_i R(e_i, x, y, e_i)$$

Scalar curvature

$$\tau = \sum_i \varepsilon_i \rho(e_i, e_i)$$

Ricci operator

$$g(\rho(x), y) := \rho(x, y) \text{ for a unit } x \text{ and } y.$$

Skew-symmetric operator

$$g(R(x, y)z, w) = R(x, y, z, w) \text{ for } x, y \in T_p \mathcal{M}.$$

Jacobi operator

$$R_x = R(\cdot, x)x \text{ for a unit } x.$$

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension n .

- ① ∇ denotes the Levi-Civita connection.
- ② $R(x, y) = [\nabla_x, \nabla_y] - \nabla_{[x, y]}$ is the curvature operator.

Ricci tensor

$$\rho(x, y) = \sum_i \varepsilon_i R(e_i, x, y, e_i)$$

Scalar curvature

$$\tau = \sum_i \varepsilon_i \rho(e_i, e_i)$$

Ricci operator

$$g(\rho(x), y) := \rho(x, y) \text{ for a unit } x \text{ and } y.$$

Skew-symmetric operator

$$g(R(x, y)z, w) = R(x, y, z, w) \text{ for } x, y \in T_p \mathcal{M}.$$

Jacobi operator

$$R_x = R(\cdot, x)x \text{ for a unit } x.$$

Szabó operator

$$R'_x = \nabla_x R(\cdot, x)x \text{ for a unit } x.$$

Generalized
locally
symmet-
ric
spaces

Sandra
Gavino
Fernández

Algebraic
conditions

Geometry
of the
Jacobi
operator

\mathbb{C} -spaces
and
 \mathfrak{P} -spaces

Geometry
of the
skew-
symmetric
curvature
operator

\mathfrak{D} -spaces
and
 \mathfrak{S} -spaces

- (\mathcal{M}, g) is locally symmetric \Leftrightarrow local geodesic symmetries are isometries

- (\mathcal{M}, g) is locally symmetric \Leftrightarrow local geodesic symmetries are isometries

$$- > (\mathcal{M}, g) \text{ is locally symmetric } \Leftrightarrow \nabla R = 0$$

- (\mathcal{M}, g) is locally symmetric \Leftrightarrow local geodesic symmetries are isometries
 - $>$ (\mathcal{M}, g) is locally symmetric $\Leftrightarrow \nabla R = 0$
 - $>$ $\dim \mathcal{M} = 3$
 (\mathcal{M}, g) locally symmetric $\Leftrightarrow \nabla \rho = 0$

- (\mathcal{M}, g) is locally symmetric \Leftrightarrow local geodesic symmetries are isometries

- $>$ (\mathcal{M}, g) is locally symmetric $\Leftrightarrow \nabla R = 0$

- $>$ $\dim \mathcal{M} = 3$

- (\mathcal{M}, g) locally symmetric $\Leftrightarrow \nabla \rho = 0$

$$W = \mathcal{A} \oplus \mathcal{B} \oplus \mathcal{C}^\perp$$

- (\mathcal{M}, g) is locally symmetric \Leftrightarrow local geodesic symmetries are isometries

$$- > (\mathcal{M}, g) \text{ is locally symmetric } \Leftrightarrow \nabla R = 0$$

$$- > \dim \mathcal{M} = 3$$

$$(\mathcal{M}, g) \text{ locally symmetric } \Leftrightarrow \nabla \rho = 0$$

$$W = \mathcal{A} \oplus \mathcal{B} \oplus \mathcal{C}^\perp$$

Einstein-like manifolds (Gray, 1978)

- (\mathcal{M}, g) is locally symmetric \Leftrightarrow local geodesic symmetries are isometries

$$- > (\mathcal{M}, g) \text{ is locally symmetric } \Leftrightarrow \nabla R = 0$$

$$- > \dim \mathcal{M} = 3$$

$$(\mathcal{M}, g) \text{ locally symmetric } \Leftrightarrow \nabla \rho = 0$$

$$W = \mathcal{A} \oplus \mathcal{B} \oplus \mathcal{C}^\perp$$

Einstein-like manifolds (Gray, 1978)

- \mathcal{A} : $(\nabla_X \rho)(X, X) = 0$,

- (\mathcal{M}, g) is locally symmetric \Leftrightarrow local geodesic symmetries are isometries

$$- > (\mathcal{M}, g) \text{ is locally symmetric } \Leftrightarrow \nabla R = 0$$

$$- > \dim \mathcal{M} = 3$$

$$(\mathcal{M}, g) \text{ locally symmetric } \Leftrightarrow \nabla \rho = 0$$

$$W = \mathcal{A} \oplus \mathcal{B} \oplus \mathcal{C}^\perp$$

Einstein-like manifolds (Gray, 1978)

- \mathcal{A} : $(\nabla_X \rho)(X, X) = 0$,
- \mathcal{B} : $(\nabla_X \rho)(Y, Z) = (\nabla_Y \rho)(X, Z)$,

- (\mathcal{M}, g) is locally symmetric \Leftrightarrow local geodesic symmetries are isometries

$$- > (\mathcal{M}, g) \text{ is locally symmetric } \Leftrightarrow \nabla R = 0$$

$$- > \dim \mathcal{M} = 3$$

$$(\mathcal{M}, g) \text{ locally symmetric } \Leftrightarrow \nabla \rho = 0$$

$$W = \mathcal{A} \oplus \mathcal{B} \oplus \mathcal{C}^\perp$$

Einstein-like manifolds (Gray, 1978)

- \mathcal{A} : $(\nabla_X \rho)(X, X) = 0$,
- \mathcal{B} : $(\nabla_X \rho)(Y, Z) = (\nabla_Y \rho)(X, Z)$,
- \mathcal{C}^\perp : $\nabla_X(\rho)(Y, Z) = \frac{1}{(n+2)(n-1)} \{nX(\tau)g(Y, Z) + \frac{n-2}{2}[Y(\tau)g(X, Z) + Z(\tau)g(X, Y)]\}.$

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is **locally symmetric**.

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is locally symmetric.
- ② (\mathcal{M}, g) is in class \mathcal{A} .

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is locally symmetric.
- ② (\mathcal{M}, g) is in class \mathcal{A} .
- ③ (\mathcal{M}, g) is in class \mathcal{C}^\perp .

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is locally symmetric.
- ② (\mathcal{M}, g) is in class \mathcal{A} .
- ③ (\mathcal{M}, g) is in class \mathcal{C}^\perp .
- ④ (\mathcal{M}, g) is in class $\mathcal{A} \oplus \mathcal{C}^\perp$.

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is locally symmetric.
- ② (\mathcal{M}, g) is in class \mathcal{A} .
- ③ (\mathcal{M}, g) is in class \mathcal{C}^\perp .
- ④ (\mathcal{M}, g) is in class $\mathcal{A} \oplus \mathcal{C}^\perp$.

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is locally symmetric.
- ② (\mathcal{M}, g) is in class \mathcal{A} .
- ③ (\mathcal{M}, g) is in class \mathcal{C}^\perp .
- ④ (\mathcal{M}, g) is in class $\mathcal{A} \oplus \mathcal{C}^\perp$.

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is **locally conformally flat**.

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is locally symmetric.
- ② (\mathcal{M}, g) is in class \mathcal{A} .
- ③ (\mathcal{M}, g) is in class \mathcal{C}^\perp .
- ④ (\mathcal{M}, g) is in class $\mathcal{A} \oplus \mathcal{C}^\perp$.

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is locally conformally flat.
- ② (\mathcal{M}, g) is in class \mathcal{B} .

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is locally symmetric.
- ② (\mathcal{M}, g) is in class \mathcal{A} .
- ③ (\mathcal{M}, g) is in class \mathcal{C}^\perp .
- ④ (\mathcal{M}, g) is in class $\mathcal{A} \oplus \mathcal{C}^\perp$.

Theorem

For a 3-dimensional Walker manifold (\mathcal{M}, g) are equivalent:

- ① (\mathcal{M}, g) is locally conformally flat.
- ② (\mathcal{M}, g) is in class \mathcal{B} .
- ③ (\mathcal{M}, g) is in class $\mathcal{B} \oplus \mathcal{C}^\perp$.

1 Algebraic conditions

2 Geometry of the Jacobi operator

- \mathcal{C} -spaces and \mathfrak{P} -spaces

3 Geometry of the skew-symmetric curvature operator

Let (\mathcal{M}, g) be a Riemannian manifold.

Let (\mathcal{M}, g) be a Riemannian manifold.

- 1 (\mathcal{M}, g) is a \mathfrak{C} -space if the Jacobi operator has **constant eigenvalues** along every geodesic.

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{C} -space if the Jacobi operator has constant eigenvalues along every geodesic.
- ② (\mathcal{M}, g) is a \mathfrak{P} -space if the Jacobi operator has **parallel eigenspaces** along every geodesic.

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{C} -space if the Jacobi operator has constant eigenvalues along every geodesic.
- ② (\mathcal{M}, g) is a \mathfrak{P} -space if the Jacobi operator has parallel eigenspaces along every geodesic.

Theorem (Berndt-Vanhecke, 1992)

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{C} -space if the Jacobi operator has constant eigenvalues along every geodesic.
- ② (\mathcal{M}, g) is a \mathfrak{P} -space if the Jacobi operator has parallel eigenspaces along every geodesic.

Theorem (Berndt-Vanhecke, 1992)

(\mathcal{M}, g) is **locally symmetric** if and only if (\mathcal{M}, g) is

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{C} -space if the Jacobi operator has constant eigenvalues along every geodesic.
- ② (\mathcal{M}, g) is a \mathfrak{P} -space if the Jacobi operator has parallel eigenspaces along every geodesic.

Theorem (Berndt-Vanhecke, 1992)

(\mathcal{M}, g) is **locally symmetric** if and only if (\mathcal{M}, g) is simultaneously **\mathfrak{C} -space** and **\mathfrak{P} -space**.

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{C} -space if the Jacobi operator has constant eigenvalues along every geodesic.
- ② (\mathcal{M}, g) is a \mathfrak{P} -space if the Jacobi operator has parallel eigenspaces along every geodesic.

Theorem (Berndt-Vanhecke, 1992)

(\mathcal{M}, g) is locally symmetric if and only if (\mathcal{M}, g) is simultaneously \mathfrak{C} -space and \mathfrak{P} -space.

Let (\mathcal{M}, g) be a Lorentzian manifold.

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{C} -space if the Jacobi operator has constant eigenvalues along every geodesic.
- ② (\mathcal{M}, g) is a \mathfrak{P} -space if the Jacobi operator has parallel eigenspaces along every geodesic.

Theorem (Berndt-Vanhecke, 1992)

(\mathcal{M}, g) is locally symmetric if and only if (\mathcal{M}, g) is simultaneously \mathfrak{C} -space and \mathfrak{P} -space.

Let (\mathcal{M}, g) be a Lorentzian manifold.

- ① (\mathcal{M}, g) is a **timelike \mathfrak{C} -space (\mathcal{TC})** if the Jacobi operator has **constant eigenvalues** along every timelike geodesic.

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{C} -space if the Jacobi operator has constant eigenvalues along every geodesic.
- ② (\mathcal{M}, g) is a \mathfrak{P} -space if the Jacobi operator has parallel eigenspaces along every geodesic.

Theorem (Berndt-Vanhecke, 1992)

(\mathcal{M}, g) is locally symmetric if and only if (\mathcal{M}, g) is simultaneously \mathfrak{C} -space and \mathfrak{P} -space.

Let (\mathcal{M}, g) be a Lorentzian manifold.

- ① (\mathcal{M}, g) is a timelike \mathfrak{C} -space ($T\mathfrak{C}$) if the Jacobi operator has constant eigenvalues along every timelike geodesic.
- ② (\mathcal{M}, g) is a **timelike \mathfrak{P} -space ($T\mathfrak{P}$)** if the Jacobi operator has **parallel eigenspaces** along every timelike geodesic.

Theorem

Let (\mathcal{M}, g) be a Lorentzian manifold.

Theorem

Let (\mathcal{M}, g) be a Lorentzian manifold.

(\mathcal{M}, g) is **locally symmetric** $\Leftrightarrow (\mathcal{M}, g)$ is simultaneously a

Theorem

Let (\mathcal{M}, g) be a Lorentzian manifold.

(\mathcal{M}, g) is **locally symmetric** $\Leftrightarrow (\mathcal{M}, g)$ is simultaneously a **timelike \mathfrak{C} -space** and a **timelike \mathfrak{P} -space**.

Theorem

Let (\mathcal{M}, g) be a Lorentzian manifold.

(\mathcal{M}, g) is locally symmetric $\Leftrightarrow (\mathcal{M}, g)$ is simultaneously a timelike \mathfrak{C} -space and a timelike \mathfrak{P} -space.

$T\mathfrak{C} \cap T\mathfrak{P}$ is equivalent to

Theorem

Let (\mathcal{M}, g) be a Lorentzian manifold.

(\mathcal{M}, g) is locally symmetric $\Leftrightarrow (\mathcal{M}, g)$ is simultaneously a timelike \mathfrak{C} -space and a timelike \mathfrak{P} -space.

$T\mathfrak{C} \cap T\mathfrak{P}$ is equivalent to

$$\nabla_x R_{xyyx} = 0, \quad \forall x \text{ timelike and } y \perp x$$

Theorem

Let (\mathcal{M}, g) be a Lorentzian manifold.

(\mathcal{M}, g) is locally symmetric $\Leftrightarrow (\mathcal{M}, g)$ is simultaneously a timelike \mathfrak{C} -space and a timelike \mathfrak{P} -space.

$T\mathfrak{C} \cap T\mathfrak{P}$ is equivalent to

$$\nabla_x R_{xyyx} = 0, \quad \forall x \text{ timelike and } y \perp x$$

This is equivalent to $\nabla R = 0$ (García-Río, Vázquez-Abal, 1993).

Theorem

Let (\mathcal{M}, g) be a Lorentzian manifold.

(\mathcal{M}, g) is locally symmetric $\Leftrightarrow (\mathcal{M}, g)$ is simultaneously a timelike \mathfrak{C} -space and a timelike \mathfrak{P} -space.

$T\mathfrak{C} \cap T\mathfrak{P}$ is equivalent to

$$\nabla_x R_{xyyx} = 0, \forall x \text{ timelike and } y \perp x$$

This is equivalent to $\nabla R = 0$ (García-Río, Vázquez-Abal, 1993).

Remark

Previous theorem is **false** in other signatures.

Theorem

Let (\mathcal{M}, g) be a Lorentzian manifold.

(\mathcal{M}, g) is locally symmetric $\Leftrightarrow (\mathcal{M}, g)$ is simultaneously a timelike \mathfrak{C} -space and a timelike \mathfrak{P} -space.

$T\mathfrak{C} \cap T\mathfrak{P}$ is equivalent to

$$\nabla_x R_{xyyx} = 0, \quad \forall x \text{ timelike and } y \perp x$$

This is equivalent to $\nabla R = 0$ (García-Río, Vázquez-Abal, 1993).

Example

A strict walker 4-dimensional manifold of signature $(2, 2)$:

$g = 2dx^1 \circ dx^3 + 2dx^2 \circ dx^4 + b(x^3)dx^4 \circ dx^4$, is always $T\mathfrak{C}$ and $T\mathfrak{P}$ but not locally symmetric ($\nabla R = 0 \Leftrightarrow b^{(3)}(x^3) = 0$).

(\mathcal{M}, g) timelike \mathfrak{C} -space $\Leftrightarrow R_\gamma$ has constant eigenvalues along γ
for any timelike γ

(\mathcal{M}, g) timelike \mathfrak{C} -space $\Leftrightarrow R_\gamma$ has constant eigenvalues along γ
 for any timelike γ
 $\Leftrightarrow \nabla_{\gamma'} \text{Tr} R_\gamma^{(k)} = 0$ for any timelike γ

(\mathcal{M}, g) timelike \mathfrak{C} -space $\Leftrightarrow R_\gamma$ has constant eigenvalues along γ
for any timelike γ

$$\Leftrightarrow \nabla_{\gamma'} \text{Tr} R_\gamma^{(k)} = 0 \text{ for any timelike } \gamma$$

$$\Leftrightarrow \nabla_X \text{Tr} R_X^{(k)} = 0, \text{ for all } X$$

(\mathcal{M}, g) timelike \mathfrak{C} -space $\Leftrightarrow R_\gamma$ has constant eigenvalues along γ
 for any timelike γ
 $\Leftrightarrow \nabla_{\gamma'} \text{Tr} R_\gamma^{(k)} = 0$ for any timelike γ
 $\Leftrightarrow \nabla_X \text{Tr} R_X^{(k)} = 0$, for all X

Lemma (Characterization)

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension 3. (\mathcal{M}, g) is a **timelike \mathfrak{C} -space** if and only if:

(\mathcal{M}, g) timelike \mathfrak{C} -space $\Leftrightarrow R_\gamma$ has constant eigenvalues along γ
 for any timelike γ
 $\Leftrightarrow \nabla_{\gamma'} \text{Tr} R_\gamma^{(k)} = 0$ for any timelike γ
 $\Leftrightarrow \nabla_X \text{Tr} R_X^{(k)} = 0$, for all X

Lemma (Characterization)

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension 3. (\mathcal{M}, g) is a timelike \mathfrak{C} -space if and only if:

$$(1) \quad \nabla_X \rho(X, X) = 0,$$

(\mathcal{M}, g) timelike \mathfrak{C} -space $\Leftrightarrow R_\gamma$ has constant eigenvalues along γ
 for any timelike γ
 $\Leftrightarrow \nabla_{\gamma'} \text{Tr} R_\gamma^{(k)} = 0$ for any timelike γ
 $\Leftrightarrow \nabla_X \text{Tr} R_X^{(k)} = 0$, for all X

Lemma (Characterization)

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension 3. (\mathcal{M}, g) is a timelike \mathfrak{C} -space if and only if:

- (1) $\nabla_X \rho(X, X) = 0$,
- (2) $\sum_i \varepsilon_i R(e_i, X, X, e_i) \nabla_X R(e_i, X, X, e_i) = 0$,

(\mathcal{M}, g) timelike \mathfrak{C} -space $\Leftrightarrow R_\gamma$ has constant eigenvalues along γ
for any timelike γ

$$\Leftrightarrow \nabla_{\gamma'} \text{Tr} R_\gamma^{(k)} = 0 \text{ for any timelike } \gamma$$

$$\Leftrightarrow \nabla_X \text{Tr} R_X^{(k)} = 0, \text{ for all } X$$

Lemma (Characterization)

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension 3. (\mathcal{M}, g) is a timelike \mathfrak{C} -space if and only if:

$$(1) \quad \nabla_X \rho(X, X) = 0,$$

$$(2) \quad \sum_i \varepsilon_i R(e_i, X, X, e_i) \nabla_X R(e_i, X, X, e_i) = 0,$$

$\forall X \in T\mathcal{M}$, where $\{e_i\}$ is an orthonormal basis.

(\mathcal{M}, g) timelike \mathfrak{C} -space $\Leftrightarrow R_\gamma$ has constant eigenvalues along γ
for any timelike γ

$$\Leftrightarrow \nabla_{\gamma'} \text{Tr} R_\gamma^{(k)} = 0 \text{ for any timelike } \gamma$$

$$\Leftrightarrow \nabla_X \text{Tr} R_X^{(k)} = 0, \text{ for all } X$$

Lemma (Characterization)

Let (\mathcal{M}, g) be a Lorentzian manifold of dimension 3. (\mathcal{M}, g) is a timelike \mathfrak{C} -space if and only if:

$$(1) \quad \nabla_X \rho(X, X) = 0,$$

$$(2) \quad \sum_i \varepsilon_i R(e_i, X, X, e_i) \nabla_X R(e_i, X, X, e_i) = 0,$$

$\forall X \in T\mathcal{M}$, where $\{e_i\}$ is an orthonormal basis.

(1) and (2) are the Ledger conditions of order 3 and 5.

Lemma

Let (\mathcal{M}, g) be a Lorentzian manifold. (\mathcal{M}, g) is a **timelike \mathfrak{P} -space** if and only if:

Lemma

Let (\mathcal{M}, g) be a Lorentzian manifold. (\mathcal{M}, g) is a timelike \mathfrak{P} -space if and only if:

$$R_{\gamma} R'_{\gamma} - R'_{\gamma} R_{\gamma} = 0$$

Lemma

Let (\mathcal{M}, g) be a Lorentzian manifold. (\mathcal{M}, g) is a timelike \mathfrak{P} -space if and only if:

$$R_{\gamma} R'_{\gamma} - R'_{\gamma} R_{\gamma} = 0$$

for all timelike geodesic γ .

Lemma

Let (\mathcal{M}, g) be a Lorentzian manifold. (\mathcal{M}, g) is a timelike \mathfrak{P} -space if and only if:

$$R_\gamma R'_\gamma - R'_\gamma R_\gamma = 0$$

for all timelike geodesic γ .

The Jacobi and the Szabó operators commute for timelike geodesics if and only $R_X R'_X - R'_X R_X = 0$ **for all X** .

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a **timelike \mathfrak{C} -space**

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a timelike \mathfrak{C} -space if and only if it is **locally symmetric**.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a timelike \mathfrak{C} -space if and only if it is locally symmetric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f of constant scalar curvature is a **timelike \mathfrak{P} -space**

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a timelike \mathfrak{C} -space if and only if it is locally symmetric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f of constant scalar curvature is a timelike \mathfrak{P} -space if and only if it is **locally symmetric** or a strictly Walker metric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a timelike \mathfrak{C} -space if and only if it is locally symmetric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f of constant scalar curvature is a timelike \mathfrak{P} -space if and only if it is locally symmetric or a **strictly Walker metric**.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a timelike \mathfrak{C} -space if and only if it is locally symmetric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f of constant scalar curvature is a timelike \mathfrak{P} -space if and only if it is locally symmetric or a strictly Walker metric.

– > A 3-dimensional Walker manifold \mathcal{M}_f of constant scalar curvature is a timelike \mathfrak{P} -space if and only if it is curvature recurrent.

- 1 Algebraic conditions
- 2 Geometry of the Jacobi operator
- 3 Geometry of the skew-symmetric curvature operator
 - \mathfrak{D} -spaces and \mathfrak{T} -spaces

Let (\mathcal{M}, g) be a Riemannian manifold.

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{D} -space if the skew-symmetric curvature operator has constant eigenvalues along each unit circle.

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{D} -space if the skew-symmetric curvature operator has constant eigenvalues along each unit circle.
- ② (\mathcal{M}, g) is a \mathfrak{T} -space if there is a **parallel Jordan basis** for the skew-symmetric curvature operator along each unit circle.

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{D} -space if the skew-symmetric curvature operator has constant eigenvalues along each unit circle.
- ② (\mathcal{M}, g) is a \mathfrak{T} -space if there is a parallel Jordan basis for the skew-symmetric curvature operator along each unit circle.

Theorem (Ivanov-Petrova, 1997)

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{D} -space if the skew-symmetric curvature operator has constant eigenvalues along each unit circle.
- ② (\mathcal{M}, g) is a \mathfrak{T} -space if there is a parallel Jordan basis for the skew-symmetric curvature operator along each unit circle.

Theorem (Ivanov-Petrova, 1997)

(\mathcal{M}, g) is **locally symmetric** if and only if (\mathcal{M}, g) is

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{D} -space if the skew-symmetric curvature operator has constant eigenvalues along each unit circle.
- ② (\mathcal{M}, g) is a \mathfrak{T} -space if there is a parallel Jordan basis for the skew-symmetric curvature operator along each unit circle.

Theorem (Ivanov-Petrova, 1997)

(\mathcal{M}, g) is **locally symmetric** if and only if (\mathcal{M}, g) is simultaneously **\mathfrak{D} -space** and **\mathfrak{T} -space**.

Let (\mathcal{M}, g) be a Riemannian manifold.

- ① (\mathcal{M}, g) is a \mathfrak{D} -space if the skew-symmetric curvature operator has constant eigenvalues along each unit circle.
- ② (\mathcal{M}, g) is a \mathfrak{T} -space if there is a parallel Jordan basis for the skew-symmetric curvature operator along each unit circle.

Theorem (Ivanov-Petrova, 1997)

(\mathcal{M}, g) is locally symmetric if and only if (\mathcal{M}, g) is simultaneously \mathfrak{D} -space and \mathfrak{T} -space.

(\mathcal{M}, g) is a \mathfrak{D} -space if and only if $\nabla_{c'} \text{Tr} R(c', \nabla_{c'} c')^{(k)} = 0$ for all unit circle c .

(\mathcal{M}, g) is a \mathfrak{T} -space if and only if $\nabla_{c'} R(c', \nabla_{c'} c')$ and $R(c', \nabla_{c'} c')$ commute for all unit circle c .

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a \mathfrak{D} -space

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a \mathfrak{D} -space if and only if it is **locally symmetric** or a strictly Walker metric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a \mathfrak{D} -space if and only if it is locally symmetric or a **strictly Walker metric**.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a \mathfrak{D} -space if and only if it is locally symmetric or a strictly Walker metric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f of constant scalar curvature is a \mathfrak{T} -space

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a \mathfrak{D} -space if and only if it is locally symmetric or a strictly Walker metric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f of constant scalar curvature is a \mathfrak{T} -space if and only if it is **locally symmetric** or a strictly Walker metric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a \mathfrak{D} -space if and only if it is locally symmetric or a strictly Walker metric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f of constant scalar curvature is a \mathfrak{T} -space if and only if it is locally symmetric or a **strictly Walker metric**.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f is a \mathfrak{D} -space if and only if it is locally symmetric or a strictly Walker metric.

Theorem

A 3-dimensional Walker manifold \mathcal{M}_f of constant scalar curvature is a \mathfrak{T} -space if and only if it is locally symmetric or a strictly Walker metric.

$$\{\mathfrak{D} - \text{space} \cap \mathfrak{T} - \text{space}\} \not\supseteq \{\nabla R = 0\}.$$

Einstein-like Walker manifolds and generalized locally symmetric spaces

Sandra Gavino Fernández



Department of Geometry and Topology
University of Santiago de Compostela

July 11th, 2009

Work in progress with Miguel Brozos Vázquez