

# Some remarks on the null sectional curvature

Alma L. Albuje  
(joint work with Stefan Haesen)

V International Meeting on Lorentzian Geometry  
Martina Franca, Taranto, Italy

July, 8-11, 2009

# Null sectional curvature

- On a Lorentzian manifold the sectional curvature function is only defined on the Grassmannian of non-degenerate planes.

# Null sectional curvature

- On a Lorentzian manifold the sectional curvature function is only defined on the Grassmannian of non-degenerate planes.
- It is well-known that it can only be continuously extended to the Grassmannian of all planes if the space has constant sectional curvature.

# Null sectional curvature

- On a Lorentzian manifold the sectional curvature function is only defined on the Grassmannian of non-degenerate planes.
- It is well-known that it can only be continuously extended to the Grassmannian of all planes if the space has constant sectional curvature.
- Harris (1982) introduced what is called the null sectional curvature of a degenerate plane:

# Null sectional curvature

- On a Lorentzian manifold the sectional curvature function is only defined on the Grassmannian of non-degenerate planes.
- It is well-known that it can only be continuously extended to the Grassmannian of all planes if the space has constant sectional curvature.
- Harris (1982) introduced what is called the null sectional curvature of a degenerate plane:

## Null sectional curvature

- Let  $(M^n, g)$  be a Lorentzian manifold, and  $\pi = \text{span}\{v, w\} \subset T_p M$  a degenerate plane at  $p \in M$  with  $g(v, v) = 0$  and  $g(w, w) > 0$ .

# Null sectional curvature

- On a Lorentzian manifold the sectional curvature function is only defined on the Grassmannian of non-degenerate planes.
- It is well-known that it can only be continuously extended to the Grassmannian of all planes if the space has constant sectional curvature.
- Harris (1982) introduced what is called the null sectional curvature of a degenerate plane:

## Null sectional curvature

- Let  $(M^n, g)$  be a Lorentzian manifold, and  $\pi = \text{span}\{v, w\} \subset T_p M$  a degenerate plane at  $p \in M$  with  $g(v, v) = 0$  and  $g(w, w) > 0$ .
- The null sectional curvature of  $\pi$  with respect to  $v$  is given by

$$K_v(p, \pi) = \frac{R(v, w, w, v)}{g(w, w)}.$$

# Null sectional curvature

- $K_v(p, \pi)$  is independent of the choice of the spacelike vector  $w \in \pi$ , but it depends quadratically on the null vector  $v \in \pi$ .

# Null sectional curvature

- $K_v(p, \pi)$  is independent of the choice of the spacelike vector  $w \in \pi$ , but it depends quadratically on the null vector  $v \in \pi$ .
- Therefore,  $K_v(p, \pi)$  is not a function on the Grassmannian of all degenerate tangent planes to  $M$ .



# Null sectional curvature

- $K_v(p, \pi)$  is independent of the choice of the spacelike vector  $w \in \pi$ , but it depends quadratically on the null vector  $v \in \pi$ .
- Therefore,  $K_v(p, \pi)$  is not a function on the Grassmannian of all degenerate tangent planes to  $M$ .
- In the case of a time-oriented Lorentzian manifold, we can normalize  $K_v(p, \pi)$  obtaining a well-defined function:

# Null sectional curvature

- $K_v(p, \pi)$  is independent of the choice of the spacelike vector  $w \in \pi$ , but it depends quadratically on the null vector  $v \in \pi$ .
- Therefore,  $K_v(p, \pi)$  is not a function on the Grassmannian of all degenerate tangent planes to  $M$ .
- In the case of a time-oriented Lorentzian manifold, we can normalize  $K_v(p, \pi)$  obtaining a well-defined function:

## $U$ -normalized null sectional curvature

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold, and let  $U \in TM$  be a globally defined timelike vector field.

# Null sectional curvature

- $K_v(p, \pi)$  is independent of the choice of the spacelike vector  $w \in \pi$ , but it depends quadratically on the null vector  $v \in \pi$ .
- Therefore,  $K_v(p, \pi)$  is not a function on the Grassmannian of all degenerate tangent planes to  $M$ .
- In the case of a time-oriented Lorentzian manifold, we can normalize  $K_v(p, \pi)$  obtaining a well-defined function:

## $U$ -normalized null sectional curvature


- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold, and let  $U \in TM$  be a globally defined timelike vector field.
- The  $U$ -normalized null sectional curvature at  $p \in M$  of the degenerate plane  $\pi \subset T_p M$  is defined as

$$K^U(p, \pi) = K_v(p, \pi),$$

with  $v \in (C_U M)_p = \{v \in T_p M \mid g(v, v) = 0 \text{ and } g(v, U_p) = -1\}$ .

# Null sectional curvature

- $K_v(p, \pi)$  is independent of the choice of the spacelike vector  $w \in \pi$ , but it depends on  $v$ .
- Therefore,  $K_v(p, \pi)$  is independent of the choice of the degenerate tangent plane  $\pi$ .
- In the case of a spacelike vector  $U_p$ , we can normalize  $K_v(p, \pi)$  to obtain



## $U$ -normalized null sectional curvature

- Let  $(M^n, g)$  be a spacetime manifold, and let  $U \in TM$  be a spacelike vector.
- The  $U$ -normalized null sectional curvature  $K^U(p, \pi)$  is defined as the null sectional curvature of the degenerate plane  $\pi$  spanned by  $U_p$  and any spacelike vector  $w \in \pi$ .

old, and let  $d$ .

$M$  of the

$$K^U(p, \pi) = K_v(p, \pi),$$

with  $v \in (C_U M)_p = \{v \in T_p M | g(v, v) = 0 \text{ and } g(v, U_p) = -1\}$ .

# Null sectional curvature

- $K_v(p, \pi)$  is independent of the choice of the spacelike vector  $w \in \pi$ , but it depends quadratically on the null vector  $v \in \pi$ .
- Therefore,  $K_v(p, \pi)$  is not a function on the Grassmannian of all degenerate tangent planes to  $M$ .
- In the case of a time-oriented Lorentzian manifold, we can normalize  $K_v(p, \pi)$  obtaining a well-defined function:

## $U$ -normalized null sectional curvature

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold, and let  $U \in TM$  be a globally defined timelike vector field.
- The  $U$ -normalized null sectional curvature at  $p \in M$  of the degenerate plane  $\pi \subset T_p M$  is defined as

$$K^U(p, \pi) = K_v(p, \pi),$$

with  $v \in (C_U M)_p = \{v \in T_p M \mid g(v, v) = 0 \text{ and } g(v, U_p) = -1\}$ .

- $K^U(p, \pi)$  is a purely Lorentzian concept.

# A geometrical interpretation for $K^U(p, \pi)$

**Why  $K^U(p, \pi)$  is called a “curvature”?**

# A geometrical interpretation for $K^U(p, \pi)$

## Why $K^U(p, \pi)$ is called a “curvature”?

- Until the moment, besides the resemblance in formulas with the classical concept of sectional curvature there does not seem to be a geometrical motivation why to call it as such.

# A geometrical interpretation for $K^U(p, \pi)$

## Why $K^U(p, \pi)$ is called a “curvature”?

- Until the moment, besides the resemblance in formulas with the classical concept of sectional curvature there does not seem to be a geometrical motivation why to call it as such.
- We give a geometrical interpretation of the null sectional curvature in terms of the difference in length of two spacelike geodesics constructed from the degenerate plane.



# A geometrical interpretation for $K^U(p, \pi)$

## Why $K^U(p, \pi)$ is called a “curvature”?

- Until the moment, besides the resemblance in formulas with the classical concept of sectional curvature there does not seem to be a geometrical motivation why to call it as such.
- We give a geometrical interpretation of the null sectional curvature in terms of the difference in length of two spacelike geodesics constructed from the degenerate plane.
- This interpretation is inspired in the one given by Levi-Civita (1917) for the sectional curvature in the Riemannian case.

# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .

# A geometrical interpretation for $K^U(p, \pi)$


- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.

# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:

# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:




# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:




# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:



# A geometrical interpretation for $K^U(p, \pi)$


- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:






# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:




# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:




# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:




# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:



# A geometrical interpretation for $K^U(p, \pi)$

- Let  $(M^n, g)$  be a time-oriented Lorentzian manifold,  $U \in TM$  a globally defined unitary timelike vector field and  $\pi = \text{span}\{v, w\}$  a degenerate plane at  $p \in M$  such that  $v \in (C_U M)_p$  and  $g(w, w) > 0$ .
- Since  $K^U(p, \pi)$  is independent of the choice of the spacelike vector  $w$ , we can assume without loss of generality that  $w$  is a unit vector.
- We make the following construction of an infinitesimal parallelogramoid at  $p \in M$ , having as edges spacelike and null geodesics:



and we prove that

$$K^U(p, \pi) \simeq 6 \frac{a' - a}{ab^2}.$$

# Isotropic and spatially constant null sectional curvature

- The  $U$ -normalized null sectional curvature is said to be **isotropic** if it is only a point function, i.e.,  $K^U(p, \pi) = K^U(p)$  for all null planes  $\pi \subset T_p M$ .

# Isotropic and spatially constant null sectional curvature

- The  $U$ -normalized null sectional curvature is said to be **isotropic** if it is only a point function, i.e.,  $K^U(p, \pi) = K^U(p)$  for all null planes  $\pi \subset T_p M$ .

## Proposition 1 (Harris, Koch-Sen)

Let  $(M^n, g)$ ,  $n \geq 3$ , be a time-orientable Lorentzian manifold and  $U$  a globally defined unitary timelike vector field. Then, the  $U$ -normalized null sectional curvature is isotropic if and only if the curvature tensor satisfies

i)  $R(X, Y)Z = k(X \wedge_g Y)Z$ ,  $\forall X, Y, Z \in U^\perp$ ,

ii)  $R(X, U)U = \mu X$ ,

with  $k, \mu : M \rightarrow \mathbb{R}$  and  $K^U \equiv k + \mu$ .

# Isotropic and spatially constant null sectional curvature

- The  $U$ -normalized null sectional curvature is said to be **isotropic** if it is only a point function, i.e.,  $K^U(p, \pi) = K^U(p)$  for all null planes  $\pi \subset T_p M$ .

## Proposition 1 (Harris, Koch-Sen)

Let  $(M^n, g)$ ,  $n \geq 3$ , be a time-orientable Lorentzian manifold and  $U$  a globally defined unitary timelike vector field. Then, the  $U$ -normalized null sectional curvature is isotropic if and only if the curvature tensor satisfies

i)  $R(X, Y)Z = k(X \wedge_g Y)Z$ ,  $\forall X, Y, Z \in U^\perp$ ,

ii)  $R(X, U)U = \mu X$ ,

with  $k, \mu : M \rightarrow \mathbb{R}$  and  $K^U \equiv k + \mu$ .

- If the  $U$ -normalized null sectional curvature  $K^U$  is isotropic, it is said to be **spatially constant** if it is constant on the space orthogonal to the chosen timelike vector field  $U$ , i.e.,  $X[K^U] = 0$ , for every  $X \in U^\perp$ .



# Isotropic and spatially constant null sectional curvature

## Theorem 1 (Harris, Karcher, Koch-Sen)

Let  $(M, g)$  be an  $n(\geq 4)$ -dimensional Lorentzian manifold and let  $U$  be a unitary timelike vector field on  $M$ . Suppose that the  $U$ -normalized null sectional curvature is non-zero, isotropic and spatially constant. Then,  $g$  is locally a Robertson-Walker metric.

# Isotropic and spatially constant null sectional curvature

## Theorem 1 (Harris, Karcher, Koch-Sen)

Let  $(M, g)$  be an  $n(\geq 4)$ -dimensional Lorentzian manifold and let  $U$  be a unitary timelike vector field on  $M$ . Suppose that the  $U$ -normalized null sectional curvature is non-zero, isotropic and spatially constant. Then,  $g$  is locally a Robertson-Walker metric.

- The condition  $n \geq 4$  is necessary in the above result.

# Isotropic and spatially constant null sectional curvature

## Theorem 1 (Harris, Karcher, Koch-Sen)

Let  $(M, g)$  be an  $n(\geq 4)$ -dimensional Lorentzian manifold and let  $U$  be a unitary timelike vector field on  $M$ . Suppose that the  $U$ -normalized null sectional curvature is non-zero, isotropic and spatially constant. Then,  $g$  is locally a Robertson-Walker metric.

- The condition  $n \geq 4$  is necessary in the above result.
- In fact, Palomo (2007) showed that the 3-dimensional sphere  $\mathbb{S}^3$  endowed with a certain Lorentzian metric has non-zero constant null sectional curvature, whereas the distribution  $U^\perp$  is not integrable and hence is not a Robertson-Walker space.

# Isotropic and spatially constant null sectional curvature

## Theorem 1 (Harris, Karcher, Koch-Sen)

Let  $(M, g)$  be an  $n(\geq 4)$ -dimensional Lorentzian manifold and let  $U$  be a unitary timelike vector field on  $M$ . Suppose that the  $U$ -normalized null sectional curvature is non-zero, isotropic and spatially constant. Then,  $g$  is locally a Robertson-Walker metric.

- The condition  $n \geq 4$  is necessary in the above result.
- In fact, Palomo (2007) showed that the 3-dimensional sphere  $\mathbb{S}^3$  endowed with a certain Lorentzian metric has non-zero constant null sectional curvature, whereas the distribution  $U^\perp$  is not integrable and hence is not a Robertson-Walker space.
- However, we can prove that the above theorem can be extended to the 3-dimensional case under some extra suitable assumptions: we will ask the space to be conformally flat.

# Isotropic and spatially constant null sectional curvature

## Lemma 1

A time-oriented, 3-dimensional Lorentzian manifold  $(M, g)$ , with non-vanishing isotropic null sectional curvature is conformally flat if, and only if,

$$\nabla^\perp[k] = 0,$$

and

$$\nabla_\xi U = -\frac{U[k]}{2(k + \mu)}\xi^\perp + g(\xi, U)\nabla^\perp[\ln(k + \mu)],$$

with  $\xi \in TM$ ,  $\xi^\perp$  its projection on the 2-space perpendicular to  $U$  and  $\nabla^\perp[f]$  the projection of the gradient of a function  $f$  on this 2-space.

# Isotropic and spatially constant null sectional curvature

## Lemma 1

A time-oriented, 3-dimensional Lorentzian manifold  $(M, g)$ , with non-vanishing isotropic null sectional curvature is conformally flat if, and only if,

$$\nabla^\perp[k] = 0,$$

and

$$\nabla_\xi U = -\frac{U[k]}{2(k+\mu)}\xi^\perp + g(\xi, U)\nabla^\perp[\ln(k+\mu)],$$

with  $\xi \in TM$ ,  $\xi^\perp$  its projection on the 2-space perpendicular to  $U$  and  $\nabla^\perp[f]$  the projection of the gradient of a function  $f$  on this 2-space.

*Proof.*  $(M^3, g)$  is conformally flat if and only if the Schouten tensor

$$S = \text{Ric} - \frac{\tau}{4}g$$

satisfies

$$(\nabla_Z S)(X, Y) = (\nabla_Y S)(X, Z).$$

# Isotropic and spatially constant null sectional curvature

## Lemma 1

A time-oriented, 3-dimensional Lorentzian manifold  $(M, g)$ , with non-vanishing isotropic null sectional curvature is conformally flat if, and only if,

$$\nabla^\perp[k] = 0,$$

and

$$\nabla_\xi U = -\frac{U[k]}{2(k+\mu)}\xi^\perp + g(\xi, U)\nabla^\perp[\ln(k+\mu)],$$

with  $\xi \in TM$ ,  $\xi^\perp$  its projection on the 2-space perpendicular to  $U$  and  $\nabla^\perp[f]$  the projection of the gradient of a function  $f$  on this 2-space.

*Proof.*  $(M^3, g)$  is conformally flat if and only if the Schouten tensor

$$S = \text{Ric} - \frac{\tau}{4}g$$

satisfies

$$(\nabla_Z S)(X, Y) = (\nabla_Y S)(X, Z).$$

- The proof follows easily by observing that by Proposition 1

$$\text{Ric} = (k - \mu)g + (k + \mu)U^\flat \otimes U^\flat,$$

being  $U^\flat$  the one-form metrically equivalent with  $U$ .

# Isotropic and spatially constant null sectional curvature

- As a first result we can prove the following,

## Corollary 1

Every time-oriented, conformally flat, 3-dimensional Lorentzian manifold  $(M, g)$  with non-vanishing isotropic null sectional curvature can be locally and isometrically embedded as a quasi-umbilical hypersurface in a 4-dimensional pseudo-Euclidean space.



# Isotropic and spatially constant null sectional curvature

- As a first result we can prove the following,

## Corollary 1

Every time-oriented, conformally flat, 3-dimensional Lorentzian manifold  $(M, g)$  with non-vanishing isotropic null sectional curvature can be locally and isometrically embedded as a quasi-umbilical hypersurface in a 4-dimensional pseudo-Euclidean space.

*Proof.*

- The Gauss equation for a hypersurface in a 4-dimensional pseudo-Euclidean space can be written in general as

$$R(X, Y, Z, W) = \varepsilon (h(X, W)h(Y, Z) - h(X, Z)h(Y, W))$$

for any  $X, Y, Z, W \in TM$ , being  $h$  the second fundamental form of the hypersurface and  $\varepsilon$  the signature of the normal direction.

# Proof of Corollary 1

- Using the expression of the Riemann curvature tensor given in Proposition 1, the Gauss equation is satisfied for

$$h(X, Y) = \alpha g(X, Y) + (\alpha + \beta)g(U, X)g(U, Y),$$

with  $k = \varepsilon\alpha^2$  and  $\mu = \varepsilon\alpha\beta$ .

# Proof of Corollary 1

- Using the expression of the Riemann curvature tensor given in Proposition 1, the Gauss equation is satisfied for

$$h(X, Y) = \alpha g(X, Y) + (\alpha + \beta)g(U, X)g(U, Y),$$

with  $k = \varepsilon\alpha^2$  and  $\mu = \varepsilon\alpha\beta$ .

- On the other hand, since  $M$  is conformally flat, the Schouten tensor is a Codazzi tensor which basically means that  $h$  satisfies the Codazzi equation for a hypersurface in a pseudo-Euclidean space.

# Proof of Corollary 1

- Using the expression of the Riemann curvature tensor given in Proposition 1, the Gauss equation is satisfied for

$$h(X, Y) = \alpha g(X, Y) + (\alpha + \beta)g(U, X)g(U, Y),$$

with  $k = \varepsilon\alpha^2$  and  $\mu = \varepsilon\alpha\beta$ .

- On the other hand, since  $M$  is conformally flat, the Schouten tensor is a Codazzi tensor which basically means that  $h$  satisfies the Codazzi equation for a hypersurface in a pseudo-Euclidean space.
- Finally, from the expression for the second fundamental form we conclude that  $M$  is quasi-umbilical.

# Isotropic and spatially constant null sectional curvature

- Our main result extends Theorem 1 to the 3-dimensional case,

# Isotropic and spatially constant null sectional curvature

- Our main result extends Theorem 1 to the 3-dimensional case,

## Theorem 2

Let  $(M, g)$  be a 3-dimensional Lorentzian manifold. Then,  $(M, g)$  is conformally flat and has non-vanishing spatially constant, isotropic null sectional curvature if, and only if,  $(M, g)$  is locally a Robertson-Walker space.

# Isotropic and spatially constant null sectional curvature

- Our main result extends Theorem 1 to the 3-dimensional case,

## Theorem 2

Let  $(M, g)$  be a 3-dimensional Lorentzian manifold. Then,  $(M, g)$  is conformally flat and has non-vanishing spatially constant, isotropic null sectional curvature if, and only if,  $(M, g)$  is locally a Robertson-Walker space.

*Proof.*

- From

$$\nabla_{\xi} U = -\frac{U[k]}{2(k + \mu)} \xi^{\perp} + g(\xi, U) \nabla^{\perp} [\ln(k + \mu)],$$

it follows easily that the vector field  $U$  is irrotational, geodesic and shear-free.

# Isotropic and spatially constant null sectional curvature

- Our main result extends Theorem 1 to the 3-dimensional case,

## Theorem 2

Let  $(M, g)$  be a 3-dimensional Lorentzian manifold. Then,  $(M, g)$  is conformally flat and has non-vanishing spatially constant, isotropic null sectional curvature if, and only if,  $(M, g)$  is locally a Robertson-Walker space.

*Proof.*

- From

$$\nabla_{\xi} U = -\frac{U[k]}{2(k + \mu)} \xi^{\perp} + g(\xi, U) \nabla^{\perp} [\ln(k + \mu)],$$

it follows easily that the vector field  $U$  is irrotational, geodesic and shear-free.

- Therefore, we can apply the Frobenius theorem, concluding that the distribution  $U^{\perp}$  is integrable.



# Proof of Theorem 2

- Moreover, the second fundamental form  $\hat{h}$  of the spacelike 2-surfaces  $\Sigma$  of the distribution is given by

$$\hat{h} = -\frac{U[k]}{2(k + \mu)}\hat{g},$$

being  $\hat{g}$  the induced metric on these surfaces.

# Proof of Theorem 2

- Moreover, the second fundamental form  $\hat{h}$  of the spacelike 2-surfaces  $\Sigma$  of the distribution is given by

$$\hat{h} = -\frac{U[k]}{2(k + \mu)}\hat{g},$$

being  $\hat{g}$  the induced metric on these surfaces.

- Therefore, these surfaces are totally umbilical in the 3-dimensional Lorentzian space  $M$  and, by the Gauss equation, their curvature  $\hat{K}$  is

$$\hat{K} = k + \left( \frac{U[k]}{2(k + \mu)} \right)^2.$$

# Proof of Theorem 2

- Moreover, the second fundamental form  $\hat{h}$  of the spacelike 2-surfaces  $\Sigma$  of the distribution is given by

$$\hat{h} = -\frac{U[k]}{2(k+\mu)}\hat{g},$$

being  $\hat{g}$  the induced metric on these surfaces.

- Therefore, these surfaces are totally umbilical in the 3-dimensional Lorentzian space  $M$  and, by the Gauss equation, their curvature  $\hat{K}$  is

$$\hat{K} = k + \left( \frac{U[k]}{2(k+\mu)} \right)^2.$$

- Because of the assumption that  $K^U = k + \mu$  is spatially constant, it follows that  $\hat{K}$  is constant on  $\Sigma$ .

# Proof of Theorem 2

- Moreover, the second fundamental form  $\hat{h}$  of the spacelike 2-surfaces  $\Sigma$  of the distribution is given by

$$\hat{h} = -\frac{U[k]}{2(k+\mu)}\hat{g},$$

being  $\hat{g}$  the induced metric on these surfaces.

- Therefore, these surfaces are totally umbilical in the 3-dimensional Lorentzian space  $M$  and, by the Gauss equation, their curvature  $\hat{K}$  is

$$\hat{K} = k + \left( \frac{U[k]}{2(k+\mu)} \right)^2.$$

- Because of the assumption that  $K^U = k + \mu$  is spatially constant, it follows that  $\hat{K}$  is constant on  $\Sigma$ .
- Summing up, the integral curves of  $U$  are geodesics whose perpendicular spaces integrate to form 2-dimensional surfaces of constant curvature.

# Proof of Theorem 2

- Moreover, the second fundamental form  $\hat{h}$  of the spacelike 2-surfaces  $\Sigma$  of the distribution is given by

$$\hat{h} = -\frac{U[k]}{2(k+\mu)}\hat{g},$$

being  $\hat{g}$  the induced metric on these surfaces.

- Therefore, these surfaces are totally umbilical in the 3-dimensional Lorentzian space  $M$  and, by the Gauss equation, their curvature  $\hat{K}$  is

$$\hat{K} = k + \left( \frac{U[k]}{2(k+\mu)} \right)^2.$$

- Because of the assumption that  $K^U = k + \mu$  is spatially constant, it follows that  $\hat{K}$  is constant on  $\Sigma$ .
- Summing up, the integral curves of  $U$  are geodesics whose perpendicular spaces integrate to form 2-dimensional surfaces of constant curvature.
- Then, the metric of  $M$  can be locally be written as

$$ds^2 = -dt^2 + f^2(t) \{ dr^2 + \Sigma^2(r, \kappa) d\theta^2 \},$$

with  $\Sigma(r, \kappa) = \sin r, r$  or  $\sinh r$  if  $\kappa = 1, 0$  or  $-1$ , respectively.

**THANK YOU VERY MUCH  
FOR YOUR ATTENTION!**