Gauge Alignment and Real-Space Locality in Higher-Rank Periodic One-Dimensional Hamiltonians

Analytic controls, numerical methodology, and evidence boundaries

Project Koios

Calculated-result technical report (unsealed)

Abstract

Matrix-valued tight-binding representations of an isolated composite subspace are not unique: a momentum-dependent unitary change of retained basis preserves the represented spectrum while changing individual Fourier hopping blocks and, consequently, the error incurred by finite-range truncation. This report develops a controlled periodic one-dimensional study of that distinction for retained ranks $r \in \{3, 4, 8, 16\}$ and noncommuting gauge-circuit depths $L \in \{1, 2, 4, 8\}$. The parent Hamiltonian, isolated retained subspace, and gauge attack are analytic. Four prescribed retained-space gauge alignments—identity, global-unitary, pointwise-unitary, and known-inverse-oracle—are compared using a common closed-loop transport of the analytic reference frame. Odd construction meshes support finite-periodic matrix Fourier transforms, while a disjoint shifted mesh separates interpolation error from deliberate finite-support truncation error. The design retains rank-normalized block tails, Parseval and Hermiticity defects, operator and ordered-spectrum errors, perturbation bounds, and mandatory mesh-refinement evidence.

Several conclusions follow analytically from the construction: the retained subspace has an exact uniform exterior-isolation energy $2.10E_{\mathrm{ref}}$; the gauge attack preserves the retained projector and spectrum; pointwise-unitary and known-inverse-oracle alignment recover the diagonal reference representation in exact arithmetic; and that recovered field has Fourier support only for $|R| \leq 2$. These are properties by construction, not new numerical findings. The maintained producer, typed evidence records, bounded producer-independent reference analysis of analytic-model and attack core invariants, bounded direct-sum all-gauge Fourier controls, and independent global/pointwise Procrustes-selection and constant-frame-transport slices are implemented and software-tested. A complete 16-case producer calculation has also been run locally and saved as canonical reusable result bytes for the fixed nine-figure inventory. Those snapshots are visibly marked "CALCULATED RESULT—UNSEALED" and are presented as calculated, integrity-admitted results available for scientific review. This status does not itself establish scientific validity, seal evidence, or confer publication acceptance. Their cache integrity and calculation-infrastructure identity are recorded; full independent recomputation of every serialized byte is not an ordinary production requirement.

Keywords: composite bands; gauge covariance; Procrustes alignment; parallel transport; matrix Fourier transform; hopping locality; truncation error; scientific evidence

1 Introduction

Localized real-space Hamiltonians are routinely obtained by Fourier transforming a matrix-valued reciprocal-space representation. For an isolated composite band subspace, the represented eigenvalues

are invariant under momentum-dependent unitary rotations of the retained basis, but the matrix entries and their Fourier coefficients are not. This familiar gauge freedom underlies both the usefulness and the interpretive difficulty of Wannier and tight-binding representations [1, 2, 3]. A short-ranged representation can be advantageous for interpolation and reduced modeling [4], yet spectral equivalence of the untruncated field alone does not imply that two gauges respond equally to a common finite-range truncation.

The present study isolates this issue from material-specific electronic-structure uncertainties. It does not run Quantum ESPRESSO, Wannier90, or a spread minimization. It instead supplies an analytic, finite-dimensional parent Hamiltonian with an exactly isolated retained subspace and then applies a deterministic higher-rank gauge circuit. The attacked representation has the same pointwise spectrum and projector as the reference representation. Any difference in its hopping locality is therefore a representation effect rather than a change of the underlying retained operator.

The central scientific question is:

> For a fixed isolated retained operator, how do retained rank, noncommuting gauge-circuit depth, and a predeclared gauge-alignment family affect matrix-valued hopping locality and the error of deliberate finite-support truncation?

This question is intentionally narrower than the problem of finding optimally localized Wannier functions. The study compares four declared feasible alignment families rather than searching an unrestricted localization functional. In particular, the analytic known inverse is retained only as an oracle control. It cannot support a claim about practical recovery from unknown data.

1.1 Contributions and organization

The report makes five methodological contributions.

(i) It defines a rank-generic analytic family with an exact, rank-independent exterior- isolation certificate and a connected noncommuting gauge attack.

(ii) It distinguishes common reference-frame transport from retained-space gauge alignment and fixes the left/right action conventions needed to compare them.

(iii) It separates identity, global-unitary, pointwise-unitary, and known-inverse-oracle gauge alignments from interpolation, truncation, and total *error channels*.

(iv) It defines an odd-mesh finite-periodic Fourier and support analysis with a disjoint shifted evaluation mesh and explicit mesh-refinement gates.

(v) It binds the scientific stages to immutable typed requests and fail-closed evidence records, while keeping independent recomputation and publication outside the producer.

Section 2 summarizes the scientific background. Sections 3–8 define the methodology. Section 9 states the analytic and implementation results currently established. Section 10 discusses their interpretation and limitations. Section 11 concludes. Detailed numerical controls, the Formal theorem crosswalk, and record ownership appear in the appendices.

1.2 Evidence status

This is a pre-retention technical report. The authoritative protocol fixes the coefficients, meshes, tolerances, schemas, resource limits, and future evidence policy. The maintained direct runner is explicitly development-only and writes no retained scientific evidence. The numerical figures in Section 9.3 come from one replaceable full-grid producer execution and are retained as report snapshots. Their canonical result bytes have passed manifest, digest, typed-round-trip, and complete-inventory integrity admission, but they are not backed by exhaustive independent recomputation, measured resource qualification, digest-gated rendering, or an evidence seal. The calculated figures are scientifically reviewable while unsealed. Cache admission establishes byte and inventory integrity, not physical validity or scientific validation, and neither admission nor presentation automatically confers publication acceptance.

2 Background and scientific setting

2.1 Represented operators and gauge covariance

Let $\mathcal{H}_{\mathrm{amb}}$ be a finite ambient state space and let $F(k) : \mathbb{C}^r \to \mathcal{H}_{\mathrm{amb}}$ be an orthonormal frame for a rank-r retained subspace at reciprocal coordinate k. For an ambient Hermitian operator $H_p(k)$, the represented retained operator is

$$H_F(k) = F(k)^\dagger H_p(k) F(k). \tag{1}$$

A retained-coordinate gauge transformation acts from the right, $F(k) \mapsto F(k)U(k)$ with $U(k) \in U(r)$, and therefore

$$H_F(k) \mapsto H_{FU}(k) = U(k)^\dagger H_F(k) U(k). \tag{2}$$

Equation (2) preserves the unordered pointwise spectrum and the ambient projector $P(k) = F(k)F(k)^\dagger$. It does not preserve individual matrix entries or Fourier coefficients.

This distinction is fundamental. A physical model is identified by its state spaces, operator, geometry, units, and energy reference. A finite matrix is a representation of that operator in a declared basis. Comparing matrices without first aligning their state spaces and basis conventions can confound a change of representation with a change of model.

2.2 Alignment versus localization

The orthogonal or unitary Procrustes problem selects the unitary factor that best aligns two frame matrices in a Frobenius norm [5]. Its polar factor can be computed from an SVD or polar decomposition [6, 7]. Pointwise Procrustes alignment supplies one unitary per reciprocal point. A global-unitary restriction instead supplies one matrix for the entire reciprocal route. These are different feasible sets, not different numerical accuracies for the same optimization problem.

Neither alignment is equivalent to minimizing a Wannier spread functional. The present protocol therefore does not claim to reproduce maximally localized Wannier functions. Its purpose is diagnostic: it asks how much locality is lost under a known gauge attack and how much is recovered under precisely declared alignment information.

2.3 Fourier locality and finite support

For a periodic represented field $H(k)$, continuum hopping blocks are

$$T(R) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ikR} H(k)\, dk, \qquad R \in \mathbb{Z}. \tag{3}$$

Rapid decay of $\|T(R)\|$ supports accurate finite-range approximation. Analytic periodicity often implies rapid Fourier decay, but a nonlinear momentum-dependent gauge rotation can generate an infinite harmonic inventory even when the reference field has finite support. Thus "same bands" and "same truncation behavior" are distinct statements.

A sampled finite transform introduces two additional distinctions. First, discrete coefficients are a finite representation of sampled data and may contain aliasing from unresolved continuum harmonics. Second, deliberately setting blocks outside a chosen support to zero is a model-reduction operation. The protocol keeps continuum behavior, finite-mesh interpolation, and deliberate truncation separate.

2.4 A gentle methods primer

A frame is a matrix whose columns are an orthonormal basis for the retained subspace. The subspace fixes the projector, but it does not fix those columns: if $A(k)$ is a frame and $Q(k)$ is unitary, then $A(k)Q(k)$ spans the same subspace. This is the frame, or gauge, ambiguity. The multiplication is on the right because $Q(k)$ mixes retained columns; left multiplication would instead act in the ambient state space. A represented operator consequently changes by unitary conjugation, even though its spectrum and ambient projector do not.

To align a candidate frame $A(k)$ to a reference frame $B(k)$, the right-action Procrustes problem minimizes

$$\|A(k)Q(k) - B(k)\|_{\mathrm{F}}^2 = \|A(k)\|_{\mathrm{F}}^2 + \|B(k)\|_{\mathrm{F}}^2 - 2\,\mathrm{Re}\,\mathrm{Tr}(Q(k)^* A(k)^* B(k))\,. \tag{4}$$

This expansion explains why the cross matrix is $A(k)^*B(k)$: it is the matrix paired with the right-acting $Q(k)^*$ in the trace. The reversed product corresponds to a different orientation, and products in the ambient space solve a different problem. In the notation used below, $A = C$ is the attacked candidate and $B = R$ is the reference. When $A(k)^*B(k) = Q(k)P(k)$ is an exact polar factorization with $P(k)$ positive semidefinite, $Q(k)$ minimizes the local objective; positive definiteness supplies the usual uniqueness condition.

The four alignment fields differ by what may vary with k. Identity alignment fixes $Q(k) = I$. Global alignment permits one constant Q for every point. Local, or pointwise, alignment permits a separately selected $Q(k)$ at each point. The known inverse is a special supplied local field satisfying $A(k)Q(k) = B(k)$ exactly in the analytic control; it is not information available in an ordinary inverse problem. These are nested information restrictions, not four error definitions.

Fourier locality begins only after a represented field and a gauge have been fixed. On a finite periodic mesh the transform is exact finite algebra for the sampled field. A single constant unitary conjugation passes through coefficient extraction, reconstruction, and support truncation. A momentum-dependent unitary generally does not: it can preserve every pointwise eigenvalue and Frobenius norm while redistributing weight among Fourier modes. Thus finite-mesh coefficient identities do not by themselves prove continuum Fourier decay, absence of aliasing, or convergence as the mesh is refined.

Finally, the report keeps the residual ledger explicit. Interpolation compares the full finite reconstruction with the direct field; deliberate truncation compares the support-truncated and full reconstructions; total error compares the truncated field with the direct field. Their fields add with a fixed sign convention, but their scalar maxima or RMS values need only satisfy norm inequalities. Gauge choice, finite Fourier representation, and these error channels are therefore related stages, not interchangeable notions of "alignment error."

3 Analytic model

3.1 Geometry, state spaces, and units

The selected one-dimensional specialization uses the dimensionless unit lattice and reduced Brillouin torus $\mathbb{T} = [-\pi, \pi)$. For retained rank $r \geq 3$, the ambient state space is $\mathbb{C}^{2r}$ with the retained coordinates ordered before the exterior coordinates. The canonical embedding $E_r : \mathbb{C}^r \to \mathbb{C}^{2r}$ and projector are

$$E_r = \begin{pmatrix} I_r \\ 0 \end{pmatrix}, \qquad P_0 = E_r E_r^\dagger. \tag{5}$$

All energies are expressed in a declared dimensionless unit with positive reference scale $E_{\mathrm{ref}} = 1$ during numerical evaluation. Formulas retain E_{ref} to expose the normalization.

The constant-projector control uses the analytic frame $F_r(k) = E_r$ and identity ambient sewing. The state-space map is therefore known directly rather than inferred from eigenvectors. Internal crossings among retained energies are permitted because the complete retained projector, not a sequence of individually labeled bands, is the scientific object.

3.2 Rank-normalized parent spectrum

For retained coordinate $a \in \{1, \ldots, r\}$, define

$$x_{r,a} = \frac{2a - r - 1}{r - 1} \in [-1, 1], \qquad c(k) = \frac{3}{4}\cos k + \frac{1}{4}\cos 2k. \tag{6}$$

The retained and exterior energies are

$$\frac{\epsilon_a(k)}{E_{\mathrm{ref}}} = 0.40 x_{r,a} + 0.25\,(1 + 0.20 x_{r,a})\,c(k), \tag{7}$$

$$\frac{\epsilon_a^{\mathrm{ext}}(k)}{E_{\mathrm{ref}}} = 3.00 + 0.30 x_{r,a} + 0.10 c(k). \tag{8}$$

Let $D_r(k) = \mathrm{diag}(\epsilon_1(k), \ldots, \epsilon_r(k))$ and define the ambient parent

$$H_p^{(r)}(k) = \begin{pmatrix} D_r(k) & 0 \\ 0 & D_{\mathrm{ext}}(k) \end{pmatrix}. \tag{9}$$

The represented reference operator is exactly $F_r^\dagger H_p^{(r)} F_r = D_r$.

Proposition 1 (Exact exterior isolation)**.** *For every selected finite rank and every $k \in \mathbb{T}$, the exterior spectrum lies at least $2.10 E_{\mathrm{ref}}$ above the retained spectrum. The lower bound is attained at $k = 0$.*

Proof. For fixed k, the retained maximum occurs at $x = 1$ and the exterior minimum at $x = -1$. Using Equations (7)–(8), their separation is $(2.30 - 0.20 c(k)) E_{\mathrm{ref}}$. Since $|c(k)| \leq 1$, the separation is at least $2.10 E_{\mathrm{ref}}$, and $c(0) = 1$ gives equality. □

The proposition is an analytic property of the finite model. A sampled minimum is retained separately as a numerical reconstruction check and is never relabeled as the continuum certificate.

3.3 Deterministic noncommuting gauge attack

Adjacent retained coordinates are mixed by real Givens rotations. For layer ℓ and pair start j, define

$$y_{r,j} = \frac{2j - r}{r - 1}, \qquad s_\ell(k) = \frac{3}{4}\sin k + \frac{1}{4}(-1)^\ell \sin 2k, \tag{10}$$

$$\theta_{\ell j}(k) = 0.35(-1)^{\ell+j}\,(1 + 0.15 y_{r,j})\, s_\ell(k). \tag{11}$$

Layer ℓ acts on the disjoint adjacent pairs with $j \equiv \ell \pmod 2$. Factors within one layer commute; neighboring layers generally do not. Increasing layer order is part of the definition. The depth-L product is denoted $G_{r,L}(k) \in SO(r)$.

The attacked frame and represented operator are

$$F'_{r,L}(k) = F_r(k) G_{r,L}(k), \qquad H'_{r,L}(k) = G_{r,L}(k)^\dagger D_r(k) G_{r,L}(k). \tag{12}$$

One layer is an incomplete-connectivity baseline; two or more alternating layers cover every adjacent edge and form a connected chain. Increasing depth changes the representation while leaving the retained operator and projector invariant.

Proposition 2 (Exact covariant equivalence)**.** *For every r, L, and k, the attacked frame is orthonormal, its ambient projector is P_0, and $H'_{r,L}(k)$ has the same unordered spectrum as $D_r(k)$.*

Proof. Each Givens factor is orthogonal, hence $G_{r,L}^\dagger G_{r,L} = I_r$. Substitution into Equation (12) proves frame orthonormality and projector equality. The represented operators are related by a unitary similarity, which proves spectral equivalence. □

The nonlinear entries of $G_{r,L}$ generally contain harmonics beyond those present in the angle functions. Consequently the attacked represented field is analytic and periodic but is not assigned a finite Fourier support by construction.

4 Gauge-alignment methodology

4.1 Canonical source and target orientation

Let $R(k) = F_r(k)$ be the reference, $C(k) = F'_{r,L}(k)$ the attacked candidate, and $H_C(k) = H'_{r,L}(k)$ the represented candidate operator. Every prescribed gauge alignment acts from the right:

$$\widehat{F}_U(k) = C(k)U(k), \qquad \widehat{H}_U(k) = U(k)^\dagger H_C(k) U(k). \tag{13}$$

Returning $U^\dagger$ while retaining Equation (13) would be a convention error, not an equivalent implementation.

For cross-covariance $X = C^\dagger R$ with complete SVD $X = L\Sigma V^\dagger$, the admitted unitary polar factor is

$$\mathcal{P}(X) = LV^\dagger. \tag{14}$$

It is selected only when

$$\sigma_{\min}(X) > \tau_{\mathrm{sv}}(X), \qquad \tau_{\mathrm{sv}}(X) = \max(10^{-12}, 10^{-10}\sigma_{\max}(X)). \tag{15}$$

Equality is deficient. No arbitrary null-space completion is introduced.

4.2 Four prescribed gauge alignments

Identity, wire value `identity`. $U_{\mathrm{id}}(k) = I_r$. This control measures the attacked representation as supplied.

Global-unitary, wire value `global-unitary`. One matrix minimizes the equal-weight objective

$$J_{\mathcal{S}}(U) = \frac{1}{|\mathcal{S}|r} \sum_{k \in \mathcal{S}} \|C(k)U - R(k)\|_{\mathrm{F}}^2. \tag{16}$$

The selection sample is exactly the final construction mesh $\mathcal{K}_{2049}$. The selected unitary is reused on the coarser construction meshes and shifted evaluation mesh without refitting.

Pointwise-unitary, wire value `pointwise-unitary`. At every construction point, $U_{\mathrm{pt}}(k) = \mathcal{P}(C(k)^\dagger R(k))$. Selection is repeated separately on each construction mesh. Selected pointwise unitaries are not interpolated to the shifted evaluation mesh; the direct recovered reference there is the analytic $D_r(k)$.

Known-inverse-oracle, wire value `known-inverse-oracle`. $U_{\mathrm{oracle}}(k) = G_{r,L}(k)^{-1} = G_{r,L}(k)^\dagger$. This action performs no optimization. It checks construction, orientation, application, and Fourier conventions. It is not a practical estimator.

The four names identify gauge alignments, not diagnostic channels. "Error channel" is reserved for the later interpolation, truncation, and total residual decomposition.

4.3 Alignment integrity evidence

For every selected Procrustes factor, the producer retains all singular values, the applied cutoff, strict rank margin, condition number when defined, and the objective recomputed from the original frames. It also evaluates

$$\varepsilon_{\mathrm{unit}} = \|U^\dagger U - I_r\|_{\mathrm{F}}, \qquad \varepsilon_{\mathrm{stat}} = \frac{\|U^\dagger X - X^\dagger U\|_{\mathrm{F}}}{\sqrt{r}}, \tag{17}$$

plus the covariance-objective identity and the negative part of the Hermitian polar factor. These checks distinguish a nominally returned matrix from a scientifically admitted alignment. The bounded independent selection reference reconstructs $X_{\mathrm{global}} = 2049^{-1} \sum_j G(q_j)^\dagger$ and evaluates its right polar factor as $X_{\mathrm{global}}(X_{\mathrm{global}}^\dagger X_{\mathrm{global}})^{-1/2}$ through an Hermitian eigendecomposition rather than the producer SVD solver. It uses $U_j = G(q_j)^\dagger$ directly for the pointwise controls and recomputes the retained rank, residual, integrity, and selected-unitary evidence.

Proposition 3 (Pointwise and oracle recovery)**.** *For the constant-projector construction, pointwise-unitary and known-inverse-oracle alignment both recover $F_r(k)$ and $D_r(k)$ in exact arithmetic.*

Proof. Here $C(k) = F_r G_{r,L}$ and $R(k) = F_r$, so $C(k)^\dagger R(k) = G_{r,L}(k)^\dagger$. This matrix is unitary and has all singular values equal to one. Its unique full polar factor is itself, which equals the known inverse. Substitution into Equation (13) gives the claim. □

The two actions remain separate in software because one reconstructs a factor from sampled frames while the other receives the generating inverse. Their agreement tests an implementation contract; it is not an empirical discovery.

5 Common closed-loop frame transport

Transport is applied to the analytic reference frame and is common to all four gauge alignments. It is not a fifth alignment and is not appended only to the pointwise case. This separation prevents transport defects from being misattributed to gauge-selection quality.

On the ordered construction mesh, let $Y_n = F_r(k_n)$ and set $\widetilde{F}_0 = Y_0$. At each neighbor link, form

$$O_n = \widetilde{F}_n^\dagger Y_{n+1} = L_n \Sigma_n R_n^\dagger, \qquad Q_n = R_n L_n^\dagger, \qquad \widetilde{F}_{n+1} = Y_{n+1} Q_n. \tag{18}$$

Then $\widetilde{F}_n^\dagger \widetilde{F}_{n+1}$ is Hermitian positive semidefinite. The strict rank rule in Equation (15) applies to every link. A deficient link terminates dependent transport; later links are unattempted rather than fabricated.

After traversing the period, write the endpoint mismatch as

$$\mathcal{H} = Z \operatorname{diag}(e^{i\gamma_1}, \ldots, e^{i\gamma_r}) Z^\dagger, \qquad \gamma_a \in (-\pi, \pi]. \tag{19}$$

The branch choice is part of the method. It assigns $+\pi$ to eigenvalue -1 and thereby fixes the distributed root. Closed frames are

$$F_n^{\text{closed}} = \widetilde{F}_n Z \operatorname{diag}(e^{-in\gamma_1/M}, \ldots, e^{-in\gamma_r/M}) Z^\dagger. \tag{20}$$

If $\widetilde{F}_M = \widetilde{F}_0 \mathcal{H}$, then $F_M^{\text{closed}} = F_0^{\text{closed}}$. For the constant reference frame and identity sewing, the exact model predicts trivial transport and zero endpoint phases; the numerical transport remains valuable as an action-orientation and closure control. The bounded independent transport reference instantiates that exact prediction directly: every overlap has singular spectrum $(1, \ldots, 1)$, every closure phase is zero, and all three retained closure right actions are I_r. It compares the link and closure evidence without invoking the transporter.

The phase tuple is an unordered multiset scientifically. Deterministic sorting is a wire convention, not a continuation of individual band labels through degeneracies.

6 Finite-periodic Fourier and locality methodology

6.1 Construction and evaluation meshes

The three construction meshes are

$$\mathcal{K}_M = \left\{ -\pi + \frac{2\pi n}{M} : n = 0, \ldots, M-1 \right\}, \qquad M \in \{513, 1025, 2049\}. \tag{21}$$

All are odd, centered, and half open. The direct evaluation mesh is

$$\mathcal{K}_{\text{eval}} = \left\{ -\pi + \frac{2\pi(m+1/2)}{8191} : m = 0, \ldots, 8190 \right\}. \tag{22}$$

It is disjoint from every construction mesh and cannot influence a fit, support, tolerance, or reporting rule.

For odd M, the canonical translation inventory is $\mathcal{R}_M = \{-(M-1)/2, \ldots, (M-1)/2\}$. There is no self-inverse Nyquist singleton; the canonical disposition is `absent-on-odd-mesh`.

6.2 Discrete transform, diagnostic ranges, and support policy

For one gauge alignment a, the finite-periodic transform is

$$T_{a,M}(R) = \frac{1}{M} \sum_{k \in \mathcal{K}_M} e^{-ikR} H_a(k), \qquad H_{a,M}(k) = \sum_{R \in \mathcal{R}_M} e^{ikR} T_{a,M}(R). \tag{23}$$

Only the forward transform carries $1/M$. The complete coefficient inventory is used for reconstruction and tail calculations. The public result retains the bounded norm inventory $|R| \leq 128$ rather than serializing all dense coefficient arrays.

The declared integer ranges serve two scientifically distinct roles. In the hopping-diagnostic inventory, a range partitions the unchanged coefficient field into included and omitted blocks for measurement only. In the retained-support inventory, the same integer interval defines a new field by setting exterior coefficients to zero. Numerically identical ranges therefore remain non-interchangeable: the first operation is diagnostic, whereas the second is an irreversible model-reduction action.

The declared maximum ranges are

$$\Omega_S = \{R \in \mathbb{Z} : |R| \leq S\}, \qquad S \in \{0, 1, 2, 4, 8, 16, 32, 64, 128\}. \tag{24}$$

Every range contains inversion pairs. All nine ranges are evaluated in both typed inventories; no first-passing or plot-selected range is promoted.

For the reference diagonal field, Equations (7) and (3) imply an exact support certificate.

Proposition 4 (Reference support)**.** *The continuum Fourier coefficients of $D_r(k)$ vanish for every $|R| > 2$.*

Proof. Every diagonal entry of D_r is a linear combination of 1, $\cos k$, and $\cos 2k$. These harmonics have Fourier indices only at 0, ± 1, and ± 2. □

Corollary 1. *In exact arithmetic, pointwise-unitary and known-inverse-oracle alignment recover a represented field with continuum support $|R| \leq 2$.*

This is an analytic certificate. Finite-transform reconstruction and binary64 residuals still require numerical checks.

6.3 Tail and error diagnostics

For hopping-diagnostic cutoff S, define the omitted quadratic weight, its square root, and the block-ℓ^1 tail from the unchanged coefficients:

$$W_{a,M}(S) = \sum_{R \notin \Omega_S} \|T_{a,M}(R)\|_{\mathrm{F}}^2, \tag{25}$$

$$N_{a,M}(S) = \sqrt{W_{a,M}(S)}, \tag{26}$$

$$L_{a,M}(S) = \sum_{R \notin \Omega_S} \|T_{a,M}(R)\|_{\mathrm{F}}. \tag{27}$$

Cross-rank comparisons retain $W/(rE_{\mathrm{ref}}^2)$ and $N/(\sqrt{r}E_{\mathrm{ref}})$ in addition to raw values. This predeclared normalization prevents matrix dimension alone from determining a trend.

For retained support Ω_S, the shifted-mesh full interpolant, truncated interpolant, and direct analytic reference define

$$e_{\text{interp}}(k) = H_{a,M}(k) - H_a^{\text{direct}}(k), \tag{28}$$

$$e_{\text{trunc}}^{(S)}(k) = H_{a,M}^{(S)}(k) - H_{a,M}(k), \tag{29}$$

$$e_{\text{total}}^{(S)}(k) = H_{a,M}^{(S)}(k) - H_a^{\text{direct}}(k). \tag{30}$$

The pointwise identity $e_{\text{total}} = e_{\text{interp}} + e_{\text{trunc}}$ is checked directly. Scalar maxima of the three error channels are not added: their maximizing points can differ, and cancellation in the total error is reported rather than called convergence.

For each residual, the study records maximum and RMS Frobenius operator error and maximum and RMS ordered-spectrum error. Ordered spectra are compared as sorted multisets rather than by tracking eigenvectors through crossings. At every point, the finite-dimensional bounds

$$\max_i |\lambda_i(H_{\text{approx}}) - \lambda_i(H_{\text{reference}})| \le \|e\|_2 \le \|e\|_{\text{F}} \tag{31}$$

are checked [8]. Qualified saturation ratios are reported only when their denominators exceed the frozen floor.

7 Numerical study design

The complete matrix contains 16 rank/depth cases, three construction meshes, four gauge alignments, and nine supports. Before failure dispositions, this corresponds to 192 mesh/alignment combinations and 1728 support rows. The global-unitary fit occurs once per case at $M = 2049$ and is reused; it is not counted as a new fit on coarser or evaluation meshes.

Figure 1 summarizes the scientific flow. Solid arrows denote fields used by later calculations. Dashed arrows denote common transport evidence that is correlated into the aggregate result but does not redefine a gauge alignment.

7.1 Refinement and qualification

The smaller meshes are mandatory refinement controls. Blockwise comparisons cover both adjacent mesh pairs. The final scalar inventory uses the mandatory $1025 \to 2049$ pair and the absolute-relative rule

$$|q_f - q_c| \le \tau_{\text{abs}} + \tau_{\text{rel}} \max(|q_c|, |q_f|), \tag{32}$$

with a denominator floor for the separately reported relative change. The shared `ScalarRefinementAnalyzer` owns this arithmetic; the higher-rank diagnostic inventory owner defines which scientific quantities enter the ledger.

A case is numerically qualified only if its mandatory construction, model, transport, alignment, Fourier, interpolation, and refinement gates pass. A global-unitary aggregate may instead carry the bounded disposition `aggregate-rank-deficient`; this makes only global-unitary downstream values inapplicable. Pointwise rank deficiency, documented backend failure, alignment integrity failure, or oracle integrity failure invalidates the complete case.

7.2 Reasonable accuracy scales and computational precision

A numerical tolerance is an acceptance standard, not a prediction that every residual will stop at that value. For normalized binary64 calculations near unit scale, machine epsilon is approximately




Figure 1: Scientific flow of one higher-rank case. Gauge alignment is performed before Fourier transformation and deliberate truncation.

2.22×10^{-16}. Matrix products, Fourier sums over up to 2049 points, eigensolvers, and repeated factorizations normally accumulate more error than one machine epsilon. Consequently, residuals of order 10^{-12} are a reasonable strict standard for structural identities in this calculation; 10^{-11} to 10^{-10} are reasonable standards for Fourier, spectrum, transport, and recovery comparisons; and the complete-interpolant operator gate is deliberately looser at 10^{-9}. The scientific effect-resolution scale is 10^{-6}, several orders of magnitude above all of these numerical gates. Table 4 gives the diagnostic-specific values and normalizations.

Some present controls nevertheless produce measured residuals of order 10^{-16}, and one decomposition defect is smaller still. Those values arise where an analytic identity, finite support, or nearly exact cancellation is represented especially faithfully in binary64 arithmetic. They show that the measured residual reached the roundoff scale; they do not establish 16 or more reliable decimal digits, justify a 10^{-18} acceptance tolerance, or improve the scientific resolution below 10^{-6}. Exact floating-point zero is reported separately because it is a representational outcome, not evidence of infinite accuracy. In logarithmic figures, 10^{-18} is only an axis-placement device that keeps exact zeros and exceptionally small positive residuals visible; all values at that scale are interpreted as computational-precision or roundoff-scale results.

7.3 Claim-resolution rules

The protocol does not infer a locality effect from visual separation alone. For a qualified case, gauge-induced locality extension at $S = 2$ is resolved only if the identity tail is at least the frozen 10^{-6} scale, the oracle tail is at most 10^{-10}, and the identity total ordered-spectrum maximum error after $S = 2$ truncation is at least $10^{-6} E_{\mathrm{ref}}$. Rank or depth differences are descriptive unless their

normalized difference exceeds both 10^{-6} and ten times the larger observed refinement change. These rules are fixed before execution and do not assert that an effect will be found.

No monotonic trend with rank or depth is assumed. Noncommuting products can produce cancellation, and a finite ladder cannot establish an asymptotic law.

8 Diagnostic and evidence methodology

8.1 Metric-explicit diagnostics

Every diagnostic names its norm and normalization. Frame and unitary residuals use $\|\cdot\|_{\mathrm{F}}/\sqrt{r}$ when rank-normalized; operator residuals use $\|\cdot\|_{\mathrm{F}}/(\sqrt{r}E_{\mathrm{ref}})$. Maximum-absolute-entry defects used internally by a backend are not relabeled as Frobenius diagnostics. Raw and normalized quantities remain available for different scientific questions.

The transform checks include representative count, uniqueness, ordering, inversion closure, reconstruction, Hermitian pairing, Parseval equality, nested-support monotonicity, and the explicit odd-mesh Nyquist disposition. These are integrity checks, not alternative definitions of locality.

8.2 Fail-closed numerical outcomes

Rank deficiency, integrity failure, and backend failure have different meanings and remain distinct typed outcomes. Documented backend failures are converted into stable codes for nonfinite arithmetic, nonconvergence, or validation failure. Runtime exception messages, tracebacks, host paths, and fabricated numerical placeholders are excluded. Downstream values that were not scientifically established are absent, not zero.

8.3 Typed stage ownership

The maintained producer decomposes one gauge-alignment analysis into recovery, Fourier, interpolation, locality, and assembly stages. Each stage binds a complete immutable request and exposes a zero-argument execution. Stage evidence retains its exact source request. Assembly rejects same-shaped but uncorrelated arrays, a replacement request, wrong mesh coordinates, mismatched case identity, wrong gauge alignment, or inconsistent selection provenance.

Stage owner	Scientific responsibility	Evidence boundary
`fields.py`	Construct one immutable represented operator and frame field.	No parallel uncorrelated dictionaries.
`recovery.py`	Compare fields with analytic or solver-owned references.	Frame and operator recovery remain separate.
`fourier.py`	Apply the PhysKit odd-mesh transform and structural checks.	Coefficients are not silently normalized or reindexed.
`interpolation.py`	Evaluate complete Fourier series and direct fields on the shifted mesh.	Evaluation coordinates are immutable and disjoint.
`locality.py`	Evaluate every support and the three error channels.	No preferred support is selected.
`assembly.py`	Correlate stages and construct the public mesh result.	No scientific recomputation during assembly.

Canonical typed JSON bytes and SHA-256 digests establish record identity and integrity; one explicit SHA-256 identifies the selected scientific calculation infrastructure. They do not establish scientific correctness. The bounded references bypass the corresponding producer actions, while rendering consumes saved results without calculating new scientific quantities.

9 Results established before retained execution

9.1 Analytic results

Table 1 collects the results proved from the selected formulas. They require no numerical execution and must not be reported as empirical observations.

Result	**Statement**	**Interpretive boundary**
Exterior isolation	$\Delta_{\mathrm{iso}} = 2.10 E_{\mathrm{ref}}$ for every selected rank.	Exact finite-model certificate; not a material band gap.
Gauge covariance	Attack preserves projector and pointwise spectrum.	Does not preserve matrix entries or hopping locality.
Pointwise recovery	Pointwise polar factor equals $G^{\dagger}_{r,L}$.	Exact constant-projector property; not evidence for noisy or unknown frames.
Oracle recovery	Known inverse restores F_r and D_r.	Construction control, not a fitted solution.
Reference support	$T_{D_r}(R) = 0$ for $\lvert R\rvert > 2$.	Continuum statement; finite-transform defects remain numerical.
Hermitian pairing	$T(-R) = T(R)^{\dagger}$.	Exact continuum identity tested separately on each finite mesh.

Table 1: Analytic results available before numerical execution.

The exact results establish a controlled contrast. Identity and global-unitary alignment are not assigned finite support by construction, while pointwise-unitary and oracle alignment recover a field whose exact support is known. Whether the selected finite attack produces a *resolved* tail or truncation effect at each (r, L) remains a numerical question governed by the frozen thresholds.

9.2 Implementation and software-verification results

The maintained Python producer implements the analytic family, global and pointwise Procrustes selection, common PhysKit transport, odd-mesh finite-periodic transforms, shifted interpolation, support diagnostics, refinement, immutable typed records, and canonical serialization. The former monolithic gauge-alignment analysis has been split into cohesive request-bound scientific owners. A separate bounded reference action now recomputes all four gauge-aligned Fourier blocks by direct sums on every construction mesh. It compares 930 reconstruction, Parseval, Hermitian-pairing, retained-block, shifted-interpolation, support-tail, and oracle exact-support quantities without invoking producer analysis actions. Here producer-independent means a direct-formula NumPy path that bypasses the corresponding scientific actions, not a separate numerical-library stack; NumPy/LAPACK and typed quantity/comparison boundaries remain shared. A third bounded action reconstructs the global and pointwise Procrustes selections without the producer model or SVD solver and compares 57 retained rank, residual, integrity, and selected-unitary quantities. A fourth bounded action directly reconstructs 45 link-rank and closure quantities for the constant-frame transport on all three meshes without calling the producer transporter. Gauge-alignment wire values are descriptive and strictly mapped from a plain domain enum; obsolete opaque values and the former ambiguous key are rejected.

At the time of this report revision, the focused higher-rank package suite passes 78 tests, the calculation-local structural suite passes 4 tests, and the complete bounded package suite passes 4715

tests with 23 environment- or historical-tree-dependent skips and 89 explicitly expensive deselections. Ruff, strict source mypy, strict Sphinx, `compileall`, LaTeX builds, and `git diff -check` also pass. These results establish current software behavior and document buildability only. They do not constitute independent numerical verification of a scientific result.

9.3 Calculated numerical snapshot: scope and admission boundary

The fixed report renderer was exercised on all 16 rank/depth cases. It assembles immutable plotting arrays from the final producer objects and renders the nine views declared before this full-grid execution. The calculation completed all requested producer paths and wrote canonical typed case bytes to a reusable calculation cache. A separate integrity admission binds their manifest digest and checks canonical round trips and complete inventory. The admitted manifest SHA-256 is `2bba6a5a208945018dc462a142412e0d6cfaba436a58fec03a1cd33cc48d4673`; it records calculation-infrastructure SHA-256 `4db05dadca1818469484b132ba0a7776c0a9281ee956e6cf94490104de0a70dd`. That infrastructure identity binds the explicitly selected scientific producer and numerical collaborator sources, not plotting or report sources. Presentation changes can therefore regenerate all nine views from the admitted result bytes without repeating the scientific calculation. Cache admission is not an independent scientific verification record. Every panel therefore carries the neutral status label *CALCULATED RESULT—UNSEALED*. The numerical values below are calculated, integrity-admitted results available for scientific review. They are neither illustrative synthetic data nor automatically publication-accepted findings; the unsealed status and disclosed coverage boundaries remain part of their interpretation. As discussed above, logarithmic panels use a display floor of 10^{-18} solely for axis placement; exact zeros remain distinct and are marked explicitly where they occur. The floor is neither an acceptance standard nor a claim of accuracy beyond binary64 computational precision.

9.3.1 $S = 2$ alignment–locality plane

Figure 2 places the final-mesh normalized frame-recovery defect on the horizontal axis and the normalized omitted-block norm beyond the hopping-diagnostic cutoff $S = 2$ on the vertical axis. Four gauge panels each retain all 16 cases; rank is encoded by marker, depth by color, and the paired $M = 1025$ and $M = 2049$ marks make mesh movement visible without converting it to a percentage near zero. Across the final mesh, identity and global-unitary tails span 8.02×10^{-4} to 9.52×10^{-2}, while the pointwise and oracle tails span approximately 1.38×10^{-16} to 2.80×10^{-16}. Identity and global frame defects remain between 2.30×10^{-1} and 1.26, whereas pointwise and oracle frame defects are at binary64 roundoff. The global and identity locality ordinates coincide to about 10^{-16} because a constant right unitary conjugates every Fourier block and preserves its Frobenius norm; this is a feasible-set invariant, not evidence that the global fit recovered the reference frame.

9.3.2 Hopping diagnostics and Fourier-block locality

Figure 3 evaluates the unchanged Fourier coefficients against every declared hopping-diagnostic cutoff. These inventories diagnose omitted block weight; they do not themselves apply a support-truncation policy. Identity and global-unitary tails decrease across several cutoffs before reaching the roundoff floor, whereas the pointwise and oracle tails reach that floor at $S = 2$, consistently with the analytic support certificate. No cutoff is omitted because its curve appears visually redundant. At the fixed interior anchor $(r, L) = (8, 4)$, the identity tail decreases from 1.52×10^{-1} at $S = 0$ to 1.21×10^{-2} at $S = 2$, 6.64×10^{-6} at $S = 8$, and 2.23×10^{-11} at $S = 16$ before reaching the numerical floor near




Figure 2: Calculated, unsealed alignment–locality plane for all 16 cases at hopping-diagnostic cutoff $S = 2$. The axes report absolute normalized operands; no identity-to-oracle ratio is formed from roundoff-scale denominators.

$S = 32$. The global-unitary values agree at the displayed precision. The pointwise and oracle tails are already about 2×10^{-16} at $S = 2$.

Figure 4 resolves the same mechanism coefficient by coefficient for the fixed $(8, 4)$ anchor. Identity and global-unitary retain a decaying sequence beyond $|R| = 2$; pointwise and oracle show only the $R = 0, 1, 2$ analytic blocks above roundoff. The plot is a finite $M = 2049$ coefficient inventory, so the apparent floor is numerical and does not establish a continuum decay rate.

9.3.3 Interpolation, truncation, and total-error channels

Figure 5 applies each declared retained-support policy and displays the resulting three error channels as separate, non-stacked curves. The complete-interpolant error is support independent and stays near binary64 roundoff at the anchor. It is therefore much smaller than truncation over most of the support path, so the truncation and total curves nearly coincide and one can obscure the other if drawn with identical markers. The revised figure draws total as a thin solid green curve and truncation as a thicker dashed orange curve with open square markers, making both visible where their numerical values overlap. For identity and global-unitary, truncation and total errors fall across the full support path; for pointwise and oracle they reach the numerical floor at $S = 2$. At $(8, 4)$, the identity normalized total ordered-spectrum maximum error decreases from 3.19×10^{-1} at $S = 0$ to 2.31×10^{-2} at $S = 2$, 8.87×10^{-6} at $S = 8$, and 3.42×10^{-11} at $S = 16$. The pointwise and oracle values at $S = 2$ are about 10^{-15}. The bottom row confirms the residual-field decomposition numerically; over the complete calculated grid, the largest normalized decomposition defect is 3.96×10^{-17}. These curves are not stacked because the residual fields add pointwise but their separately maximized norms need not.

9.3.4 Frozen claim operands and refinement

Figure 6 reports three absolute $S = 2$ locality-extension operands and one supporting global-frame diagnostic, preserving every rank/depth cell. The tail operands use the hopping-diagnostic inventory, whereas the total ordered-spectrum operand uses the retained-support analysis at the same integer range. Equal numerical cutoffs therefore do not make the two inventories interchangeable. In this calculated snapshot, all 16 cells satisfy each displayed inequality: identity tails are at least 10^{-6}, oracle tails are at most 10^{-10}, identity total ordered-spectrum errors are at least 10^{-6}, and global




Figure 3: Calculated, unsealed hopping-tail diagnostics across the declared cutoff inventory. Rows denote retained rank, columns denote gauge alignment, and line style and color denote circuit depth. The ordinate is computed from unchanged coefficients; it is not a truncation error.




Figure 4: Calculated, unsealed nonnegative Fourier-block norms at the fixed interior anchor. The vertical line marks the exact reference support boundary $|R| = 2$.




Figure 5: Calculated, unsealed non-stacked trajectories for interpolation, truncation, and total error at $(r, L) = (8, 4)$. Columns are gauge alignments; the first two rows contain all three channels for operator and ordered-spectrum error. The bottom row is the separate residual-field decomposition-identity diagnostic, not a fourth error channel.

frame residuals are at least 10^{-6}. The identity $S = 2$ total ordered-spectrum error ranges from 1.64×10^{-3} to 2.03×10^{-1}, and the global frame residual ranges from 2.30×10^{-1} to 1.25. All 16 cells satisfy each of the three available frozen claim operands. The global-frame panel exposes restricted-feasible-set behavior but is not the protocol's fourth frozen operand, the global represented-operator residual, which the current public result does not retain. The complete frozen locality-extension rule therefore remains unresolved. Neither the supporting panel nor the unsealed figure can assign a retained claim disposition.




Figure 6: Calculated, unsealed locality-extension operands and supporting global-frame matrix. Tail panels report hopping diagnostics; the ordered-spectrum panel reports a retained-support consequence. Color encodes the logarithm of the absolute normalized operand, and green or red cell borders denote whether the displayed condition is met. Missing, failed, aggregate-rank-deficient, and inapplicable states remain categorical rather than being colored as numerical zero.

Figure 7 keeps the $M = 1025$ and $M = 2049$ values paired and also shows each absolute change divided by its applied absolute–relative tolerance. The largest gate utilization in the calculated run is 5.55×10^{-6}, for an interpolation-spectrum diagnostic; every displayed utilization is far below one. This supports internal mesh stability of the producer output but does not replace independent recomputation of either side of the pair.

9.3.5 Rank/depth scaling, recovery controls, and sampled Pareto sets

Figure 8 compares ranks only at fixed depths and depths only at fixed ranks through paired $S = 2$ total-operator and total-spectrum heatmaps. In the identity and global-unitary panels, this finite snapshot shows larger errors for deeper circuits and generally smaller normalized errors at larger rank. For example, the identity ordered-spectrum matrix decreases down each fixed-depth column as r increases, while increasing across each fixed-rank row as L increases. This is a description of the selected finite grid, not a monotonicity theorem: the complete support paths contain visible nonmonotonic structure, and additional ranks or depths could reverse a finite trend. Pointwise and oracle $S = 2$ panels are dominated by roundoff and cannot support rank/depth effect ratios.

Figure 9 separates frame recovery from represented-operator construction integrity. Identity and global-unitary do not recover the reference frame, while pointwise and oracle frame recovery is at roundoff. The lower-row represented-operator recovery values compare each aligned field with its separately declared direct construction; identity and global therefore also pass this integrity check at zero. Those lower panels must not be read as claiming that identity or global alignment recovered D_r.




Figure 7: Calculated, unsealed 1025 → 2049 refinement diagnostics. The dashed line in the paired-value panel is equality; the dashed line in the utilization panel is the frozen pass boundary.

Finally, Figure 10 marks sampled nondominated supports within each case and gauge alignment. One joint mask is computed from the three-objective tuple $(2S+1,\ E^{\mathrm{total}}_{\mathrm{operator}},\ E^{\mathrm{total}}_{\mathrm{spectrum}})$ and reused in both displayed error columns. Identity and global-unitary have seven to nine efficient supports per case because additional blocks trade higher support cost for lower error. Pointwise has three to five and the oracle three to six. Beyond $S = 2$, the control errors are roundoff-scale, so exact floating-point differences can leave several supports nondominated under the tolerance-free rule. Pareto membership therefore selects no preferred support, does not pool gauge feasible sets, and does not turn the oracle into a practical optimizer.

9.4 Current numerical-result boundary

Canonical result bytes for all 16 cases have been admitted by manifest and case-file integrity checks. The implemented bounded references do not independently cover every support-truncated and total-error field or every refinement value, so cache admission is not itself scientific validation. The figures must distinguish calculated values from analytic certificates and preserve unresolved, failed, missing, and inapplicable states. The calculated figures are scientifically reviewable, but remain unsealed. Publication acceptance remains a separate human decision and is not automatically conferred by cache admission, reviewability, or presentation; changing presentation does not require a new calculation.

10 Discussion

10.1 What the controlled comparison can establish

The construction isolates representation dependence. Since the parent, projector, and untruncated retained spectrum are held fixed, a resolved difference in finite-support error would demonstrate that short-range compressibility depends on retained-space gauge within this finite analytic family. It would not demonstrate that the parent model is physically realistic or that one gauge is universally optimal.




Figure 8: Calculated, unsealed rank/depth scaling at the predeclared retained support $S = 2$. Rows are gauge alignments; columns distinguish total operator and total ordered-spectrum error. Numeric annotations retain the absolute normalized values.




Figure 9: Calculated, unsealed final-mesh gauge-recovery controls. The two rows answer different questions: reference-frame recovery above and direct represented-field construction integrity below. Exact zeros are marked "0" rather than being identified with the 10^{-18} logarithmic display floor.

Pointwise-unitary and oracle recovery form a strong implementation control because their exact target is known. Global-unitary alignment is scientifically different: one matrix cannot generally undo a momentum-dependent noncommuting attack. A nonzero qualified global-frame residual is a supporting measure of the limitation of that restricted feasible set, not solver failure. Only the separately frozen maximum normalized represented-operator residual resolves the protocol's global-alignment claim, and that operand is not retained by the current public result. Conversely, a small residual of either declared kind at one finite case would not imply equivalence of global and pointwise alignment in general.

10.2 Interpreting rank and depth

Retained rank changes matrix dimension and the number of adjacent mixing edges. Circuit depth changes the ordered product of generally noncommuting rotations. The selected $\sqrt{r}$ and r normalizations make cross-rank magnitudes interpretable without removing all rank dependence. Raw values remain necessary because normalization itself answers a different question.

Depth is not a scalar attack amplitude. Additional layers can reinforce or cancel existing harmonics, so locality need not vary monotonically with L. The finite set $L \in \{1, 2, 4, 8\}$ supports controlled comparisons only; it does not define a continuum or asymptotic-depth limit.

10.3 Transport and topological scope

The common closed-loop transport tests numerical continuity, endpoint sewing, and branch conventions. In the implemented control the exact reference projector is constant and topologically trivial. Closure phases are therefore transport diagnostics, not polarization, Wannier centers, or a topological invariant. The distributed closure is a smooth-gauge construction, not a localization optimum.




Figure 10: Calculated, unsealed sampled support–error Pareto diagnostics. Rows are gauge alignments; columns use total operator and ordered-spectrum error. Marked supports are nondominated only within their own case and gauge alignment.

A moving-projector or topologically nontrivial study would require patching, sewing, and obstruction contracts not present here. Such a study cannot be inferred by replacing the constant frame with backend eigenvectors while retaining the present evidence semantics.

10.4 Higher dimensions and nonorthonormal bases

Fiberwise equations such as $F'U$ and $U^\dagger H'U$ do not intrinsically select a spatial dimension. Meshes, Fourier indices, inversion closure, and support geometry do. The present Fourier and locality actions are therefore deliberately one-dimensional. A 2-D or 3-D study must define reviewed tensor or nonorthogonal reciprocal meshes, multi-index representatives, complete point-group or inversion orbits, and a physical support metric. No placeholder implementation is supplied.

Likewise, the current frames are orthonormal and their gauge group is $U(r)$. A nonorthonormal basis requires a positive-definite fiber metric, metric adjoint, generalized isometry group, conditioning evidence, and a metric-aware Procrustes objective. Treating that change as an optional parameter of the present action would hide a new mathematical contract.

10.5 Reproducibility and evidence lifecycle

The direct lifecycle is calculate once → verify cache integrity → render many times. The producer owns scientific calculation; bounded references provide independent controls for selected formulas and algorithms; and the renderer owns presentation only.

The calculation-local `workflow/` directory is historical provenance and remains outside the maintained runtime and tests. `run_01_calculation.py` composes the maintained owners explicitly and writes one canonical cache record per case plus a manifest containing `calculation_infrastructure_sha256`. `run_02_verification.py` checks exact SHA-256 digests, canonical typed round trips, case identities, and the complete 16-case inventory. `run_03_render.py` renders the fixed views repeatedly from the matching admitted cache without recalculating the grid. Cache integrity does not establish scientific validity, and none of these entrypoints publishes or releases a result.

10.6 Limitations

The principal limitations are:

1. the parent is synthetic, finite dimensional, and dimensionless;
2. only the one-dimensional constant-projector, real-orthogonal specialization is implemented;
3. the oracle assumes knowledge unavailable in an ordinary material calculation;
4. bounded core-invariant, all-gauge Fourier, global/pointwise Procrustes, and constant-frame transport reference analyzers are implemented; they are producer-path independent under a shared NumPy/LAPACK stack but do not independently recompute every saved quantity;
5. the finite rank/depth ladder supports no asymptotic law;
6. no 2-D, 3-D, nonorthonormal, moving-subspace, or topological result is available; and
7. passing tests does not establish scientific validation.

11 Conclusion

This report defines a practical methodological account of a controlled higher-rank gauge-locality study without overstating its evidence status. The analytic family cleanly separates a physical retained operator from its finite matrix representation. A deterministic noncommuting gauge circuit can change matrix-valued hopping locality while preserving the retained projector and pointwise spectrum; the finite change shown here is a calculated, integrity-admitted observation available for scientific review, while remaining unsealed and separate from publication acceptance. Four predeclared gauge alignments probe different information constraints, and a common transport control prevents frame-sewing effects from being conflated with gauge selection.

The exact construction already establishes a uniform isolation energy, covariant spectral equivalence, pointwise and oracle recovery, and finite reference support. The software producer implements the planned finite numerical analysis with typed immutable evidence and fail-closed outcomes. The replaceable full-grid calculated snapshot exhibits the expected contrast between long-tailed identity/global representations and finite-support pointwise/oracle controls, together with stable $1025 \to 2049$ producer diagnostics. It also shows finite rank/depth structure and sampled support–error tradeoffs. Those observations are integrity-admitted and scientifically reviewable, but remain unsealed; the implemented bounded controls and disclosed limitations define the present evidence boundary, and no automatic publication acceptance follows.

The saved canonical results now support fast presentation work without recalculation. Further scientific review may add targeted independent controls where they materially affect a claim, while publication acceptance remains a separate human decision.

A Formal theorem crosswalk and proof sketches

A.1 Pinned revision and reading rule

This appendix summarizes exact finite-dimensional results mechanized in the separate Project Koios–Formal repository. The metadata and declarations were inspected at the exact merged revision

`4902864c19b412320e56c984fe9cc98a4ad3bd89`

("Merge pull request #42: Formalize finite-mesh unitary Procrustes hierarchy"). At that revision, the Formal theorem manifest records every identifier cited below as *proved; no admissions*. The reported axiom sets contain only `propext`, `Classical.choice`, and `Quot.sound`. This is a revision-pinned cross-reference, not a claim that the present runtime produced or checked a Formal certificate.

A.2 Right-action alignment hierarchy

Let K be a finite mesh, let $A_k, B_k \in \mathbb{C}^{m\times r}$, and define

$$J_K(U) = \sum_{k\in K} \|A_k U_k - B_k\|_{\mathrm{F}}^2. \tag{33}$$

Suppose exact witnesses satisfy $A_k^* B_k = Q_k P_k$, with Q_k unitary and $P_k \succeq 0$.

Proposition 5 (Exact finite-mesh Procrustes hierarchy)**.** *The field $Q = (Q_k)$ globally minimizes Equation (33) among all pointwise unitary fields. It is no worse than any constant unitary field and no worse than identity. If every $P_k \succ 0$, equality with the minimum forces equality with Q. An exact known-inverse field O satisfying $A_k O_k = B_k$ has zero objective, is a global minimizer, and equals Q under the same positive-definiteness hypothesis.*

Proof sketch. Equation (4) reduces minimization to maximizing a real trace. For the factorization $A_k^* B_k = Q_k P_k$, write any competitor relative to Q_k; a unitary cannot increase $\mathrm{Re}\,\mathrm{Tr}(W P_k)$ above $\mathrm{Tr}(P_k)$. Positive definiteness makes equality rigid. Summing the local inequalities proves the finite-mesh statement. A known inverse makes every squared norm vanish; nonnegativity and uniqueness then give the final assertions. These are precisely the roles of `BG-ALIGNMENT-THM-041`, `BG-ALIGNMENT-THM-044`, `BG-ALIGNMENT-THM-045`, and `BG-ALIGNMENT-THM-052`–`BG-ALIGNMENT-THM-058`. □

Table 2: Exact alignment results used in the proof sketch.

Formal identifier	Mechanized result and boundary
`BG-ALIGNMENT-THM-041`	Expands the right-action objective; the cross matrix is exactly A^*B.
`BG-ALIGNMENT-THM-044`	A supplied positive-semidefinite polar witness gives a global local minimum.
`BG-ALIGNMENT-THM-045`	A positive-definite polar factor makes that minimizer unique.
`BG-ALIGNMENT-THM-052`	Pointwise polar minima jointly minimize the unweighted finite-mesh sum.
`BG-ALIGNMENT-THM-053`	Equality with the summed minimum forces field equality when every positive factor is positive definite.
`BG-ALIGNMENT-THM-054`	The pointwise polar field is no worse than any constant unitary field.
`BG-ALIGNMENT-THM-055`	The pointwise polar field is no worse than identity.
`BG-ALIGNMENT-THM-056`	A supplied exact known inverse has zero summed objective.
`BG-ALIGNMENT-THM-057`	That zero-objective known inverse is a global minimizer.
`BG-ALIGNMENT-THM-058`	Under pointwise positive definiteness, the known inverse equals the unique polar-minimizer field.

A.3 Fixed and pointwise gauges under finite Fourier analysis

Proposition 6 (Finite Fourier equivariance and its boundary)**.** *On an exact finite abelian momentum group, conjugation by one fixed unitary commutes with Fourier coefficient extraction, full reconstruction, exact support truncation, truncated reconstruction, and the full-minus-truncated residual. A constant pointwise field reduces to this case. By contrast, if a varying pointwise unitary changes the matrix field at one momentum, at least one Fourier coefficient changes, even though weighted pointwise Frobenius power remains invariant.*

Proof sketch. Fixed conjugation is a complex-linear map, so it passes through finite sums and through the coefficient mask. The residual statement follows by subtraction. Exact finite Fourier inversion makes equality of all coefficients equivalent to equality of fields, which proves the varying-gauge contraposition. Frobenius invariance is pointwise unitary invariance summed with the caller's weights. The corresponding identifiers are `BG-FOURIER-THM-043`, `BG-FOURIER-THM-044`, `BG-FOURIER-THM-045`, `BG-FOURIER-THM-046`, `BG-FOURIER-THM-047`, `BG-FOURIER-THM-052`, `BG-FOURIER-THM-053`, and `BG-FOURIER-THM-054`. □

Table 3: Finite Fourier identities and truncation bounds.

Formal identifier	Mechanized result and boundary
`BG-FOURIER-THM-043`	Fixed-unitary conjugation commutes with coefficient extraction.

Formal identifier	Mechanized result and boundary
BG-FOURIER-THM-044	Fixed-unitary conjugation commutes with full finite reconstruction.
BG-FOURIER-THM-045	Fixed-unitary conjugation commutes with an exact coefficient mask.
BG-FOURIER-THM-046	The same holds for support-truncated reconstruction.
BG-FOURIER-THM-047	The same holds for the full-minus-truncated residual.
BG-FOURIER-THM-052	A constant pointwise unitary field is the fixed-unitary case.
BG-FOURIER-THM-053	A pointwise change of the field forces a change in at least one finite Fourier coefficient.
BG-FOURIER-THM-054	Arbitrary pointwise unitary conjugation preserves weighted pointwise Frobenius power, not coefficient support.
BG-FOURIER-THM-027	Matrix Frobenius power satisfies finite Parseval with the declared normalization.
BG-FOURIER-THM-028	Full-minus-truncated matrix Frobenius power equals group cardinality times omitted coefficient power.
BG-FOURIER-THM-068	The Hilbert-valued deliberate-truncation power has the same omitted-tail identity.
BG-FOURIER-THM-069	Specializes that identity to entrywise matrix Frobenius power.
BG-FOURIER-THM-070	Hilbert-valued truncation RMS equals the square root of omitted coefficient power on a nonempty group.
BG-FOURIER-THM-071	Its squared RMS equals omitted coefficient power.
BG-FOURIER-THM-072	Matrix Frobenius truncation RMS equals the square root of omitted matrix coefficient power.
BG-FOURIER-THM-073	Its squared RMS equals omitted matrix coefficient power.
BG-FOURIER-THM-074	Each pointwise truncation norm is bounded by the omitted coefficient one-norm tail.
BG-FOURIER-THM-075	The finite maximum norm obeys the same one-norm-tail bound.

The Parseval and tail statements follow from finite-character orthogonality; the maximum bound is the triangle inequality with unit-modulus character phases. They are exact for the declared finite transform. They do not identify a coefficient index with a physical continuum range without a separate mesh and representation binding.

A.4 Residual-channel composition

Formal uses four fields—reference, interpolated, converted, and truncated—and three component residuals. Its exact identity and norm consequences are:

Formal identifier	Mechanized result
BG-HILBERT-THM-022	Total residual equals interpolation plus conversion plus deliberate-truncation residual fields.
BG-HILBERT-THM-023	The pointwise total norm is at most the sum of the three component norms.
BG-HILBERT-THM-024	The finite maximum total norm is at most the sum of the three component maxima.
BG-HILBERT-THM-025	The finite RMS total norm is at most the sum of the three component RMS values.

Proof sketch. The field identity telescopes. The pointwise estimate is the triangle inequality; the maximum and RMS estimates are the corresponding norms on finite function spaces. For the

three-field ledger in Equation (30), set Formal's converted field equal to its interpolated field, so the conversion channel is zero. Formal's reference-minus-approximation orientation is the negative of this report's approximation-minus-reference orientation; the exact field identity changes sign while all stated norm bounds are unchanged. □

A.5 Mechanization status and exclusions

Item	Status at the pinned Formal revision	Consequence for this report
Exact finite theorems above	Lean-accepted; manifest status *proved; no admissions*.	Supports only the stated conditional algebraic implications.
Runtime SVD and polar selection	**Excluded**: no floating-point SVD, rank cutoff, null-space completion, or optimizer branch is authenticated.	Producer integrity and rank gates remain numerical obligations.
Mesh authentication	**Excluded**: no runtime coordinate array, mesh identity, weight, or source digest is bound.	The construction and shifted meshes remain report and runtime data.
Continuum convergence	**Excluded**: no aliasing estimate, mesh-limit theorem, continuum Fourier-decay theorem, or physical-range identification is proved.	Refinement evidence cannot be promoted by the finite identities alone.
Runtime certificates	**Excluded**: no request/result wire, parser, retained-byte binding, or checked certificate exists; the checked certificate count is zero.	This report and its cache contain no Formal certificate.
Scientific acceptance	**Excluded**: kernel checking does not establish physical adequacy, numerical validation, novelty, publication acceptance, or release status.	Scientific review and any publication decision remain separate human actions.

B Frozen numerical inventory

Quantity	Frozen value
Retained ranks	$3, 4, 8, 16$
Circuit depths	$1, 2, 4, 8$
Ambient dimension	$2r$
Construction meshes	$513, 1025, 2049$ points
Global-unitary selection mesh	2049 points only
Shifted evaluation mesh	8191 points
Gauge alignments	identity; global-unitary; pointwise-unitary; known-inverse-oracle
Supports	$S = 0, 1, 2, 4, 8, 16, 32, 64, 128$ with $\lvert R\rvert \le S$
Singular-value cutoff	$\max(10^{-12}, 10^{-10}\sigma_{\max})$; equality deficient
Principal phase branch	$(-\pi, \pi]$
Exact isolation certificate	$2.10E_{\mathrm{ref}}$
Mandatory final refinement	$1025 \to 2049$
Arithmetic	binary64

C Frozen tolerance crosswalk

The authoritative protocol prevails if this explanatory table ever differs from the selected schema.

Table 4: Frozen one-dimensional constant-projector numerical tolerances.

Diagnostic	Normalization	Tolerance
momentum and endpoint coordinates	radian	10^{-14}
frame orthonormality, gauge/alignment unitarity, projector idempotency	raw Frobenius	10^{-12}
Procrustes stationarity, objective identity, and negative polar part	dimensionless	10^{-12}
smooth/attacked projector equality	raw Frobenius	10^{-12}
direct ordered-spectrum equivalence	E_{ref}	10^{-11}
pointwise/oracle frame recovery	$\sqrt{r}$	10^{-11}
pointwise/oracle represented-operator recovery	$\sqrt{r}E_{\mathrm{ref}}$	10^{-11}
endpoint sewing and primitive closure	normalized Frobenius	10^{-11}
maximum closure-phase magnitude	radian	10^{-10}
Fourier reconstruction and Hermitian pairing	$\sqrt{r}E_{\mathrm{ref}}$	2×10^{-11}
relative Parseval defect	dimensionless	2×10^{-11}
error decomposition and support monotonicity	dimensionless	2×10^{-11}
maximum perturbation-bound violation	E_{ref}	10^{-11}
final interpolant operator error	$\sqrt{r}E_{\mathrm{ref}}$	10^{-9}
final interpolant ordered-spectrum maximum error	E_{ref}	10^{-10}
retained-block refinement, $\lvert R\rvert \leq 128$	$\sqrt{r}E_{\mathrm{ref}}$ per block	10^{-10}

D Result and failure semantics

Disposition	**Meaning**	**Downstream consequence**
`qualified`	Required scientific and integrity gates pass.	Declared downstream values are present.
`aggregate-rank-deficient`	Global covariance fails the strict rank gate.	Only global-unitary downstream values are absent.
`pointwise-rank-deficient`	At least one pointwise covariance fails the strict gate.	The complete constant-projector case is invalid.
`alignment-numerically-unresolved`	Documented backend operation fails or becomes nonfinite.	No selected factor or locality values are fabricated.
`alignment-integrity-failed`	Returned factor fails unitarity, stationarity, objective, or PSD checks.	Complete case is invalid.
`oracle-integrity-failed`	Analytic inverse fails construction or recovery checks.	Complete case is invalid.

E Prospective retained-results presentation

A reviewed result presentation should preserve:

1. one qualification table for all 16 cases and four gauge alignments;
2. global and pointwise rank-margin summaries;

3. common transport and endpoint-closure diagnostics;
4. normalized block-norm decay for every declared support;
5. separate interpolation, truncation, and total operator and spectrum errors;
6. the complete $1025 \to 2049$ scalar-refinement ledger;
7. rank/depth comparisons with the frozen resolution rule;
8. the result-manifest SHA-256; and
9. the single `calculation_infrastructure_sha256` provenance identity.

No table should replace inapplicable or unresolved entries with zero. No plot should select a preferred support unless a separately reviewed decision rule is added before execution.

References

[1] Nicola Marzari and David Vanderbilt. Maximally localized generalized Wannier functions for composite energy bands. *Physical Review B*, 56(20):12847–12865, 1997.

[2] Ivo Souza, Nicola Marzari, and David Vanderbilt. Maximally localized wannier functions for entangled energy bands. *Physical Review B*, 65:035109, 2001.

[3] Nicola Marzari, Arash A. Mostofi, Jonathan R. Yates, Ivo Souza, and David Vanderbilt. Maximally localized wannier functions: Theory and applications. *Reviews of Modern Physics*, 84:1419–1475, 2012.

[4] Jonathan R. Yates, Xinjie Wang, David Vanderbilt, and Ivo Souza. Spectral and Fermi surface properties from Wannier interpolation. *Physical Review B*, 75(19):195121, 2007.

[5] Peter H. Schönemann. A generalized solution of the orthogonal Procrustes problem. *Psychometrika*, 31(1):1–10, 1966.

[6] Nicholas J. Higham. Computing the polar decomposition—with applications. *SIAM Journal on Scientific and Statistical Computing*, 7(4):1160–1174, 1986.

[7] Gene H. Golub and Charles F. Van Loan. *Matrix Computations*. Johns Hopkins University Press, Baltimore, 4 edition, 2013.

[8] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, Cambridge, 2 edition, 2013.