The Physician and Sportsmedicine

ISSN: (Print) (Online) Journal homepage: https://www.tandfonline.com/loi/ipsm20

The Impact of Non-Operative Hip and Core Injuries on National Football League Athlete Performance

Joseph S. Tramer, Toufic R Jildeh, Joshua P Castle, Patrick Buckley, Caden Nowak & Kelechi R. Okoroha

To cite this article: Joseph S. Tramer, Toufic R Jildeh, Joshua P Castle, Patrick Buckley, Caden Nowak & Kelechi R. Okoroha (2021): The Impact of Non-Operative Hip and Core Injuries on National Football League Athlete Performance, The Physician and Sportsmedicine, DOI: 10.1080/00913847.2021.1976603

To link to this article: https://doi.org/10.1080/00913847.2021.1976603

Accepted author version posted online: 06 Sep 2021.

Submit your article to this journal

View related articles

View Crossmark data

Full Terms & Conditions of access and use can be found at
https://www.tandfonline.com/action/journalInformation?journalCode=ipsm20

Check for updates

Publisher: Taylor & Francis & Informa UK Limited, trading as Taylor & Francis Group

Journal: *The Physician and Sportsmedicine*

DOI: 10.1080/00913847.2021.1976603

The Effect of Hip and Core Injuries on National Football League Athlete Performance

Joseph S. Tramer MD[a]*, [1]Toufic R Jildeh MD[a], Joshua P Castle MD[a], Patrick Buckley BS[a], Caden Nowak BS[b], Kelechi R. Okoroha MD[c]

[a]Henry Ford Hospital, Department of Orthopaedic Surgery, 2799 W. Grand Blvd, Detroit, MI 48202

[b]Wayne State University School of Medicine, 540 E Canfield St, Detroit, MI 48201

[c] Mayo Clinic Sports Medicine Center, 600 Hennepin Ave, Suite 310, Minneapolis, MN 55403

joe.tramer@gmail.com; touficjildeh@gmail.com; jcastle1@hfhs.org; pbuckle3@hfhs.org; caden.nowak@wayne.edu; krokoroha@gmail.com

***Corresponding Author:**

author to whom all correspondence should be addressed

Department of Orthopaedic Surgery

2799 W. Grand Blvd, Detroit, MI 48202

joe.tramer@gmail.com

P: (216) 536-2904 F: 313-916-0475

ACCEPTED MANUSCRIPT

Institutional Review Board: This project did not require review by the institutional review board.

ACCEPTED MANUSCRIPT

The Impact of Non-Operative Hip and Core Injuries on National Football League Athlete Performance

ACCEPTED MANUSCRIPT

Abstract

Objectives: Hip and core injuries are common in National Football League (NFL) athletes; however, the impact following injury remains unclear. The goal of this manuscript is to determine the impact of nonoperative hip and core injuries on return to play and performance.

Methods: NFL athletes who sustained a hip or core injury treated non-operatively between 2010-2016 were identified. Offensive and defensive power ratings were calculated for each player's injury season and 2 seasons before and after to assess longitudinal impact. A matched control group without an identified hip and/or core injury were assembled for comparison.

Results: A total of 41 offensive and 71 defensive players with nonoperative hip or core injury were analyzed. All athletes returned to play; offensive and defensive players missed 4.0 ± 5.2 and 3.1 ± 2.6 games after injury, respectively. Offensive players played fewer cumulative career games returning from core injury versus hip (23.5 ± 20.6 vs 41.0 ± 26.4). Defensive players played fewer games (58.1 ± 41.1 versus 37.4 ± 27.1, $p<0.05$) with lower defensive power rating (133.9 ± 128.5 versus 219.8 ± 212.2, $p<0.05$) cumulatively after hip or core injury. Additionally, two years following injury, defensive players played fewer games compared to controls (9.5 ± 7.0 versus 10.9 ± 6.8, $p<0.05$). Following hip injury specifically, NFL defenders played fewer games (39.8 ± 27.9 vs 61.9 ± 38.8; $p<0.05$) and had a lower defensive power rating (145.9 ± 131.7 vs 239.0 ± 205.9; $p<0.05$) compared to before injury.

Conclusion: Overall, NFL players return to play following nonoperative hip and core injuries. Defensive players played in fewer games following hip or core injury compared to controls; offensive players were unaffected. Hip injuries have a greater impact on performance compared to core injuries in defensive athletes; offensive players played fewer games upon return from core injury.

ACCEPTED MANUSCRIPT

Key Terms: Muscle injuries, football, groin pain, hip/pelvis/thigh, core

ACCEPTED MANUSCRIPT

Introduction

Injuries in the National Football League (NFL) have been shown to have a significant financial burden to both players and organizations.[1, 2] A systematic review of athletes entering the NFL combine with a musculoskeletal injury that required orthopedic intervention are uniformly at risk for decreased performance and fewer games, the effects of which resonate throughout an NFL player's career.[3, 4] Although knee and shoulder injury account for the highest incidence of time loss injuries among NFL players, they are at significant risk for injuries to other anatomic regions, particularly the hip and core.[5, 6, 7, 8]

Hip and core injuries in the NFL account for approximately 10% of NFL injuries; the majority are classified as muscle strains. [9, 10] In fact, hip flexor strains occurred 268 times over a 10 year period in the NFL, and were the most common hip injury. [10] Football players are particularly vulnerable to core injuries such as rib fractures or contusions, back stiffness as well as abdominal/oblique strains, and athletic pubalgia due to the physical nature of the game. [11, 12] Prior studies have examined the effect of surgical intervention on groin injuries, showing that operative intervention for athletica pubalgia led to significantly shorter career length and games per season compared to matched controls,[13] however the impact of nonoperative hip and core injuries on NFL player performance has not been elucidated.

The purpose of our investigation was to determine the impact of nonoperative hip and core injuries on the performance in NFL athletes. We sought to determine if these injuries had a significant impact on return to play rates, as well as offensive and defensive player productivity. It was hypothesized that both offensive and defensive NFL players who sustained a nonoperative hip or core injury would return to play but play fewer games and experience worse performance following their return as compared to healthy controls matched for age, position, and NFL experience.

ACCEPTED MANUSCRIPT

Methods

Data were collected retrospectively for this study; internal review board exemption was granted since all material is freely accessible. All NFL players who sustained a hip or core injury between 2010-2016 seasons were collected from publicly available sources such as injury reports, player profiles, team pages and game summaries using methods previously utilized in other studies.[13, 14, 15, 16] Indexed hip injuries included adductor strain, hip flexor strain, labral tears, hip inflammation or impingement while core injuries included oblique, abdominal or intercostal strains, as well as back spasms or stiffness. Injuries were cross referenced with public websites including official team pages, official press releases, and personal player websites, using methodology that has been utilized by previous studies using publicly available databases.[13, 14, 15, 16] The NFL mandates that teams quickly and accurately disseminate injury information to the league office, their opponents, local and national media, and the team's broadcast partners. Return-to-play (RTP) data was further confirmed by identifying gaps in reported statistics. Fractures, disc herniations, and any player who was reported to have undergone surgery for their injury were not included in the investigation.

Demographic data surrounding the injury were recorded such as BMI, age at injury, season in NFL that injury occurred and position. The total number of games played before and after the reported injury date were also recorded. Performance data was collected for each player including total games played, total yards (passing, rushing or receiving) and touchdowns for offensive players and total tackles, sacks and interceptions for defensive players. Offensive and defensive power ratings were also calculated from these variables, which have both been previously described and validated outcomes measures of a football player's overall contribution to the team's offense or defense.[14, 17, 18] They are calculated with the following formulas:

ACCEPTED MANUSCRIPT

Offensive power rating (OPR) = [(total yards/10) + (touchdowns/6)]

Defensive power rating (DPR) = [total tackles + (total sacks x 2) + (total interceptions x 2)]

Overall career data as well as data in the two years prior to and following the date of injury were collected in order to better assess longitudinal changes in performance. Power ratings are intended to calculate productivity over a season, not individual games; therefore, the individual statistics were also evaluated on a per game basis to account for games missed. A control group was assembled by 2:1 matching injured athletes with healthy rostered NFL players that did not sustain a documented hip and/or groin injury by age, position, performance data, and NFL experience within three seasons in accordance to previously established methodology. [13, 14, 15] Control players were chosen based on an index date which was matched with the study player's date of injury. This control group was established to compare performance before and after hip/core injury. Additionally, if players did not participate in the season prior to the index injury season they were excluded. Furthermore, athletes injured at the end of the season who were unable to return-to-play during the index season were excluded from the study. When calculating time missed from injury, bye weeks were not a factor, only the number of games missed. Power rating has been previously demonstrated in other investigations to be a valid and effective method for quantifying overall player performance;[14] however, since several studies demonstrated offensive power ratings are not an effective performance metric for offensive linemen (OL) or kickers (K), only offensive skill players (wide receivers [WR], running backs [RB], tight ends [TE], and quarterbacks [QB]) were included for offensive power ratings.[17, 18]

Statistical Analysis

All continuous data are reported as mean ± standard deviation, while categorical data are reported as counts and column percentages. Comparisons between injured and control players

ACCEPTED MANUSCRIPT

are performed using independent 2-sample t-tests for continuous variables and using chi-square tests for categorical variables. Comparisons between pre- and post- data are performed using paired t-tests if the variable is normally distributed, and Wilcoxon signed rank tests if the variable is non-normally distributed. Statistical significance is set at $p<0.05$. All analyses are performed using SAS 9.4 (SAS Institute Inc., Cary, NC, USA).

Results

A total of 112 (41 offensive and 71 defensive) NFL players were identified as having sustained a hip or core injury between 2010 and 2016 that also matched our inclusion criteria. The offensive group included 15 wide receivers, 10 offensive linemen, 6 tight ends, 5 running backs, 4 quarterbacks and 1 kicker. Defensive players included 33 defensive backs, 19 linebackers, 10 defensive ends, and 9 defensive tackles. A 2:1 matched control group consisting of 224 players was compared to the study group and demonstrated no statistical differences in age, years of NFL experience, position, or BMI (Table 1). Pre-injury performance did not significantly differ between the injured group and healthy controls. No players sustained injuries requiring retirement.

	Offensive Injury (N=41)	Offensive Control (N=82)	P-Value	Defensive Injury (N=71)	Defensive Control (N=142)	P- Value
Mean Age, y	25.7 ± 3.1	25.9 ± 2.9	0.71	26.8 ± 2.8	26.6 ± 2.8	0.65
BMI (kg/m^2)	30.6 ± 4.6	30.8 ± 4.7	0.79	30.0 ± 3.6	30.6 ± 4.4	0.33
Mean NFL experience (seasons)	6.9 ± 3.0	7.4 ± 3.2	0.41	7.3 ± 2.9	7.5 ± 2.3	0.65

Table 1: Offensive and Defensive Injury and Control group characteristics

ACCEPTED MANUSCRIPT

All players sustaining a core or hip injury were able to return to play following their injuries. Eighty-three percent of offensive players and 86% of defenders were able to return to play in the same season as the injury, the remainder of the players were able to return in the following season. Injuries occurred mostly in wide receivers (37% of offensive injuries) and defensive backs (46% of defensive injuries) and there was a total of 81 hip injuries and 31 core injuries overall. Eleven (9.8%) of all injured players had documented re-injury during the study period. Offensive players missed 4.0 ± 5.2 games when suffering a hip or core injury; defensive players missed 3.1 ± 2.6 games. Players with hip injuries missed of 2.9 ± 2.3 games; athletes with core injuries missed 4.8 ± 5.8 games.

Regarding performance, offensive skilled players' experienced a decrease in career power ratings following a hip or core injury compared to pre-injury (341.8 ± 550.6 vs. 205.7 ± 306.9, P=.03) (Table 2) while control players demonstrated no difference (350 ± 708.9 vs 335 ± 680.5; p=0.38). There were no significant differences in performance between the injured offensive skill group and healthy matched controls in the first two years after injury or cumulative post-injury performance. (Table 3)

	Before Injury	After Injury	Difference	P-value
Games Played	49.3 ± 40.1	35.0 ± 25.7	-14.3	.06
Yards Per Game	37.6 ± 59.3	35.0 ± 62.2	-2.6	.17
Touchdowns Per Game	0.27 ± 0.44	0.26 ± 0.49	-0.01	.30
Power Rating[α]	341.8 ± 550.6	205.7 ± 306.9	-136.2	**.03***

Table 2: Cumulative career performance of offensive players before and after hip or core injury (N=41)
[α] Power Rating only includes offensive skill players (WR, RB, TE, QB; N=30) [K & OL excluded]
* Denotes statistically significant difference (P<.05)

ACCEPTED MANUSCRIPT

	Two years following injury			Cumulative post-injury performance		
	Injured	Controls	P-Value	Injured	Controls	P- value
Games Played	17.8 ± 10.6	21.4 ± 12.2	.11	35.0 ± 25.7	44.5 ± 30.3	.09
Yards per game	41.8 ± 66.9	47.7 ± 78.3	.36	35.0 ± 62.2	40.6 ± 70.8	.33
Touchdowns per game	0.32 ± 0.54	0.35 ± 0.64	.43	0.26 ± 0.49	0.29 ± 0.50	.37
Power Rating[α]	106.7 ± 178.5	143.5 ± 233.3	.21	205.7 ± 306.9	335.8 ± 680.5	.11

Table 3: Performance of offensive players who sustained a hip or core injury (N=41) and healthy matched controls (N=82) in both the first two years and cumulative career statistics following hip or core injury.
[α] Power Rating only includes offensive skill players (WR, RB, TE, QB; N=30 injured; N=60 controls) [K & OL excluded]

Defensive players experienced a significant decrease in games played (58.1 ± 41.1 vs 37.4 ± 27.1, p<0.01) and power rating (219.8 ± 212.2 vs. 133.9 ± 128.8; p<0.01) compared to preinjury (Table 4). When comparing to healthy controls, defensive players participated in significantly fewer games at 2 years post injury (9.5 ± 7.0 versus 10.9 ± 6.8, p<0.05) while no significant difference was found for cumulative career games played (37.5 ± 26.9 vs. 41.4 ± 25.5, P=0.16). Power ratings did not significantly differ between injured and control defensive players at 2 years or for cumulative performance following injury (Table 5).

	Before Injury	After Injury	Difference	P-value
Games Played	58.1 ± 41.1	37.4 ± 27.1	-20.7	**<.01***
Tackles per game	3.1 ± 1.8	2.9 ± 1.6	-0.2	.13
Sacks per game	0.10 ± 0.15	0.09 ± 0.18	-0.01	.40
Interceptions per game	0.05 ± 0.07	0.05 ± 0.07	0.0	.26

ACCEPTED MANUSCRIPT

Power Rating	219.8 ± 212.2	133.9 ± 128.5	-85.9	**<.01***

Table 4: Cumulative career performance of defensive players before and after hip or core injury (N=71)
* Denotes statistically significant difference (P<.05)

	Two years following injury			Cumulative post-injury performance		
	Injured	Controls	P-Value	Injured	Controls	P- value
Games Played	9.5 ± 7.0	10.9 ± 6.8	**.04***	37.5 ± 26.9	41.4 ± 25.5	.16
Tackles per game	3.4 ± 2.0	3.2 ± 2.1	.29	2.9 ± 1.6	3.0 ± 1.7	.29
Sacks per game	0.12 ± 0.24	0.12 ± 0.20	.48	0.09 ± 0.18	0.10 ± 0.16	.37
Interceptions per game	0.05 ± 0.09	0.04 ± 0.08	.20	0.05 ± 0.07	0.05 ± 0.07	.29
Power Rating	68.3 ± 63.9	81.9 ± 62.5	.07	141.1 ± 108.9	148.4 ± 130.5	.22

Table 5: Performance of defensive players who sustained a hip or core injury (N=71) and healthy matched controls (N=142) in both the first two years and cumulative career statistics following hip or core injury.
* Denotes statistically significant difference (P<.05)

Games played and performance analysis was further stratified by either hip or core injury. Following hip injury, defensive athletes played in fewer games (39.8 ± 27.9 vs 61.9 ± 38.8; P<0.05) and had a lower defensive rating (145.9 ± 131.7 vs 239.0 ± 205.9; P<0.05) compared to their career before injury. Conversely, offensive skilled players experienced a significant decrease in power rating compared to their career before injury (395.4 ± 567.4 vs 190.1 ± 327.6; P<0.05). When compared to controls, both offensive skilled players and defensive players with

ACCEPTED MANUSCRIPT

either hip or core injuries did not experience a significant decrease in cumulative career power ratings following injury.

Discussion

The present investigation demonstrated a 100% return-to-play rate after hip or core injury for NFL players. Defensive players with a either a hip or core injury demonstrated significantly fewer games played and both offensive and defensive players experienced decreased power ratings post-injury compared to before injury. Compared to healthy controls, defensive athletes played in significantly fewer games in the two years following injury but returned to near baseline at long term career follow up. Defensive players who sustained hip injuries alone had significantly decreased games played and power ratings in their post-injury careers compared to prior. Offensive players who sustained a core injury had significantly fewer games played in their cumulative career following injury compared to healthy controls.

Hip and groin injuries occur frequently in football players and are a leading cause of injury and lost playing time during both practice and games. [19] In a descriptive epidemiologic study of 23,806 NFL injuries from 1997 to 2006, Feeley et al. found that injuries around the hip comprised 3.1% of total injuries, with muscle strains being the most common. Wide receivers (39.6% of offensive injuries) and defensive backs (47.1% of defensive injuries) were among the most affected by these injuries. [10] The present study found a nearly identical incidence of injury, with wide receivers accounting for 37% of offensive injuries and defensive backs 46% of defensive injuries, with all athletes able to RTP.

High rates of hip and groin injury have also been documented in National Collegiate Athletic Association (NCAA) football players. Kerbel et al reviewed the incidence of hip and groin injuries across multiple sports and found that football players sustained 62.7 hip or groin

ACCEPTED MANUSCRIPT

injuries per 100,000 athlete exposures, with the majority of reported injuries to either the adductor or hip flexor musculature.[20] Specifically investigating muscular strains about the hip, Eckard et al reviewed groin injuries in NCAA athletes and found that hip flexor and adductor muscles strains occurred at a rate of 1.93 and 1.43 per 10,000 athletic exposures in football players, respectively.[21] The present cohort similarly revealed groin or hip flexor strain/tear as among the most common injury type, comprising 56.1% of offensive and 67.6% of defensive injuries. Eckard et al also found 7.4% and 6.9% recurrence rate of hip flexor and adductor strains during the study period, compared to 9.8% of all athletes in our cohort with a documented re-injury.[21] Adductor strains, specifically, have the potential to alter the mechanics of the anterior musculature about the hip leading to increased risk of future rectus strains or athletic pubalgia.[22] Adductor injuries have been shown to cause significant increases in missed time from sport in other professional leagues. Long term performance and career longevity, however, were not altered. [23] These findings suggest that although muscular strains may appear benign, athletes with these injuries are at risk for future strains and time away from sport. Considering groin injuries cause significant time missed in the NFL and other professional sports, prevention is worth investigating. A study by Tyler et al revealed a dedicated physical therapy program targeted at adductor strengthening was beneficial in decreasing the incidence of groin strains in National Hockey League players; a similar training regimen may be beneficial in the NFL as well. [24]

Little is known about the effect of nonoperative core injuries such as back spasm, core stiffness and oblique strain on NFL players. A study utilizing a convenience sample of 267 NCAA football players with lumbar spine injuries found that 59.6% of athletes reporting muscular low back pain or contusion were able to return to sport in less than 24 hours and 72.6%

ACCEPTED MANUSCRIPT

of players who did not return in <24 hours returned within a week of injury.[25] Rebolledo et al examined 214 NFL athletes at the combine and found that 50% had a history of abdominal core or lower extremity muscular injury which was also associated with lower levels of Vitamin D.[9] The effect of back and core injury has also been examined in other professional sports, as Camp et al found that core and back injury accounted for 12% of injuries resulting in missed time in Major League Baseball (MLB).[26] Likewise, Marshall et al found that core injuries in MLB pitchers lead to 47.0 ± 5.6 days on the injured list with fewer strikeouts per nine innings in the year of injury.[16] In the our study, athletes sustaining a core injury missed an average of 4.8 ± 5.9 games following these injuries and all athletes were able to RTP without significant performance detriment. Additionally, offensive players who suffered a core injury played significantly fewer games throughout the remainder of their post-injury career compared to healthy controls, which could be partially attributed to similarities in athletic demand between a quarterback and MLB pitcher. While return to play rates are high and the current literature does not report significant changes in football performance, Domb et al reviewed disability claims in retired NFL athletes and found that 63 of 65 former players reported symptomatic lumbar spine strain. [27] It is important to note that, while core, and more specifically lumbar spine strains may not be associated with any surgical pathology, the repetitive stress can have implications on the future health of the athlete.

Specifically evaluating hip injuries in the NFL, Feeley et al noted that athletes lost an average of 12.3 days, with a median loss time of 4 days following a variety of hip injuries, such as muscle strains, contusions, and intra-articular pathology.[10] In the aforementioned study by Marshall et al, pitchers with hip injuries spent an average of 37.7 ± 8.1 days on the injured list and starting pitchers suffered from decreased fastball velocity during their initial season

ACCEPTED MANUSCRIPT

following injury.[16] Additionally, injuries around the hip joint have also been shown to lead to missed time in professional basketball, as Jackson et al reviewed 24 years of data and found that hip injury accounted for 14.6% of all athletic injuries and lead to an average of 6.3 ± 10.2 days missed. [28] In our investigation, defensive players were more affected by hip injury than core injuries, with individual decreases in DPR following a hip injury but not a core injury, as well as fewer games played in the two years following injury compared to healthy controls.

Although both offensive skilled players and defensive players saw a decrease in power ratings compared to their own previous performance, neither group experienced a significant decline in performance compared to the control group following a hip or core injury. In a 2017 review of player injuries at the NFL combine, Beaulieu-Jones et al. demonstrated that among NFL athletes entering the draft, defensive players with an injury history played fewer games in their first two years in the league compared to healthy controls (22.2 ± 7.9 vs 24.2 ± 7.6 games; p=0.001). Similarly, our study demonstrated that defensive players who suffered a hip or core injury played in fewer games the two years following compared to healthy defensive controls. Additionally, a systematic review by Wang et al looked at 32 studies pertaining to injury surveillance and monitoring at the NFL combine and found that overall, defensive linemen and defensive backs appeared to have their game participation most affected by an injury history compared to offensive athletes.[3] This is likely due to the inherently violent nature of defensive play as compared to offensive play, which predisposes defensive players to more energy transfer and higher impact, and thus higher injury rate. The defensive back position involves more efficient energy transfer in the kinetic chain during movements like backpedaling, quick reactionary changes in direction, and regular high-speed contact, which may help to explain the higher incidence of hip injuries among defensive players. However, it is important to note that in

ACCEPTED MANUSCRIPT

the present investigation performance decline following an injury was not significantly different for offensive skilled or defensive players when comparing to matched controls, thus while athletes may expect some decrease in individual performance, it will not statistically exceed other NFL players of similar age and position.

Our study is not without limitations. Player injury reports were collected from publicly available internet sources which may be subject to inaccuracies. However, the NFL mandates that teams provide accurate and prompt injury information to the league as well as the media. Similar to previous investigations, injuries reports were confirmed with additional publicly available injury reports. [13, 14, 15] With pressure to stay in the game and continue playing, these injuries may also be unreported, as players might not disclose injuries to the team, medical staff and media. Furthermore, while game statistics are an important indicator of player performance, there are many unmeasured contributions not captured, thus true "performance" of a player cannot be completely summarized by these statistics alone. Our use of player power rating attempts to mitigate this issue by generating a weighted value that emphasizes commonly agreed upon significant performance variables.[17] These power ratings, however, do not capture kicker or offensive linemen contributions, therefore these players were excluded from power rating analysis. In addition, hip and core injuries include a broad spectrum of pathologies, and public injury reports often do not indicate the specific diagnosis. We are unable to access clinical exam findings, subjective reports from the athlete, imaging results, and attempted conservative treatment modalities. Future studies should aim to identify specific features that favor successful nonoperative management of common hip and core injuries and correlate with performance.

Conclusion

ACCEPTED MANUSCRIPT

Overall, 100% of NFL players in our study returned to play following nonoperative hip and core injuries. Defensive players were more affected post-injury compared to offensive players. Hip injuries have a greater impact on performance compared to core injuries in defensive athletes; offensive athletes played fewer games upon return from core injury but were able to maintain their performance. At long term follow up, both offensive and defensive players were able to return to near baseline games played and power ratings.

ACCEPTED MANUSCRIPT

References

1. Kluczynski MA, Kelly WH, Lashomb WM, et al. A Systematic Review of the Orthopaedic Literature Involving National Football League Players. Orthop J Sports Med. 2019 Aug;7(8):2325967119864356. doi: 10.1177/2325967119864356. PubMed PMID: 31457068; PubMed Central PMCID: PMCPMC6702781.
2. Beaulieu-Jones BR, Rossy WH, Sanchez G, et al. Epidemiology of Injuries Identified at the NFL Scouting Combine and Their Impact on Performance in the National Football League: Evaluation of 2203 Athletes From 2009 to 2015. Orthop J Sports Med. 2017 Jul;5(7):2325967117708744. doi: 10.1177/2325967117708744. PubMed PMID: 28812033; PubMed Central PMCID: PMCPMC5529031.
3. Wang D, Weiss LJ, Abrams M, et al. Athletes With Musculoskeletal Injuries Identified at the NFL Scouting Combine and Prediction of Outcomes in the NFL: A Systematic Review. Orthop J Sports Med. 2018 Dec;6(12):2325967118813083. doi: 10.1177/2325967118813083. PubMed PMID: 30560142; PubMed Central PMCID: PMCPMC6293380.
4. Gibbs DB, Lynch TS, Gombera MM, et al. Preexisting Rotator Cuff Tears as a Predictor of Outcomes in National Football League Athletes. Sports Health. 2016 May/Jun;8(3):250-254. doi: 10.1177/1941738116636602. PubMed PMID: 26945020; PubMed Central PMCID: PMCPMC4981069.
5. Chambers CC, Lynch TS, Gibbs DB, et al. Superior Labrum Anterior-Posterior Tears in the National Football League. Am J Sports Med. 2017 Jan;45(1):167-172. doi: 10.1177/0363546516673350. PubMed PMID: 27793805.
6. Mannava S, Frangiamore SJ, Murphy CP, et al. Prevalence of Shoulder Labral Injury in Collegiate Football Players at the National Football League Scouting Combine. Orthop J

ACCEPTED MANUSCRIPT

Sports Med. 2018 Jul;6(7):2325967118783982. doi: 10.1177/2325967118783982. PubMed PMID: 30046633; PubMed Central PMCID: PMCPMC6055272.

7. Read CR, Aune KT, Cain EL, Jr., et al. Return to Play and Decreased Performance After Anterior Cruciate Ligament Reconstruction in National Football League Defensive Players. Am J Sports Med. 2017 Jul;45(8):1815-1821. doi: 10.1177/0363546517703361. PubMed PMID: 28467122.

8. Cinque ME, Hannon CP, Bohl DD, et al. Return to Sport and Performance After Anterior Cruciate Ligament Reconstruction in National Football League Linemen. Orthop J Sports Med. 2017 Jun;5(6):2325967117711681. doi: 10.1177/2325967117711681. PubMed PMID: 28680893; PubMed Central PMCID: PMCPMC5480637.

9. Rebolledo BJ, Bernard JA, Werner BC, et al. The Association of Vitamin D Status in Lower Extremity Muscle Strains and Core Muscle Injuries at the National Football League Combine. Arthroscopy. 2018 Apr;34(4):1280-1285. doi: 10.1016/j.arthro.2017.10.005. PubMed PMID: 29275983.

10. Feeley BT, Powell JW, Muller MS, et al. Hip injuries and labral tears in the national football league. Am J Sports Med. 2008 Nov;36(11):2187-95. doi: 10.1177/0363546508319898. PubMed PMID: 18641370.

11. McAdams TR, Deimel JF, Ferguson J, et al. Chondral Rib Fractures in Professional American Football: Two Cases and Current Practice Patterns Among NFL Team Physicians. Orthop J Sports Med. 2016 Feb;4(2):2325967115627623. doi: 10.1177/2325967115627623. PubMed PMID: 26925425; PubMed Central PMCID: PMCPMC4753355.

ACCEPTED MANUSCRIPT

12. Knapik DM, Gebhart JJ, Nho SJ, et al. Prevalence of Surgical Repair for Athletic Pubalgia and Impact on Performance in Football Athletes Participating in the National Football League Combine. Arthroscopy. 2017 May;33(5):1044-1049. doi: 10.1016/j.arthro.2017.01.024. PubMed PMID: 28284723.

13. Jack RA, Evans DC, Echo A, et al. Performance and Return to Sport After Sports Hernia Surgery in NFL Players. Orthop J Sports Med. 2017 Apr;5(4):2325967117699590. doi: 10.1177/2325967117699590. PubMed PMID: 28451612; PubMed Central PMCID: PMCPMC5400145.

14. Jildeh TR, Okoroha KR, Taylor KA, et al. Effect of Concussions on the Performance of Running Backs and Wide Receivers in the National Football League. Am J Sports Med. 2019 Aug 6:363546519864581. doi: 10.1177/0363546519864581. PubMed PMID: 31386551.

15. Okoroha KR, Kadri O, Keller RA, et al. Return to Play After Revision Anterior Cruciate Ligament Reconstruction in National Football League Players. Orthop J Sports Med. 2017 Apr;5(4):2325967117698788. doi: 10.1177/2325967117698788. PubMed PMID: 28451611; PubMed Central PMCID: PMCPMC5400221.

16. Marshall NE, Jildeh TR, Okoroha KR, et al. Implications of Core and Hip Injuries on Major League Baseball Pitchers on the Disabled List. Arthroscopy. 2018 Feb;34(2):473-478. doi: 10.1016/j.arthro.2017.08.298. PubMed PMID: 29225021.

17. McHale KJ, Rozell JC, Milby AH, et al. Outcomes of Lisfranc Injuries in the National Football League. Am J Sports Med. 2016 Jul;44(7):1810-7. doi: 10.1177/0363546516645082. PubMed PMID: 27166291.

ACCEPTED MANUSCRIPT

18. Nwachukwu BU, Bedi A, Premkumar A, et al. Characteristics and Outcomes of Arthroscopic Femoroacetabular Impingement Surgery in the National Football League. Am J Sports Med. 2018 Jan;46(1):144-148. doi: 10.1177/0363546517729163. PubMed PMID: 28949248.

19. Feeley BT, Kennelly S, Barnes RP, et al. Epidemiology of National Football League training camp injuries from 1998 to 2007. Am J Sports Med. 2008 Aug;36(8):1597-603. doi: 10.1177/0363546508316021. PubMed PMID: 18443276.

20. Kerbel YE, Smith CM, Prodromo JP, et al. Epidemiology of Hip and Groin Injuries in Collegiate Athletes in the United States. Orthop J Sports Med. 2018 May;6(5):2325967118771676. doi: 10.1177/2325967118771676. PubMed PMID: 29780846; PubMed Central PMCID: PMCPMC5952296.

21. Eckard TG, Padua DA, Dompier TP, et al. Epidemiology of Hip Flexor and Hip Adductor Strains in National Collegiate Athletic Association Athletes, 2009/2010-2014/2015. Am J Sports Med. 2017 Oct;45(12):2713-2722. doi: 10.1177/0363546517716179. PubMed PMID: 28745561.

22. Meyers WC, Foley DP, Garrett WE, et al. Management of severe lower abdominal or inguinal pain in high-performance athletes. PAIN (Performing Athletes with Abdominal or Inguinal Neuromuscular Pain Study Group). Am J Sports Med. 2000 Jan-Feb;28(1):2-8. doi: 10.1177/03635465000280011501. PubMed PMID: 10653536.

23. Patel BH, Okoroha KR, Jildeh TR, et al. Adductor injuries in the National Basketball Association: an analysis of return to play and player performance from 2010 to 2019. Phys Sportsmed. 2020 Mar 30:1-8. doi: 10.1080/00913847.2020.1746978. PubMed PMID: 32202444.

ACCEPTED MANUSCRIPT

24. Tyler TF, Nicholas SJ, Campbell RJ, et al. The effectiveness of a preseason exercise program to prevent adductor muscle strains in professional ice hockey players. Am J Sports Med. 2002 Sep-Oct;30(5):680-3. doi: 10.1177/03635465020300050801. PubMed PMID: 12239001.

25. Makovicka JL, Patel KA, Deckey DG, et al. Lower Back Injuries in National Collegiate Athletic Association Football Players: A 5-Season Epidemiological Study. Orthop J Sports Med. 2019 Jun;7(6):2325967119852625. doi: 10.1177/2325967119852625. PubMed PMID: 31245431; PubMed Central PMCID: PMCPMC6582304.

26. Camp CL, Conti MS, Sgroi T, et al. Epidemiology, Treatment, and Prevention of Lumbar Spine Injuries in Major League Baseball Players. Am J Orthop (Belle Mead NJ). 2016 Mar-Apr;45(3):137-43. PubMed PMID: 26991566.

27. Domb BG, Carter C, Finch NA, et al. Whole-Person Impairment in Younger Retired NFL Players: The Orthopaedic Toll of a Professional Football Career. Orthop J Sports Med. 2014 May;2(5):2325967114534824. doi: 10.1177/2325967114534824. PubMed PMID: 26535333; PubMed Central PMCID: PMCPMC4555538.

28. Jackson TJ, Starkey C, McElhiney D, et al. Epidemiology of Hip Injuries in the National Basketball Association: A 24-Year Overview. Orthop J Sports Med. 2013 Aug;1(3):2325967113499130. doi: 10.1177/2325967113499130. PubMed PMID: 26535239; PubMed Central PMCID: PMCPMC4555488.

ACCEPTED MANUSCRIPT

	Offensive Injury (N=41)	Offensive Control (N=82)	P-Value	Defensive Injury (N=71)	Defensive Control (N=142)	P- Value
Mean Age, y	25.7 ± 3.1	25.9 ± 2.9	0.71	26.8 ± 2.8	26.6 ± 2.8	0.65
BMI (kg/m^2)	30.6 ± 4.6	30.8 ± 4.7	0.79	30.0 ± 3.6	30.6 ± 4.4	0.33

Mean NFL experience (seasons)	6.9 ± 3.0	7.4 ± 3.2	0.41	7.3 ± 2.9	7.5 ± 2.3	0.65

Table 1: Offensive and Defensive Injury and Control group characteristics

	Before Injury	After Injury	Difference	P-value
Games Played	49.3 ± 40.1	35.0 ± 25.7	-14.3	.06
Yards Per Game	37.6 ± 59.3	35.0 ± 62.2	-2.6	.17
Touchdowns Per Game	0.27 ± 0.44	0.26 ± 0.49	-0.01	.30
Power Rating[α]	341.8 ± 550.6	205.7 ± 306.9	-136.2	**.03***

Table 2: Cumulative career performance of offensive players before and after hip or core injury (N=41)
[α] Power Rating only includes offensive skill players (WR, RB, TE, QB; N=30) [K & OL excluded]
* Denotes statistically significant difference (P<.05)

	Two years following injury			Cumulative post-injury performance		
	Injured	Controls	P-Value	Injured	Controls	P- value
Games Played	17.8 ± 10.6	21.4 ± 12.2	.11	35.0 ± 25.7	44.5 ± 30.3	.09
Yards per game	41.8 ± 66.9	47.7 ± 78.3	.36	35.0 ± 62.2	40.6 ± 70.8	.33
Touchdowns per game	0.32 ± 0.54	0.35 ± 0.64	.43	0.26 ± 0.49	0.29 ± 0.50	.37
Power Rating[α]	106.7 ± 178.5	143.5 ± 233.3	.21	205.7 ± 306.9	335.8 ± 680.5	.11

Table 3: Performance of offensive players who sustained a hip or core injury (N=41) and healthy matched controls (N=82) in both the first two years and cumulative career statistics following hip or core injury.
[α] Power Rating only includes offensive skill players (WR, RB, TE, QB; N=30 injured; N=60 controls) [K & OL excluded]

	Before Injury	After Injury	Difference	P-value
Games Played	58.1 ± 41.1	37.4 ± 27.1	-20.7	**<.01***
Tackles per game	3.1 ± 1.8	2.9 ± 1.6	-0.2	.13

ACCEPTED MANUSCRIPT

Sacks per game	0.10 ± 0.15	0.09 ± 0.18	-0.01	.40
Interceptions per game	0.05 ± 0.07	0.05 ± 0.07	0.0	.26
Power Rating	219.8 ± 212.2	133.9 ± 128.5	-85.9	**<.01***

Table 4: Cumulative career performance of defensive players before and after hip or core injury (N=71)
* Denotes statistically significant difference (P<.05)

	Two years following injury			Cumulative post-injury performance		
	Injured	Controls	P-Value	Injured	Controls	P- value
Games Played	9.5 ± 7.0	10.9 ± 6.8	**.04***	37.5 ± 26.9	41.4 ± 25.5	.16
Tackles per game	3.4 ± 2.0	3.2 ± 2.1	.29	2.9 ± 1.6	3.0 ± 1.7	.29
Sacks per game	0.12 ± 0.24	0.12 ± 0.20	.48	0.09 ± 0.18	0.10 ± 0.16	.37
Interceptions per game	0.05 ± 0.09	0.04 ± 0.08	.20	0.05 ± 0.07	0.05 ± 0.07	.29
Power Rating	68.3 ± 63.9	81.9 ± 62.5	.07	141.1 ± 108.9	148.4 ± 130.5	.22

Table 5: Performance of defensive players who sustained a hip or core injury (N=71) and healthy matched controls (N=142) in both the first two years and cumulative career statistics following hip or core injury.
* Denotes statistically significant difference (P<.05)

ACCEPTED MANUSCRIPT