

ASTA

CERTIFICATE OF TYPE TESTS

15. 1982-1983 1982-1983

Certificate No. 17248

SUPPLEMENT: 2000 AFAADP'S 1998 FIVE-YEAR PLAN, by the Southern Training System.
www.afaadp.com Southern system - 4 W and PE (missing)

THE UNIVERSITY OF TEXAS AT AUSTIN

[illegible]

TESTED BY: Institut „Protest für ein besseres Gesundheitswesen“ (IPGW)
Landsberger Allee 576A
12205 Berlin, Germany

BPA, Dresden GmbH
Gartenstraße 10, 10179 Berlin, Germany
Friedrichstraße 10
10117 Berlin, Germany

doi:10.1017/S0022292408004616 Published online by Cambridge University Press

The scenarios constructed in accordance with the description, strategy and propaganda instrument in the campaign has been subjected to the review of growing trend in accordance with:

ISSN 0264-3758/02 \$10.00+.00. © 2002 Blackwell Science Ltd, *Journal of Clinical Pharmacy and Therapeutics*, 27, 1-10

The results are shown in the record of Proving Tests and the enclosures attached hereto. The values obtained and the general performance are considered to comply with the above Standards and to fully represent required to the manufacturer as stated below.

Five ratings assigned by the manufacturer and proved by test are (Pages 1 and 2).

The amount of Money Back applied only to the approved items. The responsibility for conformity of any purchased items to the same dimensions with their listed costs with the Manufacturer.

The book contains: 10 pages, 1 diagram, 12 photographs, 11 photographs, 8 drawings and an index.

Only complete reproduction of this Certificate, or reproduction of this page accompanied by any payment or other fee stated on the original, shall constitute an authentic reproduction. For further information, contact the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400.



Abstract: *Chlamydia trachomatis* is the most common bacterial sexually transmitted infection (STI) in the United States. The purpose of this study was to determine the prevalence of *C. trachomatis* in a community-based sample of men and women in the United States. A cross-sectional study was conducted in 1998-1999 in a community-based sample of men and women in the United States. The prevalence of *C. trachomatis* was 1.1% (95% CI 0.4-2.8%) in men and 1.1% (95% CI 0.4-2.8%) in women. The prevalence of *C. trachomatis* was significantly higher in men than in women (1.1% vs. 0.4%, $P = 0.001$). The prevalence of *C. trachomatis* was significantly higher in men than in women (1.1% vs. 0.4%, $P = 0.001$). The prevalence of *C. trachomatis* was significantly higher in men than in women (1.1% vs. 0.4%, $P = 0.001$).

100