

## Twodimensional XY magnets with annealed nonmagnetic impurities

A. N. Berker and David R. Nelson

Citation: [Journal of Applied Physics](#) **50**, 1799 (1979); doi: 10.1063/1.327173

View online: <http://dx.doi.org/10.1063/1.327173>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jap/50/B3?ver=pdfcov>

Published by the [AIP Publishing](#)

---

### Articles you may be interested in

[Thermoelectric properties of two-dimensional topological insulators doped with nonmagnetic impurities](#)

J. Appl. Phys. **116**, 013706 (2014); 10.1063/1.4886176

[Spin waves in a two-dimensional non-ferromagnetic electron liquid with electron impurity states in a magnetic field](#)

Low Temp. Phys. **24**, 489 (1998); 10.1063/1.593629

[Critical phenomena in the twodimensional XY magnet Fe\(100\) on W\(100\)](#)

J. Appl. Phys. **79**, 4984 (1996); 10.1063/1.361923

[Twodimensional solitons in the classical Heisenberg antiferromagnet with nonmagnetic impurities](#)

J. Appl. Phys. **79**, 5368 (1996); 10.1063/1.361329

[Magnetic resonance of twodimensional Heisenberg antiferromagnets with nonmagnetic impurities](#)

J. Appl. Phys. **73**, 7010 (1993); 10.1063/1.352413

---

The advertisement features a blue background with a film strip graphic on the left. The text is in white and orange. The main headline reads 'Not all AFMs are created equal' in orange, followed by 'Asylum Research Cypher™ AFMs' in white, and 'There's no other AFM like Cypher' in orange. Below this is the website 'www.AsylumResearch.com/NoOtherAFMLikeIt' in white. In the bottom right corner is the Oxford Instruments logo, which consists of the word 'OXFORD' above 'INSTRUMENTS' inside a square frame, with the tagline 'The Business of Science®' below it.

**Not all AFMs are created equal**

**Asylum Research Cypher™ AFMs**

**There's no other AFM like Cypher**

[www.AsylumResearch.com/NoOtherAFMLikeIt](http://www.AsylumResearch.com/NoOtherAFMLikeIt)

**OXFORD**  
INSTRUMENTS  
*The Business of Science®*

# Two-dimensional XY magnets with annealed non-magnetic impurities<sup>(a)</sup>

A. N. Berker<sup>(b)</sup> and David R. Nelson<sup>(c)</sup>

Department of Physics, Harvard University, Cambridge, Massachusetts 02138

The global phase diagram of a vectorial generalization of the Blume-Emery-Griffiths model is obtained using Migdal's approximate renormalization procedure. Classical two-component spins and non-magnetic impurities populate a triangular lattice, with nearest-neighbor interactions. The phase diagram in thermodynamic field space is divided into magnetic and impurity-rich phases by a first-order surface of discontinuous impurity concentrations, terminating in an Ising-type critical line. The magnetic region is further divided into a high-temperature paramagnetic phase and a low-temperature Kosterlitz-Thouless phase. The exponent  $\eta = 1/4$  of the pure system is preserved at the higher-order surface separating these two phases. This surface terminates in a line of critical end-points on the first-order surface, and, consequently, no tricritical point occurs for any values of the model parameters. However, the Ising critical line and the line of critical end-points approach each other in a certain limit, yielding an effective tricritical phase diagram. Within the Kosterlitz-Thouless phase, lines of constant  $\eta$  bunch together as the effective tricritical point is approached, in apparent agreement with tricritical scaling. This is also a model for superfluidity and phase separation in helium films.

PACS numbers: 75.10.Hk, 64.60.Kw, 05.70.Jk, 67.70.+n

The two-dimensional XY model of magnetism, consisting of two-component spins with short-ranged interactions, has attracted much interest. Conventional long-range order, namely non-zero magnetization, is rigorously ruled out [1]. Nevertheless, low-temperature spin-wave theory [2] yields algebraic, rather than exponential, decay of correlations, and high-temperature series expansions [3] suggest a phase transition at finite temperature. Experiments now seem to indicate that a detailed theory due to Kosterlitz and Thouless [4] is substantially correct, as explained below. At all temperatures below the transition, the magnetization is zero, but the susceptibility is infinite. At the transition, a universal jump discontinuity in the spin-wave stiffness (and, hence, in the spin-wave velocity) is predicted [5]. Although verification of this result in magnets is lacking, the analogous [5] universal jump in the superfluid density of helium films was recently reported [6]. In this paper, the above picture is extended by allowing for annealed, non-magnetic impurities. We find no XY tricritical point, in contrast to the three-dimensional situation.

We studied a vectorial generalization of the Blume-Emery-Griffiths (BEG) model [7], with Hamiltonian

$$-\frac{K}{k_B T} = J \sum_{\langle ij \rangle} \vec{s}_i \cdot \vec{s}_j + K \sum_{\langle ij \rangle} |\vec{s}_i|^2 |\vec{s}_j|^2 - \Delta \sum_i |\vec{s}_i|^2. \quad (1)$$

The spins  $\vec{s}_i$  are two-component classical vectors of length unity or zero, located at the sites  $i$  of a triangular lattice. The first two sums in (1) are over all nearest-neighbor pairs of sites. The states  $|\vec{s}_i| = 1$  represent the XY spins, interacting with the coupling constant  $k_B T J$ . The state  $\vec{s}_i = 0$  allows for the occupation of site  $i$  by a non-magnetic impurity. The coupling  $k_B T K$  measures the spin-impurity interaction, and  $k_B T \Delta$  is the impurity chemical potential. The above is also a model for superfluidity and phase separation in helium films [8].

The diluted XY model (1) was treated [8,9] using a renormalization scheme due to Migdal [10] and Kadanooff [11]. This approximate procedure yields good




Fig. 1. End-point phase diagram of the diluted XY model at  $K/J = 1$ . The full lines indicate higher-order transitions. The boundary of the coexistence regions is shown with the dashed line. The Kosterlitz-Thouless phase is labeled KT.

agreement [12] with the Kosterlitz-Thouless theory [4] of the pure system, obtained in the limit  $\Delta \rightarrow -\infty$  in (1).

Our results are exhibited by phase diagrams in the variables temperature ( $J^{-1}$ ) and impurity concentration

$$x = 1 - \langle |\vec{s}_i|^2 \rangle, \quad (2)$$

at given  $K/J$ . The phase diagram for  $K/J=1$  is shown in Fig. 1. The line of XY transitions drops to lower temperature with increasing impurity concentration, until it terminates on the phase-separation boundary, at a critical end-point  $E_s$  at  $x = 0.02$ . As  $K/J$  is increased, the phase-separation (Ising-type) critical point C moves to temperatures higher and higher than the XY transitions, the end-point  $E_s$  moves closer to  $x=0$ , and, thus, phase-separation dominates the phase diagram more and more. As  $K/J$  is decreased, the end-point slides up the phase-separation boundary, but never actually reaches the tip. Thus, there is never a tricritical point. This is in contrast to the re-normalization-group treatments [13] of the two-dimensional scalar BEG model [7], and, indeed, to one's expectations for the Hamiltonian (1) in three dimensions. (In the latter case, the tricritical phase diagram observed experimentally in bulk  $^3\text{He}$ - $^4\text{He}$  mixtures [14] is expected.) On the other hand, the end-point and the phase-separation critical point do come very close together at small  $K/J$ . For  $K/J = 0$ , shown in Fig. 2, they are respectively at  $J^{-1} = 0.6581$  and  $J^{-1} = 0.6615$ . Thus, an effective tricritical phase diagram is obtained. In all cases, the ordered (Kosterlitz-Thouless) phase accommodates a small (compared to similar situations [14] in three dimensions) amount of impurities before phase separation occurs. The upper-limit concentration is  $x(E_s) = 0.12$ , for  $K/J = 0$ .

Also of interest is the critical exponent  $\eta$ , defined by the algebraic decay of correlations in the Kosterlitz-Thouless phase,

$$\langle \vec{s}(\vec{r}) \cdot \vec{s}(\vec{0}) \rangle \sim 1/r^\eta. \quad (3)$$




Fig. 2. Effective tricritical phase diagram of the diluted XY model at  $K/J = 0$ . The full lines indicate higher-order transitions. The boundary of the coexistence region is shown with the dashed line.




Fig. 3. End-point phase diagram of the diluted XY model at  $K/J = 1$ . The dark full line and the line of open circles (ooo) respectively indicate higher-order and first-order transitions. The dark circle (●) marks the isolated critical point, and E labels the end-point. Within the Kosterlitz-Thouless (KT) phase, constant  $\eta$  curves are drawn with light full lines and labeled by  $32\eta$ . The chemical potential variable (see Ref. 8) is

$$D = \{\Delta + \ln \int_0^{2\pi} (d\phi/2\pi) \exp[4J(\cos\phi - 1)]\}/K.$$

At the transition temperature in the pure system, this exponent [4] equals  $\frac{1}{4}$ , a value intimately related to the size of the jump in the spin-wave stiffness [5]. Lines of constant  $\eta$  are shown in the temperature-chemical potential phase diagrams of Figs. 3 and 4. The pure system result  $\eta = \frac{1}{4}$  is preserved along the line of XY transitions until phase separation occurs. In the  $K/J = 0$  case of Fig. 4, several lines of constant  $\eta$  seem to converge onto the effective tricritical point  $T_{\text{eff}}$  formed by the closely situated end-point and critical point. The precise value of  $\eta$  at  $T_{\text{eff}}$  thus appears indeterminate, depending instead on the path of approach! This peculiar behavior is anticipated [8] from tricritical scaling theory [15] together with the invariance property of  $\eta$  under rescaling.

#### REFERENCES

- (a) Research supported in part by the National Science Foundation under Grant No. DMR77-10210.
- (b) Supported in part by an IBM Postdoctoral Fellowship.
- (c) Supported in part by a Junior Fellowship from the Harvard Society of Fellows.
1. N. D. Mermin and H. Wagner, Phys. Rev. Lett. **17**,




Fig. 4. Effective tricritical phase diagram of the diluted XY model at  $K/J = 0$ . The dark full line and the line of open circles (ooo) respectively indicate higher-order and first-order transitions. The actual end-point and critical point are both within the dark circle at the effective tricritical point  $T_{\text{eff}}$ . Within the Kosterlitz-Thouless (KT) phase, constant  $\eta$  curves are drawn with light full lines and labeled by  $32\eta$ . As predicted by tricritical scaling theory, the  $\eta=1/2$  to  $5/32$  curves apparently converge onto  $T_{\text{eff}}$ .

- 1133 (1966); P. C. Hohenberg, Phys. Rev. **158**, 383 (1967); N. D. Mermin, J. Math. Phys. **3**, 1061 (1967).
2. F. J. Wegner, Z. Phys. **206**, 465 (1967); V. L. Berezinskii, Zh. Eksp. Teor. Fiz. **59**, 907 (1970) [Sov. Phys. - JETP **32**, 493 (1971)].
3. H. E. Stanley, Phys. Rev. Lett. **20**, 150, 589 (1968); M. A. Moore, *ibid.* **23**, 861 (1969); W. J. Camp and J. P. Van Dyke, J. Phys. C **8**, 336 (1975).
4. J. M. Kosterlitz and D. J. Thouless, J. Phys. C **6**, 1181 (1973); J. M. Kosterlitz, J. Phys. C **7**, 1046 (1974).
5. D. R. Nelson and J. M. Kosterlitz, Phys. Rev. Lett. **39**, 1201 (1977).
6. I. Rudnick, Phys. Rev. Lett. **40**, 1454 (1978); D. J. Bishop and J. D. Reppy, Phys. Rev. Lett. **40**, 1727 (1978).
7. M. Blume, V. J. Emery, and R. B. Griffiths, Phys. Rev. A **4**, 1071 (1971).
8. A. N. Berker and D. R. Nelson, Harvard University preprint (1978). This paper contains a more complete account and the technical details of the calculations reported here.
9. Similar calculations were independently done, for a square instead of triangular lattice, by J. L. Cardy and D. J. Scalapino, University of California, Santa Barbara, preprint (1978).
10. A. A. Migdal, Zh. Eksp. Teor. Fiz. **69**, 1457 (1975) [Sov. Phys. - JETP **42**, 743 (1976)].
11. L. P. Kadanoff, Ann. Phys. (N.Y.) **100**, 359 (1976); Rev. Mod. Phys. **49**, 267 (1977).
12. J. V. José, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson, Phys. Rev. B **16**, 1217 (1977).
13. T. W. Burkhardt, Phys. Rev. B **14**, 1196 (1976); A. N. Berker and M. Wortis, *ibid.* **14**, 4946 (1976); T. W. Burkhardt, H. J. F. Knops, and M. den Nijs, J. Phys. A **9**, L179 (1976); J. Adler, A. Aharony, and J. Oitmaa, J. Phys. A **11**, 963 (1978).
14. G. Ahlers, in *The Physics of Liquid and Solid Helium, Part I*, edited by K. H. Benneman and J. B. Ketterson (Wiley, New York, 1976).
15. E. K. Riedel, Phys. Rev. Lett. **28**, 675 (1972).