

Frustration, Chaos, and Order in the $d=1$ Ising Spin Glass with Long-Range Power-Law Interactions

E. Can Artun^{1,2} and A. Nihat Berker^{2,3}

¹*TUBITAK Research Institute for Fundamental Sciences (TBAE), Gebze, Kocaeli 41470, Turkey*

²*Faculty of Engineering and Natural Sciences, Kadir Has University, Cibali, Istanbul 34083, Turkey*

³*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

The $d = 1$ Ising spin glass with long-range power-law interactions Jr^{-a} is studied for all a by a renormalization-group transformation that simultaneously projects local ferromagnetism and antiferromagnetism. In the ferromagnetic case, $J > 0$, a finite-temperature ferromagnetic phase occurs for $0.75 < a < 2$. The transition temperature monotonically decreases between these two limits. At $a = 2$, the phase-transition temperature discontinuously drops to zero and for $a > 2$ there is no ordered phase above zero temperature, as predicted by rigorous results. On approaching $a = 0.75$ from above, the phase-transition temperature diverges to infinity, meaning that, at all temperatures above zero, the system is ferromagnetically ordered. Thus, the equivalent-neighbor interactions regime is entered before ($a \neq 0$) the neighbors become equivalent, namely before the interactions become equal for all separations. The critical exponents are calculated, from a large recursion matrix, as a function of a . For the antiferromagnetic case, $J < 0$, all triplets of spins have competing interactions and this highly frustrated system does not have an ordered phase. In the spin-glass phase, where all couplings for all separations are randomly ferromagnetic or antiferromagnetic (with probability p), a finite-temperatures phase diagram is obtained with order in the presence of frustration. Transient chaos is found within the ferromagnetic phase.

I. ORDERING IN ONE-DIMENSION: LONG-RANGE INTERACTIONS

Whereas systems with finite-range interactions do not order in one dimension, certain systems with long-range interactions do order. The archetypical example are the Ising models with power-law interactions, Jr^{-a} , which show varigated ordering behavior for ferromagnetic interactions. For antiferromagnetic interactions, the system incorporates saturated frustration and spin-glass phenomena, in the absence of quenched randomness. All of these behaviors are seen in our renormalization-group study.

The model that we study is defined by the Hamiltonian

$$-\beta\mathcal{H} = \sum_{r_1 \neq r_2} J |r_1 - r_2|^{-a} s_{r_1} s_{r_2} \quad (1)$$

where $\beta = 1/k_B T$ is the inverse temperature, r_1 and r_2 designate the sites on the one-dimensional system, at each site there is an Ising spin $s_i = \pm 1$ and the sums are over all sites in the system. For ferromagnetic and antiferromagnetic systems, the two-spin interactions J are $J = |J| > 0$ and $J = -|J| > 0$, respectively. For the spin-glass system, for each two spins, their interaction is randomly ferromagnetic (with probability $1 - p$) or antiferromagnetic (with probability p).

II. METHOD: LONG-RANGE RENORMALIZATION GROUP

We solve this system with Niemeyer and van Leeuwen's two-cell cluster approximation. The renormalization-group transformation is constructed by first choosing




FIG. 1. Renormalization-group cells for $d = 1$. This cell structure projects both local ferromagnetism and antiferromagnetism, and therefore also spin-glass order.

cells on the $d = 1$, as shown in Fig. 1. Each of our cells has three spins. Secondly, for each cell, a cell-spin is defined as the sign of the sum of the three spins in the cell,

$$s'_{r'} = \text{signum}(s_{r-2} + s_{r-1} + s_r) \quad (2)$$

where the signum function returns the sign of its argument, primes denote the renormalized system, and $r' = r/b$, where $b = 3$ is the length-rescaling factor. The renormalized interactions are obtained from the conservation of the partition function Z ,

$$Z = \sum_{\{s\}} e^{-\beta\mathcal{H}(\{s\})} = \sum_{\{s'\}} \sum_{\{\sigma\}} e^{-\beta\mathcal{H}(\{s'\}, \{\sigma\})} = \sum_{\{s'\}} e^{-\beta\mathcal{H}'(\{s'\})} = Z', \quad (3)$$

where the summed variable σ represents, for each cell, the four states that give the same cell-spin value. Thus, the renormalized interactions are obtained from

$$e^{-\beta\mathcal{H}'(\{s'\})} = \sum_{\{\sigma\}} e^{-\beta\mathcal{H}(\{s'\}, \{\sigma\})}. \quad (4)$$

The two-cell cluster approximation of Niemeyer and van Leeuwen consists in carrying out this transformation

for two cells, including the 6 intracell interactions and the 9 intercell interactions. A recursion relation is obtained for each renormalized interaction,

$$J'_{r'} = (1/2) \ln[R(s'_0 = +1, s'_{r'} = +1)/R(s'_0 = +1, s'_{r'} = -1)], \quad (5)$$

where

$$R(s'_0, s'_{r'}) = \sum_{\sigma_0, \sigma_{r'}} e^{-\beta \mathcal{H}_{0r'}}, \quad (6)$$

where the unrenormalized two-cell Hamiltonian contains the 6 intracell interactions and the 9 intercell interactions between the 6 spins in cells 0 and r' .

III. RESULTS: FINITE-TEMPERATURE FERROMAGNETIC PHASE IN $d = 1$

Calculated phase diagram of the $d = 1$ long-range ferromagnetic Ising model with interactions Jr^{-a} , is shown in Fig. 2, in terms of temperature $1/J$ and interaction range a . A finite-temperature ferromagnetic phase occurs for $0.75 < a < 2$. The transition temperature monotonically decreases between these two limits. At $a = 2$, the phase-transition temperature discontinuously drops to zero and for $a > 2$ there is no ordered phase above zero temperature, as predicted by rigorous results. On approaching $a = 0.75$ from above, the phase-transition temperature diverges to infinity, meaning that, at all temperatures above zero, the system is ferromagnetically ordered. Thus, the equivalent-neighbor interactions regime is entered before ($a \neq 0$) the neighbors become equal for all separations.

Specific heat critical exponent α and correlation-length critical exponent ν , continuously varying as a function of interaction range a , for the finite-temperature ferromagnetic phase transition are also shown in Fig. 2. These critical exponents are calculated from the recursion relations $J'_1, \dots, J'_n = \text{funct}(J_1, \dots, J_n)$. Convergence is obtained by calculation up to $n = 3$. The largest eigenvalue, $\lambda = b^y$ of the derivative matrix of these recursion relations at the critical point gives the correlation-length critical exponent $\nu = 1/y$ and the specific heat critical exponent $\alpha = 2 - d/y = 2 - 1/y$.

The antiferromagnetic, overly frustrated without randomness, system does not have a finite-temperature phase transition, but the spin-glass system, where all couplings for all separations are randomly ferromagnetic or antiferromagnetic (with probability p), does have finite-temperature spin-glass phase transitions and chaos inside the spin-glass phase, as seen in Fig. 3.




FIG. 2. Calculated phase diagram of the $d = 1$ long-range ferromagnetic Ising model with interactions Jr^{-a} . Top panel: Phase diagram in temperature $1/J$ and interaction range a . A finite-temperature ferromagnetic phase occurs for $0.75 < a < 2$. The transition temperature monotonically decreases between these two limits. At $a = 2$, the phase-transition temperature discontinuously drops to zero and for $a > 2$ there is no ordered phase above zero temperature, as predicted by rigorous results. On approaching $a = 0.75$ from above, the phase-transition temperature diverges to infinity, meaning that, at all temperatures above zero, the system is ferromagnetically ordered. Thus, the equivalent-neighbor interactions regime is entered before ($a \neq 0$) the neighbors become equal, namely before the interactions become equal for all separations. Bottom panel: Specific heat critical exponent α and correlation-length critical exponent ν as a function of interaction range a for the finite-temperature ferromagnetic phase transition. The antiferromagnetic, overly frustrated without randomness, system does not have a finite-temperature phase transition, but the spin-glass system, where all couplings for all separations are randomly ferromagnetic or antiferromagnetic, does have finite-temperature spin-glass phase transitions and chaos inside the spin-glass phase, as seen in Fig. 3.

IV. RESULTS: FINITE-TEMPERATURE SPIN-GLASS PHASE IN $d = 1$

The spin-glass system, where all couplings for all separations are randomly ferromagnetic or antiferromagnetic (with probability p), does have finite-temperature spin-glass phase transitions and chaos inside the spin-glass phase, as seen in Fig. 3.




FIG. 3. Calculated finite temperature phase diagram of the $d = 1$ long-range Ising spin-glass system, where all couplings for all separations are randomly ferromagnetic or antiferromagnetic (with probability p). Bottom panel: Chaos inside the spin-glass phase in $d = 1$.

V. CONCLUSION

ACKNOWLEDGMENTS

Support by the Academy of Sciences of Turkey (TÜBA) is gratefully acknowledged.

-
- [1] D. J. Thouless, Long-Range Order in One-Dimensional Ising Systems, *Phys. Rev.* **187**, 732 (1969).
 - [2] D. Ruelle, *Statistical Mechanics Rigorous Results* (Benjamin, New York, 1969).
 - [3] R. B. Griffiths, Rigorous Results and Theorems, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and M. S. Green (Academic, New York, 1972), Vol. 1.
 - [4] M. Aizenman and C. Newman, Discontinuity of the Percolation Density in One-Dimensional $1/[X - Y]^2$ Percolation Models, *Comm. Math. Phys.* **107**, 611 (1986).
 - [5] M. Aizenman, J. Chase, L. Chase, and C. Newman, Discontinuity of the Magnetization in One-Dimensional $1/|x - y|^2$ Ising and Potts Models, *J. Stat. Phys.* **50**, 1 (1988).
 - [6] G. Grinstein, A. N. Berker, J. Chalupa, and M. Wortis, Phys. Exact Renormalization Group with Griffiths Singularities and Spin-Glass Behavior: The Random Ising Chain, *Phys. Rev. Lett.* **50**, 1 (1988).
 - [7] T. Niemeyer and J. M. J. van Leeuwen, *Physica (Utr.)* **71**, 17 (1974).
 - [8] J. M. J. van Leeuwen, Singularities in the Critical Surface and Universality for Ising-Like Spin Systems, *Phys. Rev. Lett.* **34**, 1056 (1975).
 - [9] S. R. McKay, A. N. Berker, and S. Kirkpatrick, Spin-glass behavior in frustrated Ising models with chaotic renormalization-group trajectories, *Phys. Rev. Lett.* **48**, 767 (1982).
 - [10] S. R. McKay, A. N. Berker, and S. Kirkpatrick, Amorphously packed, frustrated hierarchical models: Chaotic rescaling and spin-glass behavior, *J. Appl. Phys.* **53**, 7974 (1982).
 - [11] S. R. McKay and A. N. Berker, *J. Appl. Phys.*, Chaotic Spin Glasses: An Upper Critical Dimension, *J. Appl. Phys.* **55**, 1646 (1984).
 - [12] A. N. Berker and S. R. McKay, Hierarchical Models and Chaotic Spin Glasses, *J. Stat. Phys.* **36**, 787 (1984).
 - [13] A. N. Berker and M. Wortis, Blume-Emery-Griffiths-Potts Model in Two Dimensions: Phase Diagram and Critical Properties from a Position-Space Renormalization Group, *Phys. Rev. B* **14**, 4946 (1976).