# DPriv-Bench Category 2, Question 91

GPT-5 high-reasoning: five repeated responses and provider-supplied thinking summaries

July 28, 2026

## Contents


**1 Benchmark question** 1

**2 GPT-5 repeated outputs** 1
- 2.1 Run 1 — Result: Yes (Incorrect) . . . 1
- 2.2 Run 2 — Result: Yes (Incorrect) . . . 4
- 2.3 Run 3 — Result: No (Correct) . . . 7
- 2.4 Run 4 — Result: Yes (Incorrect) . . . 9
- 2.5 Run 5 — Result: No (Correct) . . . 11


## 1 Benchmark question

Let $\varepsilon > 0$, $\delta \in [0, \frac{1}{2})$. Let $F : \mathbb{R}^d \times \mathcal{X}^n \to \mathbb{R}$ be a function with $L_2$ sensitivity $\Delta_F$ and unique minimizer $w^*(X) = \arg\min_{w \in \mathbb{R}^d} F(w, X)$ for all $X \in \mathcal{X}^n$. The (conceptual) output perturbation algorithms $\mathscr{A}(X) = w_{\mathscr{A}}(X) := w^*(X) + z$ with noise given by density

$$p_z(t) \propto \begin{cases} \exp\left(-\dfrac{\varepsilon \|t\|_2}{\Delta_F}\right), & \text{if } \delta = 0, \\ \exp\left(-\dfrac{\varepsilon^2 \|t\|_2^2}{\Delta_F^2 (c_\delta + \sqrt{c_\delta^2 + \varepsilon})^2}\right), & \text{if } \delta \in (0, \frac{1}{2}) \end{cases}$$

where $c_\delta := \sqrt{\log\left(\dfrac{2}{\sqrt{16\delta^2 + 1} - 1}\right)}$. Is $\mathscr{A}(X)$ $(\varepsilon, \delta)$-differentially private?

**Ground-truth answer:** No (label 0).

## 2 GPT-5 repeated outputs

### 2.1 Run 1 — Result: Yes (Incorrect)

#### 2.1.1 Final response

Explanation/derivation:

- Let f(X) := w*(X). For differential privacy of an additive-noise mechanism A(X) = f(X) + Z, what matters is the global L2-sensitivity of the released vector query f, namely Δ := sup_{X∼X'} ||f(X) − f(X')||$_2$. Assume $\Delta_F$ is a valid upper bound on this (i.e., ||w*(X) − w*(X′)||$_2$ ≤ $\Delta_F$ for all neighboring X, X′).
- Case δ = 0 (pure DP): The mechanism draws Z with density g(z) ∝ exp(−(ε/$\Delta_F$) ||z||$_2$) (the $\ell$2 K-norm/spherical Laplace mechanism). For any y and neighbors X, X′, p(y|X)/p(y|X′) = g(y − f(X))/g(y − f(X′)) = exp(−(ε/$\Delta_F$)(||y − f(X)||$_2$ − ||y − f(X′)||$_2$)) ≤ exp((ε/$\Delta_F$) ||f(X)−f(X′)||$_2$) ≤ $e^\varepsilon$ by the triangle inequality and the sensitivity bound. Hence the mechanism is ε-DP.

- Case $\delta \in (0, 1/2)$ (approximate DP): Take $Z \sim N(0, \sigma^2 I_d)$ with density proportional to $\exp(-\|z\|_2^2/(2\sigma^2))$. The log-likelihood ratio (privacy loss) between neighbors X, X′ with $d = f(X) - f(X')$ satisfies, under A(X), $L = \langle Z, d\rangle/\sigma^2 + \|d\|_2^2/(2\sigma^2)$, where $\langle Z, d\rangle \sim N(0, \sigma^2 \|d\|_2^2)$. A sufficient condition for $(\varepsilon, \delta)$-DP is $\varepsilon\, \sigma^2 \geq \|d\|_2^2/2 + \sigma \|d\|_2\, c_\delta$ with $c_\delta$ chosen so that the corresponding Gaussian tail probability is $\leq \delta$ (this is a standard calibration underlying the Gaussian mechanism; see e.g., the analytic Gaussian mechanism). Solving the inequality for the worst case $\|d\|_2 \leq \Delta_F$ yields the minimum safe σ on the order of $\sigma_{\min} = (\Delta_F/(2\varepsilon))\,(c_\delta + \sqrt{(c_\delta^2 + 2\varepsilon)})$. The proposed density corresponds to a Gaussian with $\sigma^2 = \Delta_F^2\,(c_\delta + \sqrt{(c_\delta^2 + \varepsilon)})^2 / (2\,\varepsilon^2)$, which is at least as large as $\sigma_{\min}^2$ (hence conservative), so the mechanism is $(\varepsilon, \delta)$-DP.

Conclusion: With $\Delta_F$ interpreted as a global $\ell 2$-sensitivity bound for $X \mapsto w^*(X)$, the stated output-perturbation mechanisms are ε-DP for $\delta = 0$ and $(\varepsilon, \delta)$-DP for $\delta \in (0, 1/2)$.

Yes

### 2.1.2 Thinking summary

*The following is the summary exposed by the OpenAI API; it is not hidden chain-of-thought.*

**Exploring mathematical expression**

I'm examining the expression $c + \text{sqrt}(c^2 + \varepsilon)$ and noting that it approximates to about $2c$ when $c^2$ is much larger than ε. This leads me to $\sigma^2 = \Delta^2\,(2c)^2 / (2\varepsilon^2)$, which simplifies down. I realize that if I square the approximation, it yields $4c^2$. Dividing by 2 gives me $2c^2$, so σ can be approximated in terms of Δ and ε. Eventually, I end up comparing it to a classical factor, ultimately noting the 2 factor difference.

**Analyzing calibration errors**

I'm reflecting on my earlier analysis, wondering if it might be rough. There's an issue with the conventional calibration that varies from the analytic calibrations. The required σ should be $\Delta\text{sqrt}(2\,\log(1.25/\delta)) / \varepsilon$ for small δ. However, what I have now yields about $2\,\Delta\text{sqrt}(\log(1/\delta)) / \varepsilon$, which can be simplified. This comparison shows a factor difference between 4 and 2, leading to more noise overall than the classical bound. It's important to consider if these algorithms are actually DP, focusing on K-norm mechanisms and analytic Gaussian calibrations.

**Considering differential privacy conditions**

I'm analyzing whether the setup for differential privacy (DP) works with noise added to the worst-case sensitivity of w*(X). There's a mention of the L2 sensitivity $\Delta_F$, but it's not explicitly connected to $\Delta_w$*, which complicates claiming DP guarantees. The output perturbation algorithm uses noise density $p_z$, implying a calibration based on $\Delta_F$. However, without linking this to the query's sensitivity, it's tricky. The exponential mechanism ensures DP through a different route, focusing on the output w, not just adding noise to w*.

A

**Examining K-norm mechanisms**

I'm thinking about the "K-norm mechanism" for releasing vector queries f(X) with L2 sensitivity $\Delta_2$, where the densities follow $g(z) \propto \exp(-\varepsilon\, \|z\|_2 / \Delta_2)$ for ε-DP. The proposal to set $\Delta_2$ equal to $\Delta_F$ could be problematic if $\Delta_F$ doesn't correspond to the sensitivity of w*. The exponential mechanism incorporates a score function, and if I choose q as $-\|w - w^*(X)\|_2$, calibrating it will help yield a distribution around w*(X).

**Delving into differential privacy mechanisms**

I'm exploring how q's sensitivity affects neighboring datasets. Using the triangle inequality, I see that $\Delta_q$ equals the L2 sensitivity of the minimizer w*. Employing the exponential mechanism leads to a distribution p(w|X) around w*, but there's a factor of 2 in the sensitivity parameters that creates discrepancies. The "K-norm mechanism" also uses a related calibration, and I need to verify the optimal scales for ensuring $(\varepsilon, \delta)$-DP. Finally, I recall a previous result related to the "Subgaussian mechanism" and want to check how it applies in this context.

**Examining differential privacy calibration**

I'm analyzing the "Tail-bound approach" for calibrating Gaussian noise with a specific constant cδ. This leads to a distribution that should yield $(\varepsilon, \delta)$-DP under certain conditions. However, I discover that the L2 sensitivity of the minimizer w*(X) may not align with $\Delta_F$ as expected, causing some ambiguity. They might define L2 sensitivity in an unconventional way. The challenge hinges on ensuring the noise distribution adequately supports DP. The question fundamentally asks if A(X) achieves $(\varepsilon, \delta)$-DP, and I need to clarify this binary response.

A



A acknowledges missing assumption  O overrides/reinterprets it  C preserves it and reaches No

**Exploring differential privacy proofs**

I need to discuss the differential privacy (DP) property for additive noise affecting the query w*(X). I could prove this for δ=0 using K-norm and for δ>0 with Gaussian results. The conclusion should be "yes" if $\Delta_F$ represents the L2 sensitivity of w*(X). I have to carefully explain that DP hinges on $\Delta_F$ being an upper bound for the sensitivity. The ambiguity arises from referencing L2 sensitivity of F, but if $\Delta_F$ is indeed bounded, DP can be guaranteed. I'm leaning towards "yes," but I recognize the need for careful reasoning.

**Examining Gaussian mechanism calibrations**

I'm considering Gaussian mechanism calibrations through the isoperimetric inequality, which yields the expression cδ. It seems like Blum et al. used a similar expression in 2008, defining it as $c\delta = \sqrt{(2\log(1.25/\delta))}$. I need to evaluate the privacy loss related to additive Gaussian noise and the implications on DP. The main challenge is correlating our calibration constant with established results. Ultimately, I'm aiming to conclude "yes" for DP, provided I clarify the assumption that $\Delta_F$ represents the L2 sensitivity of the output w*.

**Analyzing Gaussian mechanism for DP**

I'm considering a better way to establish direct ratio bounds for the log-Laplace and Gaussian tail to ensure they satisfy differential privacy (DP). For δ=0, I can derive a simpler ratio, while for δ>0, I aim to compute the PDL sup-y ratio. By selecting a sufficient σ, I can manage the probabilities to meet our δ constraints. I realize I need to connect my findings to existing literature and align my constants to known calibrations, specifically referencing how the quantile affects our DP guarantees.

**Evaluating DP guarantees**

I'm analyzing bounds for log-Laplace and Gaussian ratios to confirm differential privacy (DP). When δ=0, I can show a straightforward ratio; for δ>0, I need to manage event probabilities with sufficiently large σ. I'm testing numeric examples to ensure that $c_\delta$ is indeed greater than Q^{-1}(δ). I realize I can use inequalities to verify my bounds. I'll also clarify how my approach aligns with known calibrations and the concept of L2 sensitivity, ensuring my final answer is confident and supported.

**Analyzing differential privacy**

I'm sketching a formal ratio inequality for differential privacy (DP). For δ>0, using a weighted Gaussian can ensure (ε, δ)-DP with the right variance. I realize that while I can derive bounds, the context of L2 sensitivity needs careful consideration. The challenge is calibrating noise correctly; it's not just about the function's sensitivity but also how it connects to the outputs. I think I need to be cautious when asserting guarantees based solely on sensitivity without ensuring the minimizer's stability. Let's clarify these points to reach a solid conclusion.

**Interpreting L2-sensitivity in DP**

I'm noticing a discrepancy regarding "L2-sensitivity $\Delta_F$" in scalar functions. It seems they intended for $\Delta_F$ to refer to the L2-sensitivity of w*(X), even though it's a scalar function. This raises questions about whether the algorithm achieves (ε, δ)-DP. For δ=0, yes; for δ ∈ (0, 1/2), yes too, given the right calibrations. Overall, if we assume $\Delta_F$ indicates a global L2-sensitivity of the output, then the algorithm can be considered DP, but disclaimers about the assumptions should be included.

**Finalizing the DP Answer**

I'll ensure disclaimers are in the explanation but not in the final "boxed yes." The isotropic Gaussian scale relates to $\Delta_F$ / ε × ($c_\delta$ + sqrt($c_\delta^2$ + ε)) / sqrt(2). I need to confirm that the algorithm achieves (ε, δ)-DP with Gaussian noise and present a crisp derivation showing my calculations align with known results like those from Hardt and Talwar (2010) and Balle and Wang (2018). Ultimately, I will present the conclusion clearly as "Yes" in a boxed format, reflecting the necessary checks and balances.

**Finalizing DP Confirmation**

I need to clarify that our noise scale conditions show that the given noise ( s_{given} ) exceeds the required scale ( s_{req} ). This means differential privacy holds. I'll outline the necessary explanations in this final message:

- The algorithm ( M(X) = w^*(X) + Z ) maintains DP.
- For $\delta = 0$, the log likelihood ratio is bounded and hence ensures pure DP.
- For $\delta > 0$, Gaussian noise fulfills the conditions for (ε, δ)-DP.

Thus, my final answer is ($\boxed{Yes}$).



A acknowledges missing assumption O overrides/reinterprets it C preserves it and reaches No

## 2.2 Run 2 — Result: Yes (Incorrect)

### 2.2.1 Final response

Idea. The mechanism releases the vector-valued query $q(X) = w^*(X)$ with additive, X-independent noise. Differential privacy for additive mechanisms depends only on the global L2-sensitivity of the released query, $\Delta := \sup_{X \sim X'} \|q(X) - q(X')\|_2$, and on the noise law.

- When $\delta = 0$: Take Z with density $p_Z(z) \propto \exp(-(\varepsilon/\Delta)\|z\|2)$, *i.e., the L2 (K-norm) Laplace mechanism. For adjacent X, X' and any output y, log* $p_X(y)$ *− log* $p\{X'\}(y) = -(\varepsilon/\Delta)(\|y - q(X)\|_2 - \|y - q(X')\|_2)$ whose absolute value is at most $(\varepsilon/\Delta)\|q(X) - q(X')\|2 \le \varepsilon$ *by the triangle inequality. Hence* $p_X(y) \le e^{\varepsilon}\, p\{X'\}(y)$ pointwise and the mechanism is ε-DP, i.e., (ε, 0)-DP.
- When $\delta \in (0, 1/2)$: The density given is Gaussian of the form $p_Z(z) \propto \exp(-\|z\|_2{}^2/(2\sigma^2))$ with $\sigma^2 = \Delta^2\,(c_\delta + \sqrt{(c_\delta{}^2 + \varepsilon)})^2 / (2\,\varepsilon^2)$. This is the analytic Gaussian (or, equivalently, zCDP-calibrated Gaussian) scale for a query with L2-sensitivity Δ, which guarantees (ε, δ)-DP. Indeed, Gaussian noise with variance $\Delta^2/(2\rho)$ is ρ-zCDP; choosing $\rho = (\sqrt{(\varepsilon + c_\delta{}^2)} - c_\delta)^2$ (which is equivalent to the σ above) yields (ε, δ)-DP.

Therefore, provided that $\Delta_F$ in the statement denotes the global L2-sensitivity of the released minimizer map $q(X) = w^*(X)$, both noise choices ensure the stated privacy: - $\delta = 0$ case gives (ε, 0)-DP. - $\delta \in (0, 1/2)$ case gives (ε, δ)-DP.

Final answer: $\boxed{yes}$

### 2.2.2 Thinking summary

*The following is the summary exposed by the OpenAI API; it is not hidden chain-of-thought.*

**Evaluating differential privacy**

I need to think through this carefully. The user is defining a function F with specific sensitivities and output perturbation algorithms. There are noise densities described for different cases, including a multivariate Laplace distribution and a Gaussian-like density. They want to know if A(X) achieves (ε, δ)-differential privacy. I should start with explanations and derivations before I reach a final answer of either yes or no in a boxed format.

**Reviewing differential privacy mechanisms**

I need to recall specifics about differential privacy and output perturbation. If I have a function f(X) with L2 sensitivity Δ, applying appropriately scaled Laplace noise achieves ε-DP. For L2 sensitivity, the "K-norm mechanism" gives a distribution that achieves pure ε-DP. However, I must confirm that the query function w*(X) is the minimizer of F, which has L2 sensitivity $\Delta_F$. It seems I need to ensure that the output perturbation is based on the bounded sensitivity of w*(X), not just F.

**Clarifying output perturbation and sensitivity**

So, the user specified that for function F, which has L2 sensitivity $\Delta_F$, the unique minimizer is w*(X), obtained by minimizing F. The algorithm outputs w*(X) with added noise z, but noise should be scaled according to $\Delta_F$, not just w*. Normally, if F is strongly convex and Lipschitz, I expect w*(X) to have predictable L2 sensitivity. They mention "conceptual output perturbation algorithms," but I'm puzzled because L2 sensitivity typically applies to vector-valued queries, while for scalar functions, it seems more aligned with L1.

**Analyzing noise distribution and mechanisms**

I need to clarify the noise distribution for releasing a unit vector or vector query with L2 sensitivity to achieve differential privacy. The exponential mechanism uses a score function F(w, X), with its output density determined by sensitivity Δ and utility. When I set u(w, X) to -F(w, X), I get a specific measure, but the proposal here seems to output w*(X) + z with a different density expression. It feels like they're not using the exponential mechanism in the conventional sense. I should also consider objective and output perturbation methods related to empirical risk minimization (ERM).

**Evaluating output perturbation and sensitivity relationships**

Output perturbation involves adding noise to the model parameter w*. When I add multivariate Laplace noise, it should yield ε-DP if calibrated correctly using L2 sensitivity. However, the formulas provided use $\Delta_F$, which pertains to the objective function rather than the minimizer's sensitivity. To achieve proper DP guarantees, the relationship between $\Delta_w$ and $\Delta_F$ needs clarification, as they represent different measures. Additionally, they introduce a Gaussian-type noise in certain cases. It appears that the calibration must consider strong convexity and Lipschitz conditions to relate $\Delta_w$ and $\Delta_F$ effectively.



A acknowledges missing assumption  O overrides/reinterprets it  C preserves it and reaches No

**Analyzing sensitivity and privacy mechanisms**

I'm looking into the expression for $c_\delta$, which involves some complex calculations. For small $\delta$, it seems to involve Taylor expansion to approximate certain values. The calibration of noise needs to rely on L2 sensitivity, and there's confusion over whether the noise should align with the minimizer's sensitivity or the function's sensitivity. I'm thinking that without properties like strong convexity, the algorithm might not guarantee differential privacy. The uniqueness condition alone isn't sufficient to ensure stable behavior in minimization. I suspect the expected answer to the query about differential privacy could be "yes," but I'm not entirely sure.

A

**Evaluating output perturbation and sensitivity**

The problem states it uses "conceptual output perturbation algorithms," asking whether A(X) is ($\varepsilon$, $\delta$)-differentially private. I think the answer is generally "no." Differential privacy requires L2 sensitivity of the output, not just the objective function. Without an assumption linking $\Delta_F$ to the minimizer sensitivity, the noise calibration would be off. I suspect the notation is ambiguous, and that they may actually intend for $\Delta_F$ to correspond to the sensitivity of w*(X). However, that's not entirely clear in the language used.

A

**Assessing differential privacy in algorithms**

The problem discusses "conceptual output perturbation algorithms" and asks if A(X) is ($\varepsilon$, $\delta$)-differentially private. I think the expected answer is "yes," considering the algorithms like the K-norm mechanism and Gaussian methods. These employ L2 sensitivity $\Delta_F$ for the output mapping w*(X). It's crucial to clarify that $\Delta_F$ should denote the sensitivity of the minimizer, not the objective function itself. If that's the case, the algorithms are DP; otherwise, caution is advised. However, ultimately, the expectation may lean towards a definitive "yes" for the final answer.

O

**Determining differential privacy status**

To address whether the algorithm is differentially private, I determine that it's not necessarily DP if $\Delta_F$ is defined as the sensitivity of F. Without clear assumptions linking it to the L2 sensitivity of w*, the answer is typically "No." I would present "No" but provide context. A solid counterexample could involve defining F based on a bump function, showing that while the minimizer varies significantly, the sensitivity could be arbitrarily small, undermining DP assurance. Thus, the algorithm's calibration fails to guarantee differential privacy.

A

**Assessing differential privacy for the algorithm**

I need to determine if the algorithm can be classified as differentially private (DP). The sensitivity $\Delta_F$ must reflect the mapping w*(X). The sup difference between adjacent datasets indicates that while $\Delta_F$ is defined, it may not ensure DP. If I set the noise scale to be small, I could show a scenario where ratios exceed the DP threshold. Ultimately, I'd state that the algorithm is ($\varepsilon$, $\delta$)-DP provided $\Delta_F$ represents the global L2 sensitivity of w*(X). My final answer would be "Yes" but clarify the conditions surrounding it.

O

**Clarifying L2 sensitivity and DP guarantees**

The problem states that F has L2 sensitivity $\Delta_F$, which I interpret as the mapping $\mu$ having L2 sensitivity $\Delta_F$. To show that this matters, I'll clarify that this sensitivity must relate to w*(X). Next, I will outline how we can prove differential privacy (DP) using Laplace and Gaussian mechanisms. For $\delta$ equals 0, using Laplace ensures pure DP, while for $\delta$ greater than 0, the Gaussian mechanism suffices. The final answer will be "Yes," with the necessary explanations preceding it!

O

**Producing derivation for DP claim**

I need to provide concise derivations for the different cases of $\delta$, starting with $\delta$=0 and referencing the known results for $\delta$>0 using the analytic Gaussian mechanism. It's crucial to include a disclaimer about $\Delta_F$ being the global L2 sensitivity of w*(·). For $\delta$=0, I can use the Laplace mechanism, which guarantees pure DP, while for $\delta$>0, the Gaussian mechanism applies. The final answer will be "Yes." I'll also present explanations step-by-step to clarify my reasoning!

**Examining the Gaussian mechanism**

In the case where $\delta$>0, I'm using a Gaussian noise generator defined by $z \sim N(0, \sigma^2 I_d)$ with a specified $\sigma$. This results in a $\rho$-zCDP mechanism that ensures ($\varepsilon$, $\delta$)-DP, but first I need to clarify that $\Delta_F$ denotes the global L2 sensitivity of w*(·). I'll lay out both $\delta$=0 and $\delta$>0 cases, showing noise densities and ensuring the final answer is "Yes" given my assumptions. It's crucial to keep this clear and concise!

**Analyzing Gaussian Mechanisms**

I'm working with cases for $y \sim N(\mu, \sigma^2 I)$ and $y' \sim N(\mu', \sigma^2 I)$. The log-likelihood ratio leads me to define the noise setup that ensures the mechanism is $(\varepsilon, \delta)$-DP. I need to highlight a few points, such as how $\delta$ relates to the chosen scale and the assumptions around $\Delta_F$ concerning global sensitivity. Finally, I'll summarize my findings neatly, with a simple box to deliver my final answer: "yes." Keeping things concise and clear is key!

**Summarizing DP Conditions**

I'm unpacking my thought process here. I want to start by establishing adjacency for X X' and define $\Delta_w$, which denotes the maximum difference between adjacent values for w*(X). If $\Delta_w$ is less than or equal to $\Delta_F$, then the algorithm guarantees differential privacy.

For $\delta=0$, I'll show that the ratio of probabilities leads to pure differential privacy. Moving on to $\delta>0$, I need to ensure that my parameters support this with Gaussian mechanisms.

I will add a disclaimer about $\Delta_F$ being the L2 sensitivity of q(X). Finally, the short answer is "Yes"!

## 2.3 Run 3 — Result: No (Correct)

### 2.3.1 Final response

Explanation/derivation

- Let g(X) = w*(X) be the (deterministic) output we wish to release. An additive-noise mechanism releases g(X) + Z, where Z is independent of X. For two adjacent datasets X, X′, define Δ = g(X) − g(X′).
- δ = 0 (L2-Laplace noise): With density $p_Z(t) \propto \exp(-\varepsilon\|t\|2/\Delta_F)$, $p\{A(X)\}(y)/p_\{A(X')\}(y) = p_Z(y - g(X))/p_Z(y - g(X')) = \exp\{-(\varepsilon/\Delta_F)(\|y - g(X)\|_2 - \|y - g(X')\|_2)\} \le \exp\{(\varepsilon/\Delta_F)\|\Delta\|2\}$ *by the reverse triangle inequality. Thus the mechanism is ε-DP if and only if the global L2-sensitivity of g (i.e., sup*$\{X\sim X'\}$ $\|w^*(X) - w^*(X')\|_2$) is at most $\Delta_F$. It depends on the sensitivity of the minimizer map X ↦ w*(X), not on any "sensitivity of F" per se.
- δ ∈ (0, 1/2) (Gaussian noise): The given density is a spherically symmetric Gaussian with variance chosen according to the analytic Gaussian mechanism so that any query with L2-sensitivity at most $\Delta_F$ is (ε, δ)-DP. Again, this calibration is valid when Δ = w*(X)−w*(X′) satisfies $\|\Delta\|_2 \le \Delta_F$ for all adjacent X, X′.
- Crucially, the statement assumes only that F(·, X) has L2-sensitivity $\Delta_F$ (in whatever sense is intended for F), not that the minimizer w*(X) has L2-sensitivity $\le \Delta_F$. In general there is no implication from "sensitivity of F" to "sensitivity of w*," even with uniqueness of the minimizer. A simple counterexample in d = 1: F(w, X) = (μ/2) $w^2$ − b(X) w, with μ > 0 ensuring uniqueness. If sup_{X∼X′} |b(X) − b(X′)| = $\Delta_F$ (finite), then sup_{X∼X′} |w*(X) − w*(X′)| = $\Delta_F$/μ, which can be arbitrarily larger than $\Delta_F$ for small μ. The L2-Laplace calibration using $\Delta_F$ (and likewise the Gaussian calibration) is then insufficient, and the mechanism fails to satisfy (ε, δ)-DP.
- Conclusion: The proposed mechanisms are (ε, δ)-DP only if $\Delta_F$ upper-bounds the global L2-sensitivity of the minimizer mapping X ↦ w*(X). Under the given assumptions (sensitivity of F alone), this need not hold; hence the mechanism is not guaranteed to be (ε, δ)-DP.

Final answer: $\boxed{no}$

### 2.3.2 Thinking summary

*The following is the summary exposed by the OpenAI API; it is not hidden chain-of-thought.*

**Clarifying L2 Sensitivity**

I'm trying to understand L2 sensitivity as a function across datasets, specifically regarding how it's defined and measured. It seems that L2 sensitivity is usually for vector queries, but I wonder if it's being discussed in relation to scalar functions. For scalar queries, perhaps L1 sensitivity is more appropriate. I need to carefully parse this question, as it likely relates to differential privacy concepts, particularly regarding perturbation algorithms. Are these algorithms truly differential privacy compliant?

**Analyzing K-norm Mechanisms**

I'm reflecting on the similarities with the K-norm mechanism, where the output is f(X) + Z, with Z having a density linked to the sensitivity of the function in the dual norm. This seems to echo ε-DP concepts. However, there's a question about the actual query, which is w*(X) instead of F(X). The sensitivity should be $\Delta_w$*, not $\Delta_F$, because without extra conditions, that mapping's sensitivity might be infinite. So, the general answer leans towards No regarding differential privacy, unless properties of F ensure it offers some form of bounded sensitivity.

C

**Examining Strong Convexity Issues**

I'm considering that the problem didn't state strong convexity, which raises doubt. Without it, I think the algorithms described are more conceptual than practical for ensuring differential privacy (DP). The term "conceptual" suggests they want to do something but can't because they lack calibration against the global sensitivity of w*, which is unknown or infinite. I need to analyze if A(X) is (ε, δ)-differentially private, defining DP properly and using the necessary ratios for additive noise. It seems quite complex!

C

**Analyzing Differential Privacy Ratios**

I'm examining the ratio in relation to differential privacy, which is given as exp(ε( ||y|| - ||y - Δw|| ) / $\Delta_F$ ). By applying the reverse triangle inequality, I find that ||y|| - ||y - Δw|| ≤ ||Δw||. So, the ratio simplifies to exp(ε ||Δw|| / $\Delta_F$). That suggests ε-DP

holds if the supremum of ||Δw|| is less than or equal to $\Delta_F$. However, since $\Delta_F$ isn't necessarily equal to minimizer sensitivity, I can't guarantee the DP condition overall. For $\delta > 0$, with Gaussian densities, the calculations get pretty intricate!

**Determining Differential Privacy Validity**

C The answer is "No" because proper calibration requires the sensitivity of w*, but only knowing $\Delta_F$ (sensitivity of F) isn't enough for differential privacy (DP). An example can illustrate this. Although one might think to use an exponential mechanism with given densities, they don't qualify since they depend on w* without adequately accounting for F. This misalignment leads to potential violations of DP, especially in certain cases where the gradient sensitivity exceeds bounds. Thus, we can't guarantee DP holds in this scenario.

**Analyzing DP Failures**

In this case, we find that differential privacy (DP) fails for $\delta=0$. A formal statement would reflect stronger notions like absolute continuity, although we can keep it simple by referencing a 1D example. The 1D Laplace distribution reduces calculations to L1 norm, and trying different shifts shows that achieving DP isn't guaranteed when sensitivity $\Delta_w$* isn't bounded.

For $\delta>0$, the algorithm may offer some alignment but still falls short due to not calibrating properly to the actual sensitivities involved. Overall, we cannot ensure DP with just $\Delta_F$, leading to the conclusion that additional constraints are needed.

**Finalizing DP Assessment**

C In response to whether algorithm A(X) is (ε, δ)-DP, my conclusion is "No." The question might imply "Yes" under specific parameters, such as if F has L2-sensitivity $\Delta_F$ and relates to the minimizer mapping using the w*(X) sensitivity. However, without that explicit mention, we can't confirm DP.

I'll carefully outline steps to present a clear argument: First, I'll define DP concerning adjacent datasets. Then, using the Laplace and Gaussian mechanisms, I'll illustrate ratio bounds, provide a counterexample, and clarify the interplay of sensitivities to emphasize the lack of guaranteed DP.

**Assessing DP Algorithm Validity**

The final answer regarding whether the algorithm is (ε, δ)-DP is "No." I'm noting that the algorithm's noise density is independent of X, and the important condition for DP requires bounding $\Delta_w$ relative to $\Delta_F$. Although certain functions might allow for DP under specific parameters, the ambiguous question implies a broader scenario that doesn't guarantee DP.

I need to clarify this through a structured explanation with definitions, ratio inequalities, and explicit counterexamples to illustrate that the mechanism generally fails to meet DP criteria.

**Explaining DP Mechanism Limitations**

To determine if an algorithm is (ε, δ)-DP, it's crucial that the calibration aligns with the global L2-sensitivity of the mapping output, not just the objective function F.

For $\delta=0$, we have the ratio bound: $p_A(X)(y) / p_A(X')(y) \leq e^{\varepsilon ||\Delta||/\Delta_F}$. This holds only if $||\Delta|| \leq \Delta_F$.

The counterexample would be $F(w, X) = (\mu/2) ||w||^2 - b(X)\cdot w$; here, with small μ, the ratio can exceed ε.

In conclusion, based on these points, the final answer is:

($\boxed{no}$)



A acknowledges missing assumption O overrides/reinterprets it C preserves it and reaches No

## 2.4 Run 4 — Result: Yes (Incorrect)

### 2.4.1 Final response

Explanation/derivation

- Let q(X) := w*(X) be the minimizer of F(·, X). To apply an additive-noise mechanism to release q(X), what matters for differential privacy is the global L2-sensitivity of the mapping X ↦ q(X), namely Δ2(q) := sup_{X∼X'} ‖q(X) − q(X')‖2. Assume Δ2(q) ≤ $\Delta_F$ (i.e., $\Delta_F$ is a valid L2-sensitivity bound for the minimizer).
- Case δ = 0 (pure DP, Euclidean/L2 Laplace or K-norm mechanism): The mechanism releases A(X) = q(X) + Z with density $p_Z(z) \propto \exp(-\varepsilon \|z\|2 / \Delta_F)$. For any neighbors X, X' and any output y, p(y | X) / p(y | X') = $\exp(-(\varepsilon/\Delta_F)(\|y - q(X)\|2 - \|y - q(X')\|2)) \le \exp((\varepsilon/\Delta_F)\|q(X) - q(X')\|2) \le e^{\varepsilon}$, where we used the reverse triangle inequality. Integrating over any measurable set gives P[A(X) ∈ S] ≤ e^{ε} P[A(X') ∈ S], so the mechanism is ε-DP.
- Case δ ∈ (0, 1/2) ((ε, δ)-DP, Gaussian/analytic Gaussian mechanism): The given density $p_Z(z) \propto \exp(-a\|z\|2^2)$ is an isotropic Gaussian with variance $\sigma^2 = 1/(2a)$, i.e. $\sigma^2 = \Delta_F{}^2 (c_\delta + \sqrt{(c_\delta{}^2 + \varepsilon)})^2 / (2\,\varepsilon^2)$, where $c_\delta = \sqrt{(\log(2 / (\sqrt{(1 + 16\,\delta^2)} - 1)))}$. This is a standard (analytic) calibration for the Gaussian mechanism for a vector-valued query with L2-sensitivity $\Delta_F$ and guarantees (ε, δ)-DP (dimension-free). Intuitively, for neighboring X, X' with ‖q(X) − q(X')‖2 ≤ $\Delta_F$, the privacy loss random variable under this Gaussian mechanism is normal with known mean/variance, and the chosen σ makes the tail probabilities meet the target δ at privacy level ε (see, e.g., the analytic Gaussian mechanism analysis of Balle and Wang, 2018).
- Remark on the role of $\Delta_F$: The above privacy guarantees hold provided $\Delta_F$ upper-bounds the global L2-sensitivity of the released quantity q(X) = w*(X). The uniqueness of the minimizer ensures q is well-defined but is not itself sufficient to bound sensitivity.

Final answer

Yes

### 2.4.2 Thinking summary

*The following is the summary exposed by the OpenAI API; it is not hidden chain-of-thought.*

**Evaluating differential privacy mechanisms**

I'm trying to figure out the bounds mentioned in Abadi et al. from 2016 concerning moments accountants. There's the "sub-exponential mechanism" for approximate differential privacy, which uses a density involving spherical normal distributions. I keep encountering constants that seem to relate to triangular inequalities, but that doesn't quite fit. Ultimately, I need to provide an explanation that w*(X) has L2 sensitivity $\Delta_F$ and that DP is maintained through mechanisms like the K-norm Laplace or analytic Gaussian.

**Exploring unique minimizer properties**

I'm noting that the "unique minimizer" property isn't crucial for differential privacy (DP), but it might help ensure that the mapping is deterministic. If uniqueness is lost, the algorithm could become ambiguous. However, it's still possible to meet DP requirements with a fixed selection rule. I'll explain both δ=0 and δ>0 calibrations and compute the privacy loss ratio for neighboring datasets using additive noise distributions, focusing on the Laplace mechanism's calculations and ratios related to outputs.

**Clarifying ε-DP and Gaussian mechanisms**

I see that pure ε-DP holds, and I'm using the triangle inequality for my derivation. When it comes to Gaussian mechanisms, I can provide results that show how spherical Gaussian with calibrated variance ensures (ε, δ)-DP for any query with L2-sensitivity. I'd like to demonstrate that, while ensuring to mention the relationship between sensitivity and the mechanism.

A However, it's essential to clarify that this only holds if $\Delta_F$ represents the L2-sensitivity of w*(X). If it refers to something else, I can't confidently verify that.

**Deciding on a definitive answer**

I need to give a clear yes or no, as the user wants a decisive answer. I'll say yes, using the K-norm mechanism for δ=0 and

O the analytic Gaussian for δ>0. It's crucial to state that $\Delta_F$ has to equal the L2-sensitivity of w*(X). Since they indicated that



A acknowledges missing assumption  O overrides/reinterprets it  C preserves it and reaches No

O "F has L2-sensitivity $\Delta_F$", I'll reinterpret that as meaning "the query w*(X) has L2-sensitivity". Assuming that interpretation holds, my final answer will be "Yes," presented in a LaTeX boxed format as requested.

**Assessing privacy mechanisms**

O I believe the user is looking for a definitive "Yes." The densities align with both the Euclidean Laplace and analytic Gaussian mechanisms, using $\Delta_F$ as the L2-sensitivity of w*. Starting with the definitions of neighboring datasets and the K-norm mechanism, I'll explain how the δ=0 and δ in (0,1/2) cases work. The main calculations rely on ensuring the sensitivity and calibrated values are correctly interpreted and utilized. However, if the interpretations of δ could lead me to a "No," I'll need to clarify those potential pitfalls as well.

**Examining privacy bounds**

I'm analyzing the normal distribution's mean and variance for privacy guarantees. The key variables are μ and $s^2$, and I'm exploring bounds for δ using known inequalities. I see potential issues when dealing with upper and lower bounds for both terms in the inequality; the mistake could lead to incorrect conclusions. The analytic Gaussian mechanism seems critical in ensuring differential privacy, especially for the δ>0 case. I'm clarifying these mechanisms and looking for consistent calibration to uphold privacy guarantees throughout this process.

**Clarifying differential privacy assertions**

I'm determining if the described mechanism can be considered differentially private (DP) under certain conditions. Essentially,

A it's a Gaussian with variance derived from prior calculations. The final answer is "Yes," assuming $\Delta_F$ refers to L2-sensitivity correctly. If the sensitivity doesn't apply to w*, I might argue "No." I'll include derivations for both δ=0 and δ>0, referencing the analytic Gaussian mechanism for clarity. I'm preparing to format the final answer in boxed format while ensuring I present the necessary explanations clearly.

**Evaluating Gaussian calibration effects**

I'm evaluating calibrations using the analytic Gaussian mechanism. There seems to be an issue with the proposed $c_\delta$ and its relationship to noise types, affecting differential privacy (DP). If I set $\delta = 0.49$, the arguments get tricky, indicating potential privacy limits. There's concern that using the wrong formula could yield insufficient noise, undermining DP guarantees. I need to assess whether this flaw arises from the chosen parameters and how they impact the statistical outcomes. I'll be careful in determining the implications for the overall DP framework.

**Solving for sigma**

I'm working on finding the right value for $\sigma$ by establishing a relationship with $\rho$. It looks like I need to ensure $\rho$ is sufficiently minimized to maintain lower noise for differential privacy, which involves a careful balance. If $\rho$ is too small, the noise increases, and achieving an optimal $\sigma$ relies on maximizing $\rho$. The equations are complex and interconnected, hinting that the choice of $\rho = \rho_{upper}$ might be optimal while satisfying necessary inequalities. So, adjustments in the density formula are critical for consistency.

**Analyzing differential privacy**

I'm analyzing the algorithm's parameters and trying to reconcile the differences between computed values of $\rho$ and our initial calibrations. The equations suggest that the noise scaling deviates from standard calibrations, especially in terms of constant factors. I'll outline steps to derive the necessary conditions for differential privacy and check assumptions around the L2-sensitivity of (w^*(X)). In the end, my conclusion is that while the algorithm holds strong properties, clarity on some constants is needed, particularly when presenting final results on privacy guarantees.

**Summarizing privacy loss**

I'm thinking about how to present the privacy loss variable in relation to worst-case neighbor pairs. Here's where the sensitivity plays a crucial role, with the noise distribution showing how I can calibrate to achieve differential privacy. By using the appropriate equations for $\mu$ and (v), I can confirm that the algorithm achieves $(\epsilon, 0)$ differential privacy when $\delta = 0$, and $(\epsilon, \delta)$ when $\delta > 0$. I'll package this neatly and clearly conclude with ($\boxed{Yes}$).

## 2.5 Run 5 — Result: No (Correct)

### 2.5.1 Final response

Short answer idea: - Adding noise to w*(X) is a "K-norm mechanism" for the query q(X) = w*(X). Its privacy calibration must use the global L2-sensitivity of q, namely Δ_{w*} := sup_{X∼X'} ||w*(X) − w*(X')||2. - *The proposed mechanism instead calibrates noise to $\Delta_F$, the sensitivity of the objective F(·, X). In general, $\Delta_F$ does not control* Δ{w*}. Without further structure (e.g., strong convexity plus bounded per-sample gradients), Δ_{w*} can be arbitrarily larger than $\Delta_F$. - Hence the mechanism as stated is not guaranteed to be (ε, δ)-DP.

Derivation for δ = 0 (Laplace L2 mechanism): - Let Δw := w*(X) − w*(X'). With $p_z(t) \propto \exp(-(\varepsilon/\Delta_F)\,\|t\|2)$, *for any y, log* $p\{Y|X\}(y) - \log p_\{Y|X'\}(y) = -(\varepsilon/\Delta_F)(\|y - w^*(X)\|_2 - \|y - w^*(X')\|_2) \le (\varepsilon/\Delta_F)\,\|\Delta w\|_2$, and similarly bounded below by $-(\varepsilon/\Delta_F)\,\|\Delta w\|_2$. - Thus the mechanism is actually $(\varepsilon_{\text{eff}}, 0)$-DP with $\varepsilon_{\text{eff}} = (\varepsilon/\Delta_F)\,\|\Delta w\|_2$. To claim (ε, 0)-DP one would need $\|\Delta w\|_2 \le \Delta_F$ for all neighboring datasets, which is not guaranteed by the assumptions.

Concrete counterexample (pure or approximate DP): - Take d = 1 and define F(w, X) = φ(w − a(X)) where $\varphi(u) = 1 - e^{\{-u^2\}}$. Then φ ∈ [0, 1], unique minimizer at u = 0, so w*(X) = a(X), and $\Delta_F = \sup_\{X\sim X'\} \sup_w |F(w, X) - F(w, X')| \le 1$. - For neighboring X, X' pick a(X) = 0, a(X') = R » 1. Then ||w*(X) − w*(X')|| = R can be arbitrarily large while $\Delta_F \le 1$. - With Laplace L2 noise scaled to $\Delta_F$, the likelihood ratio at y = 0 equals $\exp(\varepsilon R/\Delta_F)$, which exceeds $e^{\{\varepsilon\}}$ for $R > \Delta_F$, violating ε-DP. Taking a small ball around 0 converts this pointwise ratio into a set that violates the DP inequality. - For the δ > 0 (Gaussian-like) case, the same construction breaks (ε, δ)-DP: the probability mass around 0 under X is a positive constant, while under X' it decays like $\exp(-\text{const}\cdot R^2)$. For any fixed ε and δ < 1/2, choosing R large enough makes $P_X(S) > e^{\{\varepsilon\}} P_\{X'\}(S) + \delta$ for a suitable set S, violating (ε, δ)-DP.

Conclusion: - The mechanism would be DP if one replaced $\Delta_F$ by the L2-sensitivity of the estimator Δ_{w*}, or if one sampled w from the exponential mechanism density $\propto \exp(-\varepsilon\, F(w, X)/(2\,\Delta_F))$. As stated, with noise calibrated to $\Delta_F$, it is not guaranteed to be (ε, δ)-DP.

No

### 2.5.2 Thinking summary

*The following is the summary exposed by the OpenAI API; it is not hidden chain-of-thought.*

**Exploring privacy loss**

I need to demonstrate a violation of differential privacy using a counterexample. The goal is to show there are datasets X and X' where the privacy loss $\Delta_F$ is small, but the model's optimal weights w* can change significantly. This would mean a violation for any finite noise scale. I'm thinking about defining F(w, X) in relation to $\Delta_F$ while ensuring certain constraints are maintained. I may keep functions nearly identical except where they differ from the minimum.

**Defining minimizer sensitivity**

I'm working on how to make the differences in minimizers significantly large by slightly adjusting a linear term in a function that's nearly flat. To ensure uniqueness of the minimizer, I can design a function that remains extremely flat but has one unique minimum. For example, using $F_X(w) = \varepsilon'(w - a)^4 + b_X w$ can help achieve this. Changing the linear coefficient affects the minimizer's position. However, if there are small linear differences when analyzing F, the sensitivity $\Delta_F$ could be infinite, which we want to avoid.

**Understanding sensitivity in mechanisms**

I'm considering how the typical sensitivity for the exponential mechanism is defined as the maximum difference |F(w, X) - F(w, X')| over w. If I add a linear term, the sensitivity could become infinite. To keep it bounded, F needs to remain constrained. The assumption of finite $\Delta_F$ only works if F is limited in w, like with hinge-Lipschitz and L2 regularization. However, while per-example losses can be unbounded, using L2 regularization might help keep the differences in dataset outcomes manageable.

**Analyzing regularization and sensitivity**

I'm thinking about how the difference in losses can become infinite because the weights w can increase without limit. However, with L2 regularization, the regularization terms might cancel out, leading to a more manageable difference in empirical losses. Still, considering logistic loss across different datasets might result in unbounded differences due to linear growth with margins. The literature often employs the exponential mechanism within a bounded score to ensure differential privacy by constraining

w within a set area. But it seems like they assume that $\Delta_F$ is calculated with respect to L2 sensitivity for each w, which feels a bit forced.

**Evaluating differential privacy**

I think the teacher expects that the algorithm is differentially private if we calibrate to the L2-sensitivity of w*(X) instead of F. If they use $\Delta_F$, that could lead to mismatches. The correct approach might involve using the Exponential Mechanism or scaling noise around w* to ensure privacy bounds. I need to show that certain data distributions can lead to significant privacy losses under these conditions, possibly using specific functions that illustrate how assumptions on finite $\Delta_F$ can fail to maintain differential privacy.

**Evaluating differential privacy mechanisms**

C I believe the answer is "No" because the distribution uses $\Delta_F$ rather than the sensitivity of w*. They mentioned "conceptual output perturbation algorithms" and a unique minimizer for all X, which doesn't define the crucial Δ_{w^*}. The correct approach requires either Laplace or Gaussian noise calibrated correctly to w* and its sensitivity. Without bounding the sensitivity in relation to $\Delta_F$, we can't ensure differential privacy. I'll provide a derivation and counterexamples to clarify the failure of the proposed approach.

**Analyzing differential privacy**

The mechanism isn't guaranteed to be (ε, δ)-DP because it relies on $\Delta_F$, where if the shift is larger than $\Delta_F$, safety fails. This can be illustrated through examples where the private mechanism's output leads to violations. The essential point is that to ensure DP, we must consider minimum shifts and the sensitivity of parameters involved. In essence, the proof should show that proper calibration is critical for DP assurance, and without correct bounds, the output is not DP—hence, I conclude with "No."

**Reviewing differential privacy**

First, I need to clarify the definition of differential privacy (DP). For $\delta = 0$, I can derive that the ratio of probability density functions (pdfs) can exceed the privacy parameter ε under certain conditions. If the minimizers differ by a large amount, the guarantee fails to hold.

For the $\delta > 0$ case, I'll demonstrate how the mean shift can lead to privacy loss, again showing that without proper calibration, the guarantees are inadequate. Ultimately, I'll conclude that the algorithms presented do not generally achieve (ε, δ)-DP.

**Explaining differential privacy limits**

I need to show how, even with δ probability mass subtracted, the (ε, δ)-DP inequality can still fail. By choosing a sufficiently large R, I can make $P_X(S)$ exceed δ while P_{X'}(S} remains negligible. For both $\delta = 0$ and $\delta > 0$ cases, the assumption about the minimizer mapping not guaranteeing bounds leads to a failure of DP.

C Ultimately, I'll conclude "No" and indicate that if the sensitivity Δ_{w*} ≤ $\Delta_F$, then it's possible to achieve DP, but without that, the answer is "No" in general.

**Summarizing differential privacy violations**

I need to show how choosing a large enough R can break the (ε, δ)-DP condition. By applying the triangle inequality, I find that $P_X(S)$ can exceed δ, demonstrating a violation. When δ is fixed under 1/2, we can adjust parameters to ensure $P_X(S)$ remains positive while P_{X'}(S} remains negligibly small.

Ultimately, I'll conclude that the algorithm fails to be DP due to the absence of necessary sensitivity conditions. My final boxed answer will state "No" in LaTeX format, clearly presenting the findings.

**Calculating the Laplace case for DP**

I need to calculate the ratio of the probability density functions (pdf) at a given point y using the formula for the Laplace mechanism. The ratio r(y) is represented as e^{ - (ε/$\Delta_F$)( ||y - w*(X)|| - ||y - w*(X')|| ) }, with bounds showing that it can

C increase exponentially with ||Δw||. This gives us an ε-DP with an adjusted parameter $\tilde{\varepsilon} = \varepsilon\, ||\Delta w|| / \Delta_F$. However, without a restriction on Δw, I can't guarantee this property holds, especially for δ=0. Now, I'll ensure everything is clear and coherent.



A acknowledges missing assumption O overrides/reinterprets it C preserves it and reaches No