

Computing Maximum Hamiltonian Paths in Complete Graphs with Tree Metric

Wojciech Rytter^{*1,2} and Bartosz Szreder¹

¹ Dept. of Mathematics, Computer Science and Mechanics,
University of Warsaw, Warsaw, Poland, [rytter,szreder]@mimuw.edu.pl

² Dept. of Math. and Informatics, Copernicus University, Toruń, Poland

Abstract. We design a linear time algorithm computing the maximum weight Hamiltonian path in a weighted complete graph K_T , where T is a given undirected tree. The vertices of K_T are nodes of T and $weight(i, j)$ is the distance between i, j in T . The input is the tree T and two nodes $u, v \in T$, the output is the maximum weight Hamiltonian path between these nodes. The size n of the input is the size of T (however the total size of the complete graph K_T is quadratic with respect to n). Our algorithm runs in $\mathcal{O}(n)$ time. Correctness is based on combinatorics of alternating sequences. The problem has been inspired by a similar (but much simpler) problem in a famous book of Hugo Steinhaus on elementary mathematical problems.

1 Introduction

The maximum Hamilton cycle and path problems are generally NP-hard, see [2,3]. We introduce an interesting class of graphs for which these problems are solvable in linear time. Although it is rather of small practical importance, it is combinatorially and algorithmically quite interesting.

In his famous book “*One Hundred Problems in Elementary Mathematics*” Hugo Steinhaus as Problem 65 asked for the value $\max(n)$ of a maximum Hamiltonian path in the graph K_n with weights of edges between a pair of vertices (i, j) given by $|i - j|$.

In other words the weights of edges correspond to the metric of a simple path of nodes, a trivial case of an undirected tree. In this paper we extend this to arbitrary tree with positive weights on edges. In case of a metric given by a simple path there are very elementary closed formulas for the total weight of maximum Hamiltonian paths in graphs implied by this metric.

Lemma 1. [H. Steinhaus, see [1]]

If n is even then $\max(n) = \frac{n^2-2}{2}$, otherwise $\max(n) = \frac{n^2-3}{2}$.




Fig. 1. A longest Hamiltonian path \mathcal{P} when T is the line with $n = 9$ nodes and unit cost edges. According to Lemma 1 we have $weight(\mathcal{P}) = \frac{n^2-3}{2} = 39$. This path starts in a centroid and ends in its neighbor. Maximum paths between arbitrary pairs of nodes (usually not adjacent ones) are more complicated.

In this paper we extend this to a more complicated problem of constructing in linear time a maximum Hamiltonian path between any given pair of nodes (with a metric implied by an arbitrary tree).

Assume T is an undirected tree with the set of nodes $\{1, 2, \dots, n\}$. By $dist_T(u, v)$ we denote the length of the shortest path between u and v in T . Let K_T be the complete graph K_n with weights of edges given by

$$weight(u, v) = dist_T(u, v).$$

We define formally our problem as follows:

input: given a tree T with n nodes and two different nodes $u, v \in T$;
output: the maximum weight Hamiltonian path in K_T from u to v .

Our main result is a linear time algorithm solving this problem. The main tools are *centroids* of a tree and *alternating sequences*: colored sequences in which adjacent elements are of different colors. In the next sections we discuss them in detail.

2 Alternating sequences \Rightarrow maximum Hamiltonian paths

Assume we have a coloring \mathcal{C} of a set of n elements. We represent a coloring as a partition of this set into color classes:

$$\mathcal{C} = (C_1, C_2, \dots, C_k)$$

We say that a sequence γ over this set is \mathcal{C} -*alternating* iff it is a permutation of all n elements of this set and no two adjacent elements of the sequence are of the same color.

* Supported by grant no. N206 566740 of the National Science Centre.

Example 1. Let

$$\mathcal{C} = (\{1, 2\}, \{3, 4, 5\}, \{6, 7, 8, 9, 10, 11\}),$$

then as an alternating sequence we can take

$$\gamma = (6 \ 4 \ 7 \ 1 \ 8 \ 3 \ 9 \ 2 \ 10 \ 5 \ 11).$$

Let's project our notion of coloring onto a set of vertices of a tree. We select some node S and root the tree in this node. Then by S -coloring we mean a coloring such that the color of each node $v \neq S$ corresponds to the subtree of S containing v and the color of S is S itself. For $S \in T$ denote




Fig. 2. A tree rooted in S gives S -coloring as a family of sets $\mathcal{C} = (\{S\}, \{2\}, \{1, 5\}, \{3, 4, 6\}, \{7, 8, 9\})$. Each set represents different color and corresponds to a distinct subtree of the tree rooted in S .

by $\Delta(S)$ the sum of distances from S to all other vertices in T .

Lemma 2. *Let S be any node of T and \mathcal{C} be the S -coloring of all nodes of T . If γ is a \mathcal{C} -alternating sequence starting with u and ending with v then:*

- (a) γ is a maximum weight Hamiltonian path from u to v in K_T
- (b) $weight(\gamma) = \Delta(S) - \text{dist}(u, S) - \text{dist}(v, S)$.

Proof. We first prove the following fact:

Claim. We show that for any node S' and a path $\gamma' = x_1 x_2 \dots x_n$, where $x_1 = u$, $x_n = v$, we have

$$weight(\gamma') \leq \Delta(S') - \text{dist}(u, S') - \text{dist}(v, S')$$

Proof. Observe that the triangle inequality

$$\text{dist}(x_i, x_{i+1}) \leq \text{dist}(x_i, S') + \text{dist}(S', x_{i+1})$$

implies the following:

$$\begin{aligned} \text{weight}(\gamma') &\leq \text{dist}(x_1, S') + \text{dist}(S', x_2) + \\ &\quad + \text{dist}(x_2, S') + \text{dist}(S', x_3) + \\ &\quad \dots \\ &\quad + \text{dist}(x_{n-1}, S') + \text{dist}(S', x_n) \\ &= \Delta(S') - \text{dist}(u, S') - \text{dist}(v, S') \end{aligned}$$

This completes the proof of the claim.

Consequently a sequence γ (if it exists) satisfying the property in the assumption of Lemma 2 is of maximum weight. This result follows from definition of S -coloring: x_i and x_{i+1} are of different colors, so they reside in different subtrees of a tree rooted in S (or one of them is S , but the argument holds anyway). Because of that we now have:

$$\forall_{1 \leq i < n} \text{dist}(x_i, x_{i+1}) = \text{dist}(x_i, S) + \text{dist}(S, x_{i+1})$$

This completes the proof of the lemma.

Unfortunately the lemma gives no hint about existence and efficient location of a node S satisfying the assumption. The next two sections give tools to find good S .

3 Some useful combinatorics of alternating sequences

Assume $\mathcal{C} = \{C_1, C_2, \dots, C_k\}$ such that $|C_1| \leq |C_2| \leq \dots |C_k|$. Denote by $\max(\mathcal{C})$ the size of the largest set of elements of the same color and by $\text{rem}(\mathcal{C})$ the number of remaining elements, hence:

$$\text{rem}(\mathcal{C}) + \max(\mathcal{C}) = n.$$

For two sequences

$$\alpha = (a_1, a_2, \dots, a_p), \beta = (b_1, b_2, \dots, b_q), \quad q \leq p$$

we define the *interleave* operation as follows:

$$\text{interleave}(\alpha, \beta) = (a_1, b_1, a_2, b_2, a_3, b_3, \dots, a_p)$$

Example 2. For sequences $\alpha = (1, 2, 5, 8, 9)$ and $\beta = (6, 3, 4)$ we have $interleave(\alpha, \beta) = (1, 6, 2, 3, 5, 4, 8, 9)$.

We also introduce the operation *sequentialize* which for a sequence of disjoint sets produces the sequence of all elements of these sets starting with elements of the first set and continuing with elements of consecutive sets (order inside sets is arbitrary).

Lemma 3 (on existence of an alternating sequence).

1. A \mathcal{C} -alternating sequence exists if and only if $\max(\mathcal{C}) \leq rem(\mathcal{C}) + 1$.
2. If it exists then it can be constructed in linear time.
3. If both end-elements of the computed sequence γ are of the same color then this color occurs $rem(\mathcal{C}) + 1$ times.

Proof.

The necessity is obvious: if $\max(\mathcal{C}) > rem(\mathcal{C}) + 1$ then we do not have enough remaining elements to separate all elements of the largest color, so in any permutation two of them should be neighbors.

Assume now that $\max(\mathcal{C}) \leq rem(\mathcal{C}) + 1$ and consider the case of odd number n of elements. We can construct the required sequence γ in the following way:

Algorithm 1: AlterSeq1(\mathcal{C})

input : $\mathcal{C} = \{C_1, C_2, \dots, C_k\}$, $|C_1| \leq |C_2| \leq \dots |C_k|$
output: \mathcal{C} -alternating sequence of $C_1 \cup C_2 \cup \dots C_k$
1 Find j and a partition of $C_j = C'_j \cup C''_j$ such that
 $|C_1 \cup C_2 \cup \dots C'_j| + 1 = |C''_j \cup C_{j+1} \cup \dots C_k|$;
2 $\alpha := sequentialize(C_k, C_{k-1}, \dots C_{j+1}, C''_j)$;
3 $\beta := sequentialize(C'_j, C_{j-1}, \dots C_2, C_1)$;
4 $\gamma := interleave(\alpha, \beta)$;
5 **return** γ ;

A similar argument can be used in case of even n : now we split the coloring to get equality: $|C_1 \cup C_2 \dots \cup C'_j| = |C''_j \cup C_{j+1} \dots \cup C_k|$. The algorithm *AlterSeq1* produces \mathcal{C} -alternating sequence γ of $C_1 \cup C_2 \dots C_k$ in linear time, assuming that $\max(\mathcal{C}) \leq rem(\mathcal{C}) + 1$. This completes the proof of the lemma.

Example 3. Let

$$\mathcal{C} = (C_1, C_2, C_3, C_4) = (\{1, 2\}, \{3, 4\}, \{5, 6, 7, 8\}, \{9, 10, 11, 12, 13\})$$

Then $j = 3$, $C'_3 = \{5, 6\}$, $C''_3 = \{7, 8\}$,

$$\begin{aligned}\alpha &= (13, 12, 11, 10, 9, 8, 7), & \beta &= (6, 5, 4, 3, 2, 1) \\ \gamma &= \text{interleave}(\alpha, \beta) = (13, 6, 12, 5, 11, 4, 10, 3, 9, 2, 8, 1, 7)\end{aligned}$$

Much more complicated is the question of alternating sequences starting and ending in given nodes u, v .

Example 4. Suppose $\mathcal{C} = (\{1, 2\}, \{3, 4\})$ and we impose the condition that the sequence starts in $u = 1$ and ends with $v = 2$. Then there is no such alternating sequence. However if we strengthen the inequality from Lemma 3 to $\max(\mathcal{C}) < \text{rem}(\mathcal{C})$ then such a sequence exists.

Lemma 4. *Assume \mathcal{C} is a coloring of $n > 1$ elements, where $n > 1$, $u \neq v$ and $\max(\mathcal{C}) \leq \text{rem}(\mathcal{C}) - 1$. Then there is a \mathcal{C} -alternating sequence γ from u to v .*

Proof. The case $n \leq 3$ can be checked directly. Hence from now on we assume $n \geq 4$. This implies that $|\gamma| \geq 2$ – it will be important later as we consider two end-elements of γ (so they should exist).

Instead of elements u, v we consider their colors A, B . Hence we need a sequence which has fixed colors A, B at its ends (it is not relevant which ends). If we remove u and v from our universe then the resulting coloring \mathcal{C}' satisfies the condition from Lemma 3: we have $\max(\mathcal{C}') \leq \text{rem}(\mathcal{C}') + 1$.

Let $\gamma' := \text{AlterSeq1}(\mathcal{C}')$ be a \mathcal{C}' -alternating sequence missing two elements (with regard to \mathcal{C}) with the colors A, B . Now we insert two previously removed elements u, v with colors A and B into γ' . The main point is to show how to do it. We have several cases depending on the colors C, D of the first and the last element of γ' .

Case 1: The trivial case: $\{A, B\} \cap \{C, D\} = \emptyset$.

We insert u at arbitrary end of γ' and v at the other end of γ' , thus obtaining a desired alternating sequence γ .

Case 2: Also rather straightforward case: $(A \neq B \wedge C \neq D)$.

There is a possibility that either C or D equals either A or B , so we might be constrained with placing u or v at one end of γ' .

Case 3: $(C = D)$

Then by Lemma 3 the color C is *exhausted* and none of A, B equals C . This is in fact a special subcase of **Case 1**.

We can assume now that $C \neq D$ and $A = B$. Without loss of generality let $A = C$. So the only remaining case is as follows.

Case 4: $(A = B = C \neq D)$.

We insert one element of color A after D . We're left with one element of color A , but we cannot put it at the end of our sequence – we have to insert it in between a pair of elements of γ' such that neither of them is of color A . We can use a simple counting argument that it can be done due to the inequality $\max(\mathcal{C}) \leq \text{rem}(\mathcal{C}) - 1$. We omit technical details.

This completes the proof.

We can write an algorithm from this Lemma in the following *pseudocode*:

Algorithm 2: AlterSeq2(\mathcal{C}, u, v)

input : $\mathcal{C} = \{C_1, C_2, \dots, C_k\}$, $|C_1| \leq |C_2| \leq \dots |C_k|$, $u \in C_i$,
 $v \in C_j$
output: \mathcal{C} -alternating sequence of $C_1 \cup C_2 \cup \dots C_k$ with u, v at its
ends

- 1 $C'_i = C_i - \{u\}$;
- 2 $C'_j = C_j - \{v\}$;
- 3 $\mathcal{C}' := \{C_1, C_2, \dots, C_{i-1}, C'_i, C_{i+1}, \dots, C_{j-1}, C'_j, C_{j+1}, \dots, C_k\}$;
- 4 $\gamma' := \text{AlterSeq1}(\mathcal{C}')$;
- 5 $A := \text{color}(u)$, $B := \text{color}(v)$;
- 6 $C := \text{color}(\text{first}(\gamma'))$, $D := \text{color}(\text{last}(\gamma'))$;
- 7 **if** $(\{A, B\} \cap \{C, D\} = \emptyset)$ **then**
- 8 **return** $u \gamma' v$;
- 9 **else if** $(A \neq B \wedge C \neq D)$ **then**
- 10 **if** $(u \gamma' v)$ is an alternating sequence **then**
- 11 **return** $u \gamma' v$;
- 12 **else**
- 13 **return** $v \gamma' u$;
- 14 **else** // Assume $A = B = C \neq D$
- 15 $\gamma'' := \gamma'$ with first element removed;
- 16 $\gamma'' := u \gamma'' v$;
- 17 insert $\text{first}(\gamma')$ into γ'' without violating alternating property;
- 18 **return** γ'' ;

Example 5. We show how the algorithm works if **Case 2** applies, let

$$\mathcal{C} = (\{1\}, \{3, 4, 5, 6\}, \{7, 8, 9, 10\}), \quad u = 3, \quad v = 8.$$

After removing these elements we get coloring $\mathcal{C}' = (\{1\}, \{4, 5, 6\}, \{7, 9, 10\})$. Then after applying algorithm from Lemma 3 we obtain:

$$\text{AlterSeq1}(\mathcal{C}') = \gamma' = (7\ 5\ 9\ 4\ 10\ 1\ 6).$$

Now we have the second case from the last proof. We insert removed elements at the ends in a suitable way and get the final result:

$$\text{AlterSeq2}(\mathcal{C}, 3, 8) = (3\ 7\ 5\ 9\ 4\ 10\ 1\ 6\ 8)$$

Example 6. We show now an example when **Case 1** applies, let:

$$\mathcal{C} = (\{1\}, \{2\}, \{3, 4\}, \{5, 6, 7\}) \quad u = 1, v = 2.$$

We have $n = 7$ and $|C_4| = \max(\mathcal{C}) < \frac{n}{2}$. Then the algorithm constructs the following alternating sequence γ from 1 to 2:

$$\gamma = (1\ 5\ 3\ 6\ 4\ 7\ 2).$$

In fact there are 12 distinct alternating sequences from $u = 1$ to $v = 2$ in this case.

We are mostly interested in colorings given by S -colorings in trees. Assume until end of the paper that the smallest color corresponds to a singleton set (in case of S -colorings to $\{S\}$).

Lemma 5. *Assume \mathcal{C} is a coloring of n elements with $|C_1| = 1$ and $\max(\mathcal{C}) = \text{rem}(\mathcal{C})$. Then there exists \mathcal{C} -alternating sequence γ starting from u and ending in v if and only if at least one of u, v is of the largest color.*

Proof. In this case $\max(\mathcal{C}) = \text{rem}(\mathcal{C}) = \frac{n}{2}$. Assume C_1 consists of a single element and u is of the largest color. We provide an algorithm for constructing a proper alternating sequence in linear time.

Algorithm 3: AlterSeq3(\mathcal{C} , u , v)

input : $\mathcal{C} = \{C_1, C_2, \dots, C_k\}$, $|C_1| \leq |C_2| \leq \dots |C_k|$
output: \mathcal{C} -alternating sequence of $C_1 \cup C_2 \cup \dots C_k$ with u, v at its ends
1 **if** ($\max(\mathcal{C}) \leq \text{rem}(\mathcal{C}) - 1$) **then**
2 **return** *AlterSeq2*(\mathcal{C} , u , v);
3 **else**
4 $\alpha := \text{sequentialize}(C_k)$;
5 $\beta := \text{sequentialize}(C_{k-1}, C_{k-2}, \dots, C_1)$;
6 $\gamma := \text{interleave}(\alpha, \beta)$;
 /* Notice that γ starts with the largest color and ends with C_1 */
7 exchange the first element of γ with u ;
8 exchange the last element of γ with v ;
9 **return** γ ;

Observation. If the assumption $|C_1| = 1$ is dropped then the last lemma is false, for example if $\mathcal{C} = \{1, 2\}, \{3, 4\}$ and u, v are of a same color then $\max(\mathcal{C}) = \text{rem}(\mathcal{C})$ but there is no \mathcal{C} -alternating path starting from u and ending in v .

4 Selecting a good node S : centroids in trees

We show that S satisfying for given u, v Lemma 2 can be chosen as one of the potentially at most two nodes minimizing the separability of the graph.




Fig. 3. The nodes v_2 and v_7 are the centroids.

The measure of *separability* of a vertex v in tree T , denoted by $\beta_T(v)$, is the size of maximum component of $T - \{v\}$. A vertex v is called a *centroid* if it has minimal separability over all vertices in T

Lemma 6. [Folklore]

1. If a vertex v is a centroid then $\beta_T(v) \leq \lfloor \frac{n}{2} \rfloor$.
2. A tree with an odd number of vertices has exactly one centroid.
3. Otherwise T has two centroids u and v which are neighbours and $\Delta(u) = \Delta(v)$.

Lemma 7. [Selecting good node S]

1. Existence of at least one (of at most two) centroid S of T implies the S -coloring satisfying conditions of Lemma 4 or Lemma 5 (see Figure 4) for any chosen pair of distinct nodes u, v .
2. Let S be a centroid and \mathcal{C} be S -coloring. Then any \mathcal{C} -alternating sequence starting with u and ending with v is a maximum weight Hamiltonian path from u to v in K_T .




Fig. 4. A bicentroidal tree with centroids denoted by S, S' . There is no alternating path from u to v with respect to S -coloring, but there exists such path with respect to S' -coloring: $\gamma = (u \ 1 \ S \ 2 \ S' \ v)$.

5 Computing maximum Hamiltonian paths

We present linear time algorithm constructing maximum Hamiltonian path in K_T between two different nodes $u, v \in T$.

Lemma 8. For a given tree T we can compute in linear time:

- a) centroids of T ,
- b) $\Delta(v)$ for each $v \in T$,
- c) the distance to each centroid, for each $v \in T$.

First we present a theorem which describes how to compute the maximum value of a path. This theorem follows directly from the results in the previous two sections.

Theorem 1. *Let S be a centroid of T such that there is an alternating sequence γ with respect to S from u to v . We know such S exists. Then γ is the maximum path from u to v in K_T and its weight equals $2 \cdot \Delta(S) - \text{dist}(u, S) - \text{dist}(v, S)$.*

Corollary 1.

1. *The maximum weight of a Hamiltonian path in K_T equals $2 \cdot \Delta(S) - \text{weight}(S, v)$, where S is one of centroids of T and v is the closest neighbor of S ($\text{weight}(S, v)$ is minimal).*
2. *The maximum weight of a Hamiltonian cycle in K_T equals $2 \cdot \Delta(S)$, where S is a centroid of T .*

Now we present how to construct in linear time the maximum path γ .

Algorithm 4: MaxPath(T, u, v)

```

1 if (there are two centroids in  $T$ ) then
2   | choose centroid  $S$  such that at least one of  $u, v$  is of the largest
   | color in  $S$ -coloring;
3 else
4   |  $S$  becomes the unique centroid;
5  $\mathcal{C} := S$ -coloring of  $T$ ;
6 if ( $T$  is bicentroidal) then
7   |  $\gamma := \text{AlterSeq3}(\mathcal{C}, u, v)$ ;
8 else
9   |  $\gamma := \text{AlterSeq2}(\mathcal{C}, u, v)$ ;
10 return  $\gamma$ ;
```

Lemma 8 together with Lemma 4 and Lemma 5 directly imply the following fact.

Theorem 2. *Algorithm MaxPath(T, u, v) computes a maximum Hamiltonian path in K_T between two given nodes in linear time.*

To quickly output the length of a maximum Hamiltonian path between a given pair of nodes in K_T (without the path itself) we need to find the

(at most two) centroids of T . For each centroid S we have to compute distance to all other nodes in v . Both operations can be done in linear time. If there's only one centroid S , then we already have all necessary information to answer queries for maximum Hamiltonian paths in constant time.

When there are two centroids in T (let's say S and S') then there's one more preprocessing step involved. We need to compute size of the largest subtrees of trees rooted in S and S' and should be able to quickly recognize which vertices belong to these subtrees. This is useful in cases similar to 4 when we need to determine which centroid to use for construction of maximum Hamiltonian paths in K_T . This additional step can also be done in linear time which leads us to the following theorem.

Theorem 3. *We can preprocess a given tree in linear time to allow later queries about the value of a maximum path between two nodes in constant time.*

References

1. H. Steinhaus, *One Hundred Problems in Elementary Mathematics*, Dover Publications (September 1, 1979)
2. Alfred V. Aho, John E. Hopcroft, Jeffrey D. Ullman, *The Design and Analysis of Computer Algorithms*, Addison-Wesley (1974)
3. M. R. Garey, D. S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness*, W. H. Freeman (1979)