

Conservative level set methods for incompressible air/water flow

Chris Kees, Matthew Farthing, Clint Dawson, and Serge Prudhomme

Computational Mechanics Brown Bag Seminar
U.S. Army Engineer Research and Development Center

Nov 20, 2008



Context

- ▶ “Particle Scale Distribution of Soil Moisture”, Military Engineering Basic Research Program 6.1 (GSL), FY2.



Context

- ▶ “Particle Scale Distribution of Soil Moisture”, Military Engineering Basic Research Program 6.1 (GSL), FY2.
- ▶ “High Fidelity Vesself Effects”, Navigation Systems Research Program (CHL), FY3.



Context

- ▶ “Particle Scale Distribution of Soil Moisture”, Military Engineering Basic Research Program 6.1 (GSL), FY2.
- ▶ “High Fidelity Vesself Effects”, Navigation Systems Research Program (CHL), FY3.
- ▶ Both these projects have at their core solving **moving boundary problems** for the **incompressible Navier-Stokes** equations.



Particle Scale Project Objectives and Products

- ▶ The overall objective is to develop a better **understanding** of variably saturated granular/porous media by combining multi-scale theoretical and computational model with data.



Particle Scale Project Objectives and Products

- ▶ The overall objective is to develop a better **understanding** of variably saturated granular/porous media by combining multi-scale theoretical and computational model with data.
 - ▶ Assemble particle scale, macroscale, and multiscale mathematical models of **air/water flow** in static granular materials.



Particle Scale Project Objectives and Products

- ▶ The overall objective is to develop a better **understanding** of variably saturated granular/porous media by combining multi-scale theoretical and computational model with data.
 - ▶ Assemble particle scale, macroscale, and multiscale mathematical models of **air/water flow** in static granular materials.
 - ▶ Develop tools for simulating at **both scales** (“classical” continuum models, pore network, lattice Boltzmann).



Particle Scale Project Objectives and Products

- ▶ The overall objective is to develop a better **understanding** of variably saturated granular/porous media by combining multi-scale theoretical and computational model with data.
 - ▶ Assemble particle scale, macroscale, and multiscale mathematical models of **air/water flow** in static granular materials.
 - ▶ Develop tools for simulating at **both scales** (“classical” continuum models, pore network, lattice Boltzmann).
 - ▶ Develop tools for estimation of macroscale **constitutive relations** (homogenization/upscaling).



Particle Scale Project Objectives and Products

- ▶ The overall objective is to develop a better **understanding** of variably saturated granular/porous media by combining multi-scale theoretical and computational model with data.
 - ▶ Assemble particle scale, macroscale, and multiscale mathematical models of **air/water flow** in static granular materials.
 - ▶ Develop tools for simulating at **both scales** (“classical” continuum models, pore network, lattice Boltzmann).
 - ▶ Develop tools for estimation of macroscale **constitutive relations** (homogenization/upscaling).
 - ▶ Develop tools for **multiscale simulation** (variational multiscale methods).



High Fidelity Vessels Project Objectives and Products

- ▶ The overall objective is to develop **accurate and robust** numerical models of **breaking waves** and **moving vessels** primarily for locally adaptive, unstructured tetrahedral meshes



High Fidelity Vessels Project Objectives and Products

- ▶ The overall objective is to develop **accurate and robust** numerical models of **breaking waves** and **moving vessels** primarily for locally adaptive, unstructured tetrahedral meshes
 - ▶ Develop **Eulerian** free surface tools: level set/volume-of-fluid (free surface evolution) and eikonal equations (geometry/boundary conditions)



High Fidelity Vessels Project Objectives and Products

- ▶ The overall objective is to develop **accurate and robust** numerical models of **breaking waves** and **moving vessels** primarily for locally adaptive, unstructured tetrahedral meshes
 - ▶ Develop **Eulerian** free surface tools: level set/volume-of-fluid (free surface evolution) and eikonal equations (geometry/boundary conditions)
 - ▶ Develop **two-phase** (air/water) Navier-Stokes capability




High Fidelity Vessels Project Objectives and Products

- ▶ The overall objective is to develop **accurate and robust** numerical models of **breaking waves** and **moving vessels** primarily for locally adaptive, unstructured tetrahedral meshes
 - ▶ Develop **Eulerian** free surface tools: level set/volume-of-fluid (free surface evolution) and eikonal equations (geometry/boundary conditions)
 - ▶ Develop **two-phase** (air/water) Navier-Stokes capability
 - ▶ Develop **three-phase** (air/water/solid) capability



Two phases separated by a sharp interface



Flow equations and jump conditions

- In Ω_a and Ω_w we assume:

$$\nabla \cdot \mathbf{v}_\alpha = 0 \quad (1)$$

$$\frac{\partial \rho_\alpha \mathbf{v}_\alpha}{\partial t} + \nabla \cdot (\rho_\alpha \mathbf{v}_\alpha \otimes \mathbf{v}_\alpha - \boldsymbol{\sigma}_\alpha) = \rho_\alpha \mathbf{g} \quad (2)$$

where $\alpha = w, a$ and

$$\boldsymbol{\sigma}_\alpha = -p_\alpha \mathbf{I} + \rho_\alpha \nu_\alpha (\nabla \mathbf{v}_\alpha + \nabla \mathbf{v}_\alpha^t) \quad (3)$$



Flow equations and jump conditions

- In Ω_a and Ω_w we assume:

$$\nabla \cdot \mathbf{v}_\alpha = 0 \quad (1)$$

$$\frac{\partial \rho_\alpha \mathbf{v}_\alpha}{\partial t} + \nabla \cdot (\rho_\alpha \mathbf{v}_\alpha \otimes \mathbf{v}_\alpha - \boldsymbol{\sigma}_\alpha) = \rho_\alpha \mathbf{g} \quad (2)$$

where $\alpha = w, a$ and

$$\boldsymbol{\sigma}_\alpha = -p_\alpha \mathbf{I} + \rho_\alpha \nu_\alpha (\nabla \mathbf{v}_\alpha + \nabla \mathbf{v}_\alpha^t) \quad (3)$$

- We **don't** allow the interface to absorb mass:

$$\mathbf{v}_a \cdot \mathbf{n}_a + \mathbf{v}_w \cdot \mathbf{n}_w = 0 \text{ on } \Gamma \quad (4)$$



Flow equations and jump conditions

- In Ω_a and Ω_w we assume:

$$\nabla \cdot \mathbf{v}_\alpha = 0 \quad (1)$$

$$\frac{\partial \rho_\alpha \mathbf{v}_\alpha}{\partial t} + \nabla \cdot (\rho_\alpha \mathbf{v}_\alpha \otimes \mathbf{v}_\alpha - \boldsymbol{\sigma}_\alpha) = \rho_\alpha \mathbf{g} \quad (2)$$

where $\alpha = w, a$ and

$$\boldsymbol{\sigma}_\alpha = -p_\alpha \mathbf{I} + \rho_\alpha \nu_\alpha (\nabla \mathbf{v}_\alpha + \nabla \mathbf{v}_\alpha^t) \quad (3)$$

- We **don't** allow the interface to absorb mass:

$$\mathbf{v}_a \cdot \mathbf{n}_a + \mathbf{v}_w \cdot \mathbf{n}_w = 0 \text{ on } \Gamma \quad (4)$$

- We **do** allow the interface to resist normal stress:

$$\boldsymbol{\sigma}_a \cdot \mathbf{n}_a + \boldsymbol{\sigma}_w \cdot \mathbf{n}_w = -\mathbf{f}_\Gamma \text{ on } \Gamma \quad (5)$$



Continuum surface force approximation

- ▶ We enforce the jump conditions through a **singular** source term in the momentum equation.



Continuum surface force approximation

- ▶ We enforce the jump conditions through a **singular** source term in the momentum equation.
- ▶ We represent Γ as $\phi(x, t) = 0$ (i.e. the zero level set of some function ϕ).



Continuum surface force approximation

- ▶ We enforce the jump conditions through a **singular** source term in the momentum equation.
- ▶ We represent Γ as $\phi(x, t) = 0$ (i.e. the zero level set of some function ϕ).
- ▶ If $x(t)$ is a particle path then it remains on Γ so

$$\frac{d\phi(x(t), t)}{dt} = \frac{\partial \phi}{\partial t} + \nabla \phi \cdot \frac{dx}{dt} = \frac{\partial \phi}{\partial t} + \nabla \phi \cdot \mathbf{v} = 0 \quad (6)$$



Continuum surface force approximation

- ▶ We enforce the jump conditions through a **singular** source term in the momentum equation.
- ▶ We represent Γ as $\phi(x, t) = 0$ (i.e. the zero level set of some function ϕ).
- ▶ If $x(t)$ is a particle path then it remains on Γ so

$$\frac{d\phi(x(t), t)}{dt} = \frac{\partial \phi}{\partial t} + \nabla \phi \cdot \frac{dx}{dt} = \frac{\partial \phi}{\partial t} + \nabla \phi \cdot \mathbf{v} = 0 \quad (6)$$

- ▶ This is an equation for the level set function ϕ that we extend to all of Ω using the fluid velocity.



Two-phase Navier-Stokes with Surface Tension

$$\nabla \cdot \mathbf{v} = 0 \quad (7)$$

$$\frac{\partial \rho \mathbf{v}}{\partial t} + \nabla \cdot [\rho \mathbf{v} \otimes \mathbf{v} - \rho \nu (\nabla \mathbf{v} + \nabla \mathbf{v}^t)] = \rho \mathbf{g} - \nabla p + \mathbf{f}_\Gamma \quad (8)$$

$$\frac{\partial \phi}{\partial t} + \mathbf{v} \cdot \nabla \phi = 0 \quad (9)$$

$$\mathbf{f}_\Gamma(\phi) = \delta(\phi) \sigma_{aw} \kappa(\phi) \mathbf{n} \quad (10)$$

$$\rho(\phi) = H(\phi) \rho_w + [1 - H(\phi)] \rho_a \quad (11)$$

$$\nu(\phi) = H(\phi) \nu_w + [1 - H(\phi)] \nu_a \quad (12)$$

$$\mathbf{n} = \frac{\nabla \phi}{\|\nabla \phi\|} \quad (13)$$

$$\kappa = -\nabla \cdot \mathbf{n} \quad (14)$$



Two-phase Flow, cont'd

- $H(\phi)$ is the Heaviside function

$$H(\phi) = \begin{cases} 0 & \phi < 0 \\ 1/2 & \phi = 0 \\ 1 & \phi > 0 \end{cases} \quad (15)$$



Two-phase Flow, cont'd

- $H(\phi)$ is the Heaviside function

$$H(\phi) = \begin{cases} 0 & \phi < 0 \\ 1/2 & \phi = 0 \\ 1 & \phi > 0 \end{cases} \quad (15)$$

- $\delta(\phi)$ is the distributional derivative of H , the Dirac delta function

$$\int_{-\infty}^{\infty} \delta(\phi) f d\phi = f(0) \quad (16)$$



The Level-Set Conservation Problem

- The integral form of conservation of mass (or volume for incompressible fluids) over an interval $[t_n, t_{n+1}]$ is

$$\int_{\Omega} \rho_a H(\phi) dV \Big|_{t_n}^{t_{n+1}} + \int_{t_n}^{t_{n+1}} \int_{\partial\Omega} \rho_a H(\phi) \mathbf{v} \cdot \mathbf{n} dS = 0 \quad (17)$$



The Level-Set Conservation Problem

- The integral form of conservation of mass (or volume for incompressible fluids) over an interval $[t_n, t_{n+1}]$ is

$$\int_{\Omega} \rho_a H(\phi) dV \Big|_{t_n}^{t_{n+1}} + \int_{t_n}^{t_{n+1}} \int_{\partial\Omega} \rho_a H(\phi) \mathbf{v} \cdot \mathbf{n} dS = 0 \quad (17)$$

- Unfortunately this is not enforced by standard discretizations of

$$\frac{\partial \phi}{\partial t} + \mathbf{v} \cdot \nabla \phi = 0 \quad (18)$$

or

$$\frac{\partial \phi}{\partial t} + \nabla \cdot (\phi \mathbf{v}) = 0 \quad (19)$$



The Level-Set Conservation Problem

- The integral form of conservation of mass (or volume for incompressible fluids) over an interval $[t_n, t_{n+1}]$ is

$$\int_{\Omega} \rho_a H(\phi) dV \Big|_{t_n}^{t_{n+1}} + \int_{t_n}^{t_{n+1}} \int_{\partial\Omega} \rho_a H(\phi) \mathbf{v} \cdot \mathbf{n} dS = 0 \quad (17)$$

- Unfortunately this is not enforced by standard discretizations of

$$\frac{\partial \phi}{\partial t} + \mathbf{v} \cdot \nabla \phi = 0 \quad (18)$$

or

$$\frac{\partial \phi}{\partial t} + \nabla \cdot (\phi \mathbf{v}) = 0 \quad (19)$$

- But equation 17 doesn't uniquely define ϕ so we can't use it in a level set approach



Example: Vortex in a Box



Example: Bubble in a Box



Methods for Maintaining Conservation

- Compute a more accurate solution.



Methods for Maintaining Conservation

- ▶ Compute a more accurate solution.
- ▶ Compute the fluid mass or volume in each element and reconstruct Γ at each time step (volume-of-fluid methods).



Methods for Maintaining Conservation

- ▶ Compute a more accurate solution.
- ▶ Compute the fluid mass or volume in each element and reconstruct Γ at each time step (volume-of-fluid methods).
- ▶ Hybrid methods



Methods for Maintaining Conservation

- ▶ Compute a more accurate solution.
- ▶ Compute the fluid mass or volume in each element and reconstruct Γ at each time step (volume-of-fluid methods).
- ▶ Hybrid methods
 1. Particle/Level Set Methods



Methods for Maintaining Conservation

- ▶ Compute a more accurate solution.
- ▶ Compute the fluid mass or volume in each element and reconstruct Γ at each time step (volume-of-fluid methods).
- ▶ Hybrid methods
 1. Particle/Level Set Methods
 2. Level Set/Volume-of-Fluid Methods



Methods for Maintaining Conservation

- ▶ Compute a more accurate solution.
- ▶ Compute the fluid mass or volume in each element and reconstruct Γ at each time step (volume-of-fluid methods).
- ▶ Hybrid methods
 1. Particle/Level Set Methods
 2. Level Set/Volume-of-Fluid Methods
- ▶ Conservative level set methods: mimic volume conservation with a smoothed Heaviside function and identify Γ with $H = 1/2$.



A New Conservative Level Set Method?

- Try to “minimally” correct the level set equation to obtain conservation, which is a “low order” property: Find ϕ' such that

$$\int_{\Omega} \left[H(\phi + \phi') - H^{n+1} \right] dV = 0 \quad (20)$$



A New Conservative Level Set Method?

- Try to “minimally” correct the level set equation to obtain conservation, which is a “low order” property: Find ϕ' such that

$$\int_{\Omega} \left[H(\phi + \phi') - H^{n+1} \right] dV = 0 \quad (20)$$

- Problems:



A New Conservative Level Set Method?

- Try to “minimally” correct the level set equation to obtain conservation, which is a “low order” property: Find ϕ' such that

$$\int_{\Omega} \left[H(\phi + \phi') - H^{n+1} \right] dV = 0 \quad (20)$$

- Problems:
 1. Such a ϕ' may not exist.



A New Conservative Level Set Method?

- Try to “minimally” correct the level set equation to obtain conservation, which is a “low order” property: Find ϕ' such that

$$\int_{\Omega} \left[H(\phi + \phi') - H^{n+1} \right] dV = 0 \quad (20)$$

- Problems:
 1. Such a ϕ' may not exist.
 2. If it does, it's not unique.



A New Conservative Level Set Method?

- Try to “minimally” correct the level set equation to obtain conservation, which is a “low order” property: Find ϕ' such that

$$\int_{\Omega} \left[H(\phi + \phi') - H^{n+1} \right] dV = 0 \quad (20)$$

- Problems:

1. Such a ϕ' may not exist.
2. If it does, it's not unique.
3. The equation nonlinear and non-differentiable.



A New Conservative Level Set Method?

- ▶ Try to “minimally” correct the level set equation to obtain conservation, which is a “low order” property: Find ϕ' such that

$$\int_{\Omega} \left[H(\phi + \phi') - H^{n+1} \right] dV = 0 \quad (20)$$

- ▶ Problems:
 1. Such a ϕ' may not exist.
 2. If it does, it's not unique.
 3. The equation nonlinear and non-differentiable.
- ▶ Sounds like a great idea!



Variational Problem

$$\min_{\phi_c \in \mathcal{A}} I(\phi_c) \quad (21)$$

where

$$I(\phi_c) = \frac{1}{2} \int_{\Omega} \|\nabla \phi\|^2 dV \quad (22)$$

$$\mathcal{A} = \{\phi_c \in V_{\phi} | J(\phi_c) = 0\} \quad (23)$$

$$J(\phi_c) = \int_{\Omega} [H(\phi_{n+1} + \phi_c) - H_{n+1}] dV \quad (24)$$



A nonlinear elliptic equation

Find $\phi' \in H^1(\Omega)$ such that

$$\int_{\Omega} \{ [H(\phi_{h,n+1} + \phi^c) - \bar{H}_{n+1}] w + \epsilon \nabla \phi^c \cdot \nabla w \} dV = 0 \quad \forall w \in H^1(\Omega) \quad (25)$$

where \bar{H}_{n+1} is piecewise constant and satisfies

$$\int_{\Omega} (\bar{H}_{n+1} - H(\phi_{h,n})) dV + \int_{t_n}^{t_{n+1}} \int_{\partial\Omega} H(\phi) \mathbf{v} \cdot \mathbf{n} dS = 0 \quad (26)$$



Results: Vortex in a Box



Results: Bubble in a Box



Summary

- We formulated a mass/volume correction equation that can be added to the standard level set methods to enforce conservation.



Summary

- ▶ We formulated a mass/volume correction equation that can be added to the standard level set methods to enforce conservation.
- ▶ It applies to higher order and discontinuous finite elements as well.



Summary

- ▶ We formulated a mass/volume correction equation that can be added to the standard level set methods to enforce conservation.
- ▶ It applies to higher order and discontinuous finite elements as well.
- ▶ To do: local (embarassingly parallel) version, complete the theory for the correction equation and the nonlinear solver.

