



Observed lag embeddings | direction 1, original trial 16

T, channel 7 | same preprocessed 500-ms pre-GO epoch | fixed K = 1, m = 3


tau = 1 ms | 498 points




tau = 2 ms | 496 points




tau = 3 ms | 494 points




tau = 4 ms | 492 points




tau = 5 ms | 490 points




tau = 6 ms | 488 points




tau = 7 ms | 486 points




tau = 8 ms | 484 points




tau = 9 ms | 482 points




tau = 10 ms | 480 points




tau = 11 ms | 478 points




tau = 12 ms | 476 points




tau = 13 ms | 474 points




tau = 14 ms | 472 points




tau = 15 ms | 470 points




tau = 16 ms | 468 points




tau = 17 ms | 466 points




tau = 18 ms | 464 points



tau = 19 ms | 462 points



tau = 20 ms | 460 points




Black = observed trajectory; no fitted surface. Time samples are linked in their original order.
Identical camera and axis limits across all 20 delays. No PCA, new filtering, or decoding.


Observed lag embeddings | direction 2, original trial 10

T, channel 7 | same preprocessed 500-ms pre-GO epoch | fixed $K = 1$, $m = 3$


tau = 1 ms | 498 points




tau = 2 ms | 496 points




tau = 3 ms | 494 points




tau = 4 ms | 492 points




tau = 5 ms | 490 points




tau = 6 ms | 488 points




tau = 7 ms | 486 points




tau = 8 ms | 484 points




tau = 9 ms | 482 points




tau = 10 ms | 480 points




tau = 11 ms | 478 points




tau = 12 ms | 476 points




tau = 13 ms | 474 points




tau = 14 ms | 472 points




tau = 15 ms | 470 points




tau = 16 ms | 468 points




tau = 17 ms | 466 points




tau = 18 ms | 464 points



tau = 19 ms | 462 points



tau = 20 ms | 460 points




Black = observed trajectory; no fitted surface. Time samples are linked in their original order.
Identical camera and axis limits across all 20 delays. No PCA, new filtering, or decoding.


Observed lag embeddings | direction 3, original trial 14

T, channel 7 | same preprocessed 500-ms pre-GO epoch | fixed $K = 1$, $m = 3$


tau = 1 ms | 498 points




tau = 2 ms | 496 points




tau = 3 ms | 494 points




tau = 4 ms | 492 points




tau = 5 ms | 490 points




tau = 6 ms | 488 points




tau = 7 ms | 486 points




tau = 8 ms | 484 points




tau = 9 ms | 482 points




tau = 10 ms | 480 points




tau = 11 ms | 478 points




tau = 12 ms | 476 points




tau = 13 ms | 474 points




tau = 14 ms | 472 points




tau = 15 ms | 470 points




tau = 16 ms | 468 points




tau = 17 ms | 466 points




tau = 18 ms | 464 points



tau = 19 ms | 462 points



tau = 20 ms | 460 points




Black = observed trajectory; no fitted surface. Time samples are linked in their original order.
Identical camera and axis limits across all 20 delays. No PCA, new filtering, or decoding.


Observed lag embeddings | direction 4, original trial 17

T, channel 7 | same preprocessed 500-ms pre-GO epoch | fixed $K = 1$, $m = 3$


tau = 1 ms | 498 points




tau = 2 ms | 496 points




tau = 3 ms | 494 points




tau = 4 ms | 492 points




tau = 5 ms | 490 points




tau = 6 ms | 488 points




tau = 7 ms | 486 points




tau = 8 ms | 484 points




tau = 9 ms | 482 points




tau = 10 ms | 480 points




tau = 11 ms | 478 points




tau = 12 ms | 476 points




tau = 13 ms | 474 points




tau = 14 ms | 472 points




tau = 15 ms | 470 points




tau = 16 ms | 468 points




tau = 17 ms | 466 points




tau = 18 ms | 464 points



tau = 19 ms | 462 points



tau = 20 ms | 460 points




Black = observed trajectory; no fitted surface. Time samples are linked in their original order.
Identical camera and axis limits across all 20 delays. No PCA, new filtering, or decoding.


Observed lag embeddings | direction 5, original trial 6

T, channel 7 | same preprocessed 500-ms pre-GO epoch | fixed $K = 1$, $m = 3$


tau = 1 ms | 498 points




tau = 2 ms | 496 points




tau = 3 ms | 494 points




tau = 4 ms | 492 points




tau = 5 ms | 490 points




tau = 6 ms | 488 points




tau = 7 ms | 486 points




tau = 8 ms | 484 points




tau = 9 ms | 482 points




tau = 10 ms | 480 points




tau = 11 ms | 478 points




tau = 12 ms | 476 points




tau = 13 ms | 474 points




tau = 14 ms | 472 points




tau = 15 ms | 470 points




tau = 16 ms | 468 points




tau = 17 ms | 466 points




tau = 18 ms | 464 points



tau = 19 ms | 462 points



tau = 20 ms | 460 points




Black = observed trajectory; no fitted surface. Time samples are linked in their original order.
Identical camera and axis limits across all 20 delays. No PCA, new filtering, or decoding.


Observed lag embeddings | direction 6, original trial 9

T, channel 7 | same preprocessed 500-ms pre-GO epoch | fixed $K = 1$, $m = 3$


tau = 1 ms | 498 points




tau = 2 ms | 496 points




tau = 3 ms | 494 points




tau = 4 ms | 492 points




tau = 5 ms | 490 points




tau = 6 ms | 488 points




tau = 7 ms | 486 points




tau = 8 ms | 484 points




tau = 9 ms | 482 points




tau = 10 ms | 480 points




tau = 11 ms | 478 points




tau = 12 ms | 476 points




tau = 13 ms | 474 points




tau = 14 ms | 472 points




tau = 15 ms | 470 points




tau = 16 ms | 468 points




tau = 17 ms | 466 points




tau = 18 ms | 464 points



tau = 19 ms | 462 points



tau = 20 ms | 460 points



Black = observed trajectory; no fitted surface. Time samples are linked in their original order.
Identical camera and axis limits across all 20 delays. No PCA, new filtering, or decoding.