

Trapezoid o' trapezoid!

Problem Statement

Consider the scalar ordinary differential equation (ODE) $\dot{x} = f(t, x(t))$ starting from $x_0 = x(0)$. Then, the trajectory $x(t)$ over $t \in [0, T]$ satisfies:

$$x(t_{n+1}) = x(t_n) + \int_{\tau=t_n}^{\tau=t_{n+1}} f(\tau, x(\tau)) d\tau$$

for $t_n = nh$. Let $g(\tau) := f(\tau, x(\tau))$. Assume that g is \mathcal{C}^2 over $[0, T]$.

a

If $0 \leq a \leq b \leq T$, then prove that g satisfies:

$$\int_a^b g(\tau) d\tau = \frac{1}{2}(b-a)[g(a) + g(b)] - \frac{1}{2} \int_a^b \left[\left(\frac{b-a}{2} \right)^2 - \left(\tau - \frac{a+b}{2} \right)^2 \right] g''(\tau) d\tau$$

Hint: Use integration by parts.

b

Then, use the identity in part a) to prove that there exists a constant M s.t.

$$\left| \int_a^b g(\tau) d\tau - \frac{1}{2}(b-a)[g(a) + g(b)] \right| \leq M \frac{(b-a)^3}{12}$$

c

Deduce that the local truncation error of the trapezoidal method is $\mathcal{O}(h^3)$; i.e.:

$$x(t_{n+1}) = x(t_n) + \frac{h}{2}[f(t_n, x(t_n)) + f(t_{n+1}, x(t_{n+1}))] + \mathcal{O}(h^3)$$

d (optional)

Under possibly additional assumptions, prove that the global error for the trapezoidal method is $\mathcal{O}(h^2)$; i.e., there exists some constant C s.t.

$$|x(t_n) - x_n| \leq Ch^2 \forall t \in [0, T]$$

Solution

a

b

c

d

Problem 2: Deriving Adams-Moulton, not 1, but 2

Problem Statement

Consider a scalar ODE $\dot{x}(t) = f(t, x(t))$ with $x(0)$ as the initial point. With a step-size of $h > 0$ and $t_n = nh$ for $n \geq 0$ we have:

$$x(t_{n+1}) = x(t_n) + \int_{\tau=t_n}^{\tau=t_{n+1}} f(\tau, x(\tau)) d\tau$$

Adams-Moulton seeks to compute x_n recursively as:

$$x_{n+1} = x_n + \int_{\tau=t_n}^{\tau=t_{n+1}} g(\tau) d\tau \quad (1)$$

where $g(\tau)$ is a polynomial approximation to $f(\tau, x(\tau))$. Define $G_n := f(t_n, x_n)$. Then, the AM(2) method seeks a quadratic approximation to g that takes the values

$$g(t_{n-1}) = G_{n-1}, \quad g(t_n) = G_n, \quad g(t_{n+1}) = G_{n+1} \quad (2)$$

Consider g of the form

$$g(\tau) = G_{n-1}L_{n-1}(\tau) + G_nL_n(\tau) + G_{n+1}L_{n+1}(\tau) \quad (3)$$

where L 's are the Legendre polynomials defined by:

$$L_{n-1}(\tau) := \frac{(\tau - t_n)(\tau - t_{n+1})}{(t_{n-1} - t_n)(t_{n-1} - t_{n+1})}, \quad L_n(\tau) := \frac{(\tau - t_{n-1})(\tau - t_{n+1})}{(t_n - t_{n-1})(t_n - t_{n+1})}, \quad L_{n+1}(\tau) := \frac{(\tau - t_{n-1})(\tau - t_n)}{(t_{n+1} - t_{n-1})(t_{n+1} - t_n)}$$

a

Verify that g defined in (3) satisfies (2).

b

Compute $\int_{\tau=t_n}^{\tau=t_{n+1}} L_k(\tau) d\tau$ for $k = n-1, n, n+1$.

Solution